

# Self-induced density modulations in the free expansion of Bose-Einstein condensates

Luca Salasnich<sup>1</sup>, Nicola Manini<sup>2,3</sup>, Federico Bonelli<sup>2</sup>, Michael Korbman<sup>2</sup>, and Alberto Parola<sup>4</sup>

<sup>1</sup>*CNISM and CNR-INFM, Unità di Padova,  
Dipartimento di Fisica “Galileo Galilei”,*

*Università di Padova, Via Marzolo 8, 35131 Padova, Italy*

<sup>2</sup>*Dipartimento di Fisica, Università di Milano,  
Via Celoria 16, 20133 Milano, Italy*

<sup>3</sup>*CNISM, Unità di Milano,  
Via Celoria 16, 20133 Milano, Italy*

<sup>4</sup>*Dipartimento di Fisica e Matematica,  
Università dell’Insubria,  
Via Valleggio 11, 22100 Como, Italy*

We simulate numerically the free expansion of a repulsive Bose-Einstein condensate with an initially Gaussian density profile. We find a self-similar expansion only for weak inter-atomic repulsion. In contrast, for strong repulsion we observe the spontaneous formation of a shock wave at the surface followed by a significant depletion inside the cloud. In the expansion, contrary to the case of a classical viscous gas, the quantum fluid can generate radial rarefaction density waves with several minima and maxima. These intriguing nonlinear effects, never observed yet in free-expansion experiments with ultra-cold alkali-metal atoms, can be detected with the available setups.

PACS numbers: 03.75.Ss, 03.75.Hh, 64.75.+g

The anisotropic free expansion of a gas of <sup>87</sup>Rb atoms was the first experimental evidence of Bose-Einstein condensation in ultra-cold gases [1]. The non-ballistic free expansion observed with <sup>6</sup>Li atoms has been saluted as the first signature of superfluid behavior in a ultra-cold fermi vapor [2]. In both cases the atomic quantum gases can be described by the hydrodynamic equations of superfluids and, because the initial density profile is an inverted parabola, the free expansion is self-similar [3, 4, 5, 6, 7].

The first theoretical investigations of the free expansion into vacuum of a classical gas sphere with constant initial density dates back to the 1960’s [8, 9]: numerical analysis was needed to analyze in detail the formation of a depletion at the center and of a shock wave at the surface [10, 11]. More recently, rarefaction waves have been produced experimentally by the free expansion of an electron plasma [12].

In this Letter we show that the free expansion of a bosonic superfluid into vacuum displays intriguing nonlinear phenomena. In particular, by integrating numerically [13] the time-dependent Gross-Pitaevskii equation [14] we prove that the expansion of a repulsive Bose-Einstein condensate (BEC) of initial Gaussian density profile gives rise to self-induced density modulations, i.e. self-induced rarefaction waves. In addition, we find that the bosonic cloud produces a shock wave at the surface, that is damped by the quantum pressure of the superfluid. These nonlinear effects, which can be observed experimentally with available techniques, are strongly suppressed if a harmonic confinement along two perpendicular directions is retained and free expansion is allowed in 1D only. Observe that the formation of shock and rarefaction waves induced by an *ad hoc* external perturba-

tion has been suggested [15, 16, 17] and observed recently in BECs [18, 19] and also in non-dissipative nonlinear optics [20].

We describe the collective motion of the BEC of  $N$  atoms in terms of a complex mean-field wavefunction  $\psi(\mathbf{r}, t)$  normalized to unity and such that  $\rho(\mathbf{r}, t) = N\rho_1(\mathbf{r}, t) = N|\psi(\mathbf{r}, t)|^2$  is the number-density distribution. The equation of motion that we assume for  $\psi(\mathbf{r}, t)$  is the time-dependent Gross-Pitaevskii equation (GPE) [14]

$$i\hbar\frac{\partial}{\partial t}\psi = \left[ -\frac{\hbar^2}{2m}\nabla^2 + U + \frac{4\pi\hbar^2 a_s}{m}N|\psi|^2 \right] \psi, \quad (1)$$

where  $U(\mathbf{r}, t)$  is a confining potential that we assume to vanish at  $t \geq 0$  (thus allowing free expansion) and  $m$  is the atomic mass. The nonlinear term represents the inter-atomic interaction at a mean-field level, where  $a_s$  is the s-wave scattering length, and we consider the repulsive regime  $a_s > 0$ .

In traditional experiments with ultra-cold alkali-metal atoms [1, 2] expansion starts from an initial state coinciding with the ground state of the confined superfluid under the action of a (often anisotropic) harmonic potential. For robust interparticle interaction (large number of particles), the density profile in this initial state resembles closely a negative-curvature parabola [14]. When such density profile is taken as the initial state of a successive free non-ballistic expansion, to a very good degree of approximation it expands in a self-similar fashion, maintaining the same shape and only spreading out and scaling down its height proportionally, until a ballistic regime is reached when dilution leads to a fully non-interacting regime [3, 4, 5, 6, 7].

In the present work we discuss the much more exciting




FIG. 1: (Color on line). Four successive frames of the radial density profile (solid line) for the expansion of a strongly interacting condensate, characterized by dimensionless interaction parameter  $g = 2000$ . The “opacity” of the expanding cloud (dashed line) given by the density  $\rho_1(\mathbf{r})$  integrated along lines at a distance  $r$  from the center (rescaled by a factor 0.036). All quantities are dimensionless: lengths in units of the initial Gaussian width  $\sigma$  and time in units of  $m\sigma^2/\hbar$ .

phenomena observed in the expansion of an interacting condensate starting off as a stationary Gaussian

$$\psi(\mathbf{r}, 0) = \frac{1}{(\pi\sigma^2)^{3/4}} \exp\left(-\sum_{i=x,y,z} \frac{r_i^2}{2\sigma_i^2}\right), \quad (2)$$

with  $\sigma^3 = \sigma_x\sigma_y\sigma_z$ . Note that a Gaussian profile is readily achieved experimentally by equilibrating the BEC with a very small scattering length  $a_s$  obtained by means of the Fano-Feshbach resonance technique [14, 21] with a carefully tuned external constant magnetic field.  $a_s$  can then be set to the desired value by a sudden change in the magnetic field at the time when the harmonic trapping potential is switched off.

The isotropic case  $\sigma_x = \sigma_y = \sigma_z = \sigma$  is conceptually advantageous, as the expansion can be studied within the GPE model in its full generality as a function of a single parameter. Consider rescaling the variables of Eq. (1) as follows:  $\mathbf{r} \rightarrow \mathbf{r}/\sigma$ ,  $t \rightarrow t\hbar/(m\sigma^2)$ , and  $\psi \rightarrow \psi\sigma^{3/2}$ , to produce a dimensionless form

$$i\frac{\partial}{\partial t}\psi = \left[-\frac{1}{2}\nabla^2 + g|\psi|^2\right]\psi, \quad (3)$$

of the equation for the free expansion of a condensate expansion starting off from a Gaussian initial state of unit width. The dimensionless interaction strength  $g = 4\pi N a_s/\sigma$  is the one free parameter determining the properties of the free expansion. A gas of non interacting ( $g = 0$ ) bosons starting from the Gaussian state expands




FIG. 2: (Color on line). Characteristic times of the free expansion as a function of the interaction strength  $g$ . For  $g \geq 48.7$ , a first minimum in the radial density profile appears at the center  $r = 0$  at time  $t_{am}$  (dashed) and fills in at  $t_{dm}$  (dot-dashed). For  $g \geq 363$ , a local maximum re-forms at the center at time  $t_{aM}$  (solid) and disappears at time  $t_{dM}$ . A density maximum appears within the rarefaction region at time  $t_{eM}$  (dot-dot-dashed). Inset: the time of appearance  $t_a$  and disappearance  $t_d$  of a denser ring (sketched in the small square for  $g = 1200$ ,  $t = 1.8$ ) of visibility  $h$  (defined in Fig. 1) 1% (solid) and 5% (dashed) in the opacity profile (dashed lines of Fig. 1), as a function of  $g$ . Units as in Fig. 1.

self similarly according to the formula [22]

$$\psi(\mathbf{r}, t) = \frac{1}{\pi^{3/2}(1+t^2)^{3/2}} \exp\left(-\frac{r^2(1+it)}{2(1+t^2)}\right). \quad (4)$$

By using an efficient finite-difference Crank-Nicolson algorithm [13] we have verified that little changes affect the expansion as long as the interaction is small.  $g < 1$  tracks the weak-coupling limit where interaction only accelerates slightly the free expansion, while  $g \gg 1$  represents the strong-coupling regime, where the mean-field self-interaction term in Eq. (3) dominates the expansion for long enough to produce substantial nonlinear effects such as those sketched in Fig. 1. In particular we observe the rapid build-up of a sharp expanding spherical density wave which leaves behind a central region of depleted density. New successively formed radial ripples cross this density-depleted region. Eventually, at very long times, when the overall density has decayed enough for the nonlinear term in Eq. (3) to become negligible everywhere, the expansion recovers a bell-shaped profile.

More quantitatively, for increasing  $g$ , the rarefaction starts to be evidenced by a local minimum at the droplet center for  $g \gtrsim 48.3$ , and becomes more and more pronounced and long lived for larger interaction strength  $g$ .




FIG. 3: (Color on line). Comparison of the density profile  $\rho_1(r) = |\psi(r)|^2$  and of the velocity profile  $v(r)/r$  for the spherically symmetrical expansion of the strongly-interacting bosonic cloud. Units as in Fig. 1.

In the highly nonlinear regime, the free expansion of the bosonic cloud into vacuum develops sequences of radial density waves with minima and maxima, each starting at a characteristic time and disappearing at a later time. Figure 2 tracks a few early times in this class, as a function of  $g$ : more could be defined for larger  $g$ . Overall, the time  $t_{\text{am}}$  of appearance of the first local minimum reduces slowly as  $g$  increases, while the time  $t_{\text{dm}}$  of disappearance of all local minima and recovery of a bell-shaped profile increases rapidly with  $g$ . In between these two characteristic times several traveling density minima and maxima can be formed depending on the value of  $g$ . Generally, the number and density difference of local minima and maxima increases with the interaction strength.

In practice, the visibility of the central depletion need not be easy to appreciate by means of total opacity measurements, since such measurements address the integrated density of a generic linear section crossing the droplet at a given distance from its center. However, Fig. 1 shows that for suitably strong interaction, a sensibly higher-opacity outer ring does indeed develop. The inset of Fig. 2 tracks the time range when this inner optical-density minimum remains lower than the outside denser ring by at least 1% and 5%: it is seen that for strong enough interaction ( $g \gtrsim 215$  and  $g \gtrsim 310$  respectively), the depleted region realizes a limited but significant visibility, which improves for stronger coupling. An example of visible opacity ring is displayed in the upper right corner of Fig. 2.

To characterize the dynamics of the bosonic cloud it is

useful to introduce its local phase velocity, given by

$$\mathbf{v} = \frac{i}{2} \frac{\psi^* \nabla \psi - \psi \nabla \psi^*}{|\psi|^2}. \quad (5)$$

This velocity can be written as  $\mathbf{v}(\mathbf{r}, t) = \nabla \theta(\mathbf{r}, t)$ , where  $\theta(\mathbf{r}, t)$  is the phase of the macroscopic wave function  $\psi(\mathbf{r}, t) = \rho_1(\mathbf{r}, t)^{1/2} e^{i\theta(\mathbf{r}, t)}$  of the bosonic superfluid. From Eq. (4) one finds immediately the radial phase velocity of a spherical non-interacting Bose gas ( $g = 0$ ):

$$v = \frac{r}{2} \frac{t}{1 + t^2}. \quad (6)$$

In the interacting case ( $g > 0$ ) the nonlinear term acts as the chemical potential of a fluid of classical pressure  $P = g|\psi|^4/2$  and sound velocity  $c_s = \sqrt{g|\psi|^2}$  [14]. For  $g \ll 1$  the gas velocity follows Eq. (6) closely, while for large  $g$  deviations from Eq. (6) are substantial, as mainly the interaction term, rather than the quantum tendency to delocalization, drives the expansion. The ratio  $v/r$ , a constant as a function of  $r$  according to Eq. (6) in a non-interacting expansion, shows strong deviations induced by the nonlinear term  $g|\psi|^2$ . In practice, the phase velocity of a strongly interacting BEC approaches the local sound velocity, with larger densities implying higher velocities. Accordingly, initially the central part of the strongly repulsive bosonic superfluid accelerates and propagates faster than the periferic part: the ensuing mass flow is responsible for the formation of the rarefaction region inside the cloud, shown in Fig. 1. The matter flowing quickly out of the central region accumulates near the profile edge on top of the slower external tail, thus tending to produce a shock wave [16, 19, 20], with a BEC density profile extremely steep at the surface, approaching a step function: this is illustrated by the  $t=0.24$  panel of Fig. 1. This steep wave front survives for a brief period, after which density oscillations shoot backwards and the surface profile rapidly smoothens its density gradient. At this point, the expansion dynamics is strongly affected by these backward density oscillations, which induce a *inverted* relation between local velocity and density, as Fig. 3 illustrates: for a strongly interacting BEC, at a fixed time  $t$  the ratio  $v/r$  finds local maxima (minima) in correspondence to the local minima (maxima) of the density profile  $\rho_1$ . This inversion demonstrates the *inward* motion of the density ripples. These local minima represent the rarefaction waves produced by the surface step smoothing. This smoothing is driven by the quantum pressure term [16].

The quantum pressure  $-(|\nabla \psi|^2)/(2\rho_1)$ , which plays a negligible role in the self-similar non-ballistic expansion [7] of both Fermi and Bose superfluids with an inverted-parabola initial profile [23], becomes relevant in regularizing the shock-wave singularity, like the dissipative term in classical hydrodynamics [19]. Analogous depletion and shock-wave phenomena are indeed observed in the hydrodynamical expansion of hot classical fluids, and can




FIG. 4: (Color on line). Self-depleting radial density profile  $\rho_1 = \rho/N$  during the expansion: comparison between the strongly interacting Bose gas (GPE, solid lines) and the classical gas (NSE, dashed lines), with  $g = 10^4$  and the same initial conditions (a unit Gaussian). In the NSE, the shear viscosity coefficient  $\eta = 10^{-5}$ . Units as in Fig. 1.

be simulated e.g. by means of the Navier-Stokes equations (NSE), which depend on the (dissipative) coefficient of shear viscosity  $\eta$  [9]. For  $\eta = 0$  the irrotational ( $\nabla \wedge \mathbf{v} = \mathbf{0}$ ) NSE reduce to the Euler equations of an ideal (non-viscous) fluid. By using  $P(\rho) = g\rho_1^2/2$  as the equation of state, the Euler equations are exactly equivalent to the GPE without the quantum pressure term [14]. In Fig. 4 we compare the expanding BEC (GPE, Eq. (3)) and classical gas (NSE), for  $g = 10000$ . The NSE are solved by means of a Lagrangian finite-difference method [24]. The four successive frames displayed in Fig. 4 show that, while expanding, the interacting BEC and the classical gas produce a remarkably similar self-depletion of the central region. On the other hand, the multi-peak rarefaction structures predicted by the GPE are not reproduced by the NSE. Thus the novelty of the free expansion of a BEC with respect to that of a weakly viscous classical fluid, stands mainly in the distinct large-amplitude rarefaction waves moving inside the depletion region, which are supported by the nondissipative nature of quantum pressure.

The expansion of the BEC in an anisotropic context is also interesting. In particular, we find that a 1D expansion starting from a Gaussian state produces no shock wave and no central depletion. This applies both for an idealized 1D BEC, represented by a purely 1D GPE, and for the 3D expansion of a Gaussian wavepacket to which an harmonic confining potential is kept along two orthogonal space directions. Interestingly, if for the initial state of the 3D geometry a spherical Gaussian (2) of width equal to the confining-potential harmonic length  $[\hbar/(m\omega)]^{1/2}$  is taken, the strongly-interacting BEC shoots out rapidly in all directions, thus producing a shock wave, depletion, and rarefaction waves around the cylindrical symmetry axis. However, the total density,

integrated in the directions perpendicular to the symmetry axis, always retains a bell shape, even for very large interaction.

In conclusion, we have shown that the free expansion of a Bose-Einstein condensate reveals novel and interesting nonlinear effects, not achievable with classical viscous fluids, and which are awaiting experimental investigations.

The authors thank Paolo Di Trapani, Luciano Reatto and Flavio Toigo for useful suggestions.

- 
- [1] M. H. Anderson et al., *Science* **269**, 198 (1995).
  - [2] K.M. O'Hara, S.L. Hemmer, M.E. Gehm, S.R. Granade, and J.E. Thomas, *Science*, **298**, 2179 (2002).
  - [3] Y. Castin and R. Dum, *Phys. Rev. Lett.* **77**, 5315 (1996).
  - [4] Yu. Kagan, E.L. Surkov, G.V. Shlyapnikov, *Phys. Rev. A* **54**, R1753 (1996); **55** R18 (1997).
  - [5] P. Öhberg and L. Santos, *Phys. Rev. Lett.* **89**, 240402 (2002).
  - [6] L. Salasnich, A. Parola, and L. Reatto, *Phys. Rev. A* **72**, 025602 (2005).
  - [7] G. Diana, N. Manini, and L. Salasnich, *Phys. Rev. A* **73**, 065601 (2006).
  - [8] P. Molmud, *Phys. Fluids* **3**, 362 (1960).
  - [9] Ya. B. Zel'dovich and Yu. P. Raizer, *Physics of Shock Waves and High-Temperature Hydrodynamic Phenomena* (Dover Publications, Mineola-NY, 2002) [in Russian, Izd. Nauka, Moscow (1966)].
  - [10] V.E. Kondrashov, A.N. Polyanichev, and V.S. Fetisov, *Fluid Dynamics* **9**, 835 (1974).
  - [11] N.C. Freeman and S. Kumar, *J. Fluid Mech.* **56**, 523 (1972); N.C. Freeman, R.S. Johnson, S. Kumar, and W.B. Buch, *J. Fluid Mech.* **68**, 625 (1975).
  - [12] J.D. Moody and C.F. Driscoll, *Phys. Plasmas* **2**, 4482 (1995).
  - [13] E. Cerboneschi, R. Mannella, E. Arimondo, and L. Salasnich, *Phys. Lett. A* **249**, 495 (1998); L. Salasnich, A. Parola, and L. Reatto, *Phys. Rev. A* **64**, 023601 (2001).
  - [14] L.P. Pitaevskii and S. Stringari, *Bose-Einstein Condensation* (Oxford University Press, Oxford, 2003).
  - [15] M. Zak and I. Kulikov, *Phys. Lett. A* **307**, 99 (2003); I. Kulikov and M. Zak, *Phys. Rev. A* **67**, 063605 (2003).
  - [16] B. Damski, *Phys. Rev. A* **69** 043610 (2004); *Phys. Rev. A* **73** 043610 (2006).
  - [17] A.M. Kamchatnov, A. Gammal, and R.A. Kraenkel, *Phys. Rev. A* **69** 063605 (2004).
  - [18] Z. Dutton, M. Budde, C. Slowe, and L.V. Hau, *Science* **293**, 663 (2001).
  - [19] M.A. Hoefer, M.J. Ablowitz, I. Coddington, E.A. Cornell, P. Engels, and V. Schweikhard, *Phys. Rev. A* **74**, 023623 (2006).
  - [20] W. Wan, S. Jia, and J.W. Fleischer, *Nature Phys.* **3**, 46 (2007).
  - [21] U. Fano, *Nuovo Cimento* **12**, 156 (1935); H. Feshbach, *Ann. Phys. (N.Y.)* **5**, 357 (1958); U. Fano, *Phys. Rev.* **124**, 1866 (1961).
  - [22] R. Robinett, *Quantum Mechanics. Classical Results, Modern Systems, and Visualized Examples* (Oxford University Press, Oxford, 2006).
  - [23] N. Manini and L. Salasnich, *Phys. Rev. A* **71**, 033625

- (2005).
- [24] R.D. Richtmyer and K.W. Morton, *Difference methods for initial-value problems* (John Wiley, New York, 1967).