

E ffect of Loops on the V ibrational Spectrum  
of P ercolation N etw ork

H isao Nakanishi

H LR Z, K FA {J ulich, Postfach 1913

W -5170 J ulich, G erm any

<sup>y</sup>P resent and perm anent address: D epartm ent of P hysics, Purdue U niversity,  
W est Lafayette, Indiana 47907 U .S A .

W e study the e ffects of adding loops to a critical percolation cluster on the dif-  
fusional, and equivalently, (scalar) elastic properties of the fractal network. From  
the num erical calculations of the eigenspectrum of the transition probability m atrix,  
we find that the spectral dim ension  $d_s$  and the walk dim ension  $d_w$  change suddenly  
as soon as the oppy ends of a critical percolation cluster are connected together to  
form relatively large loops, and that the additional inclusion of successively sm aller  
loops only change these exponents little if at all. This suggests that there is a new  
universality class associated with the loop-enhanced percolation problem .

Keywords: P ercolation, D i usion, V ibration, Spectrum , A erogel

P A C S num bers: 64.60A , 82.70G , 05.40

## I. Introduction

One of the few experimentally accessible measures of the degree to which a structure is fractal [1, 2] is the spectral dimension  $d_s$  [3]. The spectral dimension was originally defined [4] from the low energy behavior of the vibrational density of states  $n(E)$  of a fractal, elastic network:

$$n(E) \propto E^{d_s/2-1}; \quad (1)$$

where  $E$  is the energy of a mode. The energy is proportional to  $\omega^2$  where  $\omega$  is the angular frequency of the vibrational mode, and writing the above asymptotic relation in terms of the density of states per unit interval in  $\omega$ , we have the relation

$$g(\omega) \propto \omega^{d_s-1}; \quad (2)$$

in which  $d_s$  has replaced the Euclidean dimension  $d$  in a more familiar equation for the phonon density of states.

The exponent  $d_s$  can be measured by various optical and neutron scattering experiments [5], and for some silica aerogels in particular, a value in the neighborhood  $1.3 \pm 0.01$  was found. Now, such a fractal is sometimes compared to the critical percolation cluster [6] since the latter is the simplest and best studied model of an equilibrium random fractal. If this comparison is made for the silica aerogel, the experimental value of  $d_s$  happens to be close to the  $d_s$  of the scalar elastic problem [4] on the critical percolation cluster in  $d = 3$ , and much larger than the corresponding vector elastic result of about  $d_s = 0.9$  [1]. (Vector elasticity is the problem where the displacement of each node in the network is a vector rather than a scalar.) Courtens [5] argued that this is not an evidence for scalar elasticity but rather that an aerogel has very few floppy endings unlike a percolation cluster. That is, most floppy endings which may form during the growth of an aerogel are connected together to form loops during the aging and (supercritical) drying processes. Thus the final structure is much more rigid with higher value of  $d_s$  although the fractal dimension  $d_f$  is essentially the same.

In this paper, we test this suggestion, but for the simpler, scalar elasticity problem, on two- and three-dimensional critical percolation clusters. In this case, each bond

connecting two (occupied) nearest neighbor sites of the cluster acts like an ideal spring of uniform spring constant in the sense that it exerts a Hooke's law force based on the difference in the (scalar) displacements of the two sites. The effects of loops are studied by adding controlled amount (and size) of loops and calculating the dynamic exponents  $d_s$  and  $d_w$  [3]. As far as we know, this is the first quantitative study of the effects of adding loops on the structural properties of a stochastic fractal.

We study this problem as a diffusion problem using the vibration {diffusion analogy} [4] for scalar elasticity. According to this correspondence, e.g., the vibrational spectrum of the network can be calculated as the spectrum of the transition probability matrix  $W$  [7, 8], where the elements  $W_{ij}$  of  $W$  is simply the hopping probability per step from node  $j$  to node  $i$  in the discrete network. Thus this matrix  $W$  contains the information on both the network topology and the dynamics of the model of diffusion. We choose for our calculations the so-called blind ant random walk [3] as a model of diffusion although the specific choice of the type of random walk is irrelevant for our results. For a blind ant, the random walker attempts to move without the knowledge of which neighbors are part of the cluster (thus available) and which ones are not, and thus when it happens to choose an unavailable neighbor, it cannot hop and must wait at its current position for the next time step.

From the point of view of the diffusion problem, the interesting quantities include the probability  $P(t)$  for a random walker to return to its starting point after  $t$  steps and the root-mean-square distance  $R(t)$  traveled by the random walk in time  $t$ . These quantities are expected to have the power-laws for asymptotically long times  $t$ :

$$P(t) \sim t^{-d_s=2} \quad (3)$$

$$R(t) \sim t^{1=d_w} : \quad (4)$$

These relations may be considered to define the exponents  $d_s$  and  $d_w$ .

We calculate these exponents by approximately diagonalizing the matrix  $W$  using the method of Ref. [9, 8]. Once the diagonalization is done, we compute two quantities, the density of eigenvalues  $n(\lambda)$  (where  $\lambda$  denotes the eigenvalues of  $W$ ) and a certain function  $\phi(\lambda)$  (which is the product of  $n(\lambda)$  and some coefficient determined when the stationary initial state distribution is expanded in terms of the eigenvectors of  $W$  [10]). These functions are expected to behave, asymptotically near  $\lambda = 1$  [8],

as

$$n(\mathbf{r}) = \sum_{\mathbf{j}} \mathbf{j} \cdot \mathbf{r}^{d_s=2-1} \quad (5)$$

$$(\mathbf{r}) = \sum_{\mathbf{j}} \mathbf{j} \cdot \mathbf{r}^{1-2=d_w} : \quad (6)$$

## II. Loop Addition

In this section, we characterize the clusters for which the calculations described in Section I are performed. The procedure is as follows: we first generate random realizations of occupied sites with the known critical probability  $p_c$  (independently of each other) on either a square or simple-cubic grid of a predetermined size (edge length  $L$ ). The percolation problem is then constructed by imposing the nearest-neighbor connectivity among the occupied sites using periodic boundary condition in all directions. We then choose only those realizations whose maximal cluster spans the grid in all directions as well as wraps in all directions, i.e., that those on which there is a path that winds around the grid (with the periodic boundaries) in each coordinate direction. The maximal, wrapping cluster generated in this way is an allowable starting configuration.

In order to add loops in a controlled manner, we first mark all perimeter sites which are neighbors to two or more occupied sites (i.e., those empty sites which would connect two or more occupied sites if they were occupied). For each of these multiple-perimeter sites, we must decide the size of the loop it closes. While it is very easy to determine the size of the smallest loop, that is not a very interesting quantity, since it may also close a much larger loop and such a larger loop may dominate the rigidity of the structure. On the other hand, the size of the largest loop itself is also not an interesting quantity since in principle there are many loops that are embedded in the interior of the cluster with little influence on the cluster's rigidity.

Thus we use the following procedure to determine the effective size of the loop a multiple-perimeter site closes: For each such site, we consider every pair of occupied neighbors in turn. For each pair, we calculate the minimum loop size by a burning method [11], spreading a fire from each end and determining the first contact of the two fires. Then we compute the maximum among these minimum loop sizes attached to the perimeter site in question. This is the appropriate loop size  $P$ , the size of the

oppest loop associated with the particular multiple-perimeter site. We then add all multiple-perimeter sites with  $P$  greater than or equal to a certain predetermined value  $P_0$  to control how floppy a loop must be for it to be closed. For example,  $P_0 = 1$  will close no loop, and  $P = 4$  will close all loops for both the square and simple cubic lattices.

In this paper, we compare the results for  $P_0 = 1, 2, 8$ , and  $6$ . A typical starting cluster and the corresponding loop-added clusters on the square lattice are shown in Fig.1. Clearly, the fuzzy and floppy endings present in the original cluster (a) become thicker and better connected successively in (b), (c), and (d) as more loops are added, giving the impression of image sharpening. The final structure in (d) with  $P_0 = 6$  appears to be a much more solid object than the original; yet, the fractal dimension of all these clusters are virtually the same, being about 1.9.

The above procedure is followed after the starting configuration is fixed. The applicable multiple-perimeter sites are ordered in a particular way (in a typewriter fashion due to the construction of the cluster), and the addition procedure applied only once in the fixed order, with no recursion. That is, once a site is determined to be either added or not, we do not later consider it again for addition and also we do not consider perimeter sites newly created by this process itself. This removes the problem of the complete filling of lakes and fjords which would occur particularly when  $P_0$  is small, but does not solve the problem of the final structure depending somewhat on the initial ordering of the perimeter sites. Physically, no recursion rule might correspond to the assumption that the formation of additional bonds during aging and drying is largely simultaneous and not sequential.

### III. Numerical Results

Our numerical results for  $n(\epsilon)$  for the density of states and  $\chi(\epsilon)$  (related to the velocity autocorrelation function of the random walk by a Laplace transform [9, 8]) are plotted in Fig.2 and Fig.3 respectively. We see that generally  $d = 3$  data appear to be behaving in a clearer way than those of  $d = 2$ . We summarize the exponent estimates for  $d_s$  and  $d_w$  in Tables I and II where the error estimates are mostly from the least squares regression and do not take into account any finite size effects or other systematic errors that may be present.

The grid sizes for the calculations shown in the figures are  $L = 100$  for the square lattice and  $L = 30$  for the simple cubic lattice (although we have also checked for the cluster size dependence using smaller grids of  $L = 50$  for the square lattice and  $L = 20$  for the simple cubic lattice). The spanning and wrapping percolation clusters (at  $p = 0.593 = p_c$  for the square and  $p = 0.312$  for the simple cubic lattice) contained about 4600 sites for  $d = 2$  and 4000 sites for  $d = 3$  before adding any loops. For  $P_o = 12$ , the size increases to about 5050 and 4400, for  $P_o = 8$ , to about 5250 and 4900, and for  $P_o = 6$ , to 6100 and 5700 for  $d = 2$  and  $d = 3$ , respectively. For each value of  $P_o$ , averages were taken over 400 independent cluster realizations for the square lattice, and for the simple cubic lattice, the number of clusters used varied slightly as 481, 480, and 479 for  $P_o = 12, 8$ , and  $6$ , respectively.

Thus, both the size of the clusters and the number of independent samples in the disorder ensemble are rather modest; however, they are comparable to the previous calculations by the same method for the  $P_o = 1$  case [8] where the results for  $p_c$  were in excellent agreement with other calculations of  $d_s$  and  $d_w$  as well as with the scaling relation  $d_s = 2d_f = d_w$  [3] (but see also [12]). The scope of the calculations were CPU time limited mainly because of the need to calculate the eigenvalues and eigenvectors very accurately (usually to 6 digits) for the largest 200 or so eigenvalues. These calculations took typically 3 hours of CPU time on one Cray Y-MP processor for each  $P_o$  for the square lattice and 6 hours for each  $P_o$  for the simple cubic lattice. Performing these calculations on workstations for long periods of time would solve the CPU limitation, but then the memory becomes the limiting factor.

Since the slope in the log-log plot of Fig.2 must equal  $d_s = 2 - 1$  (cf. Eq.(5)), it is obvious that the original percolation clusters ( $P_o = 1$ ) yield completely different values of  $d_s$  in both  $d = 2$  and  $3$  even from the case of  $P_o = 12$ . Indeed, the estimate of  $d_s$  is about 1.30 for both  $d = 2$  and  $d = 3$  if  $P_o = 1$ , but for the loop-added cases, the estimates are  $d_s = 1.7$  for  $d = 2$  and  $2.0$  for  $d = 3$ . Since the case  $P_o = 12$  contains only about 10 % more sites and visually looks very similar to the original percolation cluster (cf. Fig.1 (a) and (b)), this is a clear evidence that the structural properties are very sensitive to the addition of even a few loops. On the other hand, the slopes of the data points for the three finite values of  $P_o$  do not differ very much from each other. Thus once the larger loops are added, any further addition of smaller

loops do not appear to affect the results very much.

There does seem to be a tendency for the slope to get steeper when more loops are added. This tendency is within the statistical standard deviations for  $d = 2$  but definitely outside reasonable statistical errors for  $d = 3$ . A study of smaller clusters ( $L = 50$  for the square and  $L = 20$  for the simple cubic lattices) also shows different behaviors for  $d = 2$  and  $3$ . For the square lattice, this slope gets consistently steeper for all  $P_o = 12, 8$ , and  $6$  as  $L$  increases; however, the increase in the slope is well within the statistical standard deviation in all cases. For the simple cubic lattice, on the other hand, the slope becomes less steep for  $P_o = 12$  and steeper for  $P_o = 6$  as  $L$  increases, the changes being slightly outside of the statistical errors. These considerations suggest that the different  $d_s$  for different  $P_o < 1$  might be a real effect for  $d = 3$  but less likely to be so for  $d = 2$ . However, a more complete study, e.g., of the finite size effects is needed to definitively answer this question. In any case, the differences in the slope among the  $P_o = 12, 8$ , and  $6$  are far smaller than the difference between these values of  $P_o$  and  $P_o = 1$ .

Compared to the results for the density of states  $n(\cdot)$ , the results for  $(\cdot)$  in Fig.3 show a similar, but much less drastic relative differences in  $d_w$  between the loop-added clusters and the original percolation clusters. That is, the exponent  $d_w$  changes from about 2.9 for the original cluster to about 2.4 for the loop-added case for  $d = 2$ , a change of only about 17 % compared to the case of  $d_s$  where the change is more than 30 %. This can be understood since the addition of a few loops may not change the velocity autocorrelation of a random walker in  $t$  steps as much as its chances to return to the starting point, or put another way, the structural rigidity is much more sensitive to the addition of a few loops than the overall random walk displacement on the same network.

Again, the three-dimensional results appear to be better behaving than the two-dimensional ones. However, in both cases, the slopes in the log-log plot of Fig.3 are significantly steeper for  $P_o = 1$  than for the loop-added clusters, with a hint of a slight trend for the larger slope for  $P_o = 6$  in  $d = 3$ . Since this slope must equal  $1 - 2/d_w$  (cf. Eq.(6)), a larger slope implies a larger value of  $d_w$ . These differences among  $P_o < 1$  are, however, at least as ill-defined as for the case of  $n(\cdot)$ .

#### IV . Sum m ary

In sum m ary, we have presented the analysis of the dynam ical properties of the random walk con ned to loop-enhanced critical percolation clusters. Through the vibration{di usion analogy [4], this work has implications for the (scalar) elasticity problem on the same fractal network. From the num erical calculations of the eigenspectrum of the transition probability m atrix, we nd what appears to be the crossover to a new universality class characterized by a signi cantly larger spectral dimension  $d_s$  and signi cantly smaller walk dimension  $d_w$  as soon as the oppy ends of a critical percolation cluster are connected together to form relatively large bops. The additional inclusion of successively sm aller bops does not seem to further change the exponents  $d_s$  and  $d_w$  very m uch. This result supports the observation of Ref.[5] that the deviation between the experim ental observation of  $d_s$  of an aerogel and the previous theoretical calculations for the vector elasticity of the critical percolation cluster is due to the lack of oppy endings in the aerogel because of the chem ical reactions during aging and (supercritical) drying.

The loop-enhanced percolation problem is rem iniscent of the problem of the external surface of the critical percolation cluster. The latter can be de ned in several ways, e.g., by the percolation hull [6] and by the perim eter sites accessible to a random walk from the outside. The accessibility in that case depends on the size of the walker [13]; in a som ewhat sim ilar way, the oppiness of the loops in our problem depends on the m inim um loop size  $P_0$ . As in our case, the external perim eter problem also leads to one new universality class (in addition to the one for the hull) in two dim ensions [13]. One di erence is, however, that our problem leads to a new universality class also in three dim ensions while this is not so for the external perim eter problem .

Beyond the qualitative analogy, there m ight lurk a m ore quantitative one, since the problem of the external surface is that of successively controlling the accessibility for a particle di using from outside while our problem is that of successively controlling the reach of a particle di using on the cluster itself. This is, however, beyond the scope of this work and m ust await further research.



## Acknowledgments

I am grateful to Hans Herrmann for suggesting this problem and to him and Dietrich Staufer for helpful discussions. This work was carried out while I was visiting the HLRL Supercomputer Center at KFA {J{lich in Germany. I would like to thank the Center and Hans Herrmann for warm hospitality.

## References

- [1] B.Mandelbrot, *Fractal Geometry of Nature* (Freeman, San Francisco, 1982).
- [2] J.Feder, *Fractals* (Plenum, New York, 1988).
- [3] See, e.g., S.Havlin and D.Ben-Avraham, *Adv.Phys.* 36 (1987) 695.
- [4] S.Alexander and R.Orbach, *J.Phys.Lett. (Paris)* 43 (1982) 625.
- [5] E.Courtens, R.Vacher and E.Stoll, in *Fractals in Physics*, eds.A.Aharony and J.Feder (North-Holland, Amsterdam, 1989).
- [6] D.Stauffer and A.Aharony, *Introduction to Percolation Theory* (Taylor and Francis, London, 1992).
- [7] A.B.Harris, Y.Meir and A.Aharony, *Phys.Rev.B* 36 (1987) 8752.
- [8] H.Nakanishi, S.Muckherjee and N.H.Fuchs, *preprint* (1992).
- [9] N.H.Fuchs and H.Nakanishi, *Phys.Rev.A* 43 (1991) 1721.
- [10] D.Jacobs and H.Nakanishi, *Phys.Rev.A* 41 (1990) 706.
- [11] H.J.Herrmann, D.C.Hong and H.E.Stanley, *J.Phys.A* 17 (1984) L261.
- [12] H.Nakanishi and H.J.Herrmann, *HLRZ preprint*.
- [13] T.Grossman and A.Aharony, *J.Phys.A* 16 (1986) L745; 20 (1987) L1193.

## F I G U R E S

Fig. 1: The result of adding successively smaller loops to the critical percolation cluster is illustrated with: (a) a typical spanning and wrapping critical percolation cluster on the square lattice; (b) when all loops of loop parameter  $P = 12$  are added; (c) when all loops of  $P = 8$  are added; (d) when all loops of  $P = 6$  are added. The grid size is  $100 \times 100$ .

Fig. 2: Density of eigenvalues  $n(\lambda)$  for (a) the square lattice and (b) the simple cubic lattice. The symbols  $\circ$ ,  $\square$ ,  $\triangle$ , and  $\diamond$  correspond to  $P_0 = 1, 12, 8$ , and  $6$ , respectively. Lines drawn are linear least squares fits to the corresponding data shown here. The data for  $P_0 = 1$  are from Ref.[8].

Fig. 3: The function  $\chi(\lambda)$  for (a) the square lattice and (b) the simple cubic lattice. The symbols  $\circ$ ,  $\square$ ,  $\triangle$ , and  $\diamond$  correspond to  $P_0 = 1, 12, 8$ , and  $6$ , respectively. Lines drawn are linear least squares fits to the corresponding data shown here, and the data for  $P_0 = 1$  are from Ref.[8]

# TABLES

Table 1: Numerical estimates of the exponents  $d_s$  and  $d_w$  for loop-enhanced critical percolation cluster on the square lattice. The results for  $P_o = 1$  are from Ref.8. The error estimates are mainly from the least squares regression.

$P_o$	$d_s$		$d_w$	
1	1:30	0:02	2:86	0:04
12	1:69	0:12	2:40	0:12
8	1:66	0:10	2:42	0:12
6	1:63	0:11	2:39	0:15

Table 2: Numerical estimates of the exponents  $d_s$  and  $d_w$  for loop-enhanced critical percolation cluster on the simple cubic lattice. The results for  $P_o = 1$  are from Ref.8. The error estimates are mainly from the least squares regression.

$P_o$	$d_s$		$d_w$	
1	1:30	0:02	3:70	0:07
12	1:99	0:02	2:67	0:07
8	1:94	0:02	2:76	0:08
6	1:89	0:01	2:76	0:04