

Non-equilibrium relaxation and interface energy of the Ising model

Nobuyasu Ito

Institute for Theoretical Physics

University of Cologne

D-5000 Cologne 41, Germany

and

Computing and Information Systems Center

Japan Atomic Energy Research Institute

Tokai, Ibaraki 319-11, Japan

Abstract

From the non-equilibrium critical relaxation study of the two-dimensional Ising model, the dynamical critical exponent z is estimated to be 2.165 ± 0.010 for this model. The relaxation in the ordered phase of this model is consistent with $\exp(-t^\beta)$ behavior. The interface energy of the three-dimensional Ising model is studied and the critical exponent of the correlation length and the critical amplitude of the surface tension σ_0 are estimated to be 0.6250 ± 0.025 and 1.42 ± 0.04 , respectively. A dynamic Monte Carlo renormalization group method is applied to the equilibrium properties of the three-dimensional Ising model successfully.

1 Introduction

The study of the non-equilibrium relaxation turned out to be useful to study the critical slowing down and other dynamical aspect of the Ising models [1, 2, 3, 4, 5, 6].

For the critical relaxation, it is shown that the analysis based on the dynamical finite-size scaling theory [7] produces precise estimation of the value of β/ν , where β and ν denote the equilibrium critical exponents of the spontaneous magnetization and the correlation length, respectively, and ν denotes the dynamical critical exponent. It is estimated directly from the initial relaxation curve of the magnetization $m(t)$ which follows asymptotically $t^{-\beta/\nu}$. This method is simpler than the other previous methods which require the calculations of the two-time correlations and the estimations of the correlation times from them. In this paper, the values of β/ν for the two-dimensional Ising model is studied from the non-equilibrium relaxation. The exact values of β and ν of this model are known to be $1/8$ and 1 , respectively, and the value of ν is estimated from the value of β/ν .

The non-equilibrium relaxation analysis is useful also for the relaxation process in the ordered phase which is not yet fully understood. It has been pointed out that the equilibrium relaxation of the magnetization may be stretched exponential, that is,

$$C_m(t) \sim e^{-at}; \quad (1)$$

where $C_m(t)$ denotes the equilibrium time-correlation function of the magnetization and $a > 0$. Two phenomenologies predicts $a = 1/2$ [8] and $1/3$ [9] for the two-dimensional Ising model. In this paper, the non-equilibrium relaxation in the ordered phase is studied assuming the similar stretched exponential decay, that is,

$$m(t) - m_s \sim ae^{-bt}; \quad (2)$$

The value of β/ν for the three-dimensional Ising model was estimated to be 0.250 ± 0.002 [5]. The value of β/ν of this model has been estimated by several methods and the non-equilibrium critical relaxation provides one method to study its value. When the scale transformation with scale l is applied n times repetitively, the scale transformed magnetization is expected to follow the scaling form

$$m(t; n) = l^{-\beta/\nu} f(t l^{-\nu}) \quad (3)$$

at the critical point for large $t l^{n_z}$.

In this scaling region, the magnetization at the fixed time follows [10, 11]

$$m \sim t^{-1/n_z} : \quad (4)$$

When we use the total magnetization M instead of the magnetization per site, it behaves as

$$M \sim l^{-n(d-1)} : \quad (5)$$

This relation provides a new method of Monte Carlo renormalization group. The study of β for three-dimensional Ising model based on this scaling argument is also included.

The second problem of this paper, the interface energy and the surface tension [12]. In the recent studies of this problem [13, 14, 15, 16, 17, 18], it seems to be believed that the standard Monte Carlo simulation with single spin updating dynamics is not an appropriate method for this problem. In this paper, however, it is shown that the naive simulation works successfully and a precise study of the interface energy is made. We reconfirm the values of the critical exponent and the critical amplitude, which is important to verify the universality hypothesis about the critical amplitude [19].

The non-equilibrium relaxation study of the two-dimensional Ising model is described in the next section and the interface energy of the three-dimensional Ising model in the third section. The β analysis is given in the appendix A. The last section contains the summary.

2 Two-dimensional Ising model

The non-equilibrium relaxation phenomena is studied for the two-dimensional Ising model in this section.

2.1 Dynamics and algorithm

The Hamiltonian of the ferromagnetic Ising model on square lattices is

$$H(f, g) = K \sum_{i,j=1}^X f_i g_j \quad (K > 0; f_i = \pm 1); \quad (6)$$

where $\beta = 1/kT$ and the summation runs over all the nearest neighbor site pairs. The dimensionless constant K is used as the inverse temperature in the following. The Ising spins at the lattice points are denoted by

$$S_{(i,j)} \quad (i = 1; 2; \dots; L_x; j = 1; 2; \dots; L_y) \quad (7)$$

where L_x and L_y are assumed to be $L + 1$ and L , respectively, and L is an even integer. Skew boundary condition is applied to L_x direction and periodic boundary condition to L_y direction. There is no intrinsic dynamics of this model and the dynamics studied here is the single-spin updating stochastic dynamics with two-sublattice flip sequence and Metropolis-type transition probability. The spins are classified into two sublattices defined by

$$A = \{S_{(i,j)}; i+j = \text{even}\} \text{ and } B = \{S_{(i,j)}; i+j = \text{odd}\} \quad (8)$$

The spins in A are updated firstly and then those in B are updated. The transition probability is

$$p_M(S_0 \rightarrow S_0; S_1; S_2; S_3; S_4) = \min[1, \exp(-2\beta E(S_0; S_1; S_2; S_3; S_4))] \quad (9)$$

where S_1, S_2, S_3 and S_4 are the nearest-neighbor spins of S_0 and the local energy E denotes

$$E(S_0; S_1; S_2; S_3; S_4) = K S_0(S_1 + S_2 + S_3 + S_4) \quad (10)$$

This dynamics is efficiently vectorized and it is the reason why it is used. 64 independent systems with the same lattice are simulated simultaneously. The 64 spins on the same site of these 64 systems are coded in the same eight-byte integer variables, that is, the independent-system coding technique is employed [20, 21, 22]. The spin-updating procedure is realized by 15 logical operations and the generation of one random integer. The condition that a spin S_0 is flipped to $-S_0$ is

$$r \leq p_M(S_0 \rightarrow -S_0; S_1; S_2; S_3; S_4) \quad (11)$$

where r is a random number. The transition probability p_M is the function of local energy $E(S_0; S_1; S_2; S_3; S_4)$. The spin variable s_i which takes -1 or $+1$ value is stored in one bit of one computer word which takes 0 or 1 value and

the corresponding bit is denoted by I_i . Now the above condition is expressed by

$$I_1 = I_0 + I_2 \oplus I_0 + I_3 \oplus I_0 + I_4 \oplus I_0 + 2 \oplus IX\ 1T(r) + IX\ 2T(r) \oplus 2; \quad (12)$$

where \oplus denotes the exclusive-OR operation and $IX\ 1T(r)$ and $IX\ 2T(r)$ are the functions of random number r defined by

$$IX\ 1T(r) = \begin{cases} 1 & r \leq e^{8K} \\ 0 & \text{otherwise} \end{cases} \quad (13)$$

and

$$IX\ 2T(r) = \begin{cases} 1 & e^{8K} < r \leq e^{4K} \\ 0 & \text{otherwise} \end{cases} : \quad (14)$$

This condition should be checked by the bitwise logical operations. The following operations realize it with 14 logical operations. In the following, $\langle I;J \rangle$ denotes 2^{I+J} .

1. Make $J1$ and $J2$ so that $\langle J1;J2 \rangle = I1 + I2 + I3$. It takes five operations.
2. Take the exclusive-OR of $J1, J2$ and $J4$ with $I0$, respectively. It takes three operations. The same notations are used for the results.
3. Make $K1$ and $K2$ so that $\langle K1;K2 \rangle = I4 + IX\ 1T(r) + IX\ 2T(r)$. It takes three operations.
4. Make ID which holds the truth value of spin- \uparrow condition by

$$K1 \text{ or } J1 \text{ or } (J2 \text{ and } K2): \quad (15)$$

It takes three operations.

Further details of this updating algorithm are found in the FORTRAN routines given in the Appendix B. To do the independent simulation of the systems at the same temperature with the independent-system coding technique, the recycle algorithm of the random numbers is used by shuffling the Boltzmann factor tables[23]. The RNDTICK routine[24] are used for the random number generations.

The NEC SX3/11 of Kohn university and the MONTE-4 of Japan Atomic Energy Research Institute are used for the simulation. The performance of the simulation codes developed for the present study, named IS2DVP, on these vector processors is shown in Table 1. The MONTE-4 has four processors and they share the main memory. So the simulation speed on one processor is influenced by other jobs on other processors. The performance in Table 1 is measured with/without other jobs on other processors. The concurrent jobs were Monte Carlo simulations of two- and three-dimensional Ising models which are highly vectorized. The speed-down shown in Table 1 under multi-job environment can be regarded as the worst case. If the concurrent jobs are badly vectorized jobs or the jobs which are highly vectorized but do not generate the main memory access massively, the speed-down becomes smaller. In fact, almost no speed-down was observed when the performance of IS2DVP is measured with three lowly vectorized jobs.

The value of magnetization at MCS from all up initial condition is denoted by $m(t)$. This time-dependent magnetization of the above-mentioned dynamics is estimated by the simulation. It is estimated from the values of magnetization at t of independent Monte Carlo runs.

2.2 At the critical point

The critical point of the square-lattice Ising model is $K_c = J/k_B T_c = 0.440686793$ where the time-dependent magnetization $m(t)$ decays to zero following asymptotically to

$$m(t) \sim t^{-z} = t^{-1/8}; \quad (16)$$

where the exact values of the critical exponents of the spontaneous magnetization and the correlation length, $\nu = 1/8$ and $\beta = 1$, are used.

The simulations are listed in Table 2. The deviation of $m(t)$ due to the finite lattice is expected to be exponentially small for large system. The system-size dependence at several time are shown in Fig. 1. No size dependence is observed up to $t = 400$ MCS, although it is observed at $t = 500$ MCS. In the following, the estimated values of $m(t)$ for $L = 1500$ lattice up to 400 MCS which are shown in Fig. 2 are used for estimating the value of z .

Firstly the estimator

$$R(t; m) = t \left[\frac{m(t-1)}{m(t)} - 1 \right] \quad (17)$$

is used for this purpose. The average of this $R(t)$ over N successive t , that is,

$$R_{\text{ave.}}^N(t; N; m) = \frac{1}{N} \sum_{t=t_0}^{t_0+N-1} R(t; m); \quad (18)$$

is used to reduce the statistical fluctuation, where the mean time, $(t; N) = t_0 + (N-1)/2$. The values of $z_{\text{ave.}}^{25} = 1/(8R_{\text{ave.}}^{25})$ is plotted in Fig. 3. If the form of $m(t)$ is $t^{\alpha} (a_0 + a_1 t + a_2 t^2 + \dots)$, $R_{\text{ave.}}^N(t; m)$ is the form of $t^{\alpha} + d_1 t^{-1} + d_2 t^{-2} + \dots$. It is observed that the value of z converges to 2.165 ± 0.005 assuming that the correction to the asymptotic simple power law is analytic, which is consistent with our data.

The estimates for z using the least-square fitting analysis are shown in Fig. 4. The function of the form of $\log m(t) = a + b \log t$ is fitted to the successive fifty values of m at magnetization, that is, $m(t), m(t+1), \dots, m(t+49)$ and $m(t+49)$. The values of $z_{\text{fitting}}^{50} = 8/b$ are plotted. In the large time limit, the value of z estimated from this Fig. 4 is consistent with the estimated $z = 2.165 \pm 0.005$ from the ratio analysis.

There are many theoretical studies on this value of z [25, 26, 27, 28, 29]. The estimates for the two-dimensional Ising model are listed in Table 3 and they are plotted in Fig. 6. It is observed that the simulational values which includes the estimates obtained by temperature dependence analysis, finite-size scaling analysis, Monte Carlo renormalization group analysis, non-equilibrium relaxation analysis and other analysis of the simulational results are converging.

Above analysis assumes the form of the function $m(t)$ to be a simple power-law with simple correction terms and the results might have unknown systematic error caused by this assumption. Therefore the naive error region might be underestimated and an overestimated error region will be safer as our present conclusion. So we adopt $z = 2.165 \pm 0.010$ as our conclusion.

There is a room for the qualitatively different assumption for the critical-slowing down phenomena from above-mentioned simple power law. For example, a behavior with logarithmic correction to the naive diffusive behavior, $z = 2$, was proposed [65]. Assuming $m(t)$ to be $t^{-z} (\log t)^c$, where $-z = 1/6$, the values of $t^{1/6} m(t)$ is plotted in Fig. 5. The value of c is roughly estimated to be 0.04 from the tangent of the plot about $t = 200 \sim 400$ ($\log t = 5.3 \sim 6.0$), although the convex curvature seems to remain.

2.3 In the ordered phase

The simulations listed in Table 4 are made. Each run started from all up initial configuration. In the previous section, we observed that the value of the magnetization of $L = 1500$ lattice up to 400 MCS at the critical point can be regarded as values in the thermodynamics limit with the present accuracy. Now the simulation is made at β -critical temperature in the ordered phase and the maximum Monte Carlo step is up to 90 or 35. So the results can also be regarded as the values in the thermodynamics limit within the present accuracy.

The estimated values of magnetization are plotted in Fig. 7. The exact solution of the spontaneous magnetization,

$$m_s = [1 - \frac{1}{\sinh^4(2K)}]^{1/8}; \quad (19)$$

is used. The asymptotic slopes of these curves correspond to the value of β .

At $t = 0$ where $m = 1$, the values of $\ln[1 - \ln(m - m_s)]$ are 0.647, 0.819, 0.943 and 1.041 for $K = 0.47, 0.49, 0.51$ and 0.53 , respectively, and these values correspond to $\ln t = -1$. Therefore the tangents of the curves in Fig. 7 approach the β value from smaller side and the curves are concave.

The value of β observed from the Fig. 7 is clearly larger than $1/3$. The results for $K = 0.51$ and 0.53 are consistent with $\beta = 1/2$. For $K = 0.47$ and 0.49 , the curves do not reach $\beta = 1/2$ behavior yet, but the β tangent are already larger than $1/3$ which is observed in Fig. 8.

We cannot exclude the possibility that the curves continue to become steeper for longer time and finally the simple exponential decay ($\beta = 1$) appears. We can conclude that the conjecture $\beta = 1/3$ is not true for the non-equilibrium relaxation of this model in the presently studied time region.

3 Three-dimensional Ising model

The interface energy of the three-dimensional Ising model is studied in this section.

3.1 Monte Carlo simulation

The interface energy is estimated by the difference of the energy in the systems with and without a interface. In this study, the different boundary conditions are used to realize the constraint for the interface. One lattice has periodic boundary conditions for all of the x , y and z directions. In this case, the conformation without interface is dominant because the present simulation is started from all-up initial conformation and we take lattice size to infinity at fixed t . The other lattice has anti-periodic boundary condition in the z direction and periodic to the x and y direction. This means that the lattice is pasted into torus with anti-ferromagnetic bond with coupling constant K (< 0) for z direction. In this case, a single interface conformation is dominant because the all-up initial conformation has one interface in the case of the anti-periodic lattice.

The lattice size is specified by two integers, L and H , in the following. We use the box of cubic lattice of the size of $L \times L \times H$ for x , y and z directions. Odd L and even H are used for the purpose of vectorizing the Monte Carlo dynamics.

The total energies of periodic and anti-periodic lattices are denoted by E_p and E_{ap} . The interface energy per unit area denoted by σ_s is defined by

$$\sigma_s(K; L; H) = (E_{ap} - E_p)/L^2; \quad (20)$$

In the thermodynamic limit, the critical behavior of this interface energy is expected to be

$$\sigma_s(K) - \sigma_s(K; L=1; H=1) = \frac{2\sigma_0}{K} (1 - \frac{K_c}{K})^{2-\beta}; \quad (21)$$

where σ_0 denotes the critical amplitude of the surface tension and the factor $1=K$ is added instead of $1=K_c$ because it gives a better description of the present results and the critical behavior is essentially the same.

The simulations listed in Table 5 were made on MONTE-4 of JAERI and NEC SX3/11 of Köln University. In each simulation 64 independent simulations were run simultaneously using the independent-system coding technique[20, 21, 22] and the recycle algorithm for the random number[23]. Half of these 64 systems have the periodic boundary condition and the rest have the anti-periodic boundary condition. All the simulations here are made in the ordered phase above the surface roughening temperature. The triplet

(A, B, C) in the MCS column of Table 5 denotes the length of the simulation and the energy measurement timing. From all-up initial configuration, A MCS are made without energy calculation. The value of A is selected to be twenty or more times larger than the correlation time of the energy. Then the energy is calculated C times in every B MCS and one estimate of the interface energy is calculated using the estimates of the energy of 64 systems. These operations are repeated five times independently. If five runs can be made within the CPU time limit of one batch job, the initialization of the configuration is not made for every run but the last configuration of the previous run is used as the next initial configuration. The obtained five estimates of the interface energy are used to estimate the average, $\sigma_s(K; L; H)$, and its error.

3.2 Finite size effect

For a fixed L, the interface energy will be independent of the height H if H is larger than the interface thickness which is known to increase as $\log L$, as far as there is no interface in the periodic lattice and there is only one interface in the anti-periodic lattice. Fig. 9 shows the H dependence of the estimates of $\sigma_s(K; L; H)$. It is observed that $\sigma_s(K; L; H)$ is independent of H if H is not very small. The horizontal solid line and the broken lines show the estimate of $\sigma_s(K; L) = \sigma_s(K; L; H = 1)$ and its 1 σ error region, which is estimated from the data $H \geq 20$, that is, all the data except two small H points shown in the figure.

For every L at every K, lattices with several H are made. The values of the interface energy were estimated from the estimates for the H-independent region. The region of H used for this procedure is given in Table 6 with the values of σ_s .

Now the extrapolation to $L \rightarrow 1$ has to be made. The finite size correction to $\sigma_s(K)$ is expected to be the order of $1/L^2$ from the capillary wave contribution, that is,

$$\sigma_s(K; L) = \sigma_s(K; L = 1) + \frac{a}{L^2}; \quad (22)$$

up to the first order correction. The values of the interface energy in the thermodynamic limit $\sigma_s(K) = \sigma_s(K; L = 1)$ are estimated from the estimates of $\sigma_s(K; L)$ with this size dependence. This $1/L^2$ correction is appropriate

to our results. One example of this extrapolation is shown in Fig. 10. The estimated values of $\chi_s(K)$ are also given in Table 6.

3.3 Critical behavior

The critical exponent ν and the critical amplitude χ_0 are estimated from our estimates of χ_s . The present estimates of χ_s are plotted in Fig. 11.

The value of the critical point K_c has been estimated by several methods and usually is in the region $0.22165 - 0.22166$ [82] and this accuracy is enough precise for the present analysis. So $K_c = 0.221657$ [83, 84] is used for the present analysis.

The tangent of the consecutive two points in Fig. 11 are shown in Fig. 12 and the value of $2 - 1$ at $K = K_c$ is estimated to be $0.250(5)$ from this figure. This means that the estimate of ν is $0.6250 - 0.0025$. This values of ν is consistent with the result of a recent Monte Carlo renormalization study [82]. The previous estimates of ν are also cited in Ref. [82].

The value of χ_0 is estimated from Fig. 13 which shows the values of

$$\chi_0(\nu) = \chi_s(K) \frac{K}{2} \left(1 - \frac{K_c}{K}\right)^{1-\nu}; \quad (23)$$

from our estimates of $\chi_s(K)$ for several values of ν . From this figure and above estimate of ν , we estimate that the value of χ_0 is $1.42 - 0.04$. The estimates in the previous works are shown in Table 7 and are mostly confirmed by our more accurate data.

4 Summary

First the non-equilibrium relaxation of the two-dimensional ferromagnetic Ising model was studied. The critical relaxation exponent z was estimated to be $2.165 - 0.010$. This value is smaller than the recent other estimations for this value from the same non-equilibrium relaxation method by Munkel et al [6] and from the high-temperature expansion study by Dammann and Reger [79]. But the upward behavior of the two-dimensional data shown in Fig. 1 of Ref. [6] can be interpreted as a sample fluctuation and this might cause a overestimation. The coefficients of the high-temperature expansion given in Ref. [79] were reanalyzed by Adler [81] and an estimate consistent

with the present result was reported. The relaxation process in the ordered phase is stretched exponential. The exponent is larger than $1/3$ [9] and it is consistent with $1/2$ [8].

Secondly the critical exponent ν and the critical amplitude ϕ_0 are estimated to be 0.6250 ± 0.0025 and 1.42 ± 0.04 , respectively. The fully vectorized simulation with single spin updating dynamics can efficiently study the interface. When we used only about 20 hours of NEC SX3/11, the data were as accurate as those of Ref. [17] from 10^3 work station hours.

Finally the new Monte Carlo renormalization group method produced reasonable estimate of β with a quite small scale simulation.

The total number of updated spins are 3.19×10^{15} .

Acknowledgment

The author thanks D. Stauffer for his valuable advice and discussion. This study is partially supported by SFB 341.

Appendix A

A new Monte Carlo renormalization group method is applied to the estimation of β value of the three-dimensional Ising model. The value of the magnetization of the scale transformed configuration with scale $l = 2$ majority-rule real-space scale transformation is estimated. A 256^3 lattice is simulated up to 2000 MCS from all-up initial configuration with an appropriate Monte Carlo dynamics. Totally 45 independent simulations are made and therefore the total number of the updated spin is 1.51×10^{13} .

Following the eq. (5), the values of $\langle n; t \rangle$ defined by

$$\langle n; t \rangle = \frac{\log[M(n-1; t) - M(n; t)]}{\log l} \quad (24)$$

are studied. This $\langle n \rangle$ is expected to be $d \log 2 = 12$, where d is the lattice dimensionality, in the limit of $t \rightarrow \infty$ if the infinitely large lattice is studied. The behavior of the $\langle n; t \rangle$ is shown in Fig. 14. From this figure, the value of $\langle n; t = 1 \rangle$ is estimated to be $2.4625(1)$. In table 8, the estimated values of $\langle n \rangle$ ($\langle n; t = 1 \rangle$) are shown for 256^3 and 1024^3 lattice. The 1024^3

lattice was simulated up to 5644 MCS for one sample and the number of the updated spin is 6.06×10^{12} .

When we extrapolate these estimates of $\langle n \rangle$ are extrapolated to $n \rightarrow 1$ with

$$\langle n \rangle = \langle n \rangle_0 + c_1 n^{\alpha}; \quad (25)$$

the value of $\langle n \rangle$ behaves as shown in Fig. 15. The estimates for the 256^3 lattice given in Table 8 are used for the extrapolation. From the present preliminary data, the value of α is estimated to be 0.7(2) and the corresponding value of $\langle n \rangle_0$ is 2.492(8). this means the value of $\langle n \rangle = 0.508(8)$. If the estimate $\alpha = 0.80$ [0.85[82]] is assumed, $\langle n \rangle$ is estimated to be 2.487 [2.490] and $\langle n \rangle_0 = 0.513$ [0.510]. If we assume $\alpha = 1$, $\langle n \rangle$ and $\langle n \rangle_0$ are 2.486 and 0.514, respectively.

Appendix B

Two routines of IS2DVP code which are specific to the two-dimensional Ising model are shown. BFL2DF generates the Boltzmann factor tables and SU2DSK updates the spin configuration.

```
C*****
C IS2DVP: Monte Carlo simulation of two-dimensional Ising Model
C
C Boltzmann-factor table preparation routine
C   METROPOLIS-TYPE TRANSITION PROBABILITY
C   1992.09.19. VERSION 1.00 BY NOBUYASU ITO
C
C TK(64) R*8 : Give a inverse temperature of 64 systems
C IX1(0:IRLST), IX2(0:IRLST) I*8 : The BFT is generated in them.
C   IRLST is maximum number of the random integer.
C*****
      SUBROUTINE BFL2DF(TK, IX1, IX2, IRLST)
      PARAMETER (IWIDTH=64)
      DIMENSION TK (IWIDTH)
      REAL*8 TK
      DIMENSION IX1 (0:IRLST), IX2 (0:IRLST)
      DIMENSION IK1 (IWIDTH), IK2 (IWIDTH), I2P (IWIDTH)
```

```

      REAL*8  TNORM
C
      TNORM=DFLOAT (IRLST)
      DO 10 I=1, IWIDTH
          IK1 (I)=IDINT (TNORM*DEXP (-8.0D0*TK (I) ))
          IK2 (I)=IDINT (TNORM*DEXP (-4.0D0*TK (I) ))
10  CONTINUE
      DO 30 I=1, IWIDTH
          I2P (I)=ISHFT (1, (IWIDTH-I) )
30  CONTINUE
      DO 60 I=0, IRLST
          IX1 (I)=0
          IX2 (I)=0
60  CONTINUE
*VDIR NODEP
      DO 50 I=1, IWIDTH
      DO 40 J=0, IRLST
          IXT1=0
          IXT2=0
          IF (J.LE.IK1 (I) ) IXT1=I2P (I)
          IF (J.LE.IK2 (I) ) IXT2=I2P (I)
          I2=IEOR (IXT1, IXT2)
          I1=IAND (IXT1, IXT2)
          IX2 (J)=IOR (IX2 (J) , I2)
          IX1 (J)=IOR (IX1 (J) , I1)
40  CONTINUE
50  CONTINUE
      RETURN
      END

C*****
C IS2DVP: Monte Carlo simulation of two-dimensional Ising Model
C
C Spin configuration updating routine for
C ferromagnetic Ising model with Metropolis-type transition
C probabilities
C BFL2DF routine should be used for BFT preparation

```

```

C      1992.09.19. VERSION 1.00 BY NOBUYASU ITO
C
C  ISTEP I*8 : number of Monte Carlo sweeps
C  L1, L2 I*8: lattice size
C  IS((-L1+1):(L1*(L2+1)) I*8: spin configuration
C  IRD(L1*L2) I*8: work area
C  IRLST I*8: Maximum number of random integer
C  IX1(0:IRLST), IX2(0:IRLST) I*8: Boltzmann factor tables(BFT)
C*****
      SUBROUTINE SU2DSK(ISTEP,L1,L2,IS,IRD,IRLST,IX1,IX2)
      DIMENSION IS((-L1+1):(L1*(L2+1)))
      DIMENSION IRD(L1*L2)
      DIMENSION IX1(0:IRLST),IX2(0:IRLST)
      NSYS=L1*L2
      LS=L1
      DO 10 IMCS=1,ISTEP
      CALL RNDO2I(NSYS,IRD)
      IFIRST=1
*VDIR NODEP
      40 DO 20 I=-LS+1,0
      20 IS(I)=IS(I+NSYS)
*VDIR NODEP
      DO 30 I=NSYS+1,NSYS+LS
      30 IS(I)=IS(I-NSYS)
*VDIR NODEP
      DO 50 IJ=IFIRST,NSYS,2
      I0=IS(IJ)
      I1=IS(IJ+1)
      I2=IS(IJ-1)
      I3=IS(IJ+L1)
      K1=IEOR(I1,I2)
      K2=IAND(I1,I2)
      J2=IEOR(K1,I3)
      K3=IAND(K1,I3)
      J1=IOR(K2,K3)
      J1=IEOR(I0,J1)
      J2=IEOR(I0,J2)

```

```

      I4=IS (IJ-L1)
      I4=IEOR (I4, I0)
      IRT=IRD (IJ)
      IX1T=IX1 (IRT)
      IX2T=IX2 (IRT)
      K2=IEOR (I4, IX2T)
      K1T=IAND (I4, IX2T)
      K1=IOR (K1T, IX1T)
      ID=IOR (J1, K1)
      K4=IAND (J2, K2)
      ID=IOR (ID, K4)
      IS (IJ)=IEOR (IS (IJ) , ID)
50  CONTINUE
      IF (IFIRST.EQ.1) THEN
          IFIRST=2
          GOTO 40
      END IF
10  CONTINUE
      RETURN
      END

```

References

- [1] M . K ikuchi and Y . O kabe, J.Phys. Soc. Jpn. 55 (1986) 1359.
- [2] D . Stau er, Physica A 186 (1992) 197.
- [3] D . Stau er, Int. J. M od. Phys. C 3 (1992) 287.
- [4] G . A . K ohring and D . Stau er, Int. J. M od. Phys. C .
- [5] N . Ito, to appear in Physica A (1993).
- [6] C . M unkel and D . W . Heerm ann, J. Adler, M . G ofm an and D . Stau er,
to appear in Physica A (1993).
- [7] M . Suzuki, Phys. Lett. A 58 (1976) 435.
- [8] D . A . H use and D . S . F ischer, Phys. Rev. B 35 (1987) 6841.

- [9] H. Takano, H. Nakanishi and S. Miyashita, *Phys. Rev. B* 37 (1988) 3716.
- [10] N. Ito and M. Suzuki, *Prog. Theor. Phys.* 77 (1987) 1391.
- [11] N. Ito and M. Suzuki, *J. de Physique C* 8 (1988) 1565.
- [12] B. Widom, *Phase Transition and Critical Phenomena* ed. C. Domb and F. Green 2 (Academic Press, London, 1972) 79.
- [13] M. Hasenbusch, to be published in *J. Phys. I*.
- [14] S. Klessinger and G. Munster, preprint.
- [15] G. Munster, *Int. J. Mod. Phys. C* 3 (1992).
- [16] B. A. Berg, U. Hansmann and T. Neuhaus, preprint.
- [17] M. Hasenbusch and K. Pinn, *Physica A* 192 (1993) 343.
- [18] H. Gausterer, J. Potvin, C. Rebbi and S. Sanielevici, *Physica A* 192 (1993) 343.
- [19] K. Binder, *Phase Transition and Critical Phenomena* ed. C. Domb and J. L. Lebowitz 8 (Academic Press, London, 1983) 1.
- [20] N. Ito and Y. Kanada, *Supercomputer* 5 No. 5 (1988) 31, see also G. Bhanot, D. Duke and R. Salvador, *Phys. Rev. B* 33 (1986) 7841 and C. Michael, *Phys. Rev. B* 33 (1986) 7861.
- [21] N. Ito and Y. Kanada, *Proc. Supercomputing '90* (New York) (IEEE Computer Society Press, 1990, Los Alamitos) 753.
- [22] N. Kawashima, N. Ito and Y. Kanada, to be published in *Int. J. Mod. Phys. C*.
- [23] N. Ito, M. Kikuchi and Y. Okabe, to be published in *Int. J. Mod. Phys. C*.
- [24] N. Ito and Y. Kanada, *Supercomputer* 7 No. 1 (1990) 29.
- [25] K. Kawasaki, in *Phase Transition and Critical Phenomena* 2 ed. C. Domb and M. S. Green (Academic Press, London, 1972) 443.

- [26] P. C. Hohenberg and B. I. Halperin, *Rev. Mod. Phys.* 49 (1977) 465.
- [27] G. F. Mazenko and O. T. Valls, *Phys. Rev. B* 24 (1981) 1419.
- [28] D. P. Landau, S. Tang and S. Wansleben, *J. de Physique C* 8 (1988) 1525.
- [29] M. D. Lacasse, J. Vinals and M. Grant, to be published in *Phys. Rev. B*.
- [30] K. Kawasaki, *Prog. Theor. Phys.* 40 (1968) 706.
- [31] H. Yahata and M. Suzuki, *J. Phys. Soc. Jpn.* 27 (1969) 1421.
- [32] H. Yahata, *J. Phys. Soc. Jpn.* 30 (1971) 657.
- [33] B. I. Halperin, P. C. Hohenberg and S.-K. Ma, *Phys. Rev. Lett.* 29 (1972) 1548.
- [34] E. Stoll, K. Binder and T. Schneider, *Phys. Rev. B* 8 (1973) 3266.
- [35] C. De Dominicis, E. Brezin and J. Zinn-Justin, *Phys. Rev. B* 12 (1975) 4945.
- [36] H. C. Bolton and C. H. J. Johnson, *Phys. Rev. B* 13 (1976) 3025.
- [37] Z. Racz and M. F. Collins, *Phys. Rev. B* 13 (1976) 3074.
- [38] S.-K. Ma, *Phys. Rev. Lett.* 37 (1976) 461.
- [39] Y. Achiam, *J. Phys. A* 11 (1978) L129.
- [40] Y. Achiam and J. M. Kosterlitz, *Phys. Rev. Lett.* 41 (1978) 128.
- [41] W. Kinzel, *Z. Phys. B* 29 (1978) 361.
- [42] G. F. Mazenko, M. J. Nolan and O. T. Valls, *Phys. Rev. Lett.* 41 (1978) 500.
- [43] M. C. Yalabik and J. D. Gunton, *Prog. Theor. Phys.* 62 (1979) 1573.
- [44] Y. Achiam, *Phys. Rev. B* 19 (1979) 376.

- [45] M .Suzuki, K .Sogo, I.M atsuba, H .Ikeda, T .Chikam a and H .Takano,
Prog.Theor.Phys. 61 (1979) 864.
- [46] S.T .Chui, G .Forgacs and H .L.Frisch, Phys.Rev.B 20 (1979) 243.
- [47] S.K .M a, Phys.Rev.B 19 (1979) 4824.
- [48] M .P.N ightingale and H .W .J.B lote, Physica A 104 (1980) 352.
- [49] G .F.M azenko, M .J.Nolan and O .T .Valls, Phys.Rev.B 22 (1980)
1275.
- [50] G .Forgacs, S.T .Chui and H .L.Frisch, Phys.Rev.B 22 (1980) 415.
- [51] Y .A chiam , J.Phys.A 13 (1980) 1355.
- [52] G . F . M azenko, J. E . Hirsch, M . J. Nolan and O . T . Valls,
Phys.Rev.Lett.44 (1980) 1083.
- [53] K .Binder, J.Stat.Phys. 24 (1981) 69.
- [54] J.Tobochnik, S.Sarker and R.Cordery, Phys.Rev.Lett. 46 (1981)
1417.
- [55] R.Bausch, V.D ohm , H .K .Janssen and R.K .P .Zia, Phys.Rev.Lett.
47 (1981) 1837.
- [56] N .Jan and D .Stau er, Phys.Lett.A 93 (1982) 39.
- [57] J.C .Angles d'Auriac, R.M aynard and R.Ram m al, J.Stat.Phys. 28
(1982) 307.
- [58] H .Takano, Prog.Theor.Phys. 68 (1982) 493.
- [59] S.L.K atz, J.D .G unton and C.P.Liu, Phys.Rev.B 25 (1982) 6008.
- [60] M .C .Yalabik and J.D .G unton, Phys.Rev.B 25 (1982) 534.
- [61] N .Jan, L.L.M oseley and D .Stau er, J.Stat.Phys. 33 (1983) 1.
- [62] F.Haake and M .Lewenstein, Phys.Rev.B 27 (1983) 5868.
- [63] F.Haake, M .Lewenstein and M .W ilkens, Z.Phys.B 54 (1984) 333.

- [64] C .K alle, J.Phys.A 17 (1984) L801.
- [65] E .D om any, Phys. Rev. Lett. 52 (1984) 871.
- [66] J.O .Indekeu, A .L .Stella and J.R ogiers, Phys. Rev. B 32 (1985) 7333.
- [67] G .F .M azenko and O .T .Valls, Phys. Rev. B 31 (1985) 1565.
- [68] S.M iyashita and H .Takano, Prog. Theor. Phys. 73 (1985) 1122.
- [69] J.K .W illiam s, J.Phys. A 18 (1985) 49.
- [70] H .J.H erm ann, J. Stat. Phys. 45 (1986) 145.
- [71] L .de A rcangelis and N .Jan, J.Phys. 19 (1986) L1176.
- [72] O .F .de A lcantara Bon m , Europhys. Lett. 4 (1987) 373.
- [73] N .Ito, M .Taiji and M .Suzuki, J.Phys. Soc. Jpn. 56 (1987) 4218.
- [74] S.Tang and D .P .Landau, Phys. Rev. B 36 (1987) 567.
- [75] J.G .Zabolitzky and H .J.H erm ann, J. Com p. Phys. 76 (1988) 426.
- [76] M .M ori and Y .T suda, Phys. Rev. B 37 (1988) 5444.
- [77] J.R ogiers and J.O .Indekeu, Phys. Rev. B 41 (1990) 6998.
- [78] D .Stau er, Physica A 184 (1992) 201.
- [79] B .D am m ann and J.D .R eger, to be published in Europhys. Lett.
- [80] J.W ang, Phys. Rev. B 47 (1993) 869.
- [81] J.A dler, private com m unication.
- [82] C .F .Baillie, R .G upta, K .A .H aw ick and G .S .Paw ley, Phys. Rev. B 45 (1992) 10438.
- [83] N .Ito and M .Suzuki, J.Phys. Soc. Jpn. 60 (1991) 1978.
- [84] N .Ito, A IP C onf.P roc.248 C om puter-A ided Statistical Physics (Taipei, Taiwan, 1991) ed.C .K .H u (A IP, New York, 1990) 136.

- [85] K .B nder, Phys. Rev. A 25 (1982) 1699.
- [86] K .K .M on and D .Jasnow , Phys. Rev. A 31 (1985) 4008.
- [87] K .K .M on, Phys. Rev. Lett. 60 (1988) 2749.

Figure 1: The size-dependences of $m(t)$ at (a) $t = 10$, (b) 100, (c) 400 and (d) 500 are shown.

Figure 2: The estimated values of $m(t)$ for $L = 1500$ lattice up to 400 MCS are shown in logarithmic scale. They are regarded as those in the thermodynamic limit within the present accuracy.

Figure 3: The value of the averaged local exponent z_{ave}^{25} is shown. The magnetization obtained for $L = 1500$ lattice is used.

Figure 4: The estimates of z from the least-square fitting are shown. $(t; 50)$ denotes the mean time of the data, that is, $t + (50 - 1)/2$.

Figure 5: The values of $m(t)$ is plotted assuming the logarithmic correction.

Figure 6: The estimates for z are plotted. The points marked with \square and \triangle show simulational and other results, respectively. The horizontal axis shows the publication year except the recent preprints.

Figure 7: The non-equilibrium relaxation curves of magnetization are shown. The point marked by \square , \triangle , \circ and \times correspond to $K = 0.47, 0.49, 0.51$ and 0.53 , respectively. The tangent of the solid line is $\beta = 1/2$. Those of broken lines are $\beta = 1/3$ (gentler one) and 1 (steeper one).

Figure 8: The magnified figure of the magnetization at $K = 0.47$. The solid line has the slope of $1/3$. It is observed that the exponent β is already larger than $1/3$.

Figure 9: The height dependence of the interface energy is shown. The estimates of $\sigma_s(K; L; H)$ for $L = 21$ are plotted as an example.

Figure 10: The L dependence of the interface energy is shown. The estimates of $\sigma_s(K; L)$ at $K = 0.24$ are plotted as an example.

Figure 11: The critical behavior of the interface energy is shown. $K_s(K)$ is plotted versus $K - K_c$ in the logarithmic scale.

Figure 12: The values of the exponent $2 - 1$ from the consecutive two points are shown.

Figure 13: The values of $\phi_0(\beta)$ are plotted. $\beta = 4, 4.5, 5, 5.5$ show the estimates for $\beta_c = 0.6300, 0.6275, 0.6225$ and 0.6200 , respectively.

Figure 14: The behavior of the $\langle \phi^2(t) \rangle$ is shown. Every point denotes an averaged value of $\langle \phi^2(t) \rangle$ over successive 1000 MCS.

Figure 15: The extrapolated values of $\phi(n)$ to $n \rightarrow 1$ assuming the value of $\phi(1)$ are plotted.

machine	N_{job}	performance			
		U		U + M	
one processor of MONTE-4	0	1.31		0.77	
	1	1.1	1.2	0.6	0.7
	2	0.9	1.0	0.5	
	3	0.7	0.9	0.4	0.5
SX 3/11		1.05		0.63	

Table 1: The performance of the simulation code, IS2DVP, is shown. The simulation speed in the case of 1501 \times 1500 lattice is shown. The unit of the figures is GUPS, that is, giga(10^9) update per second. The figures in "U" and "U + M" denote the speed of the spin updating procedures only and of the spin updating procedure with magnetization calculation in every Monte Carlo sweep. The number N_{job} in MONTE-4 shows the number of concurrent jobs on other processors of MONTE-4. The concurrent jobs are highly vectorized with dense memory access. Therefore the speed in $N_{\text{job}} = 1$ case shows the worst case.

L	N _{sample}		updated spins	
1000	1:54	10 ⁵	7:69	10 ¹³
1500	1:18	10 ⁶	1:33	10 ¹⁵
2000	6:14	10 ⁴	1:23	10 ¹⁴
3000	1:54	10 ⁴	6:91	10 ¹³

Table 2: The simulations made for the non-equilibrium critical relaxation study are listed. N_{sample} denotes the number of independent runs. The simulation up to 500M CS is made for every sample from all-up initial condition.

year	method	z	reference
1968		1.75	[30]
1969	HTE	2.00 0.05	[31]
1971	HTE	2	[32]
1972	RG	2.0	[33]
1973	MC	1.90 0.10	[34]
1975	RG	1.82	[35]
1976	MC	2.30 0.30	[36]
1976	HTE	2.125 0.01	[37]
1976	MCRG	1.4 0.4	[38]
1978	RG	2.2	[39]
1978	RG	2.19	[40]
1978	RG	1.7	[41]
1978	RG	1.67	[42]
1979	MC+FSS	2.0 0.1	[43]
1979	RG	2.09	[44]
1979	RG	1.96	[45]
1979	RG	2.064 (energy)	[46]
1979	RG	1.819 (magnetization)	[46]
1979	MCRG	1.85 0.15	[47]
1980	TM+FSS	1.99	[48]
1980	RG	2.22	[49]
1980	RG	1.819	[50]
1980	RG	2.2	[51]
1980	RG	1.791	[52]
1981	MC	1.9	[53]
1981	MCRG	2.22 0.13	[54]
1981	RG	2.126	[55]
1982	MC+NER	2.1	[56]
1982	MC+FSS	2.0	[57]
1982	MC	2.2 0.1	[58]
1982	MCRG	2.23	[59]
1982	MCRG	2.08	[60]

1983	M C R G	2:12		[61]
1983	R G	2:2		[62]
1984	R G , H T E	2:32		[63]
1984	M C R G + N E R	2:14	0:02	[64]
1984		2		[65]
1985	R G	1:73		[66]
1985	R G	1:856		[67]
1985	M C	2:2		[68]
1985	M C R G	2:13	0:03	[69]
1986	Q 2 R	2:1	0:1	[70]
1986	M C R G	2:19	0:05	[71]
1987	M C + N E R	2:16	0:05	[72]
1987	M C + F S S	2:132	0:008	[73]
1987	M C + F S S	2:17	0:06	[74]
1988	Q 2 R	2:16		[75]
1988	M C	2:22		[75]
1988	M C + N E R	2:076	0:005	[76]
1990	H T E	2:34	0:03	[77]
1992	M C + N E R	2:2		[78]
1992	H T E	2:183	0:005	[79]
1993	H T E	2:21	0:005	[80]
1993	M C + N E R	2:21	0:03	[6]
1993	H T E	2:165	0:015	[81]
1993	M C + N E R	2:165	0:010	present study

Table 3: The estimates of z are listed. The H T E , R G , M C , F S S and N E R in this table are the abbreviations of high-temperature expansion, renormalization group, Monte Carlo, finite-size scaling and non-equilibrium relaxation, respectively.

K	L	T _{m ax.}	N _{sample}	updated spins
0:47	1500	90	1:02 10 ⁶	2:07 10 ¹⁴
0:49	2000	35	3:58 10 ⁵	5:02 10 ¹³
0:51	2000	35	3:58 10 ⁵	5:02 10 ¹³
0:53	2000	35	1:54 10 ⁵	2:15 10 ¹³

Table 4: The simulations made for the relaxation study in the ordered phase. K and T_{m ax.} show the inverse temperature and the Monte Carlo steps for each run.

L	K	H	M C S
11	0:23, 0:24, 0:25	6, 10, 16, 20, 26, 30	(1K , 10, 1K)
15	0:23, 0:24, 0:25	6, 10, 16, 20, 26, 30, 36, 40, 46	(1K , 10, 1K)
21	0:23, 0:24, 0:25	10, 16, 20, 30, 40, 50, 60	(2K , 10, 2K)
	0:23	80, 100, 150, 200, 250, 300, 350, 400	(2K , 10, 2K)
	0:225, 0:2275	50, 74, 100, 124, 150, 174, 200, 250	(10K , 10 ² , 1K)
25	0:23, 0:24, 0:25	10, 20, 30, 40, 50, 60, 70, 80	(2K , 10, 1K)
31	0:23, 0:24, 0:25	10, 20, 30, 40, 50, 60, 70, 80, 90	(2K , 20, 1K)
	0:235, 0:245	50, 60, 70, 80, 90	(2K , 10, 2K)
	0:225, 0:2275	50, 74, 100, 124, 150	(10K , 10 ² , 1K)
35	0:23, 0:24, 0:25	20, 40, 60, 80, 100, 120	(2K , 20, 1K)
	0:235, 0:245	50, 60, 70, 80, 90	(2K , 10, 2K)
41	0:23, 0:24, 0:25	20, 40, 60, 80, 100, 120	(2K , 10, 1K)
	0:235, 0:245	50, 60, 70, 80, 90	(2K , 10, 2K)
	0:225, 0:2275	100, 124, 150	(20K , 200, 1K)
45	0:235, 0:245	50, 60, 70, 80, 90	(2K , 10, 2K)
51	0:23, 0:24, 0:25	26, 50, 76, 100, 126, 150, 176, 200	(2K , 10, 2K)
	0:235, 0:245	50, 60, 70, 80, 90	(2K , 10, 2K)
	0:225, 0:2275	100, 124, 150	(20K , 200, 1K)
61	0:23, 0:24, 0:25	30, 60, 90, 120, 150, 180, 210	(2K , 10, 2K)
	0:225, 0:2275	100, 124, 150	(20K , 200, 500)
	0:235, 0:245	50, 60, 70, 80, 90	(2K , 10, 2K)
71	0:225, 0:2275	100, 124, 150	(20K , 200, 500)
81	0:23, 0:24, 0:25	40, 60, 80, 100, 120	(2K , 10, 2K)
	0:235, 0:245	50, 60, 70, 80, 90	(2K , 10, 2K)
101	0:225	100	(20K , 200, 400)
	0:225	124	(20K , 200, 300)
	0:225	150	(20K , 200, 250)
	0:23, 0:24, 0:25	50, 60, 70, 80, 90, 100	(2K , 10, 2K)
151	0:23, 0:24, 0:25	50, 60, 70, 80, 90	(2K , 10, 2K)
171	0:23	50, 70, 90	(2K , 10, 2K)

Table 5: The simulations made for the interface study are listed. The three numbers in M C S column show the M C S schedules and their meaning is explained in the text (1K = 1000). Totally 1:26 10^{15} spins are updated.

K	L	H	N _H	s
0.225	31	50{150	5	2.5592 (32)
	41	100{150	3	2.6342 (56)
	51	100{150	3	2.6741 (32)
	61	100{150	3	2.6977 (43)
	71	100{150	3	2.7036 (39)
	101	100{150	3	2.7234 (25)
	1	{	{	2.74 (1)
0.2275	21	50{250	8	2.8858 (56)
	31	50{150	5	2.9938 (40)
	41	100{150	3	3.0365 (51)
	51	100{150	3	3.0578 (37)
	61	100{150	3	3.0725 (49)
	71	100{150	3	3.0832 (32)
	1	{	{	3.10 (1)
0.23	11	16{30	4	2.7649 (89)
	15	16{46	7	3.0443 (30)
	21	30{400	12	3.1726 (20)
	25	30{80	6	3.2200 (28)
	31	30{90	7	3.2608 (17)
	35	40{120	5	3.2790 (35)
	41	40{120	5	3.2904 (22)
	51	76{200	6	3.3034 (13)
	61	60{150	4	3.3118 (14)
	81	40{120	5	3.32151 (95)
	101	50{100	6	3.32622 (58)
	119	50{80	4	3.32875 (66)
	151	50{90	5	3.33083 (40)
	171	50{90	3	3.33128 (57)
	1	{	{	3.335 (3)

K	L	H	N _H	s
0.235	31	50{90	5	3.5859 (17)
	35	50{90	5	3.5939 (11)
	41	50{90	5	3.6029 (18)
	45	50{90	5	3.6073 (13)
	51	50{90	5	3.6132 (14)
	61	50{90	5	3.62088 (83)
	81	50{90	5	3.62346 (67)
	1	{	{	3.630 (5)
0.24	11	16{30	4	3.5778 (22)
	15	16{46	7	3.6852 (34)
	21	20{400	13	3.7447 (14)
	25	30{80	6	3.7616 (26)
	31	30{90	7	3.7829 (19)
	35	20{120	6	3.7875 (15)
	41	20{120	6	3.7943 (13)
	51	26{200	8	3.79888 (24)
	61	30{210	7	3.80299 (95)
	81	40{120	5	3.80849 (81)
	101	50{100	6	3.81055 (57)
	151	50{90	5	3.81181 (32)
	1	{	{	3.813 (2)

K	L	H	N _H	γ_s
0.245	31	50{90	5	3.9101 (16)
	35	50{90	5	3.9141 (13)
	41	50{90	5	3.9181 (10)
	45	50{90	5	3.9202 (16)
	51	50{90	5	3.92080 (67)
	61	50{90	5	3.92691 (61)
	81	50{90	5	3.92687 (63)
	1	{	{	3.930 (2)
0.25	11	16{30	4	3.8552 (26)
	15	16{46	7	3.9297 (21)
	21	20{400	13	3.96912 (78)
	25	20{80	7	3.9839 (16)
	31	30{90	7	3.9906 (15)
	35	20{120	6	3.9929 (12)
	41	20{120	6	4.0028 (11)
	51	26{200	8	4.00478 (52)
	61	30{210	7	4.00704 (83)
	81	40{120	5	4.00900 (50)
	101	50{100	6	4.01017 (32)
	151	50{90	5	4.01138 (20)
	1	{	{	4.012 (1)

Table 6: The estimated values of γ_s for some values of K and L are shown. The column n H shows the height region over which the average is calculated. N_H is the number of different H used in the averaging operation. The values of γ_s in L = 1 denotes the extrapolated values of $\gamma_s(K;L)$ to L = 1 and they are the estimates for the interface energy in the thermodynamics limit.

year	α_0		reference
1982	1.05	0.05	[85]
1985	1.2	0.1	[86]
1988	1.58	0.05	[87]
1993	1.5	0.1	[13]
1993	1.25	1.68	[14]
1993	1.52	0.05	[16]
1993	1.221	1.492	[17]
1993	1.92	0.15	[18]
1993	1.42	0.04	present study

Table 7: The estimates of α_0 are shown with the year of publication.

n	$\langle n \rangle; 256^3$	$\langle n \rangle; 1024^3$
2	2.4625 (1)	2.4624 (2)
3	2.4738 (5)	2.4735 (5)
4	2.481 (1)	2.480 (2)
5	2.485 (3)	2.485 (5)

Table 8: The estimates of $\langle n \rangle$ are shown. The estimates from 256^3 and 1024^3 lattice coincide within the present accuracy of the estimation.