

# Surface magnetization of aperiodic Ising quantum chains

L Turban and B Berche

Laboratoire de Physique du Solide, URA CNRS n°155, Université de Nancy I,  
BP 239, F-54506 Vandœuvre les Nancy Cedex, France

**Abstract.** We study the surface magnetization of aperiodic Ising quantum chains. Using fermion techniques, exact results are obtained in the critical region for quasiperiodic sequences generated through an irrational number as well as for the automatic binary Thue-Morse sequence and its generalizations modulo  $p$ . The surface magnetization exponent keeps its Ising value,  $\nu_s = 1/2$ , for all the sequences studied. The critical amplitude of the surface magnetization depends on the strength of the modulation and also on the starting point of the chain along the aperiodic sequence.

## 1. Introduction

Since the discovery of quasicrystals [1] there has been a growing interest in the structural and physical properties of quasiperiodic systems (see e.g. [2-5]). Due to their situation, which is intermediate between periodic and random ones, these systems are particularly interesting from the point of view of their critical behaviour.

On the Penrose lattice, an approximate renormalization group study of the ferromagnetic Ising model [6] suggested a logarithmic singularity in the specific heat, i.e. ordinary two-dimensional Ising behaviour. In the same way, random and self-avoiding walks, studied via Monte-Carlo techniques, were shown to keep their usual asymptotic properties [7].

Exact results have been obtained, either for aperiodic quantum spin chains or for the corresponding layered classical counterparts ([8-12] and references therein). A condition for quasiperiodicity was established [13] in the case of aperiodic chains generated through an inflation rule. The one-dimensional lattice is quasiperiodic, i.e. it displays a Fourier spectrum with Dirac delta peaks for a set of  $Q$ -values which are not integral multiples of any single wave-vector, when the characteristic polynomial of the substitution has only one root with an absolute value greater than one.

Mainly bulk critical properties of aperiodic Ising systems have been worked out in the last years. Tracy [9] found that, although for quasiperiodically layered Ising systems the specific heat displays a logarithmic singularity, this is no longer true for some aperiodic sequences which do not satisfy the criterion for quasiperiodicity. The low energy finite-size behaviour of the spectrum of quasiperiodic Ising quantum chains was found to be in agreement with the prediction of conformal theory by Igloi [10]. The bulk critical exponents deduced from the gaps are those of the periodic system. The structure of the finite-size Ising spectrum has been carefully studied for the Fibonacci chain in [12]. Besides the conformally invariant scaling of the levels at low energy, a multifractal scaling is observed for higher ones.

In the present work we examine the surface critical behaviour of aperiodic Ising quantum chains. The surface magnetization is calculated exactly in the critical region making use of fermion techniques [14,15]. The outline of the paper is as follows. In section 2 we recall how the surface magnetization of the inhomogeneous Ising quantum chain can be obtained. The case of a quasiperiodic chain generated by an irrational number is considered in section 3. This is followed by a study of the aperiodic binary Thue-Morse sequence in section 4 and of its generalizations modulo  $p$  in section 5. The results are discussed in section 6.

## 2. Surface magnetization of the inhomogeneous Ising quantum chain

Let us consider a semi-infinite inhomogeneous Ising quantum chain with modulated interaction strength  $J_j$  and a constant transverse field  $h = 1$ . The Hamiltonian can be written as

$$H = \frac{1}{2} \sum_{j=1}^X \left( \sigma_j^z + J_j \sigma_j^x \sigma_{j+1}^x \right) \quad (2.1)$$

where the  $\sigma_j^{\alpha}$  are Pauli spin operators. This quantum chain also corresponds, in the extreme anisotropic limit, to a semi-infinite layered two-dimensional classical Ising system with modulated interactions on the bonds perpendicular to the surface.

Using the Jordan-Wigner transformation [16] the Hamiltonian is rewritten as a quadratic form in fermion operators  $c_j^+, c_j$ , which may be diagonalized through the canonical transformation [14,17]

$$c_j = \frac{1}{2} \sum_{\alpha} \left[ \epsilon_{\alpha}(j) + \epsilon_{\alpha}(j) \right] + \left[ \epsilon_{\alpha}(j) - \epsilon_{\alpha}(j) \right]^{\dagger} \quad (2.2)$$

where  $\epsilon_{\alpha}$  and  $\epsilon_{\alpha}^{\dagger}$  are new fermion operators.  $\epsilon_{\alpha}$  and  $\epsilon_{\alpha}^{\dagger}$  are the real normalized eigenvectors of two matrices leading to the excitation energies  $\epsilon_{\alpha}$  of the diagonal Hamiltonian  $H = \sum_{\alpha} (\epsilon_{\alpha}^{\dagger} \epsilon_{\alpha} - 1)$ . They are related through

$$(A + B)_{ij} = (A - B)_{ij} = \delta_{i, j+1} \quad (2.3)$$

Below the critical point, in the ordered phase, the ground-state is two-fold degenerate and the lowest excitation  $\epsilon_0$ , which corresponds to an eigenvector localized near to the surface, vanishes exponentially with the size of the system. This provides the condition for criticality [18]

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{j=1}^N (\epsilon_0)_c = 1 \quad (2.4)$$

The surface magnetization can be obtained using a simple method due to Peschel [15]. The asymptotic form of the surface two-point correlation function gives the square of the surface magnetization  $m_s$  which can be expressed as the matrix element  $\langle 0 | \sigma_1^x | j \rangle$  where  $|0\rangle$  is the vacuum of  $H$  and  $|j\rangle = \sum_{\alpha} \epsilon_{\alpha}^{\dagger} |0\rangle$  its first excited state. Rewriting  $\sigma_1^x$  in terms of diagonal fermions through (2.2), one obtains  $m_s = \langle 0 | \epsilon_0 \rangle$ . Using (2.3) with

= 0 in the ordered phase where the lowest excitation vanishes, a recursion equation for the successive components of  $\psi_0$  is obtained leading to

$$\psi_0(j) = (\lambda)^{j-1} \psi_0(1) \prod_{i=1}^{j-1} \lambda_i^{-1} \quad j > 1 \quad (2.5)$$

The surface magnetization then follows from the normalization of the eigenvector and reads

$$m_s = \frac{\sum_{i=1}^{\infty} \lambda^i \prod_{j=1}^{i-1} \lambda_j^{-1}}{\sum_{i=1}^{\infty} \lambda^i \prod_{j=1}^{i-1} \lambda_j^{-1}} \quad (2.6)$$

### 3. Quasiperiodic modulation generated by an irrational number

Let the position  $x_j$  of the  $j$ -th spin along the chain be defined as [10, 13]

$$x_j = jb + \frac{j+1}{!} (a-b) \quad (a > b) \quad (3.1)$$

where  $[x]$  is the integer part of  $x$ . When  $!$  is irrational, one obtains a succession of short and long bonds which are distributed according to a quasiperiodic sequence. With a modulation amplitude  $r$  such that  $a$ -bonds have a strength  $r$  and  $b$ -bonds a strength  $1$ , the number of modified bonds from the surface to site  $j$  is given by

$$n_j = j - \frac{j+1}{!} \quad (3.2)$$

corresponding to an asymptotic density  $\rho = 1 - !$ . The critical coupling  $c_c$  follows from (2.4) with

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{j=1}^N (\lambda_j)_c = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{j=1}^N r^{N-n_j} = 1 \quad (3.3)$$

so that

$$c_c = r^{1-!} \quad (3.4)$$

The surface magnetization (2.6) involves the sum of products

$$\sum_{i=1}^{\infty} \lambda^i \prod_{j=1}^{i-1} \lambda_j^{-1} = \sum_{j=1}^{\infty} r^{2n_j} = r^{2!} \sum_{j=1}^{\infty} \frac{c_c^{2j}}{r^{2f_{j+1}g}} \quad (3.5)$$

In the following we take  $r > 0$  without loss of generality.




Figure 1. Surface magnetization  $m_s$  of the Fibonacci Ising quantum chain with a modulation amplitude  $r = 1; 2; 3; 5$  from top to bottom.

In the last term  $n_j$ , given by (3.2), was rewritten as

$$n_j = j \frac{j+1}{!} + \frac{j+1}{!} \quad (3.6)$$

where  $f(x)$  is the fractional part of  $x$ . The eigenvector  $\psi_0$ , which is localized near to the surface in the ordered phase, extends into the bulk in the critical regime ( $! = !_c$ ). Then the last factor in (3.5) may be replaced by its averaged value [11]. Since  $!$  is irrational,  $f(j+1) = !$  is uniformly distributed over  $[0; 1]$  and the average is given by

$$\int_0^1 r^{-2f(j+1)} dx = \int_0^1 r^{-2x} dx = \frac{r^2 - 1}{r^2 \ln r^2} \quad (3.7)$$

Summing the geometrical series, one obtains the exact leading contribution in the scaling region

$$m_s^{(1)} = \frac{1}{c} \frac{\ln r^2}{r^2 - 1} \quad (3.8)$$

The surface exponent  $\nu_s = 1=2$  is the same as for an homogeneous Ising system and the critical amplitude keeps the same form for any sequence generated by an irrational. It only depends on the modulation amplitude and the critical coupling.

When  $!$  is the golden mean  $(\sqrt{5} + 1)/2$ , the modulation is distributed according to the Fibonacci sequence. Such a sequence may be obtained in various ways. It can be represented by an infinite word  $W_1 = 01001010 \dots$  where 0 stands for an a-bond and 1 for a b-bond. This word is the limit of an infinite sequence of finite words  $W_n$  obtained through the recursion relation

$$W_{n+1} = W_n W_{n-1} \quad W_0 = 1 \quad W_1 = 0 \quad (3.9)$$




Figure 2. Ratio of the Fibonacci surface magnetization  $m_s$  to its leading critical term  $m_s^{(1)}$  for  $r = 2; 3; 4; 5$  from bottom to top. The linear behaviour corresponds to a correction to scaling of order  $3=2$ .

to which is associated the characteristic polynomial  $x^2 - x - 1$  [9]. The length of successive words is given by Fibonacci numbers.

The Fibonacci sequence also follows from the inflation rule [9]

$$0 \rightarrow 01 \quad 1 \rightarrow 10 \tag{3.10}$$

giving

$$\begin{aligned} &0 \\ &01 \\ &010 \\ &01001 \\ &01001010 \\ &\vdots \end{aligned} \tag{3.11}$$

The surface magnetization of the Fibonacci quasilattice as given by (2.6) is plotted in figure 1 for different values of the modulation amplitude  $r$ . The ratio  $m_s/m_s^{(1)}$  shown in figure 2 as a function of  $1 - \lambda_c/\lambda$  indicates that the corrections to scaling are of order  $3=2$  in this case.

#### 4. The binary Thue-Morse sequence

Let us now consider the binary Thue-Morse sequence  $f_j^{(2)} = 0$  or  $1$  ( $j = 1; 2; 3; \dots$ ) which is not quasiperiodic [19] and sometimes called automatic since it can be generated




Figure 3. Surface magnetization of the binary Thue-Morse aperiodic chain with a modulation amplitude  $r = 1; 2; 3; 5$  from top to bottom.

through a 2-automaton. It may be deduced from the binary representation of non-negative integers [20]

$$0 \ 1 \ 10 \ 11 \ 100 \ 101 \ 110 \ 111 \ \dots \quad (4.1)$$

by counting the sum of their digits modulo 2

$$01101001 \ \dots \quad (4.2)$$

or through the recursion

$$W_{n+1} = W_n \ W_{n-1} \ W_n \ W_{n-1} \quad (4.3)$$

It also results from the substitution

$$0! \ 01 \quad 1! \ 10 \quad (4.4)$$

leading to

$$\begin{aligned} &0 \\ &01 \\ &0110 \\ &01101001 \\ &\underline{0110100110010110} \\ &\dots \end{aligned} \quad (4.5)$$

The aperiodic chain is constructed through the correspondences  $0! \rightarrow r$  and  $1! \rightarrow r$ . It is clear from (4.4) that the asymptotic density of modified bonds is  $\rho_1 = 1/2$  and, according to (3.3),

$$\lambda_c = r^{1/2} \quad (4.6)$$




Figure 4. Ratio of the binary Thue-Morse surface magnetization  $m_s$  to its leading critical contribution  $m_s^{(1)}$  for a modulation amplitude  $r = 2;3;4;5$  from top to bottom. The corrections to scaling, due to the higher regularity of the sequence, are much weaker than for the Fibonacci sequence.

This value of the density is also reached for any finite sequence with an even length so that the number of modified bonds satisfies

$$n_{2k} = k \tag{4.7}$$

The substitution (4.4) is such that the binary Thue-Morse sequence is reproduced by selecting the sequence of its odd elements ( $f_{2k+1}^{(2)} = f_{k+1}^{(2)}$ ) as indicated in the last line of (4.5). As a consequence

$$n_{2k+1} = k + f_{k+1}^{(2)} \quad k = 0;1;2 \dots \tag{4.8}$$

The sum involved in the surface magnetization (2.6) can then be written as follows

$$\sum_{j=1}^D \sum_{i=1}^E \lambda^i = \sum_{k=1}^D \frac{\lambda^{4k}}{c} + \sum_{k=0}^D \frac{\lambda^{2(2k+1)}}{c} r^{1-2f_{k+1}^{(2)}} \tag{4.9}$$

Near the critical point the last term may be replaced, as above, by its average over the infinite sequence in which 0 and 1 occur with the same probability. Then

$$r^{1-2f_{k+1}^{(2)}} = \frac{1}{2} (r + r^{-1}) \tag{4.10}$$

and the exact leading critical behaviour reads

$$m_s^{(1)} = \frac{2^D r^D}{r+1} \left( 1 - \frac{c}{2} \right)^{\#_{1=2}} \tag{4.11}$$

The surface magnetization obtained numerically using (2.6) is plotted in figure 3 for different values of  $r$  and the ratio  $m_s = m_s^{(1)}$  is shown in figure 4. As a result of the higher regularity of the Thue-Morse sequence, the corrections to scaling are much weaker than for the Fibonacci sequence.

5. Thue-Morse sequences modulo  $p$ 

The results of the last section can be generalized by considering Thue-Morse sequences  $f_j^{(p)}$  defined modulo  $p$ . These sequences are generated by taking the sum modulo  $p$  of the digits of non-negative integers written in base  $p$ . With  $p=3$ , for example, one obtains

$$0 \ 1 \ 2 \ 10 \ 11 \ 12 \ 20 \ 21 \ 22 \ 100 \ 101 \ 102 \ \dots \quad (5.1)$$

and the sequence  $f_j^{(3)}$  reads

$$012120201120 \ \dots \quad (5.2)$$

$f_j^{(p)}$  also follows from the substitution

$$0 \rightarrow 012 \ \dots; \ 1 \rightarrow 1! \ 12 \ \dots; \ 2 \rightarrow 2! \ 20 \ \dots; \ \dots; \ p-1 \rightarrow (p-1)! \ (p-1)0 \ \dots; \ p \rightarrow p^2 \quad (5.3)$$

which for  $p=3$  leads to

$$\begin{aligned} &0 \\ &012 \\ &012120201 \\ &\underline{012120201120201012201012120} \\ &\dots \end{aligned} \quad (5.4)$$

One associates a modified coupling on the chain with any nonzero digit along the sequence, i.e. we use the correspondence

$$0 \rightarrow 1; 2; \dots; \ p-1 \rightarrow r \quad (5.5)$$

Then, according to (5.3), the asymptotic density of modified bonds is  $\rho_1 = (p-1)/p$  and the critical coupling  $\beta_c = r^{-(p-1)/p}$ . The asymptotic density is also obtained when the number of bonds  $n_j$  is any multiple  $pk$  of  $p$  so that

$$n_{pk} = pk \rho_1 = k(p-1) \quad (5.6)$$

One may check on the last line of (5.4) that  $f_{pk+1}^{(p)} = f_{k+1}^{(p)}$  whereas  $f_{pk+2}^{(p)}; f_{pk+3}^{(p)} \dots$  are deduced from  $f_{k+1}^{(p)}$  through circular permutations on  $0; 1; 2; \dots; \ p-1$ . Then, the number of modified bonds after  $pk+q$  steps can be written as

$$n_{pk+q} = k(p-1) + m_{k+1}(q) \quad 0 \leq q < p-1 \quad (5.7)$$

where

$$m_k(q) = \sum_{i=0}^{q-1} g_i f_k^{(p)} + i \pmod{p} \quad 0 < q < p-1; \quad m_k(0) = 0 \quad (5.8)$$




Figure 5. Surface magnetization of binary Thue-Morse aperiodic chains with  $r = 2$ , starting on the  $i$ -th digit along the sequence with  $i = 1; 2; 3; 4$ . The amplitude of the leading term given by (4.11) when  $i$  is odd, is the geometric mean of the amplitudes obtained for even values of  $i$ . In the strong coupling regime the local environment governs the behaviour of the surface magnetization.

and

$$g(i) = 1 \quad i; 0 \tag{5.9}$$

The sum in (2.6) may be splitted into  $p$  parts and rewritten as

$$\sum_{j=1}^i Y^j = \sum_{k=1}^i \frac{r^{2pk}}{r^{2q(p-1)+p}} + \sum_{q=1}^i \frac{r^{2q}}{r^{2q(p-1)+p}} \tag{5.10}$$

When  $r^{2m_{k+1}(q)}$  in the last sum may be replaced by its averaged value. On the infinite sequence,  $m_k(q)$  is equal to  $q-1$  with probability  $q-p$  and equal to  $q$  with probability  $(p-q)=p$ , so that

$$\frac{D}{r^{2m_{k+1}(q)}} = \frac{qr^{2(q-1)} + (p-q)r^{2q}}{p} \tag{5.11}$$

In the critical region, one finally obtains

$$m_s^{(1)} = \frac{p}{p} \frac{r^{1-p} - r^{-1-p}}{r - r^{-1}} \frac{1}{1 - \frac{c}{r^{2p}}} \tag{5.12}$$

for the leading contribution to the surface magnetization in agreement with (4.11) when  $p=2$ .




Figure 6. Ratio of the surface magnetization  $m_s$  to its leading term  $m_s^{(1)}$  for the binary Thue-Morse Ising chain with  $r = 2$  and different starting points  $i$  along the sequence. Although the leading amplitude is the same for all chains with odd values of  $i$ , the corrections to scaling are different.

#### 6. Discussion

In the cases studied so far the modulation was distributed on the spin chain according to a given sequence, starting with the first digit of the sequence. One expects a change in the surface magnetization amplitude for chains starting elsewhere along the sequence. The previous calculations are easily extended to the case where one begins the chains on the  $i$ -th digit.

For the Fibonacci sequence,  $n_j$  in (3.6) is changed into

$$n_j = j - 1 \frac{1}{j!} + \frac{j+1}{j!} \frac{1}{j!} \quad (6.1)$$

leading to

$$m_s^{(1)}(i) = r^{1 - f_{i-1}^{(1)}} \frac{\ln r^2}{r^2 - 1} \prod_{l=2}^i \frac{1}{2^{f_{l-1}^{(1)}}} \quad i > 1 \quad (6.2)$$

With the binary Thue-Morse sequence starting on  $i$  even, i.e. with  $f_{21}^{(2)}$ , one obtains

$$n_{2k} = k - 1 + f_{k+1}^{(2)} + f_{21}^{(2)} \quad n_{2k+1} = k + f_{21}^{(2)} \quad i = 2l \quad (6.3)$$

and then, on the surface site,

$$m_s^{(1)}(2l) = \frac{2r^{f_{21}^{(2)}}}{r+1} \prod_{l=2}^{2l} \frac{1}{2^{f_{l-1}^{(2)}}} \quad (6.4)$$

Otherwise, when  $i$  is odd,

$$n_{2k} = k \quad n_{2k+1} = k + f_{k+1}^{(2)} \quad i = 2l+1 \quad (6.5)$$

like in (4.7-8). Then the leading term is still given by (4.11) which is just the geometrical mean of the two allowed values in (6.4). The dependence of the surface magnetization on  $i$  is shown in figure 5. The local environment clearly governs the behaviour when  $c = 1/2$ . The surface magnetization is reduced when the surface site is linked to the bulk via a weak bond for  $i = 1$  and 4. Near the critical point, on the contrary, only the long-range structure of the modulation and the parity of the surface site are important.

Although the amplitude of the leading contribution to the surface magnetization is the same for all the aperiodic chains starting with odd values of  $i$  along the Thue-Morse sequence the corrections to scaling differ as shown on figure 6.

Similar results would be obtained with the Thue-Morse sequence defined modulo  $p$  with the same leading amplitude for chains starting on  $i = pl+1$  along the sequence.

For all the sequences considered here, the aperiodic modulation, either quasiperiodic or automatic, leaves the surface magnetic exponent unaffected,  $\nu_s = 1/2$ , like in the homogeneous semi-infinite Ising model. On the other hand, the critical amplitude depends on the modulation and it would be of interest to study other surface quantities in order to check whether surface critical amplitude ratios remain universal. The study of aperiodic sequences which do not lead to ordinary bulk Ising behaviour [9] would also be worthwhile.

## References

- [1] Shechtman, D., Blech, I., Gratias, D., Cahn, J.W. Phys. Rev. Lett. 53 (1984) 1951
- [2] Henley, C.J. Comments Cond. Mat. Phys. 13 (1987) 59
- [3] Janssen, T. Phys. Reports 168 (1988) 55
- [4] Guyot, P., Kramer, P., de Boissieu, M. Rep. Prog. Phys. 54 (1991) 1373
- [5] Janot, C., Dubois, J.M., de Boissieu, M. Am. J. Phys. 57 (1989) 972
- [6] Godreche, C., Luck, J.M., Orland, H. J. Stat. Phys. 45 (1986) 777
- [7] Langie, G., Igloi, F. J. Phys. A 25 (1992) L487
- [8] Luck, J.M., Nieuwenhuizen, Th.M. Europhys. Lett. 2 (1986) 257
- [9] Tracy, C.A. J. Phys. A 21 (1988) L603; J. Stat. Phys. 51 (1988) 481
- [10] Igloi, F. J. Phys. A 21 (1988) L911
- [11] Ceccatto, H.A. Phys. Rev. Lett. 62 (1988) 203; Z. Phys. B 75 (1989) 253
- [12] Henkel, M., Patkós, A. J. Phys. A 25 (1992) 5223
- [13] Bombieri, E., Taylor, J.E. J. Physique Colloq. C3 47 (1986) 19; Contemp. Math. 64 (1987) 241
- [14] Lieb, E.H., Schultz, T.D., Mattis, D.C. Ann. Phys. (N.Y.) 16 (1961) 406
- [15] Peschel, I. Phys. Rev. B 30 (1984) 6783
- [16] Jordan, P., Wigner, E.Z. Phys. 47 (1928) 631
- [17] Feuty, P. Ann. Phys. (N.Y.) 57 (1970) 79
- [18] Feuty, P. Phys. Lett. 72A (1979) 245
- [19] Axel, F., Allouche, J.P., Klemm, M., Mendès-France, M., Peyrière, J. J. Physique Colloq. C3 47 (1986) 181
- [20] Dekking, M., Mendès-France, M., van der Poorten, A. Math. Intelligencer 4 (1983) 130