

Hole-hole contact interaction in the t - J model

A. L. Chernyshev

Institute of Semiconductor Physics, 630090 Novosibirsk, Russia

A. V. Dotsenko and O. P. Sushkov

School of Physics, The University of New South Wales

P. O. Box 1, Kensington, NSW 2033, Australia

and Budker Institute of Nuclear Physics, 630090 Novosibirsk, Russia

Abstract

Using an analytical variational approach we calculate the hole-hole contact interaction on the Neel background. Solution of the Bethe-Salpeter equation with this interaction gives bound states in d - and p -waves with binding energies close to those obtained by numerical methods. At $t=J/2-3$ the bound state disappears. In conclusion we discuss the relation between short range and long range interactions and analogy with the problem of pion condensation in nuclear matter.

1 Introduction

Due to connection with high- T_c superconductivity, the problem of mobile holes in the t - J model has received considerable development during recent years. This model with less than half-filling is defined by the Hamiltonian

$$H = H_t + H_J = t \sum_{\langle nm \rangle} (c_n^\dagger c_m + H.c.) + J \sum_{\langle nm \rangle} S_n^z S_m^z + \frac{1}{2} (S_n^+ S_m + S_n^- S_m^+); \quad (1)$$

where c_n^\dagger is the creation operator of a hole with spin ($= \uparrow, \downarrow$) at site n of a two-dimensional square lattice. The operator c_n^\dagger acts in the Hilbert space where there is no double electron occupancy. The spin operator is $S_n = \frac{1}{2} c_n^\dagger \sigma c_n$. $\langle nm \rangle$ are the neighbor sites on the lattice. The Hamiltonian (1) with $t = 1$ corresponds to the t - J model, and for $t = 0$ it is the t - J_z model. Below we set $J = 1$.

At half-filling (one hole per site) the t - J model is equivalent to the Heisenberg antiferromagnet model [1, 2] which has long range Neel order in the ground state [3, 4, 5]. The problem is the behavior of a system under doping by additional holes. Strong interaction of mobile holes with localized spins makes this problem very complicated.

Intense studies have led to quite firm establishing of one particle properties in the t - J model. A single hole is an object with a complex structure due to virtual admixture of spin excitations. It has been shown both analytically [6]–[13] and numerically [14]–[18] that one hole has a ground state with a momentum of $k = (\pi/2, \pi/2)$. The energy is almost degenerate along the line $\cos k_x + \cos k_y = 0$ which is the edge of the magnetic Brillouin zone. However, even the two hole problem remains controversial because there are many sophisticated effects in such systems.

The hole can interact with spin waves. For the Neel background, the spin wave spectrum is gapless according to the Goldstone theorem. The effective coupling constant for the interaction of a hole with long wavelength spin excitations for $t=J=5$ was calculated in our work [19]. Earlier it had been done in Ref. [20] in the perturbation theory limit ($t=J=1$) and for $zS=1$ (z is the number of neighbor sites).

If interaction of the Goldstone excitation with particles exceeds a certain critical value, the system becomes unstable. This was understood a long time ago for electron-phonon interaction (lattice instability) [21]. Just because of this fact and because the hole-magnon coupling constant is large enough, any small but finite doping destroys the long range antiferromagnetic order in the t - J model. It was shown for $t=J$ in Ref. [22]. For the Hubbard model, the instability has been proven in the Hartree-Fock approximation [23]. For the t - J model, the instability of long range antiferromagnetic order for $t=J=5$ was demonstrated in our work [24]. In the same paper, we pointed out the direct analogy of this instability with pion condensation in nuclear matter (for review see Ref. [25]).

In the present work we calculate contact interaction between two holes on the Neel background. "Contact" means that it is due to exchange by spin excitations with momentum $q=0$. There is no retardation in contact interaction. We consider this interaction for both parallel and opposite directions of the holes spins. We calculate the vertex function which provides the most general description of interaction. For the most interesting case of opposite spins, there is an attraction at $t=J=2-3$ which gives a very shallow bound state. Our result for binding energy agrees with that of Ref. [26] obtained by a numerical variational method in a restricted Hilbert space. It agrees also with recent result of Monte Carlo simulation [27]. However, there is a disagreement with results of exact diagonalizations on finite-size clusters [16, 28]–[32]. Following Refs. [16, 26, 33], we suspect that the discrepancy is caused by finite-size effects in exact diagonalizations. Recent calculation [34] of the binding energy on larger clusters (up to 26 sites) and the scaling with system size indicated a critical value of $J=t$ between 0.3 and 0.5, in agreement with our number.

We base our study on results obtained in [13]. The suggested trial wave function of a single hole was of the form $\psi_k = h_k^\dagger \phi$, where ϕ is the background and h_k^\dagger is the creation operator of a dressed hole

$$\begin{aligned} h_{k\uparrow}^\dagger &= \frac{1}{\sqrt{2N}} \sum_n (1 - \delta_{n,0}) e^{-ik \cdot n} c_{n\uparrow}^\dagger + S_{n\uparrow}^+ e^{-ik \cdot n} c_{n\downarrow}^\dagger; \\ h_{k\downarrow}^\dagger &= \frac{1}{\sqrt{2N}} \sum_n (1 + \delta_{n,0}) e^{-ik \cdot n} c_{n\downarrow}^\dagger + S_{n\downarrow}^+ e^{-ik \cdot n} c_{n\uparrow}^\dagger; \end{aligned} \quad (2)$$

Here $\delta_{n,0} = +1$ for the spin-up sublattice and $\delta_{n,0} = -1$ for the spin-down sublattice, \mathbf{e} is a unit vector corresponding to one step in the lattice, N is the total number of sites. The hole energy corresponding to ansatz (2) was found in Ref. [13] by variational method

$$\epsilon_k = \epsilon_0 + \frac{0}{2} S_k; \quad S_k = \frac{0}{4} + 4t^2 (1 + y) - 4t^2 (x + y) \frac{1}{k^2}; \quad (3)$$

where

$$\epsilon_k = \frac{1}{2}(\cos k_x + \cos k_y); \quad (4)$$

Expressions for the parameters ϵ_0 , ϵ_1 , x , and y in terms of ground state correlators are presented in Ref.[13]. With the normalization of one hole per the lattice of N sites the coefficients in Eq. (2) are as follows

$$\epsilon_k = \frac{1}{2} \frac{\epsilon_0 + 2S_k}{X S_k}^{1=2}; \quad \epsilon_k; = \frac{t}{[Y S_k (\epsilon_0 + 2S_k)]^{1=2}} [(1+v) - (u+v) e^{ik}]; \quad (5)$$

The parameters X , Y , u , and v are also expressed in terms of ground state correlators [13]. In the present work we consider two backgrounds: 1) Ising background which is the ground state of the t - J_z Hamiltonian and has no quantum fluctuations ($S_n^z = \frac{1}{2}$). 2) Neel background which is the ground state of the Heisenberg model [$S = 1$ in the Hamiltonian (1)]. For these states the numerical values of parameters in Eqs. (3) and (5) are [13]

$$\begin{aligned} I: \quad & \epsilon_0 = 1.0; \quad \epsilon_1 = 1.50; \quad x = 0.00; \quad y = 0.00; \quad X = 1.0; \quad Y = 1.00; \quad u = 0.00; \quad v = 0.00; \\ N: \quad & \epsilon_0 = 1.2; \quad \epsilon_1 = 1.33; \quad x = 0.56; \quad y = 0.14; \quad X = 0.8; \quad Y = 0.72; \quad u = 0.42; \quad v = 0.12; \end{aligned} \quad (6)$$

Ansatz (2) includes states with a simple hole and states with one hopping to a nearest neighbor site ($S_n^+ d_{n+}^Y$ can be represented as $d_{n+}^Y d_n d_{n+}^Y$). Using the string picture suggested in Ref.[35], one can say that ansatz (2) includes only strings of the minimal length $L = 1$. It is quite natural for $t = 1$, but it is a rather rough assumption for the Ising background at $t = 1$. However, for the Neel background this ansatz is justified by comparison with numerical simulations. It turns out that it provides good results up to $t = 5$ (see discussion in Refs. [13, 18]). It means that the hole motion on the Neel background is mostly determined by quantum fluctuations or, in the string language, that fluctuations cut strings. As far as we understand, a similar conclusion was made in Ref.[23]. Thus we believe that the wave function we use is valid for $t = 5$.

The plan of the paper is as follows. In Sec. II we study the interaction of holes with parallel spins. In Sec. III we consider the case of the holes with opposite spins in the Ising limit and the bound state of two holes. An analysis of the Neel case is performed in Sec. IV. Sec. V presents our conclusions.

2 Interaction of the holes with parallel spins

We consider a state with two holes

$$|j;2i = h_{k_1}^Y h_{k_2}^Y |Di; \quad (7)$$

Using Eq. (2), one can easily check that the effective creation operators h_k^Y obey the usual anticommutation relations:

$$f h_{k_1}^Y; h_{k_2}^Y g = 0; \quad (8)$$

It means that in the present description the dressed holes are fermions and the two hole state (7) is well defined.

In the naive perturbation theory, the vertex corresponding to scattering $1;2 \rightarrow 3;4$ is just the matrix element $\langle h;4 | h | j;2i$. However in our case, the holes are composite objects and we must take into account that the overlapping $\langle h;4 | j;2i$ is not equal to zero. In this case the vertex is equal to

$$\langle 3;4;1;2 \rangle = \langle h;4 | h | j;2i - (\epsilon_1 + \epsilon_2) \langle h;4 | j;2i; \quad (9)$$

where ϵ_i is the single particle energy (3). Due to the energy conservation in scattering, $\epsilon_1 + \epsilon_2 = \epsilon_3 + \epsilon_4$. This procedure with subtraction is well known in the many-body perturbation theory (see, e.g., Ref. [36]). The physical meaning of Eq. (9) is especially evident at $\langle 3;4i = |j;2i$ when the interaction is just the difference between the total energy and the sum of single-particle energies.

We consider now the Ising background (the t - J_z model). Within the accepted approximation, there is no dispersion for this background [see Eqs.(3) and (6)]. Therefore, we may introduce an effective creation operator h_n^Y of a hole localized at site n . Equations (2-5) reduce to

$$\begin{aligned} h_k^Y &= \frac{2}{N} \sum_n h_n^Y \exp(ik \cdot n); \\ h_n^Y &= \frac{1}{2} (1 - \epsilon_n) d_{n+}^Y + S_n^+ d_{n+}^Y; \end{aligned} \quad (10)$$

$$\begin{aligned}
&= \frac{7}{4} S; \quad S = \frac{P}{9=16+4t^2}; \\
&= \frac{1}{2} \frac{3=2+2S}{S} \stackrel{1=2}{}, \quad = \frac{t}{[S(3=2+2S)]^{1=2}}:
\end{aligned}$$

In the coordinate space the vertex has the form

$$\langle m^0; n^0; m; n \rangle = \langle m^0; n^0 | H_{Jn} | m; n \rangle - 2 \langle m^0; n^0 | J_n | m; n \rangle; \quad (11)$$

where $|m; n\rangle = h_m^\dagger h_n^\dagger |0\rangle$ ($m \in n$ and m and n are associated with the same sublattice). One can easily check that matrix elements $\langle m^0; n^0 | H_{Jn} | m; n \rangle$ and $\langle m^0; n^0 | J_n | m; n \rangle$ are not equal to zero only for $m^0 = m$ and $n^0 = n$. (There is also a possibility of $m^0 = n$ and $n^0 = m$ which, due to anticommutation relation (8), gives the usual fermionic antisymmetrization.) Obviously, for distant sites m and n the second term in (11) exactly cancels out the first term. The interaction is not zero only for the closest possible m and n : $\langle m; n+2x; n; n+2x \rangle$ (Fig. 1a) and $\langle m; n+x+y; n; n+x+y \rangle$ (Fig. 1b). The calculation is straightforward and leads to

$$\begin{aligned}
\langle m; n+2x; n; n+2x \rangle & \quad V_k = 4t^{-3} (1 - 4^2) \frac{1}{2} 2^{-4} + 12^{-6}; \\
\langle m; n+x+y; n; n+x+y \rangle & \quad V_? = 16t^{-3} (1 - 4^2) 2^{-2} 12^{-4} + 48^{-6};
\end{aligned} \quad (12)$$

The terms with t are due to the hopping part H_t of the Hamiltonian (1), and the other terms are due to the H_J part. In calculations we used that $2 + 4^2 = 1$ and $2 = 2 + 10^2 - 8t$.

The $\langle m; n; m; n \rangle$ is in essence the potential energy in coordinate space. There is a very weak attraction at $t = 1$ which is due to configurations with closed holes minimizing H_{Ji} (see, e.g., Fig. 2). In minima at $t = 0.5$, $V_k = V_? = 0.018$. Practically within the accuracy of the calculation, it is equal to zero. At $t = 1$ the interaction (12) is repulsive. At $t = 1$, $V_k = \frac{1}{16} (t - \frac{13}{8})$ and $V_? = \frac{1}{4} (t - \frac{7}{8})$.

One can easily convert the vertex into momentum representation [Eq. (9) with $J_{2i} = h_{k_1}^\dagger h_{k_2}^\dagger |0\rangle$ and $J_{4i} = h_{k_3}^\dagger h_{k_4}^\dagger |0\rangle]$

$$\langle k_3; k_4; k_1; k_2 \rangle = \frac{8}{N} V_k (\cos^2 q_k + \cos^2 q_?) + V_? \cos q_k \cos q_? \quad (3 \neq 4) \quad 12;34: \quad (13)$$

Here $q = k_1 - k_3 = k_4 - k_2$; $12;34 = 1$ if $k_1 + k_2 = k_3 + k_4$ and $12;34 = 0$ otherwise.

We expect that the vertex (13) calculated for the Ising background provides also a reasonable estimate for the case of the Neel background. Moreover, we think that for $t = 1$, due to the arguments presented after Eq. (6), it is actually better justified for the Neel background than for the Ising one. The calculation of the quantum fluctuation correction is rather cumbersome and we omit it for the case of parallel spins. We feel, having an experience with holes of opposite spins, that the correction is not large.

3 Interaction of the holes with opposite spins on the Ising background. Energy of the bound state

Similarly to the previous case, it is convenient to use coordinate representation (10). The contact vertex is defined by Eq. (11) with the state $|m; n\rangle = h_m^\dagger h_n^\dagger |0\rangle$. For distant sites m and n , the second term in (11) exactly cancels out the first term and the interaction vanishes. As in the case of parallel spins, there is a potential-like term $\langle m; n+x; n; n+x \rangle$ (Fig. 3a). In addition, there appear effective hopping of clusters $\langle m; n+y; n; n+x \rangle$ (Fig. 3b) and $\langle m; n-x; n; n+x \rangle$ (Fig. 3c) which turn out to be equal. Simple calculation gives

$$\begin{aligned}
\langle m; n+x; n; n+x \rangle & \quad U = 4t \left((1 - 5^2 + 4^4) \frac{1}{4} 4^2 + \frac{27}{2} 4^{-4} - 12^{-6} \right); \\
\langle m; n-x; n; n+x \rangle & = \langle m; n+y; n; n+x \rangle \quad T; \\
T & = 2t \left((1 - 9^2 + 32^4) 2^{-2} (1 - 4^2) (1 - 12^2) \right);
\end{aligned} \quad (14)$$

Again, the terms with t are due to the hopping part of the Hamiltonian (1), and the other terms are due to the H_J part. To be accurate, we should say that within the wave function (10) in the t - J_2 model there are more distant configurations which contribute to hole-hole interaction (Fig. 4). However, the corresponding values of interaction potential

$$\langle m; n+2x+y; n; n+2x+y \rangle = 3 \langle m; n+3x; n; n+3x \rangle = \frac{3}{4} 4^{-4} \quad (15)$$

are very small and we neglect them.

Now we can consider the bound state of two holes. It is quite obvious that in the present approximation the wave function of such a state with the total momentum must be of the form

$$\psi_p = \frac{1}{N} \sum_{n,m} \langle r_n, r_m | h_m^\dagger h_{n\#}^\dagger \rangle \exp[i\frac{1}{2}p \cdot (r_n + r_m)] \quad (16)$$

The equation for bound state is given by variation of $\langle h_p | j(H - E) | j_p \rangle$ with respect to the coefficients $\langle i |$. If we separate the binding energy ($E = 2 + \epsilon$), the equation becomes of the form

$$\langle h_p | j(H - 2) | j_p \rangle - \langle h_p | j_p \rangle = 0 \quad (17)$$

The matrix elements $\langle h_p | j(H - 2) | j_n \rangle$ are given by Eqs. (14). In the term $\langle h_p | j_p \rangle$ we set $\langle h_p | j_n \rangle = \langle h_p | j_p \rangle$. Actually, the correction to the last equality is small, and therefore it gives only a small correction to ϵ . To avoid misunderstanding, we should note that in the effective interaction (12,14) we calculated the overlapping $\langle h_p | j_n \rangle$ exactly because 2 is much larger than the binding energy ϵ . Further calculation is straightforward. Only the coefficients $\langle i |$ corresponding to the closest positions of clusters are not equal to zero. The obtained coefficients correspond to p- and d-wave states which turn out to be degenerate. The energy is minimum at $p = 0$, and the binding energy is equal to

$$\epsilon = U - 2T = 16t^3(1 - 7^2) - \frac{1}{4} 2^2 - 18 \cdot 5^4 + 84 \cdot 6^6 \quad (18)$$

Now we would like to derive Eq. (18) using the momentum representation. For the t-J_z model, it is just another mathematical way. Nevertheless, it is useful because for the t-J model, which we are mainly interested in, the coordinate representation is not convenient. Taking $|j; 2i\rangle = h_{k_1}^\dagger h_{k_2}^\dagger |i\rangle$ and $|j; 4i\rangle = h_{k_3}^\dagger h_{k_4}^\dagger |i\rangle$ in Eq.(9) and using Eqs. (10), one can easily transform the interaction (14) into momentum representation [cf. with Eq. (13)]

$$V_{\#}(k_3; k_4; k_1; k_2) = \frac{8}{N} A_{q_1} + B (i_1 i_3 + i_2 i_4) i_{12; 34} \quad (19)$$

$$A = U - 2T = 16t^3(1 - 7^2) - \frac{1}{4} 2^2 - 18 \cdot 5^4 + 84 \cdot 6^6;$$

$$B = 4T = 8t (1 - 9^2 + 32^4) - 4^2 (1 - 4^2) (1 - 12^2);$$

Here $i_i = k_i$. The simple structure of the vertex (19) reflects the simplicity of quasiparticle ansatz (10). The quasiparticles interact only when they are at the nearest neighbor sites which gives only q_1 and i_j terms.

Let g_k be the wave function of a pair with the total momentum $p = 0$:

$$\langle r_n; r_m \rangle = \frac{1}{N^{3/2}} \sum_k g_k h_k^\dagger h_{k\#}^\dagger \exp[ik \cdot (r_n - r_m)] \quad (20)$$

This function obeys the Bethe-Salpeter (BS) equation

$$(E - 2\epsilon_k) g_k = \sum_{k^0} V_{\#}(k; k; k^0; -k^0) g_{k^0} \quad (21)$$

where the summation is carried out over the Brillouin zone ($\cos k_x^0 + \cos k_y^0 = 0$). For the Ising case $k = [0, 0]$. Using the vertex (19), one can easily find that there are two degenerate solutions of Eq. (21) corresponding to two symmetries

$$\begin{aligned} \text{d-wave: } g_k &= \frac{1}{2} (\cos k_x - \cos k_y); \\ \text{p-wave: } g_k &= 2 \sin k_x; \end{aligned} \quad (22)$$

The i_j terms cancel in the process of integration. The value of binding energy is determined by the q_1 part $\epsilon = A = U - 2T$ which exactly agrees with Eq. (18).

Plot of the hole-hole binding energy in the t-J_z model as a function of t is presented in Fig. 5. The result at $t = 0$ ($\epsilon = 1/4$) can be very easily understood. Two simple holes bound together have one less antiferromagnetic bond broken than when they are apart. As t increases, holes tend to part to gain delocalization

energy. For $t \geq 2$ there is a reasonable agreement between our result and that of Ref.[26]. However for larger t , our curve lies substantially lower than in Ref.[26]. Keeping in mind that our wave function on the Ising background (10) is not very good for $t \geq 1$ [see discussion after Eq. (6)], we would not like to insist on our result. Let us discuss the point in more detail. In our calculations for $t \geq J_2$ model we make in essence an expansion in $\frac{t}{zJ}$, where $z = 4$ is the number of neighbor sites. Therefore, one can expect it to be valid up to $t=J \approx 4$. Unfortunately at large t , there takes place a strong compensation in the hopping part of the binding energy (18) $= U - 2T - \frac{t}{2}(1 - 7^{-2})! - \frac{t}{16}$. It means that this contribution is asymptotically $1=z$. Corrections from including longer strings into the wave function (10) are $1=z$ as well. Therefore they are very important for this value. The situation is different in the t - J model and now we come to this problem.

4 Interaction of the holes with opposite spins on the Neel background. Energy of the bound state

Since we are finally interested in calculation of the binding energy, let us look at the BS equation (21). The Neel case differs from the Ising one in two ways: 1) There is a non-trivial dispersion ϵ_k [Eq.(3)]. 2) The vertex function is renormalized by quantum fluctuations. Generally speaking, the calculation of fluctuation correction to is straightforward: One should only substitute the wave functions (2) into Eq. (9). As we have pointed out in discussion after Eq. (6), the wave function (2) on the Neel background is actually more reliable than on the Ising one. Unfortunately straightforward calculation of is extremely cumbersome because there appear many complicated correlators. Thus we have to make some simplifications.

1) For the Ising case the wave function of bound state (22) is concentrated near the edge of the Brillouin zone ($k_x = 0$). We will prove that for the Neel background this concentration is even stronger. (The same conclusion can be made from the plots of wave functions presented in Ref. [26].) The reason of this effect is quite evident: The bottom of the single-hole band (3) is exactly at the edge of the Brillouin zone. Using this fact, we will consider the vertex $\chi_{\#}(k_3; k_4; k_1; k_2)$ only for momenta lying at the edge of the Brillouin zone ($k_x = 0$).

2) Another simplification is connected with calculation of background spin correlators $\langle \psi_n^+ S_m S_l^z \rangle$. We make an expansion in the parameter of the Hamiltonian (1) and restrict ourselves to the first order. Although the physical value is $\gamma = 1$, this can be quite a reasonable way to estimate the influence of quantum fluctuations. For example, for the transverse nearest neighbor correlator this gives $\langle \psi_n^+ S_{n+1} \rangle = \frac{2}{6}! \approx 0.167$ which is in good agreement with exact numerical value. In the first order in γ , it is the only correlator which differs from its Ising value.

3) The last simplification is the following. We will calculate quantum fluctuation correction only in the H_t part of the vertex. Actually, in the Ising limit there is no compensation in H_J contribution into the vertex (19) and into the binding energy (18). Therefore, one should to expect only small correction from fluctuations. It is not the case for the H_t contribution where happens a strong compensation: $= U - 2T - \frac{t}{2}(1 - 7^{-2})! - \frac{t}{16}$ (see discussion in the end of the previous section).

Calculation with the given above simplifications is still cumbersome but straightforward. Using Eqs. (2) and (9), we get the correction

$$\chi_{\#}(k_3; k_4; k_1; k_2) = -2qt \frac{8}{N} \gamma^3 (14 \gamma_q + \gamma_{k_1+k_3} + \gamma_{k_2+k_4}) + (16 \gamma^3 \gamma^3 - 56 \gamma^5) \gamma_{k_1+k_3+k_2+k_4}; \quad (23)$$

where $2q_l = \langle \psi_n^+ S_{n+1} \rangle$, $\gamma_l = \frac{1}{6}$ is the nearest neighbor correlator. Clearly, the correction (23) is repulsive. The coefficients γ and γ_l in Eq. (23) correspond to the Ising limit (10) since the expansion in γ is carried out explicitly. The γ^3 term in Eq. (23) comes from expansion of $\langle \psi_n^+ S_{n+1} \rangle$ in Eq. (9), the $\gamma^3 \gamma^3$ term comes from overlapping $\langle \psi_n^+ S_{n+1} \rangle$, and the γ^5 term comes from expansion of energy $(\epsilon_1 + \epsilon_2)$. The $(\gamma_{k_1+k_3} + \gamma_{k_2+k_4})$ part arises from interaction of the holes at next-nearest neighbor sites. Its relatively small value reflects the importance of quasiparticles interaction at the nearest neighbor sites.

Adding the quantum fluctuation correction (23) to the Ising value of vertex (19), we get contact hole-hole vertex for the t - J model

$$\begin{aligned} \chi_{\#}^0(k_3; k_4; k_1; k_2) &= \frac{8}{N} A^0 \gamma_q + B (\gamma_1 \gamma_3 + \gamma_2 \gamma_4) + \frac{C}{2} (\gamma_{k_1+k_3} + \gamma_{k_2+k_4}) \gamma_{k_1+k_3+k_2+k_4}; \\ A^0 &= A + 10 t^3 \gamma^3; \quad C = \frac{2}{3} t^3 \gamma^3; \end{aligned} \quad (24)$$

Expressions for A and B are presented in Eq. (19). In transformation from Eq. (23) to Eq. (24) we take into account the normalization condition $\gamma^2 + 4 \gamma^2 = 1$.

Let us consider two hole bound state with zero momentum $p = 0$. If we expand the single hole dispersion (3) near the band bottom

$$\epsilon_k = \epsilon_{\text{bot}} + \frac{1}{2} k^2; \quad \epsilon_k = \frac{4t^2(x+y)}{[4 + 4t^2(1+y)]^{1/2}}; \quad (25)$$

the BS equation (21) can be rewritten in the form

$$(\epsilon_k - \epsilon_{\text{bot}})g_k = \sum_{k^0} \epsilon_{\#}^0(k; k; k^0; -k^0)g_{k^0}; \quad (26)$$

where $\epsilon_{\#}^0$ is given by Eq. (24). Let us first treat t -term perturbatively. At $t = 0$, in spite of appearance of $(k_1 + k_3 + k_2 + k_4)$ term in the vertex, the solutions (22) still satisfy Eq. (26) with binding energies

$$\epsilon_p^{(0)} = a_p - A^0 - C; \quad \epsilon_d^{(0)} = a_d - A^0 + C; \quad (27)$$

The first corrections in perturbation theory are

$$\epsilon_p^{(1)} = \epsilon_p^{(0)} + \frac{3}{16} k^2; \quad \epsilon_d^{(1)} = \epsilon_d^{(0)} + \frac{1}{16} k^2; \quad (28)$$

At $t = 1$ the corrections become too large and the perturbation theory is not valid. However, general solutions of Eq. (26) are obvious:

$$\begin{aligned} \text{p-wave: } g_k & \propto \frac{\sin k_x}{k^2}; \\ \text{d-wave: } g_k & \propto \frac{\cos k_x \cos k_y}{k^2}. \end{aligned} \quad (29)$$

Let us emphasize that the factor $(\epsilon_k - \epsilon_{\text{bot}})^{-1}$ (s is the symmetry label, $s = p, d$) substantially enhances the wave function near the edge of the Brillouin zone ($k_x = 0$). Substitution of the solution (29) into the BS equation (26) gives the selfconsistency condition for ϵ_s . The binding energy vanishes at the point $t = t_s$ where a_s (27) vanishes. Near this point

$$\epsilon_s \approx \frac{a_s^2}{t - t_s^2} \quad (30)$$

To be precise, we should note that for the d-wave there is an additional weak logarithmic $(\ln \frac{1}{t - t_d})$ dependence in the coefficient in this estimation.

The numerical solutions for ϵ_s as a function of t are presented in Fig. 6 (d-wave) and Fig. 7 (p-wave). Dashed lines correspond to $\epsilon = 1$ in vertex (24) which is the physical value. There are higher orders in t which we do not take into account. To estimate their contribution, we present also results with $\epsilon = 0.5$ (solid lines). We believe that exact solution lies somewhere between $\epsilon = 0.5$ and $\epsilon = 1$. Of course the choice of $\epsilon = 0.5$ is rather arbitrary. However, we can recall the linear expansion in t for the parameters x and y which determine the single-hole dispersion (3): $x = 12q = \dots$, $y = 4q = \dots = 3$ [13]. To fit the physical values (6), we must take $q = 0.5$. Anyway, it is not very important since the interval between the curves with $\epsilon = 0.5$ and $\epsilon = 1$ in Figs. (6) and (7) is quite narrow. At most t , the d-state has lower energy. There is a good agreement between our results and results obtained by a numerical variational method [26] which are also presented in Figs. (6) and (7). Thus to our view, the bound state disappears at $t \approx 2.3$. This conclusion is in agreement with the results of recent works [27, 34].

Solving the BS equation for $p \neq 0$, one can easily derive the dispersion of the bound state. For example for small p at $t = J$

$$E_p(p) \approx 0.165p^2; \quad E_d(p) \approx 0.048p^2; \quad (31)$$

For $p \neq 0$ the solution in general case has no definite symmetry (d or p). The labels in above equation denote the symmetry of corresponding solution at $p = 0$.

5 Conclusion

We calculated the vertex function of contact interaction between two holes in the t - J model. Contact interaction is caused by exchange of spin excitations with momentum q and has no retardation. We considered this interaction for both parallel and opposite directions of the holes spins. For the most interesting case of opposite

spins, there was found an attraction at $t=J$ [2, 3] which gives a very shallow bound state (with p or d symmetry). Our result for binding energy agrees with that of recent works [26, 27, 34], but disagrees with that of exact diagonalizations on small-size clusters [16], [28]–[32]. Following Refs. [16, 26, 33], we suspect that the discrepancy is caused by finite-size effects in exact diagonalizations. Recent calculation [34] of the binding energy for larger clusters (up to 26 sites) and the scaling with system size probably confirms this point of view.

The real physical value of $t=J$ in high- T_c superconductors is $t=J/3$ (see, e.g., [37, 38, 39]). Therefore, the two hole short-range bound state obtained in the present work is probably irrelevant to high- T_c superconductivity. Moreover, we think that this bound state is by-product of the pure t - J model. Actually, the size of bound state is of the order of one lattice space $a \approx 7a_B$ (a_B is the Bohr radius). Therefore, the Coulomb hole-hole repulsion on neighbor sites is $V = e^2/(a) \approx 4\text{eV}$, where ϵ is the dielectric constant. More realistic structure of hole wave function, which takes into account oxygen ions [38], does not change this estimation. Even with the static value $\epsilon = 50$, the Coulomb repulsion $V \approx 0.1\text{ eV}$ destroys the bound state $\epsilon \approx 0.3J \approx 0.03\text{ eV}$. Actually the Coulomb repulsion is even larger because in interaction we must use the dynamic value of the dielectric constant (!) with $\epsilon \approx 0.01\text{ eV}$ which is definitely smaller than the static value.

Thus we have to introduce a short range hole-hole repulsion into the t - J model. One way to do it is to consider a t - J - V model [40, 41]. However, we do not think that the results are sensitive to the specific way of introduction of short range repulsion. We believe that the main problem is long range dynamics at distances p^{-1} . This problem includes the long range antiferromagnetic order instability [24]. Nevertheless the short range repulsion is very important.

The situation is quite similar to the pion condensation in nuclear matter [25], where both the interaction of a nucleon with the Goldstone excitation (pion) and short range nucleon-nucleon repulsion are important. At the moment we believe that suitable description of high- T_c superconductors is an effective long-range theory with spin-1/2 holes and Goldstone gapless spin-waves. (To avoid misunderstanding, we should note that long range instability destroys antiferromagnetic order and therefore Goldstone spin-waves definitely are not the physical excitations. They just provide a suitable basis set.) The single hole properties as well as hole - spin-wave vertex [19] are described by the t - J model. However besides that, we have to introduce "by hands" the contact hole-hole repulsion.

It is interesting to notice that investigation of high- T_c superconductivity could help to better understanding of pion condensation. Actually the long-range instability (pion condensation) [25] and the BCS-type nucleon-nucleon pairing [42] are usually considered as independent phenomena. However now, due to high- T_c superconductivity, we understand that they are closely related.

ACKNOWLEDGMENTS

We are grateful to V. I. Belinicher who draw our attention to contact hole-hole interaction. We wish to acknowledge stimulating discussions with him as well as with I. B. Khriplovich and I. V. Kolokolov. We are especially grateful to D. Poilblanc for supplying us with some references and communicating to us the results of work [34] prior to publication. One of us (A.L.C.) thanks the Soros-Akademgorodok Foundation for financial support.

References

- [1] J. E. Hirsch, Phys. Rev. Lett. 54, 1317 (1985).
- [2] C. Gross, R. Joynt, and T. M. Rice, Phys. Rev. B 36, 381 (1987).
- [3] J. Oitmaa and D. D. Betts, Can. J. Phys. 56, 897 (1971).
- [4] D. A. Huse, Phys. Rev. B 37, 2380 (1988).
- [5] E. Manousakis, Rev. Mod. Phys. 63, 1 (1991).
- [6] B. Shraiman and E. Siggia, Phys. Rev. Lett. 61, 467 (1988).
- [7] S. Schmitt-Rink, C. M. Varma, and A. E. Ruckenstein, Phys. Rev. Lett. 60, 2793 (1988).
- [8] C. L. Kane, P. A. Lee, and N. Read, Phys. Rev. B 39, 6880 (1989).
- [9] S. Sachdev, Phys. Rev. B 39, 1232 (1989).
- [10] C. Gross and M. D. Johnson, Phys. Rev. B 40, 9423 (1989).
- [11] J. H. Xu, C. S. Ting, and T. K. Lee, Phys. Rev. B 43, 8733 (1990).
- [12] G. Martinez and P. Horsch, Phys. Rev. B 44, 317 (1991).
- [13] O. P. Sushkov, Solid State Communications 83, 303 (1992).
- [14] S. A. Trugman, Phys. Rev. B 37, 1597 (1988).
- [15] E. Kaxiras and E. Manousakis, Phys. Rev. B 38, 866 (1988).
- [16] J. Bonca, P. Prelovsek, and I. Sega, Phys. Rev. B 39, 7074 (1989).
- [17] E. Dagotto, R. Joynt, S. Bacci, and E. Cagliano, Phys. Rev. B 41, 9049 (1990).
- [18] O. P. Sushkov, Phys. Lett. A 162, 199 (1992).
- [19] O. P. Sushkov, Physica C 206, 264 (1993).
- [20] B. Shraiman and E. Siggia, Phys. Rev. B 40, 9162 (1989).
- [21] G. Wentzel, Phys. Rev. 83, 168 (1951).
- [22] B. Shraiman and E. Siggia, Phys. Rev. Lett. 62, 1565 (1989).
- [23] A. Singh and Z. Teseanovic, Phys. Rev. B 41, 614 (1990).
- [24] O. P. Sushkov and V. V. Flambaum, Physica C 206, 269 (1993).
- [25] A. B. Migdal, Rev. Mod. Phys. 50, 107 (1978).
- [26] R. Eder, Phys. Rev. B 45, 319 (1992).
- [27] M. Boninsegni and E. Manousakis, Florida State University Preprint, FSU-SCRI-92-143, October 1992.
- [28] Y. Hasegawa and D. Poilblanc, Phys. Rev. B 40, 9035 (1989).
- [29] K. H. Luk and D. L. Cox, Phys. Rev. B 41, 4456 (1990).
- [30] E. Dagotto, J. A. Riera, and A. P. Young, Phys. Rev. B 42, 2347 (1990).
- [31] T. Itoh, M. Arai, and T. Fujiwara, Phys. Rev. B 42, 4834 (1990).
- [32] J. A. Riera, Phys. Rev. B 43, 3681 (1991).
- [33] J. Bonca, P. Prelovsek, and I. Sega, Phys. Rev. B 42, 10706 (1990).
- [34] D. Poilblanc, private communication.

- [35] L. N. Bulaevskii, E. N. Nagaev, and D. L. Khomskii, Zh. Eksp. Teor. Fiz. 54, 1562 (1968) [Sov. Phys. JETP 27, 836 (1968)].
- [36] V. A. Dzuba, V. V. Flambaum, P. G. Silvestrov, and O. P. Sushkov, in Relativistic Quantum Electrodynamics and Weak Interaction Effects in Atoms, edited by W. Johnson, P. Mohr, and J. Sucher, AIP Conference Proceedings 189, 254 (1989).
- [37] H. Eskes, G. A. Sawatzky, and L. F. Feiner, Physica C 160, 424 (1989).
- [38] V. V. Flambaum and O. P. Sushkov, Physica C 175, 347 (1991).
- [39] I. V. Belinicher and A. L. Chemshev (submitted to Phys. Rev. B).
- [40] S. Kivelson, V. Emery, and H. Q. Lin, Phys. Rev. B 42, 6523 (1990).
- [41] E. Dagotto and J. Riera, Phys. Rev. B 46, 12084 (1992).
- [42] A. Bohr and B. R. Motelson, Nuclear Structure, Vol. 2 (Benjamin, New York, 1974).

FIGURE CAPTIONS

FIG . 1. Contact interaction of the holes with parallel spins. Only the configurations with holes in the centers of square clusters are shown, but each hole hops within the corresponding cluster.

FIG . 2. Configuration with closed holes which minimizes the H_J and gives a very small attraction between holes with parallel spins.

FIG . 3. Matrix elements for the holes with opposite spins. Only the configurations with holes in the centers of square clusters are shown, but each hole hops within the corresponding cluster. Fig. a represents diagonal interaction, and Figs. b,c represent hopping of cluster.

FIG . 4. Distant configurations for interaction of opposite spin holes in the $t\text{-}J_z$ model. Only the configurations with holes in the centers of square clusters are shown, but each hole hops within the corresponding cluster.

FIG . 5. Hole-hole binding energy in the $t\text{-}J_z$ model as a function of t . The d- and p-waves are degenerate. Solid line is the result of the present work [Eq. (18)]. The circles are results from numerical variational method [26], and the triangles are results from exact diagonalizations [16]).

FIG . 6. Hole-hole d-wave binding energy in the $t\text{-}J$ model. Solid line is the result of the present work at $U = 0.5$. Dashed line is the same at $U = 1$. The circles represent the results of numerical variational method [26].

FIG . 7. Hole-hole p-wave binding energy in the $t\text{-}J$ model. Solid line is the result of the present work at $U = 0.5$. Dashed line is the same at $U = 1$. The circles represent the results of numerical variational method [26].