

A B S T R A C T

We study the electronic states of isolated fullerene anions C_{60}^n ($1 \leq n \leq 6$) taking into account the effective interaction between electrons due to exchange of intramolecular phonons. If the vibronic coupling is strong enough such an effect may overwhelm Hund's rule and lead to an ordering of levels that can be interpreted as on-ball pairing, in a manner similar to the pairing in atomic nuclei. We suggest that such effects may be sought in solutions of fulleride ions and discuss recent experimental results.

I. INTRODUCTION

The discovery of superconductivity in alkali-metal-doped fullerenes¹ K_3C_{60} and Rb_3C_{60} has raised interesting questions about the electron-phonon coupling in such compounds and its interplay with Coulomb repulsion. C_{60} is a highly symmetrical molecule i.e. it is a truncated icosahedron and its electronic lowest unoccupied molecular orbitals (LUMO) are threefold degenerate^{2;3;4}. They form a T_{1u} representation of the icosahedral group I_h . Filling the LUMO in C_{60}^n anions leads in a naive picture to narrow, partially filled bands in the bulk fullerides. The bandwidth W is determined by the hopping between the C_{60} molecules which are quite far apart and $W \approx 0.5$ eV. The coupling of some H_g phonons with electrons residing in the T_{1u} orbital has been suggested to be responsible for the superconductivity^{5;6;7}. The Coulomb repulsion also may be important on the ball⁸. Several authors⁹⁻¹² have undertaken the study of the Jahn-Teller distortion that is expected in the fullerene anions. In such calculations one considers the electrons as fast degrees of freedom and the phonon normal coordinates are treated as static¹³.

In this paper we investigate the interplay between the electronic and phononic degrees of freedom on an isolated fullerene anion. We study an effect that goes beyond the Born-Oppenheimer approximation which is the modification of electronic levels due to phonon exchange. We obtain the lifting of degeneracy by a perturbation calculation in the case of an undistorted anion. The ordering of levels can be described as "anti-Hund" rule. Our calculation is very close in spirit to the standard treatment of the electron-phonon coupling in superconducting metals. Here we argue that the energy scale of this effect may be comparable to that of the Jahn-Teller effect. This is because the phonons have high frequencies as well as medium to strong coupling to electrons. The effect we observe

may be sought by spectroscopy of solutions of fullerenes in liquid ammonia, for example. We discuss the opposite effect of Coulomb interaction, leading to Hund's rule in ordinary situations. Finally we point out that experimentally observed spectra may be at least partially explained by our crude calculation.

In section II we discuss the electron-phonon coupling, section III gives our results for the electronic levels of the anions, section IV discusses the competition with Coulomb repulsion and section V contains a brief discussion of near-IR spectroscopic measurements and EPR experiments on solutions of fullerene anions as well as our conclusions.

II. THE ON-BALL ELECTRON-PHONON INTERACTION

The electronic structure of electrons in the C_{60} molecule is well known to be given by a simple Hückel calculation. The levels are labeled⁴ by the irreducible representations (irreps) of the icosahedron group I_h . One important property has to be noted: three of the I_h irreps are the $l=0,1,2$ spherical harmonics of $SO(3)$ which do not split under the I_h group. They are commonly named A_g , T_{1u} , H_g . In addition there is also the twofold spin degeneracy.

In the ground-state of the neutral C_{60} molecule all levels up to H_u included are completely filled thus building a singlet state $j=0$. The LUMOs are the six T_{1u} states. These are occupied upon doping with extra electrons and the ground-state becomes then degenerate. One then expects the Jahn-Teller effect to distort the anion and lift this orbital degeneracy^{13;14}. We focus on another effect which goes beyond the Born-Oppenheimer approximation in the sense that nuclear motions are crucial for its very existence: the coupling of the T_{1u} electrons to the vibrational modes of the molecule (also referred to

as phonons). Phonon exchange between electrons leads to an effective electron-electron interaction that competes with Coulomb repulsion and may lead to anti-Hund ordering of energy-levels.

For simplicity we treat this effect assuming the absence of the Jahn-Teller distortion. The next logical step would be to compute first the distortion pattern of the anion under consideration, then obtain its vibrational spectrum and the electron-phonon coupling in the distorted structure and then compute again the effective electron-electron interaction. As a first investigation of electron-phonon coupling we use a perturbation scheme suited to degenerate levels we will derive an effective electron-electron interaction with the assumption that filled states lying below the T_{1u} level remain frozen so that intermediate states involve only T_{1u} excitations. Indeed the H_u $\{T_{1u}$ gap is $\sim 2\text{eV}$ whereas maximum phonon energies are $\sim 0.2\text{eV}$.

A typical electron-phonon interaction term reads:

$$W = \sum_{m_1, m_2} \sum_l X_{lm_1 m_2} \sum_{\sigma_1, \sigma_2} f_{m_1 m_2} X_{l\sigma_1} c_{m_1}^{\sigma_1} c_{m_2}^{\sigma_2} :$$

Here X_{lm} are normal coordinates, the subscript referring both to the irrep and to the row in the irrep they belong to, $c_{m_1}^{\sigma_1}$ is the creation operator for an electron with spin σ_1 in the T_{1u} ($l=1$) level, m_1 taking one of the $m = -1, 0, 1$ values, and $f_{m_1 m_2}$ are complex coefficients. The $c_{m_1}^{\sigma_1}$ operators transform as $l=1$ j_l vectors under I_h symmetries, and their conjugates $c_{m_1}^{\sigma_1}$ transform as $(-1)^{m_1+1} j_l$ vectors. The $(-1)^{m_2+1} c_{m_1}^{\sigma_1} c_{m_2}^{\sigma_2}$ products transform then as members of the $T_{1u} \otimes T_{1u}$ representation, which in the I_h group splits as:

$$T_{1u} \otimes T_{1u} = A_g + T_{1g} + H_g :$$

This selects the possible vibrational modes T_{1u} electrons can couple to. In fact, only H_g modes split the degeneracy⁷.

Let us consider a particular ν -fold degenerate multiplet of H_g modes. Their normal coordinates will be labelled X_m , m ranging from -2 to $+2$. Since H_g appears only once in the product $T_{1u} \times T_{1u}$, the interaction is determined up to one coupling constant g by the usual formula for the coupling of two equal angular momenta to zero total angular momentum :

$$W = g \sum_m (-1)^m X_m \quad (1):$$

The X_m may be chosen such that $X_m^y = (-1)^m X_{-m}$ and have the following expression in terms of phonon operators:

$$X_m = \frac{1}{\sqrt{2}} a_m + (-1)^m a_m^y \quad (2)$$

whereas the a_m are the irreducible $l=2$ tensor operators built from the $c^y c$ products according to:

$$a_m = \sum_{m_1} (1; 1; 2 | m_1; m_1; m) (-1)^{(m-m_1+1)} c_{m_1}^y c_{m+m_1} \quad (3)$$

where $(l_1; l_2; l | m_1; m_2; m)$ are Clebsch-Gordan coefficients.

We now consider a doped C_{60}^n molecule, $0 < n < 6$. Its unperturbed degenerate ground-states consist of j_0 to which n T_{1u} electrons have been added times a zero-phonon state. They span a subspace denoted by E_0 . In E_0 the unperturbed Hamiltonian H_0 reads:

$$H_0 = \sum_{m; t_{1u}} \epsilon_m^y c_m c_m + \hbar \omega \sum_m a_m^y a_m$$

where ϵ_{1u} is the energy of the T_{1u} level, $\hbar\omega$ is the phonon energy of the H_g multiplet under consideration. Within E_0 the effective Hamiltonian up to second order perturbation theory is given by:

$$H_{eff} = E_0 P_0 + P_0 W P_0 + P_0 W (1 - P_0) \frac{1}{E_0 - H_0} (1 - P_0) W P_0$$

where P_0 is the projector onto E_0 , E_0 is the unperturbed energy in this subspace which is just the number of doping electrons times ϵ_{1u} . The linear term in W gives no contribution. Using expressions (1) and (2) for W and X_m one finds:

$$H_{eff} = H_0 + \frac{g^2}{2\hbar^2} \sum_{m_1, m_2} X_m (1)^{m_1+m_2} \quad (4)$$

where we have now included spin indices. We can now use equation (3) to express H_{eff} as a function of c and c^\dagger operators and put it in normal ordered form using fermion anticommutation rules. In this process there appears a one-body interaction term which is a selfenergy term. We will henceforth omit the H_0 term which is a constant at fixed number of doping electrons.

Let us now define pair creation operators $A_{lm}^{s\ y}$ which when operating on the vacuum $|0\rangle$ create pair states of T_{1u} electrons that are eigenfunctions of L, S, L_z, S_z , where L, S are total angular momentum and spin, and L_z, S_z their z -projections. l and s can take the values 0, 1, 2 and 0, 1 respectively. This holds also if $|0\rangle$ is taken to be the singlet state $|0\rangle_i$.

$$A_{lm}^{s\ y} = \sum_{m_1, m_2} (1; 1; l; m_1; m_2; m_1 + m_2) \left(\frac{1}{2}; \frac{1}{2}; s; j_1; j_2\right) c_{m_1-1}^y c_{m_2-1}^y \quad (5)$$

The quantity $A_{lm}^{s\ y}$ is non-zero only if $(l+s)$ is even and the norm of $A_{lm}^{s\ y}|0\rangle$ is then equal to $\sqrt{2}$. The inverse formula expressing $c^\dagger c^\dagger$ products as $A^{s\ y}$ operators is:

$$c_{m_1-1}^y c_{m_2-2}^y = \sum_{l, s} (1; 1; l; m_1; m_2; m_1 + m_2) \left(\frac{1}{2}; \frac{1}{2}; s; j_1; j_2\right) A_{lm_1+m_2}^{s\ 1+2\ y} : \quad (6)$$

$A_s H_{eff}$ is a scalar, its two-body part may be written as a linear combination of diagonal $A_{lm}^s A_{lm}^s$ products whose coefficients depend only on l and s :

$$\sum_{ls,m} F(l;s) A_{lm}^s A_{lm}^s$$

The $F(l;s)$ coefficients are calculated using expressions (4), (3), (6). We then get H_{eff} in normal form :

$$H_{eff} = \frac{5g^2}{6h!} \hat{N} + A_{00}^{00y} A_{00}^{00} + \frac{1}{2} \sum_m A_{1m}^1 A_{1m}^1 + \frac{1}{10} \sum_m A_{2m}^{00y} A_{2m}^{00} : \quad (7)$$

In this formula \hat{N} is the electron number operator for the T_{1u} level; the \hat{N} term appears when bringing H_{eff} of expression (4) in normal ordered form. In our Hamiltonian formulation the effective interaction is instantaneous.

There are actually eight H_g multiplets in the vibrational spectrum of the C_{60} molecule. To take all of them into account we only have to add up their respective coefficients $5g^2=6h!$, their sum will be called \dots .

III-THE ELECTRONIC STATES OF FULLERENE ANIONS

We shall now, for each value of n between 1 and 6, find the n -particle states and diagonalize H_{eff} . The Hamiltonian to be diagonalized is that of equation (7) where the prefactor is replaced by \dots . The invariance group of H_{eff} is $I_h \times SU(2)$. The n -particle states may be chosen to be eigenstates of $L; S, L_z; S_z$ and we shall label the multiplets by $(l;s)$ couples, in standard spectroscopic notation (^2s+1L stands for (l,s)). The pair $(l;s)$ label $SO(3) \times SU(2)$ irreps which, as previously mentioned, remain irreducible under $I_h \times SU(2)$ as long as l doesn't exceed 2; for larger values of l $SO(3)$ irreps split under I_h . Fortunately enough, the relevant values of l never exceed 2. Moreover given any value

of $n, (l; s)$ multiplets appear at most once so that the energies are straightforwardly found by taking the expectation value of the Hamiltonian in one of the multiplet states. The degeneracies of the levels will then be $(2l+1)(2s+1)$. We now proceed to the construction of the states.

$n=1$: There are six degenerate 2P states $c_m^y |j_0\rangle$ whose energy is ϵ_1 .

$n=2$: There are 15 states, generated by applying $A_{lm}^{s\ y}$ operators on $|j_0\rangle$. There is one 1S state, nine 3P states and five 1D states.

$n=3$: There are 20 states. States of given $l; m; s;$ can be built by taking linear combinations of $A_{lm}^{s\ y} |j_0\rangle$ states according to:

$$\sum_{m_1; s_1}^X (l_1; 1; l; m_1; m) (s_1; \frac{1}{2}; s; j_1; j_0) A_{l_1 m_1}^{s_1\ y} c_{m_1}^y |j_0\rangle$$

These states belong to the following multiplets: $^4S, ^2P, ^2D$.

$n=4$: There are 15 states, which are obtained by applying $A_{lm}^{s\ y}$ operators on $A_{00}^{00y} |j_0\rangle$.

$n=5$: There are six 2P states which are $c_m^y A_{00}^{00y} A_{00}^{00y} |j_0\rangle$ and whose energy is ϵ_5 .

$n=6$: There is one 1S state whose energy is 0.

The corresponding energies are given explicitly in Table I and displayed in Fig. 1. It is interesting to note that the above treatment of electron-phonon interaction parallels that of pairing forces in atomic nuclei^{15,16}. Of course in the case of finite fermionic systems there is no breakdown of electron number but there are well-known "odd-even" effects that

appear in the spectrum. In our case pairing shows up in the 1S ground state for C_{60}^2 rather than 3P as would be preferred by Coulomb repulsion i.e. Hund's rule. The construction of the states above is that of the seniority scheme in nuclear physics¹⁶. We note that similar ideas have been put forward by V. Kresin some time ago, also in a molecular context¹⁷. The effective interaction that he considered was induced by core polarization.

IV - THE EFFECT OF COULOMB REPULSION

We now consider the Coulomb electron-electron interaction and assume it to be small enough so that it may be treated in perturbation theory. To get some feeling of the order of magnitude of this repulsion we use the limiting case of on-site interaction i.e. the Hubbard model. This Hamiltonian is not specially realistic but should contain some of the Hund's rule physics. The two-body interaction now reads:

$$\frac{U}{2} \sum_i c_{i\uparrow}^\dagger c_{i\uparrow}^\dagger c_{i\downarrow} c_{i\downarrow} ;$$

where the i subscript now labels the orbitals on the C_{60} molecule. The quantity U is

2-3 eV from quantum chemistry calculations¹⁸. Since level degeneracies are split at first order in perturbation theory we confine our calculation to this order and have thus to diagonalize the perturbation within the same subspace E_0 as before. In this subspace it reads:

$$W_H = U \sum_i \sum_{\mu, \nu} h_{\mu\nu} c_{i\mu}^\dagger c_{i\nu}^\dagger c_{i\nu} c_{i\mu} ;$$

where greek indices label one-particle states belonging either to j_{0i} or to the T_{1u} level.

Let us review the different parts of W_H . Note that since the j_{0i} singlet remains frozen

we have the identity: $c_{i\mu}^\dagger c_{i\mu} = 1$ if μ, ν label states belonging to j_{0i} .

{A part involving states belonging to j_0 only:

$$W_{H_1} = U \begin{matrix} X \\ i; \end{matrix} \quad h \ jij^2 h \ jij^2 c'' c'' c'_{\#} c_{\#} :$$

, belong to j_0 . This term is thus diagonal within E_0 and merely shifts the total energy by a constant that does not depend on the number of doping electrons. It won't be considered in the following.

{A part involving both states belonging to j_0 and to the T_{1u} level:

$$W_{H_2} = U^X \begin{matrix} & h & j & i & j & i & j \\ i & j & i & i & i & i & i \end{matrix} c^y \quad c \quad c^y \quad c$$

where ψ_1, ψ_2 belong to the T_{1u} level whereas ψ_3 belongs to $j = 0$. It reduces to:

$$W_{H_2} = U \begin{matrix} X \\ c^y \\ c \end{matrix} \begin{matrix} X \\ h \\ i \end{matrix} \begin{matrix} X \\ j \\ j \end{matrix} \begin{matrix} X \\ i \\ j \end{matrix}$$

The sum over j is just the density on site i for a given spin direction of all states belonging to j_0 which is built out of completely filled irreps. As a result this density is uniform and since j_0 contains 30 electrons for each spin direction it is equal to $1/2$. W_{H_2} then becomes diagonal and reads:

$$W_{H_2} = \frac{U}{2} \frac{X}{C^y C} :$$

Its contribution is thus proportional to the number of T_{1u} electrons. It represents the interaction of the latter with those of the singlet and we won't consider it in the following.

{A part involving only states belonging to the T_{1u} level:

W_{H_3} has the same form as W_H with all indices now belonging to the T_{1u} level. Whereas the interaction has a simple expression in the basis of π states, we need its matrix elements in

the basis of the T_{1u} states. There are in fact two T_{1u} triplets in the one-particle spectrum of the C_{60} molecule, the one under consideration having higher energy. To construct the latter we have first constructed two independent sets of states which transform as $x; y; z$ under I_h . These are given by:

$$|j_i\rangle = \sum_i^X e_i \cdot \mathbf{r}_i |j_i\rangle \quad \text{and} \quad |j_i^0\rangle = \sum_i^X e_i \cdot \mathbf{K}_i |j_i\rangle$$

where \mathbf{e} are three orthonormal vectors, i labels sites on the molecule, the \mathbf{r}_i are the vectors joining the center of the molecule to the sites while the \mathbf{K}_i join the centre of the pentagonal face of the molecule the site i belongs to to the site i . We assume that the bonds all have the same length. These states span the space of the two T_{1u} triplets. The diagonalization of the tight-binding Hamiltonian in the subspace of these six vectors yields then the right linear combination of the $|j_i\rangle$ and $|j_i^0\rangle$ states for the upper lying triplet. From the $x; y; z$ states one constructs $l=1$ spherical harmonics. We then get the matrix elements of W_{H_3} in the basis of T_{1u} states. As E_0 is invariant under I_h operations and spin rotations, W_{H_3} which is the restriction of W_H to E_0 is invariant too. It may thus be expressed using the A, A^\dagger operators by using formula (5) in the same way as the phonon-driven interaction and we finally get:

$$W_{H_3} = \frac{U}{40} A_{00}^{00\dagger} A_{00}^{00} + \frac{U}{100} \sum_m^X A_{2m}^{00\dagger} A_{2m}^{00} \quad (8)$$

which is the only part in W_H that we will keep. Note that there is no contribution from $l=1, s=1 A^\dagger A$ products. Indeed the Hubbard interaction is invariant under spin rotation and couples electrons having zero total S_z . As the coefficients of $A^\dagger A$ products depend solely on l and s they must be zero for $s \neq 0$. The spectrum for any number of T_{1u} electrons is now easily found: see fig 2 and table I.

V -CONCLUSION

The ordering of energy levels in the electron-phonon scheme are clearly opposite to those of Hund's rule (compare fig. 1 and fig. 2). The clear signature of what we can call "on-ball" pairing is the ground state 1S of C_{60}^2 : the two extra electrons are paired by the electron-phonon coupling. We note that the U of the Hubbard model appears divided by large factors: this is simply due to the fact that the C_{60} molecule is large. As a consequence, if $U \approx 2$ eV, Coulomb repulsion may be overwhelmed by phonon exchange. With a H_g phonon of typical energy 100 meV and coupling $O(1)$ as suggested by numerous calculations^{6;7;10}, the quantity λ may be tens of meV.

It seems to us that the cleanest way to probe this intramolecular pairing would be to look at solutions of fullerides leading to free anions such as liquid ammonia solutions or organic solvents¹⁹⁻²². EPR or IR spectroscopy should be able to discriminate between the two types of spectra. Measurements by EPR should determine whether or not the two extra electrons in C_{60}^2 are paired, for example. In near-IR spectroscopy the lowest allowed transition for C_{60}^2 should be at higher energy than that of C_{60} due to the pairing energy while in the Coulomb-Hubbard case it is at lower energy.

Present experiments^{19;20} have studied the near-IR spectra of solutions of fulleride anions prepared by electrochemical reduction. There are several peaks that do not fit a simple Hückel scheme of levels. They do not have an immediate interpretation in terms of vibrational structure^{19;20}. With our energy levels in table I, a tentative fit would lead to

80 meV assuming $U = 0$. Such a value leads to intriguing agreement with the major peaks seen for C_{60}^2 and C_{60}^3 while this is no longer the case for C_{60}^4 and C_{60}^5 .

Finally we mention that recent EPR experiments²² have given some evidence for non-Hund behaviour of the fulleride anions. While one may observe some trends similar to the results of the phonon-exchange approximation, it is clear that the model we used is very crude. The interplay with conventional Jahn-Teller effect is an important factor missing in our study and of a similar order of magnitude. In a bulk conducting solid we do not expect the previous scheme to be valid since the levels are broadened into bands.

Acknowledgements

We thank K.M.Kadish, M.T.Jones, C.A.Reed and J.W.White for informing us about their recent experimental work. We would like to thank also J.P.Blaizot, S.Doniach, C.Fabre, T.Garel and A.Rassat for discussions. We have also benefited from the help of the GDR "Fullerenes" supported by the Centre National de la Recherche Scientifique (France).

R E F E R E N C E S

- [1] For a review see A . F . H e b a r d , Physics Today, November 1992.
- [2] S . S a t p a t h y , Chem . Phys. Letters 130, 545 (1986).
- [3] R . C . H a d d o n and L . T . S c o t t , Pure Appl. Chem . 58, 137 (1986); R . C . H a d d o n , L . E . B r u s , and K . R a g h a v a c h a r i , Chem . Phys. Letters 125, 459 (1986).
- [4] G . D r e s s e l h a u s , M . S . D r e s s e l h a u s , and P . C . E k l u n d , Phys. Rev. B 45, 6923 (1992).
- [5] M . L a n n o o , G . A . B a r a , M . S c h l u t e r and D . T o m a n e k , Phys. Rev. B 44, 12106 (1991). See also K . Y a b a n a and G . B e r t s c h , Phys. Rev. B 46, 14263 (1992).
- [6] V . d e C o u l o n , J . L . M a r t i n s , and F . R e u s e , Phys. Rev. B 45, 13671 (1992).
- [7] C . M . V a r m a , J . Z a a n e n , and K . R a g h a v a c h a r i , Science 254, 989 (1991).
- [8] S . C h a k r a v a r t y and S . K i v e l s o n , Europhys. Lett. 16, 751 (1991); S . C h a k r a v a r t y , M . G e l f a n d , and S . K i v e l s o n , Science 254, 970 (1991); S . C h a k r a v a r t y , S . K i v e l s o n , M . S a l k o l a , and S . T e w a r i , Science 256, 1306 (1992).
- [9] A . A u e r b a c h , "V i b r a t i o n s and B e r r y p h a s e s of c h a r g e d B u c k m i n s t e r f u l l e r e n e", preprint (1993).
- [10] N . K o g a and K . M o r o k u m a , Chem . Phys. Letters 196, 191 (1992).
- [11] J . C . R . F a u l h a b e r , D . Y . K . K o , and P . R . B r i d d o n , Phys. Rev. B 48, 661 (1993).
- [12] F . N e g r i , G . O r l a n d i and F . Z e r b e t t o , Chem . Phys. Lett. 144, 31 (1988).
- [13] M . C . M . O ' B r i e n and C . C . C h a n c e y , Am . J . Phys. 61, 688 (1993).

- [14] J.B. Bersuker, *The Jahn-Teller Effect and Vibronic Interactions in Modern Chemistry*, Plenum Press, (1984).
- [15] P. Ring and P. Schuck, *The Nuclear Many-Body Problem*, Springer-Verlag, (1980).
- [16] J.M. Eisenberg and W. Greiner, "Nuclear Theory, Microscopic Theory of the Nucleus", North Holland, Amsterdam (1972), Vol. III, see pp. 287-317, the treatment of a single-j shell is close to our section II.
- [17] V.Z. Kresin, *J. Supercond.* 5, 297 (1992); V.Z. Kresin, V.A. Litovchenko and A.G. Panasenko, *J. Chem. Phys.* 63, 3613 (1975).
- [18] V.P. Antropov, O. Gunnarsson and O. Jepsen, *Phys. Rev. B* 46, 13647 (1992).
- [19] G.A. Heath, J.E. McGrady, and R.L. Martin, *J. Chem. Soc., Chem. Commun.* 1272 (1992).
- [20] W.K. Fullagar, I.R. Gentle, G.A. Heath, and J.W. White, *J. Chem. Soc., Chem. Commun.* 525 (1993).
- [21] D. Dubois, K.M. Kadish, S. Flanagan and L.J. Wilson, *J. Am. Chem. Soc.* 113, 7773 (1991); *ibid.* 113, 4364 (1991).
- [22] P. Bhyrappa, P. Paul, J. Stinchcombe, P.D.W. Boyd and C.A. Reed, "Synthesis and Electronic Characterization of Discrete Buckminsterfulleride salts", to appear in *JACS*, November 1993.

TABLE CAPTIONS:

Table I: The left column is the electron number. The levels are identified by their quantum numbers and the energies are obtained by straightforward perturbation theory.

FIGURE CAPTIONS:

Fig. 1: The levels of fullerene anions taking into account the phonon-mediated coupling.

Fig. 2: The energy levels of fullerene anions taking into account a Hubbard interaction.

ELECTRONIC STRUCTURE OF
FULLERENE ANIONS

L. Bergomi and Th. Jolicœur,*

Service de Physique Théorique**
C.E. Saclay
F-91191 Gif-sur-Yvette CEDEX, France

Submitted to: Physical Review B

November 1993
PACS No: 75.10J, 75.50E.

cond-mat/9309xxx
SPHT/93-119

* C.N.R.S. Research Fellow

** Laboratoire de la Direction des Sciences de la Matière du Commissariat à l'Energie Atomique