

Strong-Coupling Behavior of Two $t - J$ Chains with Interchain Single-Electron Hopping

Guang-Ming Zhang,^{*} Shiping Feng,^{**} and Lu Yu[†]

International Center for Theoretical Physics, P.O.Box 586, 34100, Trieste, Italy.

Using the fermion-spin transformation to implement spin-charge separation of constrained electrons, a model of two $t - J$ chains with interchain single-electron hopping is studied by abelian bosonization. After spin-charge decoupling the charge dynamics can be trivially solved, while the spin dynamics is determined by a strong-coupling fixed point where the correlation functions can be calculated explicitly. This is a generalization of the Luther-Emery line for two-coupled $t - J$ chains. The interchain single-electron hopping changes the asymptotic behavior of the interchain spin-spin correlation functions and the electron Green function, but their exponents are independent of the coupling strength.

PACS numbers: 75.10.Lp, 75.10.Jm

An important issue of current interest is whether the peculiar properties of one-dimensional (1D) Luttinger liquid (LL) [1], [2] will survive in two- and three-dimensions. The renormalization group (RG) studies seemed to indicate an instability of LL behavior with respect to the interchain single-electron hopping (SEH) t_{\perp} [3]. However, Anderson suggested that *one should treat the intrachain correlations exactly including the spin-charge separation before switching on SEH*. Using the asymptotic Green functions in 1D for a finite Hubbard U , he argued that SEH is an irrelevant variable and named this property as "confinement" of the 1D Hubbard model [4]. His idea has stimulated several further studies [5]–[10], most of which did not confirm his conjecture in the strict sense. SEH is indeed renormalized to zero, but e-e or e-h pair hopping is generated, which drives the coupled chains towards a strong-coupling fixed point corresponding to superconducting or density-wave states. In fact, this type of instability was studied earlier in connection with organic superconductors [11]. Nevertheless, this result is not convincing because the validity of the perturbative RG at strong-coupling fixed point with large U is questionable, as in the single-impurity Kondo problem. The Kondo physics is determined by the Wilson strong-coupling fixed point [12]. The poor-man's scaling [13] correctly directs the RG flow towards it, but the calculation can not be justified by itself [14].

In this paper, we consider two coupled $t - J$ chains, using a fermion-spin transformation, proposed recently by Feng *et al.* [15], where the charge degrees of freedom are described by spinless fermions, while the spin degrees of freedom are represented by *hard-core* bosons, which in turn, can be expressed as another type of spinless fermions via Jordan-Wigner transformation. The on-site local constraint for single occupancy is satisfied even in the mean-field approximation (MFA) and the sum rule for physical electrons is obeyed. We combine this transformation with the abelian bosonization technique [1], [16] to consider the effect of SEH on the correlation functions. After spin-charge decoupling the charge dynamics can be solved trivially, while the spin dynamics can be mapped into noninteracting spinless fermions. This strong-coupling fixed point is similar to the Luther-Emery line of the single chain problem with back scattering [17]. We confirm that the spin-charge separation by itself

does not produce Anderson confinement [8], [9]. Moreover, SEH changes the asymptotic behavior of the interchain spin-spin correlation functions and the electron Green function, but their exponents are independent of the coupling strength t_\perp .

We consider two coupled $t - J$ chains

$$H = -t_\parallel \sum_{i,\sigma} (C_{1,i,\sigma}^\dagger C_{1,i+1,\sigma} + C_{2,i,\sigma}^\dagger C_{2,i+1,\sigma} + h.c.) - \mu \sum_{i,\sigma} (C_{1,i,\sigma}^\dagger C_{1,i,\sigma} + C_{2,i,\sigma}^\dagger C_{2,i,\sigma}) \\ + 2J \sum_i (\vec{S}_{1,i} \vec{S}_{1,i+1} + \vec{S}_{2,i} \vec{S}_{2,i+1}) - t_\perp \sum_{i,\sigma} (C_{1,i,\sigma}^\dagger C_{2,i,\sigma} + h.c.), \quad (1)$$

with local constraint $\sum_\sigma C_{i,\sigma}^\dagger C_{i,\sigma} \leq 1$. Here $C_{1,i,\sigma}^\dagger (C_{2,i,\sigma}^\dagger)$ creates an electron with spin σ at site i on chain 1 (2), and $\vec{S}_{1,i} (\vec{S}_{2,i})$ is the corresponding electron spin operator; t_\parallel is the intrachain hopping and μ is the chemical potential. The fermion-spin transformation of constrained electrons [15]

$$C_{i,\uparrow} = P_i a_i S_i^- P_i^\dagger, \quad C_{i,\downarrow} = P_i a_i S_i^+ P_i^\dagger$$

can implement the spin-charge separation without additional constraints. Here a_i and a_i^\dagger are "holon" (or "electron" in the particle representation) operators, represented by spinless fermions. S_i^\pm and S_i^z are spinons or pseudo-spin operators represented by CP^1 *hard-core* bosons, different from the electron spin operators in Eq.(1). P is a projection operator removing the extra degrees of freedom in the CP^1 representation. The anticommutation relations for constrained fermions $C_{i,\sigma}$ are strictly preserved. Moreover, the local constraint is satisfied exactly. However, the projection operator P is cumbersome to handle and in many cases, for example, MFA, we can drop it, with very good results [15].

To establish notations, consider first a single $t - J$ chain

$$H_1 = -t_\parallel \sum_i (a_i^\dagger a_{i+1} + h.c.) (S_i^+ S_{i+1}^- + h.c.) - \mu \sum_i a_i^\dagger a_i \\ + 2J \sum_i (a_i^\dagger a_i) \vec{S}_i \vec{S}_{i+1} (a_{i+1}^\dagger a_{i+1}). \quad (2)$$

As shown in [15], using the above fermion-spin representation, the Jordan-Wigner transformation $S_i^+ = f_i^\dagger e^{i\pi \sum_{l<i} f_l^\dagger f_l}$, $S_i^- = (S_i^+)^+$, $S_i^z = f_i^\dagger f_i - \frac{1}{2}$, and MFA, one finds the ground state energy and gapless spinon and holon spectra, in good agreement with the exact solution [18].

However, to obtain correct exponents for correlation functions, one has to go beyond the MFA, taking into account holon-spinon interactions. Following Weng *et al.* [19], this can be done by "squeezing out" holes from the spin chain, *i.e.*, to replace $a_i^\dagger a_{i+1}(f_i^\dagger f_{i+1} + f_i f_{i+1}^\dagger)$ by $a_i^\dagger a_{i+1}$ wherever there is a hole at site i and introducing the "string operators" which in our case are given by [15]

$$C_{i,\uparrow} = [a_i e^{i\pi(N - \sum_{l>i} a_l^\dagger a_l)}][f_i e^{-i\pi \sum_{l<i} a_l^\dagger a_l}],$$

$$C_{i,\downarrow} = [a_i e^{i\pi(N + \sum_{l>i} a_l^\dagger a_l)}][f_i^\dagger e^{i\pi \sum_{l<i} a_l^\dagger a_l}].$$

In the resulting Hamiltonian, the "holon" part is free and can be easily bosonized, while the spinon part is an antiferromagnetic Heisenberg spin-1/2 chain, which can also be bosonized and reduced to a standard 1+1 quantum sine-Gordon (SG) model [20]

$$H_{1,s} = \int dx \left(\frac{v_s K_s}{2} \Pi^2 + \frac{v_s}{2K_s} (\nabla \varphi)^2 - \frac{2v_s K_s^2 k_F^2}{(2\pi\alpha)^2} \cos \sqrt{16\pi} \varphi \right), \quad (3)$$

where α is an ultraviolet cut-off, while the boson field φ describes the low-energy excitations of spinons, Π is its conjugate momentum with a commutation relation $[\varphi(x), \Pi(x')] = i\delta(x - x')$. The spinon velocity is $v_s = 2J \left[(1 - \delta)^2 - \left(\frac{\sin \delta \pi}{\pi} \right)^2 \right] \sqrt{1 + \frac{4}{\pi}}$, with δ as doping concentration. The parameter determining the exponent of the spinon correlation function is $K_s = (1 + \frac{4}{\pi})^{-1/2}$, which should be independent of δ , and our result for K_s is slightly away from the exact value derived for half-filling [20], [21]. In principle, the abelian bosonization is exact only at $J_z/J_\perp \approx 0$ for the Heisenberg spin-1/2 chain [16]. However, the exact Bethe-ansatz solution does not show any singularities for $-1 < J_z/J_\perp < 1$, and the isotropic antiferromagnetic coupling $J_z/J_\perp = 1$ is described by this fixed point [21]. On the other hand, the 1+1 SG model with $A \cos \beta \varphi$ has only one weak-coupling fixed point for small A at $\beta^2 = 8\pi$ [22]. Thus we can associate the cosine interaction (3) with this fixed point of the SG model in order to rectify K_s to be 1/2 after rescaling $\Pi \rightarrow \sqrt{K_s} \Pi$, $\varphi \rightarrow \frac{\varphi}{\sqrt{K_s}}$. Since the fixed point of SG Hamiltonian under RG for $\beta \geq 8\pi$ corresponds to the vanishing of the cosine term [22], we can easily calculate the asymptotic behavior of the spin-spin correlation functions and the electron Green functions, in good agreement with exact results [18], *e.g.*,

$$S(x_i - x_j, t) \sim \frac{\cos 2k_F(x_i - x_j)}{[(x_i - x_j)^2 - (v_h t)^2]^{\frac{1}{4}} [(x_i - x_j)^2 - (v_s t)^2]^{\frac{1}{2}}}, \quad (4)$$

$$\begin{aligned} & \langle TC_{i,\sigma}(t) C_{j,\sigma}^\dagger(0) \rangle \\ & \sim \frac{e^{ik_F(x_i - x_j)}}{[(x_i - x_j)^2 - (v_h t)^2]^{\frac{1}{16}} [(x_i - x_j) - (v_s t)]^{\frac{1}{2}} [(x_i - x_j) - (v_h t)]^{\frac{1}{2}}}, \end{aligned} \quad (5)$$

where the holon velocity is the exact value $v_h = 2t_{\parallel} \sin \delta \pi$, and $k_F = \frac{\pi}{2}(1 - \delta)$.

Now consider two coupled chains and use the MFA to decouple the interchain holon-spinon interaction. The Hamiltonian (1) is reduced to the following form $H = H_h + H_s$, and

$$\begin{aligned} H_h = & -t_{\parallel} \sum_i (a_{1,i}^\dagger a_{1,i+1} + a_{2,i}^\dagger a_{2,i+1} + h.c.) - t_{\perp} \eta_1 \sum_i (a_{1,i}^\dagger a_{2,i} + h.c.) \\ & - \mu \sum_i (a_{1,i}^\dagger a_{1,i} + a_{2,i}^\dagger a_{2,i}), \end{aligned} \quad (6)$$

$$H_s = 2J^{eff} \sum_i (\vec{S}_{1,i} \vec{S}_{1,i+1} + \vec{S}_{2,i} \vec{S}_{2,i+1}) - t_{\perp} \eta_2 \sum_i (S_{1,i}^+ S_{2,i}^- + h.c.), \quad (7)$$

where we have defined two MF order parameters η_1 and η_2 .

The holon Hamiltonian is trivially diagonalized by introducing $A_k = \frac{1}{\sqrt{2}}(a_{1,k} + a_{2,k})$ and $B_k = -\frac{1}{\sqrt{2}}(a_{1,k} - a_{2,k})$ with excitation energies $\varepsilon_k^A = -2t_{\parallel} \cos k - t_{\perp}$ and $\varepsilon_k^B = -2t_{\parallel} \cos k + t_{\perp}$, respectively, where we assume $\eta_1 \approx 1$, as will be confirmed later. The SEH splits the original holon excitation spectrum by $2t_{\perp}$ and in the low doping case for a finite value $t_{\perp} > t_{\parallel}(1 - \cos 2\delta \pi)$, only the upper band has vacancies and the lower band is fully occupied. The above condition on t_{\perp} is usually satisfied. Thus, it is easy to find the self-consistent value $\eta_2 = -(1 - \delta)$, as well as the interchain holon correlation functions using the abelian bosonization technique [1]

$$\langle e^{\mp i\pi \sum_{l < i} a_{1,l}^\dagger(t) a_{1,l}(t)} e^{\pm i\pi \sum_{l < j} a_{2,l}^\dagger(0) a_{2,l}(0)} \rangle \sim [(x_i - x'_j)^2 - (v_h t)^2]^{-\frac{1}{16}}, \quad (8)$$

$$\begin{aligned} & \langle e^{-i\frac{\pi}{2} \sum_{l < i} a_{1,l}^\dagger(t) a_{1,l}(t)} a_{1,i}(t) a_{2,j}^\dagger(0) e^{i\frac{\pi}{2} \sum_{l < j} a_{2,l}^\dagger(0) a_{2,l}(0)} \rangle \\ & \sim \frac{e^{ik_F^B(x_i - x'_j)}}{[(x_i - x'_j)^2 - (v_h t)^2]^{\frac{1}{64}} [(x_i - x'_j) - (v_h t)]^{\frac{1}{2}}}, \end{aligned} \quad (9)$$

where $k_F^B = k_F - \frac{t_\perp}{v_h}$. Due to the single occupancy constraint, the situation here is simpler than the weak-coupling case where both bands have to be considered [5]- [10].

The spinon part can be reduced to the following form:

$$H_s = \int dx \left[\frac{v_s K_s}{2} \Pi_1^2 + \frac{v_s}{2K_s} (\nabla \varphi_1)^2 + \frac{v_s K_s}{2} \Pi_2^2 + \frac{v_s}{2K_s} (\nabla \varphi_2)^2 + \frac{(1-\delta)t_\perp}{(\pi\alpha)^2} \cos(\sqrt{\pi}(\tilde{\varphi}_1 - \tilde{\varphi}_2)) \right], \quad (10)$$

where $\tilde{\varphi}(x)$ is the dual field of $\varphi(x)$, and is defined by $\frac{\partial \tilde{\varphi}}{\partial x} = \Pi$ and $\tilde{\Pi} = -\frac{\partial \varphi}{\partial x}$. Note the difference of (10) from (3), where no dual fields are involved. Introducing symmetric and antisymmetric combinations of spinon fields: $\varphi_S = \frac{1}{\sqrt{2}}(\varphi_1 + \varphi_2)$, $\varphi_A = \frac{1}{\sqrt{2}}(\varphi_1 - \varphi_2)$, (10) can be rewritten as: $H_s = H_s^S + H_s^A$, where

$$H_s^S = \int dx \left(\frac{v_s K_s}{2} \Pi_S^2 + \frac{v_s}{2K_s} (\nabla \varphi_S)^2 \right), \quad (11)$$

$$H_s^A = \int dx \left(\frac{v_s K_s}{2} \Pi_A^2 + \frac{v_s}{2K_s} (\nabla \varphi_A)^2 + \frac{(1-\delta)t_\perp}{(\pi\alpha)^2} \cos(\sqrt{2\pi} \tilde{\varphi}_A) \right). \quad (12)$$

The symmetric part H_s^S is a LL with the same parameters v_s and K_s as for a single chain. As for the antisymmetric part, a rescaling and use of self-duality of the non-interacting part lead to the following 1+1 quantum SG model

$$H_s^A = \int dx \left(\frac{v_s}{2} [\tilde{\Pi}_A^2 + (\nabla \tilde{\varphi}_A)^2] + \frac{(1-\delta)t_\perp}{(\pi\alpha)^2} \cos\left(\sqrt{\frac{2\pi}{K_s}} \tilde{\varphi}_A\right) \right). \quad (13)$$

With the corrected $K_s = \frac{1}{2}$, the coupling strength of the SG model is $\beta^2 = 4\pi$. This is nothing but a free massive Thirring model [22] with a mass gap in the excitation spectrum $\Delta_s \approx 2(1-\delta)t_\perp$. It is known that t_\perp is a relevant variable in the range $0 < \beta^2 < 8\pi$. Some years ago, Haldane [23] considered the renormalization of the Bethe ansatz equations for the massive Thirring model, equivalent to the 1+1 SG model [24]. He extracted a quantum fluctuation parameter that controlled the correlation functions of this model, and found that at $\beta^2 = 4\pi$ the renormalization of the model stops and it corresponds to a free spinless fermion field. This means that $\beta^2 = 4\pi$ is just the strong-coupling fixed point, analogous to the Toulouse limit of the single-impurity Kondo problem [25]. It is the fixed point that

controls the properties of this model in the whole region $0 < \beta^2 < 8\pi$. It is remarkable that after correcting the K_s value using the Bethe ansatz solution for the single chain, we end up *exactly at this fixed point for two coupled t - J chains*. If we did not rectify the parameter K_s in the absence of the SEH, the coupling strength of the above SG model would be at $(\beta')^2 < 4\pi$, and it should be renormalized to strong-coupling fixed point $\beta^2 = 4\pi$, while the parameter K_s is renormalized to $\frac{1}{2}$. In the end, the spinon correlation functions of the single $t - J$ chain could still have correct asymptotic behavior.

In the weak coupling approach, it was also found that the ground state of a single chain is unstable with respect to SEH and the two-coupled chains are driven to a strong-coupling fixed point corresponding to $\beta^2 = 4\pi$ with opening a gap in one of the spinon excitation spectra [9], which is equivalent to the Luther-Emery line for a single chain with back scattering. However, for a finite t_\perp , the spin and charge degrees of freedom are still coupled, and the renormalizaion process cannot be carried out in the perturbative approach. Therefore, it is not possible to calculate the correlation functions at this strong-coupling fixed point . On the contrary, within our strong-coupling approach, we end up exactly at this strong-coupling fixed point and the spinon excitations reduce to two modes, one is the original LL branch for the single chain, while the other is a massive free fermion branch, corresponding to the soliton gas of the quantum SG model. After convolution, the interchain spin-spin correlation function and electron Green function for two-coupled chains can be calculated as

$$S(x_i - x'_j, t) \sim \frac{\cos 2k_F(x_i - x'_j)}{[(x_i - x'_j)^2 - (v_h t)^2]^{\frac{1}{16}} [(x_i - x'_j)^2 - (v_s t)^2]^{\frac{1}{4}}}, \quad (14)$$

$$\begin{aligned} & \langle TC_{1,i,\sigma}(t) C_{2,j,\sigma}^\dagger(0) \rangle \\ & \sim \frac{e^{ik_F^B(x_i - x'_j)}}{\left[(x_i - x'_j)^2 - (v_h t)^2 \right]^{\frac{1}{64}} \left[(x_i - x'_j) - (v_s t) \right]^{\frac{1}{4}} \left[(x_i - x'_j) - (v_h t) \right]^{\frac{1}{2}}}. \end{aligned} \quad (15)$$

The parameter $\eta_1 \approx 1$, as mentioned earlier. As compared with (4) and (5) for a single $t - J$ chain, the SEH has generated new exponents, *independent of t_\perp* . The singularity is weaker due to the presence of a gap in one of the excitation branches. The "spinon"

exponent is $-\frac{1}{2}$ instead of -1 for the spin-spin correlation function, while it is $-\frac{1}{4}$ instead of $-\frac{1}{2}$ for the electron Green function. The singularity due to holons is also weakened because of the hybridization of two chains and the single occupancy constraint. The gap in the excitation spectrum for the antisymmetric spinon field $\tilde{\phi}$ leads to an exponential decay for $\langle \tilde{\phi}(x, t) \tilde{\phi}^\dagger(0, 0) \rangle$, but a constant contribution to the correlation functions for the fermionic fields.

To summarize, we have found strong-coupling fixed point controlling the behavior of two coupled $t - J$ chains and have calculated explicitly the interchain spin-spin correlation function and the electron Green function. Our work has reconfirmed some results of the previous weak-coupling studies, namely, (i) the spin-charge separation does not produce by itself Anderson confinement; (ii) the exponents of the interchain correlation functions are changed due to the presence SEH [8], [9]. However, there are significant differences between these two approaches. (i) The weak-coupling approach indicates the existence of a strong-coupling fixed point, but can not provide a valid calculation scheme at that fixed point. (ii) The spin-charge separation for a single chain in the weak-coupling sense in general does not guarantee the spin-charge separation for two coupled chains (except for a special g_4 -ology model [8]), so it is not possible to calculate explicitly the correlation functions. The situation here is similar to the single chain problem. The correlation exponents depend on the interaction strength in the weak-coupling limit, while it is independent of interaction strength in the strong-coupling limit; the spin-charge separation is valid in the sense of "almost complete factorization" of the wave function in the large- U limit [18]. Here we use the spin-charge separation in the same sense and it can be thus justified.

Acknowledgements. The authors are very grateful to Dr. X. Y. Zhang for several enlightening comments. They appreciate helpful discussions with A. Parola and E. Tosatti and thank A. Finkel'stein for sending Ref.9 prior to publication. They also acknowledge the support of ICTP for these studies.

REFERENCES

- * Permanent address: Pohl Institute of Solid State Physics, Tongji University, Shanghai 200092, P. R. China.
- ** Permanent address: Department of Physics, Beijing Normal University, Beijing 100875, P. R. China.
- † Permanent address: Institute of Theoretical Physics, Academia Sinica, P. O. Box 2735, P. R. China.
- [1] F. D. M. Haldane, Phys. Rev. Lett. **45**, 1358 (1980); J. Phys. C **14**, 2585 (1981).
- [2] P. W. Anderson, Phys. Rev. Lett. **64**, 1839 (1990); *ibid.* **65**, 2306 (1990).
- [3] X. G. Wen, Phys. Rev. B **42**, 6623 (1990); H. J. Schulz, Int. J. Mod. Phys. B **5**, 57 (1991).
- [4] P. W. Anderson, Phys. Rev. Lett. **67**, 3844 (1991).
- [5] M. Fabrizio, A. Parola, and E. Tossatti, Phys. Rev. B **46**, 3159 (1992).
- [6] C. Castellani, C. Di Castro, and W. Metzner, Phys. Rev. Lett. **69**, 1703 (1992).
- [7] F. V. Kusmartsev, A. Luther, and A. Nersesyan, JETP Lett. **55**, 724 (1992) and their preprint; see also V. M. Yakovenko, JETP Lett. **56**, 510 (1992).
- [8] M. Fabrizio and A. Parola, Phys. Rev. Lett. **70**, 226 (1993).
- [9] A. M. Finkel'stein and A. I. Larkin, Phys. Rev. B **47**, 10461 (1993).
- [10] D. Schmeltzer, preprint; Z. Y. Weng *et al.*, preprint.
- [11] See, *e.g.*, C. Bourbonnais and L. G. Caron, Int. J. Mod. Phys. B **5**, 1033 (1991) and references therein.
- [12] G. Toulouse, C. R. Acad. Sc. Paris B **268**, 1200 (1969); K. G. Wilson, Rev. Mod. Phys. **47**, 773 (1975).

- [13] P. W. Anderson, G. Yuval, and D. R. Hamann, Phys. Rev. B **1**, 4464 (1970).
- [14] P. Nozières, Proc. 15th Conf. Low Temp. Phys. Vol. 5, M. Krusius and M. Vuorio, eds (Amsterdam, North-Holland), p. 339.
- [15] Shiping Feng, Z. B. Su, and L. Yu, accepted by Phys. Rev. B (BBR508).
- [16] A. Luther and I. Peschel, Phys. Rev. B **9**, 2911 (1974); *ibid.* **12**, 3908 (1975); M. Fowler, J. Phys. C **13**, 1459 (1980).
- [17] A. Luther and V. J. Emery, Phys. Rev. Lett. **33**, 589 (1974).
- [18] M. Ogata and H. Shiba, Phys. Rev. B **41**, 2326 (1990).
- [19] Z. Y. Weng, D. N. Shen, C. S. Ting and Z. B. Su, Phys. Rev. B **45**, 7850 (1992).
- [20] M. P. den. Nijs, Phys. Rev. B **23**, 6111 (1981).
- [21] C. N. Yang and C. P. Yang, Phys. Rev. **150**, 321, 327 (1967); R. J. Baxter, Phys. Rev. Lett. **26**, 832 (1971).
- [22] S. Coleman, Phys. Rev. D **11**, 2088 (1975); D. J. Amit, Y. Y. Goldschmidt, and G. Grinstein, J. Phys. A **13**, 585 (1980).
- [23] F. D. M. Haldane, J. Phys. A **15**, 507 (1982).
- [24] H. Bergknoff and H. B. Thacker, Phys. Rev. D **15**, 3666 (1979).
- [25] A. M. Tselick and P. B. Wiegmann, Adv. Phys. **32**, 453 (1983).