Gauge Interactions and Bosonized Ferm ion Liquids

H.-J.Kwon, A.Houghton, and J.B.Marston Department of Physics, Brown University, Providence, RI 02912–1843 (January 20, 1994)

We investigate ferm ion liquids interacting with longitudinal and transverse abelian gauge elds via bosonization. In two spatial dimensions we obtain the ferm ion propagator for the specie class of a Coulom b plus Chern-Sim ons gauge action. We discuss the relevance of this result to the Halperin-Lee-Read theory of the = 1=2 Landau level and demonstrate how Kohn's theorem is satisfied.

05.30 Fk, 11.40.-q, 72.20 My, 73.40 Hm

cond-m at/9401041

The many body problem of ferm ions coupled to gauge elds is central to the physics of strongly correlated electronic system s. Gauge elds play an important role in the physics of the half-led Landau level [1] and may be important in the cuprate superconductors [2,3]. The elects of transverse gauge elds, which are not screened, are particularly interesting as perturbative expansions in the coupling constant break down [4]. Renormalization group analysis suggests that a novel xed point exists [5]. Bosonization in dimensions greater than one [6[9] is a powerful tool for understanding the elects of gauge elds on ferm ion liquids as the realm in which it is applicable, for low-energy excitations near the Ferm i surface, is precisely where these interactions are most important. Bosonization does not rely upon a Ferm i liquid form for the quasiparticle propagator; consequently non-trivial xed points are accessible. In an appropriate lim it, bosonization reduces com plicated m any-body H am iltonians to quadratic form s. In this paper we consider the general problem of longitudinal and transverse gauge elds coupled to a Ferm i gas in two dimensions. Specializing to the Chem-Sim ons theory studied by Halperin, Lee, and Read [1] (HLR) we show that a novel non-Ferm i-liquid xed point controls the low energy physics. Furtherm ore, by including the Landau param eter f₁ we autom atically recover K ohn's theorem [10]: the bare cyclotron frequency, not a renorm alized value, appears in the collective m ode spectrum.

For simplicity, we can ne our attention to the case of spinless ferm ions and spherical Ferm i surfaces and set h = 1. We begin with the bare Ham iltonian for ferm ions interacting with non-compact longitudinal and transverse gauge elds A (q). In the Coulomb gauge, r = A = 0 it is:

$$H = \int_{-\infty}^{Z} d^{D} x \frac{1}{2m} c^{Y}(x) (ir A)^{2} A_{0} c(x) + \frac{1}{2} \int_{-\infty}^{Z} d^{D} x d^{D} y V(x y) c^{Y}(x) c^{Y}(y) c(y) c(x)$$
(1)

where $k = k^2 = 2m$ for the case of a Ferm i gas and V (x y) represents C oulom b or other two-body interactions. Next we integrate out the high-energy Ferm i degrees of freedom with the use of the renorm alization group [11]. The resulting low-energy elective theory is expressed in terms of quasiparticles k which obey canonical anticom mutation relations and which are related to the bare ferm ion operators via the wavefunction renorm alization factor Z_k :

$$_{k} = Z_{k}^{1=2} C_{k}$$
(2)

for m om enta k restricted to a narrow shell of thickness around the Ferm i surface: $k_F = 2 < j_k j < k_F + 2$. The factor Z_k rescales the discontinuity in the quasiparticle occupancy at the Ferm i surface back to one. The operation of integrating out the high energy elds can be written explicitly by introducing the partition function:

$$Z = DA D D 6 e^{iS[A;;6]}$$
(3)

Here, 6 are the high-energy Ferm i elds that are integrated out rst. Introducing the D + 1-vector notation q (!;q), the resulting low energy elective action m ay be written as:

$$S[A;] = d^{D} x dt \quad (x)(i\theta_{t} + A_{0}) \quad (x) + \frac{1}{2m} \quad (x)(r \quad iA)^{2} \quad (x) + \frac{X}{2} \quad \frac{d!}{2} \quad 6 \quad (q) A \quad (q) A \quad (q): (4)$$

In addition there are two-body ferm ion interactions which in general include both the long range C oulom b interaction and short range Ferm i liquid type interactions which we include later. There are also irrelevent operators consisting of higher powers and/or derivatives of the gauge and Ferm i degrees of freedom which may safely be neglected in the low-energy lim it [5]. The quadratic 6 term is generated by integrating out the 6 elds deep inside the Ferm i sea. It contributes to the total diam agnetic term of (f = 2m) A(q) A(q).

We now review the main aspects of bosonization and refer the reader to previous papers [7{9] for more details. In D-spatial dimensions, the Fermi quasiparticle elds may be expressed [7] in terms of the charge boson elds as:

$$(S;x;t) = \frac{p}{\frac{1}{V}} - \frac{p}{a} e^{ik_{s} \cdot x} expfi - (S;x;t)gO(S);$$
(5)

where S labels the patch [S (;) in three dimensions] on the Ferm i surface with momentum $k_{\rm S}$. V is the volume of the system which equals $L^{\rm D}$ in D-dimensions; the factor of V is introduced to maintain canonical ferm ion anticommutation relations. Both and elds are dened inside a squat box centered on S with height in the radial (energy) direction and area $^{\rm D-1}$ along the Ferm i surface. These two scales must be small in the following sense: $k_{\rm F} >> >>$. We satisfy these limits by setting $k_{\rm F}$ =N and $k_{\rm F}$ =N where 0 < <1 and N ! 1. The quantity a in the bosonization formula Eq. (5) is a real-space cuto given by a 1= . Here $^{\rm D-1}$ (L=2)^D equals the number of states in the squat box divided by . Finally, O (S) is an ordering operator introduced [7,12] to maintain Ferm i statistics in the angular direction along the Ferm i surface. (Anticommuting statistics are obeyed autom atically in the radial direction.) The charge current in a given Ferm i surface patch S is dened by:

$$J(S;q) \qquad (S;k+q) \quad (S;k) f_{k+q}^{\gamma} = k_{q;0}^{3} n_{k}g:$$
(6)

Here (S;k) = 1 if k lies inside a squat box of dimensions D^{-1} centered at S and equals zero otherwise. The charge currents obey the abelian K ac-M oody algebra [7].

We now rewrite the e ective action Eq.(4) in terms of charge currents J:

$$S[A;a] = \frac{X \quad X \quad Z}{V} \frac{d!}{2} \quad (! \quad v_{F} \quad q \quad f_{S}) \quad a \quad (S;q) \quad a(S;q) \\ + \frac{1}{V} \frac{X}{Q} \frac{d!}{2} \frac{X}{S} \quad J \quad (S;q) \quad [A_{0}(q) + v_{S} \quad A(q)] \quad \frac{f}{2m} \quad A(q) \quad A(q) :$$
(7)

Here $v_s = k_s = m$ is the renormalized Ferm i velocity vector at patch S and the charge currents are related to the canonical boson operators a and a^y by:

$$J(S;q) = \frac{p_{1}}{j\hat{n}_{S}} q_{j}[a(S;q) (\hat{n}_{q} q) + \check{a}(S; q) (\hat{n}_{S} q)]:$$
(8)

Also (x) = 1 if x > 0 and is zero otherwise and f is the full ferm ion number density. Exchange scattering, which involves high momentum gauge elds, has not been included in the action, as it is not singular in the low -momentum lim it. We also assume that no superconducting or charge-density-wave instabilities occur. The bare gauge action, which we have not yet discussed, is in general a quadratic form :

$$S_{G}^{0}[A] = \frac{1}{2}^{Z} \frac{d^{D}q}{(2)^{D}} \frac{Z}{2} K^{0}(q) A(q) A(q) :$$
(9)

Below we will specialize to the particular case of the Chem-Sim ons action.

We proceed to evaluate the exact boson correlation function. Rather than integrate out the gauge elds to obtain an elds to obtain an elds a current-current interaction, we instead introduce a generating functional for the boson correlation function and rst integrate out the boson elds a and a . In this way we avoid the intermediate step of summing an in nite perturbation series in powers of the elds elds a rate interaction to obtain the boson propagator. We construct the generating functional by coupling elds ; to the boson elds a ; a. Then, by completing the square, we integrate out the boson elds to obtain the elds to obtain the elds can be carried out to the resulting elds elds can be carried out to yield the generating functional explicitly as a function of ; .

$$Z[;] = DA DaDa \exp fi(S[A;a] + S_G^0[A])g \exp i \frac{X X Z}{S q;q \hat{n}_S > 0} \frac{d!}{2} [(S;q)a(S;q) + (S;q)a(S;q)]$$
$$= N DA \exp fi(S_{eff}[A] + S_G^0[A])g$$

$$\exp i \frac{X}{s} \frac{Z}{(2)^{p}} \left(q g\right) \frac{Z}{2} \frac{d!}{2} \frac{p}{q} \frac{f_{B}}{f_{E}} \left[(S;q) (A_{0}(q) + v_{S} A(q)) + (S;q) (A_{0}(q) + v_{S} A(q)) \right] \frac{X}{s} \frac{X}{2} \frac{Z}{2} \frac{d!}{2} \frac{1}{s} \frac{1}{s} \left[(S;q) (A_{0}(q) + v_{S} A(q)) + (S;q) (S;q) + (S;q) (S;q) \right]$$

$$+ (S;q) (A_{0}(q) + v_{S} A(q)) = i \frac{X}{s} \frac{X}{q;q} \frac{d!}{f_{S}>0} \frac{d!}{2} \frac{1}{s} \frac{1}{v_{F} q} \frac{f_{S}}{f_{S}} \left[(S;q) (S;q) (S;q) \right]$$

$$+ (S;q) (A_{0}(q) + v_{S} A(q)) = i \frac{X}{s} \frac{X}{q;q} \frac{f_{S}}{f_{S}>0} \frac{d!}{2} \frac{1}{s} \frac{1}{v_{F} q} \frac{f_{S}}{f_{S}} \left[(S;q) (S;q) (S;q) \right]$$

$$+ (S;q) (A_{0}(q) + v_{S} A(q)) = i \frac{X}{s} \frac{X}{q;q} \frac{f_{S}}{f_{S}>0} \frac{d!}{2} \frac{1}{s} \frac{1}{v_{F} q} \frac{f_{S}}{f_{S}} \left[(S;q) (S;q) (S;q) \right]$$

$$+ (S;q) (A_{0}(q) + v_{S} A(q)) = i \frac{X}{s} \frac{X}{q;q} \frac{f_{S}}{f_{S}>0} \frac{d!}{s} \frac{1}{s} \frac{1}{s} \frac{f_{S}}{f_{S}} \frac{f_{S}}{f_{S}}$$

The boson progator is obtained by di erentiating twice:

ha (S;q)
$$a^{y}(S;q)i = \frac{2 \ln Z[;]}{(S;q)}j = 0$$
: (11)

In Eq.(10), the elective action equals:

$$S_{G} [A] = S_{eff} [A] + S_{G}^{0} [A] = \frac{1}{2}^{Z} \frac{d^{D} q}{(2)^{D}}^{Z} \frac{d!}{2} K \quad (q) A \quad (q) A \quad (q);$$
(12)

where the integrand is:

$$K (q) A (q) A (q) = K^{0} (q) A (q) A (q) + {}^{0} (q) A_{0} (q) A_{0} (q) + {}^{T} (q) A (q) A (q) :$$
(13)

Here, 0 and T are the longitudinal and transverse susceptibilities given in two spatial dimensions by:

$${}^{0}(q) = N \quad (0) \quad 1 \quad \frac{(x^{2} \quad 1) \quad \dot{x} \quad j}{p \quad x^{2} \quad 1} + i \quad \frac{(1 \quad x^{2}) \quad \dot{x} \quad j}{1 \quad x^{2}}$$
(14)

and

^T (q) =
$$v_F^2 N$$
 (0) $x^2 + (x^2 1) \dot{y} \dot{y} \dot{y} \dot{x}^2 1 + i (1 x^2) \dot{y} \dot{y} \dot{j} \frac{p}{1 x^2}$ (15)

where x $\frac{!}{v_{p} \text{ jrj}}$ and N (0) = m =2 is the quasiparticle density of states at the Ferm isurface. The boson propagator can now be determined exactly from Eqs. (10, 11, 12). Expressed in terms of the gauge propagator, D (q) = K (q) ¹, \pm is:

$$ha(S;q) a^{Y}(S;q)i = \frac{i}{! v_{F}q \quad \hat{g} + i \quad sgn(!)} + i \frac{D^{-1}}{(2^{-})^{D}} q \quad \hat{g} \quad \frac{D^{-}(q;!)}{[! v_{F}q \quad \hat{g} + i \quad sgn(!)]^{2}};$$
(16)

where (S) is the D + 1-vector (1; v_s).

Next we turn to the problem of the half-led Landau level. The HLR theory [1] describes this system as a collection of quasiparticles which obey Ferm i statistics. Each quasiparticle is a composite object consisting of the physical electron plus a ux tube of quanta ~. The attachment of ux tubes to quasiparticles is natural in a Chem-Sim ons theory. The average eld strength of the ux tubes is arranged to cancel out the external magnetic eld. Thus, at the mean-eld level, the quasiparticles behave as free ferm ions in zero net eld. Consequently, the in nite Landau degeneracy is lifted and presum ably the state is stable against two-body Coulom b interactions. The important question then is whether gauge uctuations (or equivalently quasiparticle density uctuations) modify or destroy this mean-eld state. Unlike physical M axwell electrom agnetic gauge elds, the transverse component of the statistical gauge eld is not suppressed by a numerical factor of the ne structure constant multiplied by the square of the Ferm i velocity to the speed of light ratio. In the following we assume that despite strong gauge uctuations the Ferm i surface rem ains well enough de ned for our bosonic construction to apply and later verify that this assumption is correct. Experimental evidence supports the existence of a Ferm i surface [13]. The gauge elds in the HLR theory are com pact and instantons may play an important role in the electron G reen's function [14]. For simplicity we ignore these e ects here. In the C oulom b gauge the HLR e ective action is:

$$S = \int_{-\infty}^{Z} d^{2}x \, dt \quad {}^{y} (i \, \theta_{t} + A_{0}) + \frac{1}{2m} \quad {}^{y} (r \quad iA)^{2} + \frac{A_{0}}{2 \, \sim} (r \quad A)$$

$$\frac{1}{2} \frac{1}{(2 \, \sim)^{2}} \int_{-\infty}^{Z} d^{2}x \, d^{2}y \, dt [r_{x} \quad A(x)] [r_{y} \quad A(y)] V(x \quad y) : \qquad (17)$$

At lling fraction = 1=2 we set $\sim = 2$ to cancel out the external eld. The last term in Eq. (17) is the two-body Coulomb interaction, now written in terms of the gauge eld on making use of the constraint that a ux tube is

attached to each electron [1]. We should also include Ferm i liquid interactions, which if all coe cients except f₁ are set equal to zero, contribute to the action a term of the form :

$$S_{FL} = \frac{f_1}{2V k_F^2} \sum_{s, T, q}^{L} J(s;q) k_s \neq J(T;q)$$
(18)

which is made gauge covariant by the replacement [15] k = ir ! ir A G iven the action Eq. (17) the bare gauge term Eq.(9) becomes:

$$S_{G_0} = \frac{i}{4 \sim}^{Z} \frac{d^2q}{(2\gamma)^2} \frac{\frac{d!}{2}}{\frac{d!}{2}} fA_0(q) [q \quad A(q)] \quad A_0(q) [q \quad A(q)]g$$

$$\frac{1}{2} \frac{1}{(2\gamma)^2} \frac{d^2q}{(2\gamma)^2} \frac{\frac{d!}{2}}{\frac{d!}{2}} q^2 V(q) [A(q) \quad A(q)]; \qquad (19)$$

where V (q) is the Fourier transform of V (x y). For the Coulomb interaction, V (q) = $\frac{2 e^2}{jqj}$, where $e^2 = c$ and 1=137 is the ne-structure constant. In the q-lim it of ! << jq jwe nd the gauge propagator to be given by:

D (q) (S) (S) '
$$\frac{jq}{q^2} \frac{ft_S j^2}{\frac{f}{v_F jq} j} \frac{v_F^2 N (0)}{\frac{q^2}{(2^{-\gamma})^2} [1 + N (0) V (q)]}$$
: (20)

In this limit the boson self energy has an inaginary part which rejects quasiparticle damping. For the Coulomb interaction the pole is located at ! $ie^2q^2=k_F$. If, on the other hand, V (q) is a short-range interaction, the pole is at ! $iv_F jnj^2=k_F^2$ in agreement with the RPA result [1,3]. In the opposite !-limit of ! >> jnjwe obtain instead:

D (q) (S) (S)
$$'\frac{1}{2}\frac{1^2}{q^2}\frac{v_F^2 N}{1^2(2 - \frac{1}{2})^2 m^2}$$
: (21)

The poles in the !-lim it which characterize collective excitations [8] are found at j! $j = \frac{2}{m} e^{-r}$, the cyclotron frequency for a free particle of mass m in an external magnetic eld of $B = \frac{2}{e} e^{-r}$. As it stands this result violates K ohn's theorem [10]. The correct result is obtained on including the Fermi liquid interaction, Eq. (18), in the fully quantum m echanical bosonic theory. The poles are then shifted to j! $j = 2 e_{f} \left(\frac{1}{m} + \frac{f_{1}}{4}\right)^{-r} = \frac{2}{m} e^{-r}$ in accord with K ohn's theorem. The details of the proof of this result will be given elsewhere [16]. The form of Eq.(21) is similar to that found for the super long range interaction of B ares and W en [17] and therefore can be expected to make a sm all contribution to the anom alous exponent appearing in the real space ferm ion G reen's function [9]. We can estimate from Eq. (20) the ferm ion quasiparticle self-energy [8,9] by expanding the ferm ion G reen's function in powers of D ; how ever it should be noted that there is no obvious sm all parameter to justify the expansion. G iven this caveat, to rst order in the case of a short-range interaction we obtain $Im e_{f} = j! e^{2} e^{-r}$ and for the Coulom b interaction Re $_{f} = 1$ h j! jin agreement with Ref. [1].

A sthe expansion in powers of D is unreliable, it is more useful to determ ine the quasiparticle propagator directly in real space and time. We focus on the physical case of the Coulomb interaction. Given the bosonization formula Eq.(5), the real space ferm ion G reen's function is obtained by computing the Fourier transform of the boson correlation function:

$$\ln G_{f}(S;x;t) = \ln \left(\frac{a}{x \ \ f} \ v_{F} t\right) + i \frac{d^{2}q}{(2 \)^{2}} \frac{d!}{2} \left[e^{i(q \ x \ ! \ t)} \ 1\right] D \ (q;!) \ (S) \ (S) \left[! \ v_{F} \ q \ \ f + i \ sgn\left(! \)\right]^{2} : (22)$$

To further simplify the calculation we consider only the equal time propagator by setting t = 0. In this lim it the integrals appearing in Eq. (22) can be evaluated and we obtain:

$$G_{f}(S;x) = \frac{1}{\hat{n}_{S}} \frac{c^{2}}{x_{F}} \frac{\dot{n}_{S}}{k_{F}} \frac{x}{2} \hat{n}_{S} x \hat{j};$$
 (23)

where $=\frac{(2 - \tilde{r})^2}{4} \left(\frac{v_{\rm p}}{c}\right)$. The quasiparticle pole has been destroyed by the transverse gauge interactions. Apparently the half-led Landau level system is controlled by a novel non-Fermi liquid xed point. Despite the fact that the

G reen's function has a Luttinger liquid form, the quasiparticle occupancy changes abruptly at the Ferm i surface as the momentum scale for the change is set by $\frac{c}{v_{p}} \frac{2}{k_{F}} \frac{2}{v_{p}}$! 0 as ! 0. Evidently the xed point is more like a marginal Ferm i liquid than a Luttinger liquid. The existence of a discontinuity agrees with experimental observation [13]. It is also consistent with the assumed K ac-M oody current algebra, as the derivation of the algebra itself relies upon the existence of a discontinuity in the quasiparticle occupancy [7]. The novel xed point therefore is accessible via bosonization.

B osonization of ferm ion liquids in two or three spatial dimensions coupled to gauge elds yields non-perturbative insight into strongly correlated systems. The generating functional approach is a convenient way to compute the correlation function of the interacting bosons; in fact previous results obtained by an exact resum mation of a perturbative expansion for the cases of a single Ferm i liquid interaction parameter f_0 [3] and a pure C oulom b interaction [9] are reproduced directly by the longitudinal part of Eq. (16). By applying the method to the HLR theory of the half-led Landau level we identi ed the novel non-Ferm i liquid xed point that governs the low-energy physics. K ohn's theorem is satis ed autom atically.

H.-JK.thanks YoungjaiKiem and JBM.thanks P.Nelson and R.Shankar for helpful discussions. This research was supported by the National Science Foundation through grants DMR-9008239 (A H.) and DMR-9357613 (JBM.).

- [1] B. I. Halperin, P. A. Lee and N. Read, Phys. Rev. B 47, 7312 (1993).
- [2] See, for example, G. Baskaran and P.W. Anderson, Phys. Rev. B 37, 580 (1988); I.A. eck and J.B.M. arston, Phys. Rev. B 37, 3774 (1988); L.B. Io e and A. I. Larkin, Phys. Rev. B 39, 8988 (1989).
- [3] P.A. Lee and N. Nagaosa, Phys. Rev. B 46, 5621 (1992).
- [4] B.Blok and H.Monien, Phys. Rev. B 47, 3454 (1993).
- [5] J.G an and E.W ong, Phys. Rev. Lett. 71, 4226 (1993); Chetan N ayak and Frank W ilczek, \N on-Ferm iLiquid Fixed Point in 2 + 1 D im ensions," Princeton University preprint (cond-m at/9312086).
- [6] F.D.M. Haldane, \Luttinger's Theorem and Bosonization of the Ferm i Surface," Proceedings of the International School of Physics \Enrico Ferm i," Varenna 1992 (R.Schrie er and R.A.Broglia, eds.).
- [7] A.Houghton and J.B.Marston, Phys. Rev. B 48, 7790 (1993).
- [8] A. Houghton, H.-J.Kwon, and J.B. Marston, \On the Stability and Single-Particle Properties of Bosonized Ferm iLiquids," Brown University preprint (cond-mat/9310043).
- [9] A. Houghton, H.-J. Kwon, J. B. Marston and R. Shankar, \Coulom b Interaction and the Ferm i Liquid State: Solution by Bosonization," Brown University preprint (cond-m at/9312067).
- [10] W .Kohn, Phys. Rev. 123, 1242 (1961).
- [11] R. Shankar, Physica A 177, 530 (1991) and to appear in Rev. M od. Phys.
- [12] A.Luther, Phys. Rev. B, 19, 320 (1979).
- [13] R.L.W illett, R.R.Ruel, K.W. West, and L.N.Pfei er, Phys. Rev. Lett. 71, 3846 (1993)
- [14] Yong Baek K in and X iao-G ang W en, \Instantons and the spectral function of electrons in the half- led Landau level," M IT preprint (cond-m at/9401032).
- [15] Steven H. Sim on and Bertrand I. Halperin, \F inite-W avevector E lectrom agnetic Response of Fractional Quantized Hall States, "Harvard University preprint (1993), submitted to Phys. Rev. B.
- [16] H.-J.Kwon, A.Houghton, and J.B.Marston, to be published.
- [17] P.Bares and X.G.W en, Phys. Rev. B 48, 8636 (1993).