

New perturbation theory of low-dimensional quantum liquids I: the pseudoparticle operator basis

J.M.P. Carmelo^{1,2}, A.H. Castro Neto², and D.K. Campbell²

¹ Instituto de Ciencia de Materiales, C.S.I.C., Cantoblanco, SP - 28949 Madrid, Spain

² Department of Physics, University of Illinois at Urbana-Champaign,

1110 West Green Street, Urbana, Illinois 61801-3080

(Received February 1994)

We introduce a new operator algebra for the description of the low-energy physics of one-dimensional, integrable, multicomponent quantum liquids. Considering the particular case of the Hubbard chain in a constant external magnetic field and with varying chemical potential, we show that at low energy its Bethe-ansatz solution can be interpreted in terms of the new pseudoparticle operator algebra. Our algebraic approach provides a concise interpretation of and justification for several recent studies of low-energy excitations and transport which have been based on detailed analyses of specific Bethe-ansatz eigenfunctions and eigenenergies. A central point is that the exact ground state of the interacting many-electron problem is the non-interacting pseudoparticle ground state. Furthermore, in the pseudoparticle basis, the quantum problem becomes perturbative, i.e., the two-pseudoparticle forward-scattering vertices and amplitudes do not diverge, and one can define a many-pseudoparticle perturbation theory. We write the general quantum-liquid Hamiltonian in the new basis and show that the pseudoparticle-perturbation theory leads, in a natural way, to the generalized Landau-liquid approach.

PACS numbers: 64.60.Fr, 03.65.Nk, 05.70.Jk, 72.15.Nj

I. INTRODUCTION

For more than sixty years the "Bethe ansatz" (BA) [1,2] has played a central role in the analytic solution of a variety of "integrable" [3] many-body problems in condensed matter physics and quantum field theory [1,2,3,4,5]. Initially applied to determine the ground state energies and spatial eigenfunctions, BA techniques were generalized to describe excited states, thermodynamics, and correlation functions [3,5] and remain an active subject of study today. Spurred by a conjectured relationship to the microscopic mechanism for high-temperature superconductivity [6], those particular integrable interacting electron models (such as the Hubbard Hamiltonian [4]) which exhibit "Luttinger-liquid" behavior [7] have been extensively investigated [8,9,10,11,12,13,14,15].

In particular, recent investigations have established the existence of Hubbard-model eigenstates possessing off-diagonal order (a feature essential for superconductivity) [16] and have shown that, although not themselves complete, the BA eigenstates can be extended to form a complete set of states for the one-dimensional Hubbard model [17]. Importantly, both these new results have been derived using powerful algebraic techniques, which rely on the symmetries of the Hubbard Hamiltonian. A vital and to date open issue is the extent to which these or similar algebraic techniques and symmetries can be used to describe specifically those excitations that dominate response and transport at low energies in integrable quantum liquids and related models.

In the present article, which extends our earlier results [18], we show that there is a new operator algebra associated with the low-energy Hamiltonian eigenstates of integrable quantum liquids solvable by BA [1,2,4,8,13]. We show that this algebra is expressed most naturally in terms of operators describing the "pseudoparticle" excitations introduced in several previous papers [14,15,18,19,20,21,22] and proven to dominate low-energy transport and response functions.

Whereas this previous work focused on the eigenstates and eigenfunctions of the pseudoparticles, here we are able to work in terms of algebraic operators alone and hence to

obtain a much more compact and general representation. Considering the particular case of the Hubbard chain in a magnetic field and chemical potential, we demonstrate that the familiar BA solution can be interpreted naturally in terms of the pseudoparticle basis and that the Hamiltonian can be expressed simply in terms of operators in that basis. We then establish that a pseudoparticle perturbation theory, which is also naturally described in terms of the new operator algebra, can be used to study the low-energy excitations and the response and correlation functions.

Our primary goal in the present article is to establish the utility of the pseudoparticle operator algebra for expressing the Hamiltonian and for calculating in perturbation theory the low-energy excitations of the one-dimensional Hubbard and related models. In a companion paper, which we shall henceforth call II ([23]), we will apply the perturbation theory introduced here to the study of the Virasoro operator algebras of multi-component integrable quantum liquids. Papers I and II contain a detailed exposition and extension of our earlier results [18].

In Section II we introduce the Hubbard Hamiltonian in the presence of non-zero external magnetic field and chemical potential. We discuss prior results establishing that in this case the low-energy physics is dominated by a particular class of eigenstates: these lowest-weight states (LWS) of both the spin and spin algebras [16,17,24,25,26,27,28,29,30] which refer to real rapidities [15]. We call these "LWS I" to distinguish them from the LWS associated with complex, non-real, rapidities, which we call "LWS II". In the sectors of parameter space with $U(1) \times U(1)$ symmetry the LWS II excitations have an energy gap and do not contribute to the low-energy physics. The LWS I can be described simply in terms of the pseudoparticles and the corresponding algebra. We demonstrate that in the pseudoparticle basis the ground state of the many-body problem is a "non-interacting" state, i.e., a simple Slater determinant of filled pseudoparticle levels. A central point is that for all canonical ensembles and symmetries with densities $0 \leq n \leq 1$ and spin densities $0 \leq m \leq n$ the ground state is always an LWS I. We discuss the nature of the pseudoparticle Fermi sea and the low-energy excitations, showing that these excitations can be simply understood as

pseudoparticle-pseudohole pairs and that they span the full low-energy Hilbert space. We introduce momentum-density operators and show that the operator expressions for momentum, pseudoparticle number, and pseudoparticle-density fluctuations all have the expected forms.

In Section III we relate the important rapidity numbers used in the BA to the momentum-distribution operators and use this relation to write the Hamiltonian (formally) in terms of the pseudoparticle operators.

In Section IV we discuss the (potentially important) effects of normal-ordering and show that, in terms of the pseudoparticle operators, the normal-ordered Hamiltonian in the low-energy subspace has a simple form. This permits the introduction of a systematic, non-singular pseudoparticle perturbation theory. Further, it establishes directly the universal form of the Hamiltonian for integrable multicomponent quantum liquids. These two results lead, in a natural way, to the one-dimensional Landau-liquid theory studied in Refs. [14,15,18,19,20,21,22]. Our operator representation normal-ordered with respect to the ground state leads also to the results obtained from conformal field theory [13,18], as we show in detail in paper II: the energy-momentum stress tensor and the generators of the Virasoro algebras can also be written simply in the pseudoparticle operator representation.

In Section V we present a discussion and concluding remarks. We indicate how results derived for the Hubbard model can be extended to multicomponent BA solvable models [8,13,18] in which the theory has 2 independent branches of gapless elementary excitations. We contrast the perturbative character of the interaction effects in the pseudoparticle approach with the non-perturbative character of the usual electronic basis. We explain that the universal character of the present class of one-dimensional integrable quantum liquids can be understood in terms of a straightforward generalization to pseudoparticles of Wilson's renormalization group arguments [31]: the pseudo-Fermi points of the pseudoparticles replace the particle Fermi surface and close to the pseudo-Fermi points only a few two-pseudoparticle scattering processes are relevant for the low-energy physics. Finally, we indicate briefly possible implications of our results for real quasi-one dimensional materials.

and for two-dimensional quantum systems.

II. THE PSEUDOPARTICLE OPERATOR BASIS

Consider the Hamiltonian describing the Hubbard chain in an external magnetic field H and with chemical potential μ [13,15,20,21]:

$$\hat{H} = -t \sum_j c_j^\dagger c_{j+1} + c_{j+1}^\dagger c_j + U \sum_j c_{j\uparrow}^\dagger c_{j\uparrow} c_{j\downarrow}^\dagger c_{j\downarrow} + \mu \sum_j c_{j\uparrow}^\dagger c_{j\uparrow} + H \sum_j \hat{S}_z^j; \quad (1)$$

where t , U , μ , H , and \hbar are the transfer integral, the onsite "Coulomb" interaction, the chemical potential, the magnetic field, and the Bohr magneton, respectively. The operator c_j^\dagger (c_j) creates (annihilates) a spin electron at the site j , and

$$\hat{N}_z = \frac{1}{2} \sum_j \hat{N}_z^j; \quad \hat{S}_z = \frac{1}{2} \sum_j \hat{S}_z^j; \quad (2)$$

are the diagonal generators of the $SU(2)$ spin and spin algebras, respectively [16,17,29,30]. As usual, \uparrow, \downarrow refers to the spin projections $\pm \frac{1}{2}$ when used as an operator or function index and is given by $\delta_{\sigma, \uparrow}$ otherwise. In Eqs. (2)

$$\hat{N} = \sum_j c_j^\dagger c_j = \sum_{k^0} c_{k^0}^\dagger c_{k^0} = \hat{N}(0); \quad \hat{N}(k) = \sum_{k^0} c_{k^0+k}^\dagger c_{k^0}; \quad (3)$$

is the number operator for spin electrons. The operator $\hat{N}(k)$ in Eq. (3) is the one-pair electron operator associated with the spin fluctuations. Here c_k^\dagger (c_k) is the momentum Fourier representation of the operator c_j^\dagger (c_j).

The one-dimensional Hubbard Hamiltonian (1) describes an interacting quantum system of N_\uparrow down-spin electrons and N_\downarrow up-spin electrons on a chain of N_a sites with lattice constant a . Henceforth we employ units such that $a = t = \hbar = 1$. Introducing $n = N/N_a$ and $n = n_\uparrow + n_\downarrow$, we define $(k_F = \pi n)$ and the total Fermi momentum by $k_F = \pi n/2$.

In the absence of the chemical potential and magnetic field terms the Hamiltonian (1) has $SO(4) = SU(2) \times SU(2) = Z_2$ symmetry [16,17,24,25,26,28,29,30]. When N_a is even, the operator $\hat{N}_z + \hat{S}_z$ (see Eq. (2)) has only integer eigenvalues and all half-odd integer

representations of $SU(2) \times SU(2)$ are projected out [17,28]. The two $SU(2)$ algebras $\{\hat{c}_{j\sigma}, \hat{c}_{j\sigma}^\dagger\}$ and $\{\hat{c}_{j\sigma}, \hat{c}_{j\sigma}^\dagger\}$ have diagonal generators given by Eq. (2) and off-diagonal generators [17,28]

$$\hat{S}^x = \sum_j (\hat{c}_{j\sigma}^\dagger \hat{c}_{j\sigma} - \hat{c}_{j\sigma} \hat{c}_{j\sigma}^\dagger); \quad \hat{S}^y = \sum_j (\hat{c}_{j\sigma}^\dagger \hat{c}_{j\sigma}^\dagger - \hat{c}_{j\sigma} \hat{c}_{j\sigma}); \quad (4)$$

and

$$\hat{S}^z = \sum_j \hat{c}_{j\sigma}^\dagger \hat{c}_{j\sigma}; \quad \hat{S}^y = \sum_j \hat{c}_{j\sigma}^\dagger \hat{c}_{j\sigma}; \quad (5)$$

respectively.

In the presence of both the magnetic field and chemical potential terms, the symmetry is reduced to $U(1) \times U(1)$, with \hat{S}_z and \hat{S}_z commuting with \hat{H} . Hence at any fixed value of U , the parameter space of the Hamiltonian (1) is two dimensional (dimensional for the multicomponent case when the number of subalgebras is n and the symmetry is $[U(1)]^n$). According to these values, the system has different symmetries as follows [29]: when $\mu \neq 0$ and $S_z \neq 0$ the symmetry is $U(1) \times U(1)$, for $\mu = 0$ and $S_z \neq 0$ it is $SU(2) \times U(1)$, when $\mu \neq 0$ and $S_z = 0$ it is $U(1) \times SU(2)$, and at $\mu = 0$ and $S_z = 0$ the symmetry is $SO(4)$. Note that the $U(1) \times U(1)$ symmetry sector always corresponds to two non-zero eigenvalues of the diagonal generators, whereas in sectors of higher symmetry, one or both of these eigenvalues vanish.

Although the two conserved quantum numbers are usually taken (as in Eq.(3)) to be \hat{N}_σ and \hat{N}_σ , an alternative choice is \hat{N} and \hat{N}_z , i.e. $\hat{N} = \hat{N}_\sigma + \hat{N}_\sigma = \hat{c}^\dagger(0)$ and $\hat{N}_z = \hat{N}_\sigma - \hat{N}_\sigma = \hat{c}^z(0)$, where the charge and spin one-pair electron operators read

$$\hat{c}(k) = \hat{c}^\sigma(k) + \hat{c}^\sigma(k); \quad \hat{c}^z(k) = \hat{c}^\sigma(k) - \hat{c}^\sigma(k); \quad (6)$$

The eigenvalues μ and S_z can be expressed in terms of the conserved numbers, as shown by Eq.(2). This holds true also for the eigenvalues of the diagonal generators of multicomponent integrable systems. In these systems, in addition to the chemical potential and magnetic field, the system can include other external fields associated with the additional conserved quantum numbers. Note, however, that the number of fields always equals the number of types of particles of the quantum liquid.

In the above discussion we have worked in the parameter space specified by eigenvalues $(z; S_z)$, corresponding to the canonical ensemble. Equivalently, we could choose the two external fields $\{H$ and $\mu\}$ as parameters, or in fact use a "mixed" representation. We remark that in the papers [14,15,20,21] a mixed representation involving the magnetic field H and the density n (i.e. number of particles N) was often used.

Historically, most investigations of the Hubbard chain and related models have considered the higher-symmetry sectors. For instance, in the case of the Hubbard model studies, have typically focused on the zero magnetic field ($H = 0$) case of symmetry $U(1) \times SU(2)$ (or $SO(4)$). By considering the critical point of the one-dimensional Hubbard model in a magnetic field, Frahm and Korepin [13] were able to discover interesting new features that remained hidden in previous studies of the model at zero magnetic field. For instance, in contrast to the $H = 0$ case, where the spinon gapless excitations are LWS II, at $H > 0$ (and $n \notin 1$) all gapless Hamiltonian eigenstates are LWS I and the form of the corresponding BA equations is much simpler. (The BA equations associated with the LWS II are in large number and very complicated { see, for example, Refs. [17,28] } whereas the BA equations which describe the LWS I are only two and have the same structure as the ground-state equations of Lieb and Wu [4,13,15].) However, as discussed in detail in Ref. [15], the zero magnetic-field limit of the $H > 0$ low-energy expressions provides the correct $H = 0$ exponents [13] and low-energy quantities [15]. Similarly, several papers focusing on the lowest symmetry region of parameter space have developed a pseudoparticle description and studied its relation to the Landau-liquid properties of the low-energy physics [14,15,18,19,20,21,22,23].

We consider here the $U(1) \times U(1)$ sector which refers to electronic densities $0 < n < 1$ and spin densities $0 < m < n$. This sector contains the "regular" BA states which are LWS of both algebras [17]. We concentrate our attention on the LWS I. In this $U(1) \times U(1)$ sector $SO(4)$ multiplets are generated by acting with the raising operators (4) and (5) on the corresponding LWS (for both LWS I and II). The total number of states is 4^{N_a} [17].

It is important to reiterate that our focus on the $U(1) \times U(1)$ symmetry sector, in which the gapless BA Hamiltonian eigenstates are easier to describe, does not prevent us

from obtaining relevant information about the sectors of higher symmetry. This is because, as mentioned above, the limits of the expressions derived for the physical quantities in the lowest-symmetry parameter space generally approach the correct values for the same quantities in the sectors of higher symmetry [15]. When the transition from the sector of parameter space of lowest symmetry to the higher-symmetry sector does not involve the opening of a gap in one of the two elementary excitation branches, the values in the higher symmetry sector are obtained by taking in the expressions obtained in the $U(1) \times U(1)$ sector the limiting values for the eigenvalues of the diagonal generators which characterize the higher-symmetry sectors. For example, taking the limit $S_z \rightarrow 0$ in the expressions for the physical quantities derived in the present $U(1) \times U(1)$ sector leads to the correct $S_z = 0$ expressions of the $U(1) \times SU(2)$ (or $SO(4)$) sector. Although the metal-insulator transition can cause some subtleties [20,21,29] in determining the $n = 1$ expressions { corresponding to the symmetry $SU(2) \times U(1)$ (or $SO(4)$) } these can also be obtained from the $S_z \rightarrow 0$ values.

As noted above, in the general component case, in the sectors of lowest symmetry [$U(1)$] the number of $U(1)$ subalgebras also provides the number of independent gapless branches of elementary excitations [18,29]. As we discuss below, each of these branches is characterized by a different "color" quantum number, which we shall call λ . These colors label the pseudoparticle operators and define "new" orthogonal directions in the low-energy Hilbert space. Furthermore, the numbers of pseudoparticles and pseudoholes are good quantum numbers. Therefore, at constant values of the particle-particle interaction U , we may use the numbers of pseudoparticles or the numbers of pseudoholes as alternatives to the particle numbers N associated with the operators (3), to define the parameter space.

We return now to the specific case of the Hubbard model and study in detail the operator algebra introduced in Ref. [18], which generates the Hilbert sub-space spanned by the regular BA eigenstates described by real rapidities (LWS I).

A central point is that in canonical ensembles of the $U(1) \times U(1)$ sector both the non-LWS multiplets and the LWS II have energy gaps relative to the ground state [15,18,29].

Therefore, for energies smaller than those gaps, the Hilbert subspace spanned by the regular BA LWSI coincides with the full Hilbert space of the quantum problem. The perturbation theory which we introduce in Sec. IV refers to that Hilbert space corresponding to energy scales smaller than the above gaps. We note, however, that LWSI contain both high- and low-energy states. In this section we study the entire Hilbert subspace spanned by the LWSI, but stress that it coincides with relevant, low-energy Hilbert space of the quantum problem only at energy scales smaller than the above gaps.

The Hilbert subspace spanned by the LWSI can be generated by acting on the vacuum state with the pseudoparticle algebra we introduce below. The LWSI pseudoparticles are massless in the sense of field theory (describe gapless modes); of course, they have finite static and transport "condensed-matter" masses [15,18], which are given in II (for the transport masses see Eq. (71) of II.) For completeness we note that a more general Landau-liquid theory including the LWSI states can be constructed [32]. These LWSI can also be described by pseudoparticles, but these new pseudoparticles have a mass in the field-theory sense (gap, in the present condensed-matter language).

Although the pseudoparticles associated with the LWSI are the transport carriers at low energy [15], they refer to purely non-dissipative excitations, i.e. the Hamiltonian commutes with the currents in the subspace spanned by the LWSI [15,33]. Therefore, these pseudoparticle currents give rise only to the coherent part of the conductivity spectrum, i.e. to the Drude peak [15,18,23]. The non-coherent part is associated with the massive pseudoparticles which describe the LWSI and some of the non-LWSI multiplets [32]. In the Hilbert subspace spanned by those excitations, the Hamiltonian does not commute with the current operators. In addition, and to make life more difficult (and interesting), the excited LWSI states (18) below disappear at $n = 1$ (or $H = 0$) and the corresponding massive pseudoparticle becomes massless (U(1) to SU(2) transition) [15,32,33]. Fortunately, for most of our discussion, this subtlety will be irrelevant.

The new operator algebra studied in this paper generates all the LWSI from the electronic vacuum $| \Psi_i \rangle$. At constant values of U and $S_z < 0$, the vacuum $| \Psi_i \rangle$ corresponds to the limit of

vanishing electronic density $n = 0$. This is the $S_z < 0, U(1) \times U(1)$ vacuum which refers to the limits $n_{\uparrow} = 0$ and $n_{\downarrow} = 0$ with $n_{\uparrow} = n_{\downarrow} > 1$. We note that when this vacuum is obtained from the ground state (given in Eq. (11) below) by taking that limit, in the case of some matrix elements this does not commute with the limit $n_{\uparrow} = 0$ and $n_{\downarrow} = 0$ with $n_{\uparrow} = n_{\downarrow} = 1$, which defines the $S_z = 0, U(1) \times SU(2)$, vacuum. In turn, these two limits lead to the same $S_z = 0$ vacuum when they refer to the ground states alone (see Eq. (11) below), i.e. not to the limiting value of matrix elements involving these states. This is revealed by the study of the electron - pseudoparticle canonical transformation which will be presented elsewhere [34].

The new operator algebra involves two types of pseudoparticle creation (annihilation) operators $b_q^\dagger, (b_q)$ which obey the usual fermionic algebra [18]

$$b_q^\dagger b_q^\dagger = 0; \quad b_q b_q = 0; \quad b_q^\dagger b_q = 1; \quad b_q b_q^\dagger = 0; \quad b_q^\dagger b_q^\dagger = 0; \quad b_q b_q = 0; \quad (7)$$

Here refers to the two (or , in the general case) pseudoparticle colors c and s [15,18,29]. The discrete pseudomomentum values are

$$q_j = \frac{2}{N_a} I_j; \quad (8)$$

where I_j are consecutive integers or half integers. (In the case of the Hubbard chain we have that $I_j^c = I_j$ and $I_j^s = J$ [with $j = j$], where I_j and J refers to the notation of Lieb and Wu [4].) There are N values of I_j , i.e. $j = 1, \dots, N$. An LWSI is specified by the distribution of N occupied values, which we call pseudoparticles, over the N available values. There are $N - N$ corresponding empty values, which we call pseudoholes. We emphasize that the BA wave functions vanish for double-occupied configurations of the I_j quantum numbers [5].

In the present $U(1) \times U(1)$ sector of parameter space, the numbers of pseudoholes $N - N$, such that $N > N$, are determined by the corresponding eigenvalues of the two (or) diagonal generators. In the case of the Hubbard chain we have that

$$S_z = \frac{1}{2} [N_c - N_c]; \quad S_z = \frac{1}{2} [N_s - N_s]; \quad (9)$$

where

$$N_c = N_a; \quad N_c = N; \quad N_s = N''; \quad N_s = N_{\#} : \quad (10)$$

The numbers I_j^c are integers (or half integers) for N_s even (or odd), and I_j^s are integers (or half integers) for N_s odd (or even) [4]. Since only single and zero occupancy of the values I_j are allowed, only pseudoparticles of the color c can occupy the states labeled by the numbers I_j . Therefore, the pseudoparticles have a fermionic character, as assured by the anticommuting algebra (7). For the Hubbard model the BA spatial wave function for the LWSI depends on the quantum numbers I_j through two sets of real numbers, which many authors call rapidities. Note that for the LWSII these rapidities are complex, non-real, numbers. The expression of the spatial wave functions in terms of the quantum numbers I_j requires the solution of two systems of algebraic equations which define the two rapidities as functions of the quantum numbers I_j [2,4,13,21]. Although the expression of the spatial wave function for the LWSI in terms of the quantum numbers I_j requires the solution of the above systems of equations, which constitutes a problem of considerable complexity, the description of these eigenstates in the basis of the above BA quantum numbers I_j that diagonalize the quantum liquid is much simpler. Historically, the Hamiltonian eigenstates were introduced in terms of the spatial BA wave functions [1,2]. The diagonalization of the problem leads then to the two systems of algebraic equations. In the case of LWSI these equations introduce the integer or half-integer quantum numbers I_j which describe these eigenstates.

One of the principal advantages of the algebraic approach we use in this paper is that it permits the description of the LWSI in terms of the quantum numbers I_j and does not require the spatial wave-function representation. In the basis associated with these quantum numbers the description of the LWSI does not involve the rapidity numbers. As in the case of the spatial wave functions, the expression of the energy in terms of the quantum numbers I_j involves the rapidities. In Sec. III and Appendix A we will consider the above algebraic equations for the case of the LWSI of the Hubbard chain. The introduction of

a suitable operator representation reveals that the rapidities are the eigenvalues of rapidity operators which in the pseudoparticle basis determine the Hamiltonian interaction many-pseudoparticle terms.

The above electronic vacuum state $|0\rangle$ has no electrons. At fixed values of the onsite interaction U , the ground state of a canonical ensemble characterized by eigenvalues $\epsilon_z < 0$ and $S_z < 0$ can be constructed from that vacuum. Following Refs. [15,18,20,21,29] and as shown elsewhere [30,32], it corresponds to filling symmetrically around the origin N consecutive I_j values of all colors. In addition, the ground state of canonical ensembles of all symmetries and such that $0 \leq n \leq 1$ and $0 \leq m \leq n$ are always LWSI. This holds even at the $SO(4)$ "point", where $n = 1$ and $m = 0$, and the available Hilbert space contains only one LWSI, which is the corresponding ground state; in this case, all LWS excited states are LWSII, as we discuss below. The ground state associated with a canonical ensemble of $(\epsilon_z; S_z)$ values has the form

$$|0; \epsilon_z; S_z\rangle = \prod_{c,s} \left[\prod_{q=q_c^{(+)}}^{q_c^{(-)}} b_q^s \right] |0\rangle; \quad (11)$$

where when N is odd (even) and I_j are integers (half integers) the pseudo-Fermi points are symmetric and given by

$$q_F^{(+)} = -q_F^{(-)} = \frac{1}{N_a} [N - 1]; \quad (12)$$

On the other hand, when N is odd (even) and I_j are half integers (integers) we have that

$$q_F^{(+)} = \frac{1}{N_a} N; \quad q_F^{(-)} = \frac{1}{N_a} [N - 2]; \quad (13)$$

or

$$q_F^{(+)} = \frac{1}{N_a} [N - 2]; \quad q_F^{(-)} = \frac{1}{N_a} N; \quad (14)$$

Similar expressions are obtained for the pseudo-Brioullin zones limits $q^{(\pm)}$ if we replace in Eqs. (12)–(14) N by N' .

The simple form of the ground-state expression (11) has a deep physical meaning. It reveals that in the pseudoparticle basis the ground state of the many-electron quantum problem is a "non-interacting" pseudoparticle ground state of simple Slater-determinant form. However, that the numbers I_j of the R.H.S. of Eq. (8) can be integers or half-integers for different ground states which, for example, differ by 1 in one of the eigenvalues of the diagonal generators (9), makes the problem much more involved than a simple non-interacting case. This change in the integer or half-integer character of some of the numbers I_j of two ground states, shifts all the occupied pseudomomenta and leads to the orthogonal catastrophes [6]. This is the reason for the absence of quasiparticle peaks in the single-particle spectral function. On the other hand, if we consider the evaluation of quantities which do not change the eigenvalues of the diagonal generators (9), the integer or half-integer character of the numbers I_j is, in the thermodynamic limit, irrelevant.

In some problems it is important to distinguish the two situations $q_F^{(+)} = q_F^{(-)}$ and $q_F^{(+)} \neq q_F^{(-)}$. For instance, the total momentum expression (22) below reveals that when $q_F^{(+)} \neq q_F^{(-)}$ the pseudoparticles give a finite contribution to the momentum of the ground state (11). However, in many evaluations we can assume the simplest case $q_F^{(+)} = q_F^{(-)}$ and similarly for the pseudo-Brillouin zones limits. This is because in either case $q_F^{(+)} - q_F^{(-)} = 2q_F - 2 = N_a$ and $q_F^{(+)} - q_F^{(-)} = 2q_F - 2 = N_a$, and, in addition, except for terms of order $1/N_a$, we have that $q_F^{(+)} = q_F^{(-)} - q_F$ and $q_F^{(+)} = q_F^{(-)} + q$, where

$$q_F = \frac{N}{N_a}; \quad q = \frac{N}{N_a}; \quad (15)$$

The representation (8) was used by Yang and Yang in Ref. [35]. In the thermodynamic limit we take the continuum limit $q_j \rightarrow q$ and the set of possible pseudomomenta (8) maps into a continuum domain of q values. For the calculation of many quantities we can then replace $q_F^{(-)}$ and $q_F^{(+)}$ by the values q_F and q , respectively, given in Eq. (15). For instance, in Refs. [15,20,21] that approximation was used in the evaluation of the Landau-functional expansions because this leads to the correct values for the corresponding pseudoparticle bands and f functions. In contrast, however, when we insert in that functional particular

forms of the pseudoparticle deviations which refer to changes in the numbers N_i , the integer or half-integer character of the quantum numbers I_j plays a crucial role [13,18,23].

As in the case of Landau's Fermi liquid theory [36,37], we will use the exact ground state (11) of the quantum problem as reference state. In the pseudoparticle basis we will write all operators in normal order relative to that ground state. The corresponding universal form of the normal-ordered quantum-liquid Hamiltonian will confirm that in that basis the present quantum system is a Landau liquid [14,15,19,20,21].

It is useful to know how many LW S I and LW S II there are for given numbers n_z and S_z . The total number of LW S, i.e. of regular BA Hamiltonian eigenstates, was evaluated in Ref. [17]. The following numbers refer both to canonical ensembles of the present $U(1) \times U(1)$ sector and of the $U(1) \times SU(2)$, $SU(2) \times U(1)$, and $SO(4)$ sectors when $0 \leq n \leq 1$ and $0 \leq m \leq n$. Only in the $U(1) \times U(1)$ sector do all the LW S II states have an energy gap relative to the ground state (11). The number of LW S I and LW S II is given by

$$\begin{matrix} 2 & & 3 & 2 & 3 \\ \frac{6}{4} & N_a & \frac{7}{5} & \frac{6}{4} N_s & \frac{7}{5} \end{matrix} ; \quad (16)$$

$$N_s + N_s \quad N_s$$

and

$$\begin{matrix} 2 & & 3 & 2 & 2 & & 3 & 2 & & 3 & 2 & & 3 & 2 & & 3 & 2 & 3 \\ \frac{6}{4} N_a & \frac{7}{5} & \frac{6}{4} N_a & \frac{7}{5} + \frac{6}{4} N_a & \frac{7}{5} & \frac{6}{4} N_a & \frac{7}{5} & \frac{6}{4} N_a & \frac{7}{5} & \frac{6}{4} N_a & \frac{7}{5} & \frac{6}{4} N_a & \frac{7}{5} & \frac{6}{4} N_a & \frac{7}{5} & \frac{6}{4} N_s & \frac{7}{5} \end{matrix} ; \quad (17)$$

$$N_s \quad N_s \quad N_s \quad 2 \quad N_s + 1 \quad N_s \quad 1 \quad N_s \quad 1 \quad N_s + N_s \quad N_s$$

respectively [29]. The square brackets in the above equation refer to the usual combinatoric coefficients. Adding the numbers (16) and (17), one recovers the total number of regular BA Hamiltonian eigenstates for given values n_z and S_z [17].

Since the colors and the pseudomomentum q are the only quantum numbers involved in the description of the pseudoparticles whose distributions define the LW S I, these distributions can be generated by applying to the vacuum $|j\rangle$ the \hat{N}^P creation operators b_q^\dagger . The resulting Hamiltonian eigenstates, which are LW S I and of total number given by

(16), include the ground state (11) and the excited states which can be generated from it by pseudoparticle-pseudoholes processes and have the form

$$|j_z; S_z\rangle = \sum_{c; s} \prod_{i; j=1}^{N_{ph}} [b_{q_j}^y b_{q_i}] |0; j_z; S_z\rangle; \quad (18)$$

where q_j (q_i) denotes the different locations of the pseudoparticles (pseudoholes) relative to the reference state (11) and N_{ph} is the number of pseudoparticle-pseudohole processes. SO(4) multiplets are generated by acting the raising generators (4) and (5) of the spin and spin algebras, respectively, onto the LW S I (11) and (18). As is shown in Ref. [30], in contrast to the higher-symmetry sectors, in the present U(1) × U(1) sector we have that non-LW S multiplets with values of j_z and S_z as in the corresponding canonical ensemble have an energy gap relative to the ground state (11).

The LW S I states of the form (11) and (18) and of total number given by (16) constitute a complete orthonormal basis which spans an important Hilbert subspace, which we call H_I . At energy scales smaller than the gaps for non-LW S multiplets [30] and LW S II, H_I represents the full accessible Hilbert space. We emphasize that the generators which transform the ground state (11) into the excited eigenstates (18) are products of one-pseudoparticle operators of the form $b_{q+k}^y b_q$, where $0 < |q| < q_F^{(1)}$ and $q_F^{(1)} < |q+k| < q_F^{(1)}$.

Equations (11) and (18) reveal that in the basis associated with the pseudoparticle operators b_q^y and b_q , all Hamiltonian eigenstates of the many-electron system which are LW S I have a "non-interacting" form. Since for a given canonical ensemble of values $(j_z; S_z)$ all these excitations can be generated by successive applications of the operator $b_{q+k}^y b_q$ to the ground state (11), the expression for this operator in the usual electronic basis would provide interesting information about the nature of the states (18) from the point of view of electronic configurations: while at large $2q_F$ momenta and low energy the one-pair pseudoparticle-pseudohole Hamiltonian eigenstates can correspond to multipair electronic excitations [22,34], at small values of the momentum the operators $b_{q+k}^y b_q$ are a superposition of the two one-pair electronic (or particle) operators (4) [29,34]. Fortunately, the BA solutions are naturally expressed in terms of the pseudoparticle basis. Therefore, although

we do not know, in general, the precise form of the above generators in the electronic representation, these solutions provide the expression for the Hamiltonian and other operators in the pseudoparticle basis, and we can extract relevant information about the quantum system without describing the problem in terms of electronic configurations. An important limitation, however, is that the BA solution provides the pseudoparticle expressions for some operators only. The general problem of the electron-pseudoparticle canonical transformation will be studied elsewhere [29,34]. The fact that the BA solutions are not most naturally expressed in terms of the original electronic basis is the main reason why it has been difficult, in previous work, to extract the information concerning correlation functions and matrix elements contained in the BA solutions.

Despite the non-interacting form of the Hamiltonian eigenstates (11) and (18), we find in Sec. IV that the normal-ordered Hamiltonian includes pseudoparticle interaction terms and is, therefore, a many-pseudoparticle operator. However, these pseudoparticle interactions have a pure forward-scattering, zero-momentum transfer, character. This agrees with the Landau-liquid studies of Refs. [14,15,20,21], which established this result using eigenenergies and eigenfunctions rather than operators.

The expression of the Hamiltonian (1) in H_{Γ} involves exclusively the two ($\sigma = c; s$) pseudomomentum distribution operators

$$\hat{N}^{\sigma}(q) = b_q^{\sigma\dagger} b_q^{\sigma}; \quad (19)$$

which play a key role in the present basis. The operators (19) commute with each other, i.e. $[\hat{N}^{\sigma}(q); \hat{N}^{\sigma_0}(q^0)] = 0$. Note, however, that the two (or) pseudomomentum distribution operators (19) are not the two momentum distribution electron (or particle) operators $\hat{N}^{\sigma}(k) = c_k^{\sigma\dagger} c_k^{\sigma}$. The operator $\hat{N}^{\sigma}(q) \hat{N}^{\sigma}(k)$, which has a simple form in the pseudoparticle (electronic) basis, has an involved expression in the electronic (pseudoparticle) basis [34].

Since in the pseudoparticle basis the Hamiltonian expression involves only the operators (19), it follows that in H_{Γ} the Hamiltonian commutes with these operators. These play a central role in this Hilbert subspace because all the Hamiltonian eigenstates which are LWS

They are also eigenstates of $\hat{N}_c(q)$. These LW S I are of form (11) or (18) and, therefore, obey eigenvalue equations of the form

$$\hat{N}_c(q)j_z;S_zi = N_c(q)j_z;S_zi; \quad (20)$$

where $N_c(q)$ represents the real eigenvalues of the operators (19), which are given by 1 and 0 for occupied and empty values of q , respectively.

It follows from Eq. (9) that in H_I and in the pseudoparticle basis the diagonal generators (2) of the two $U(1)$ diagonal subalgebras become

$$\hat{N}_z = \frac{1}{2} \sum_q [\hat{N}_c(q)]; \quad \hat{S}_z = \frac{1}{2} \sum_q [\hat{N}_s(q)]; \quad (21)$$

When we take the zero limit in one or several of the eigenvalues of the diagonal generators, i.e. limit $N_z \rightarrow 0$ or (and) $S_z \rightarrow 0$, we approach a boundary line sector of parameter space of higher symmetry. Following Eqs. (9) and (21), this corresponds to taking one or several of the limits $N_c \rightarrow N_c$. If, for example, we take this limit for one of the pseudoparticle branches only, since the corresponding number of pseudoholes vanishes, Eqs. (15) and (16) reveal that the number of configurations reduces to 1. This is a filled sea of pseudoparticles similar to that of the corresponding ground state (see Eqs. (11) and (15)). In particular, when we take the limit $N_c \rightarrow N_c$ for all pseudoparticle branches, Eq.(16) gives one single configuration which is the $SO(4)$ half-integer and zero-magnetic field ground state [29]. In this case there are no excited LW S I (18) and, except for the ground state, all LW S are LW S II [28,29]. On the other hand, it also follows from Eq. (16) that taking the limit $N_c \rightarrow 0$ in all pseudoparticle branches corresponds also to a single configuration. In this case all bands are empty, i.e. full of pseudoholes. This single state is the vacuum \mathcal{V}_i , which is both the electronic and pseudoparticle vacuum.

These arguments show that, when we reach a parameter space sector of higher symmetry and/or a new phase by changing the eigenvalue of one of the diagonal generators (21), the corresponding branch of pseudoparticle-pseudohole eigenstates disappears. In the case of a phase transition, the collapsing branch is not replaced by a new gapless branch of

LW S II, whereas in the "microscopic transitions" [15,29] one of the branches of LW S II becomes gapless. In the latter case the number of gapless branches remains the same, but the symmetry is higher. In the sector of parameter space of lowest symmetry we have that the number of configurations in (16) is larger than one for all colors . Moreover, all non-LW S multiplets [30] and LW S II, the study of which we have omitted from this paper, have an energy gap [15,20,29,32].

Equation (21) gives the expression of the diagonal generators (2) in the pseudoparticle basis. Combining the operator representation that we have just introduced with the information contained in the BA solution, it is straightforward to write other simple operators in the pseudoparticle basis. The momentum \hat{P} and the number of pseudoparticle operator \hat{N} , for example, read

$$\hat{P} = \sum_q q \hat{N}(q); \quad \hat{N} = \sum_q \hat{N}(q); \quad (22)$$

respectively. These expressions were derived by combining Eqs. (10) and (20) with Eqs. (8)–(10) of Ref. [20]. Note that these operators contain non-interacting pseudoparticle terms only.

To close this section, and in order to clarify the relation of our operator representation to the Landau-liquid functional of Refs. [14,15,20,21], we discuss the eigenvalues $N(q)$ of Eq. (20). From Eq. (22) we observe that these eigenvalues obey the following normalization equation

$$N = \sum_q N(q) = \frac{N_a}{2} \int_{q^{(-)}}^{q^{(+)}} dq N(q); \quad (23)$$

For instance, in the case of the ground state (11) we have that

$$\begin{aligned} N^0(q) &= (q_F^{(+)} - q); \quad 0 < q < q_F^{(+)} \\ &= (q - q_F^{(-)}); \quad q_F^{(-)} < q < 0; \end{aligned} \quad (24)$$

where $q_F^{(-)}$ is defined in Eqs. (12)–(14) and $q_F^{(+)}$ is given by similar equations with N replaced by N^- .

Using the ground state (11) as the reference state, among all the excited eigenstates of the form (18) we will be mostly interested in low-energy excitations. These involve a redistribution of a small density of pseudoparticles relative to the distribution (24). Let us introduce the normal-ordered pseudomomentum distribution operator

$$:\hat{N}(\mathbf{q}) := \hat{N}(\mathbf{q}) - N^0(\mathbf{q}); \quad (25)$$

which is such that $\langle j_z; S_z | :\hat{N}(\mathbf{q}) : | j_z; S_z \rangle = 0$. The normal-ordered operators (25) obey the eigenvalue equations

$$:\hat{N}(\mathbf{q}) : | j_z; S_z \rangle = N(\mathbf{q}) | j_z; S_z \rangle; \quad (26)$$

where $| j_z; S_z \rangle$ denotes any Hamiltonian eigenstate of form (11) or (18). Here $N(\mathbf{q}) = \hat{N}(\mathbf{q}) - N^0(\mathbf{q})$ and, therefore, $N(\mathbf{q}) = 0$ when $| j_z; S_z \rangle = | j_z; S_z \rangle$. The eigenvalues $N(\mathbf{q})$ and $N^0(\mathbf{q})$ are nothing but the pseudomomentum distributions and deviations, respectively, of the Landau-liquid theory studied in Refs. [14,15,19,20,21]. Equations (25) and (26) imply that these are the expectation values:

$$N(\mathbf{q}) = \langle j_z; S_z | \hat{N}(\mathbf{q}) | j_z; S_z \rangle; \quad (27)$$

and

$$N^0(\mathbf{q}) = \langle j_z; S_z | :\hat{N}(\mathbf{q}) : | j_z; S_z \rangle; \quad (28)$$

respectively. Therefore, the introduction of the operator algebra (7), (11), and (18) clarifies the deep reasons for the validity of the one-dimensional Landau-liquid theory, as discussed in more detail in Sec. IV.

III. RAPIDITY OPERATORS

In this section we continue to restrict our study to H_I . As discussed above, in the case of LWSI, the BA solutions of the Hubbard model lead to two systems of algebraic equations. Given the configuration of quantum numbers that describes each Hamiltonian eigenstate of

form (11) or (18) (LW S I), these equations fully define the two sets of rapidities which determine the corresponding spatial wave function. As in the case of the quantum numbers I_j , each of these types of rapidities is associated with one of the colors. In the thermodynamic limit, we can take the pseudomomentum continuum limit $q_j \rightarrow q$ and the rapidities become functions of q which we call $R(q)$. If we combine the operator representation introduced in Sec. II with the properties of the BA solution, it is straightforward to show from the relation between the rapidities $R(q)$ and the pseudomomentum distributions (19) and (25) that these rapidity functions are nothing but real eigenvalues of rapidity operators $\hat{R}(q)$ such that

$$\hat{R}(q) |j_z; S_z\rangle = R(q) |j_z; S_z\rangle \quad (29)$$

The two rapidity operators $\hat{R}(q)$ contain all information about the many-pseudoparticle interactions of the quantum-liquid Hamiltonian. There are two fundamental properties which imply the central role that the rapidity operators of Eq. (29) have in the present quantum problem :

- (a) Each of the normal-ordered rapidity operator $:\hat{R}(q):$ can be written exclusively in terms of the two pseudomomentum distribution operators (25);
- (b) The normal-ordered Hamiltonian can be written, exclusively, in terms of the two pseudomomentum distribution operators (25), but all the corresponding many-pseudoparticle interaction terms can be written in terms of the rapidity operators $:\hat{R}(q):$. It follows that in the H_I the rapidity operators commute with the Hamiltonian.

In the thermodynamic limit the BA algebraic equations for the LW S I are replaced by a system of two coupled integral equations [2,4]. The standard treatment of the BA solutions is very lengthy, e.g., for each eigenstate we have to rewrite a new set of equations. This is because the eigenvalues $R(q)$ of the rapidity operators $\hat{R}(q)$ are different for each state $|j_z; S_z\rangle$ (18).

However, within the present operator description, we can introduce a single set of two general operator equations which apply to all eigenstates. Each of these equations defines

one of the rapidity operators $\hat{R}^{\pm}(q)$ in terms of a and b summation containing functionals of the rapidity operators and pseudomomentum-distribution operators. This system of coupled equations has a unique solution which defines the expressions for the rapidity operators in terms of the pseudomomentum-distribution operators. Although the structure of these equations leads to many universal features which we will discuss later, they involve the spectral parameters which are not universal but are specific to each model. Also the formal expression for the Hamiltonian in terms of the rapidity operators is not universal. However, the corresponding normal-ordered expressions do have a universal form for the integrable multicomponent systems, as we shall discuss in Sec. IV.

For consistency with the previous notation [15,20,21], we use for the Hubbard chain the notation $\hat{R}_c(q) = \hat{K}^+(q)$ and $\hat{R}_s(q) = \hat{S}^-(q)$. The spectral parameters are the numbers $(4t-U) \sin[K^+(q)]$ and $S^-(q)$, which appear in Appendix A and in Eqs. (30)–(32) below in operator form.

We have mentioned several times that the BA solution is most naturally expressed in the pseudoparticle basis. One reflection of this is the simple expression for the Hamiltonian (1) in that basis. Combining Eq. (29) with the energy expression (4) of Ref. [21], which refers to the LWS I, the result is

$$\hat{H} = \sum_q \hat{N}_c(q) f_{2t \cos[K^+(q)] - U} = 2g \sum_z \hat{N}_z - 2 \sum_z \hat{S}_z; \quad (30)$$

where the expressions of the diagonal generators are given in Eq. (21). This is the exact expression of the Hamiltonian (1) in H_I . At energy scales smaller than the gaps for the non-LWS multiplets and LWS II, (30) gives the exact expression of that Hamiltonian in the full Hilbert space. Despite its simple appearance, the Hamiltonian (30) describes a many-pseudoparticle problem. The reason is that the expression of the rapidity operator $\hat{K}^+(q)$ in terms of the operators $\hat{N}^{\pm}(q)$ contains many-pseudoparticle interacting terms.

To specify the Hamiltonian (30) completely, we must indicate the operator equations that define the rapidity operators in terms of the pseudoparticle momentum distribution operators. In the case of the Hubbard chain these two operator equations read

$$\hat{K}(q) = \frac{2}{N_a} \sum_{q^0} \hat{N}_s(q^0) \tan^{-1} \hat{S}(q^0) - (4t=U) \sin \hat{K}(q) \quad |j_z; S_z\rangle = q |j_z; S_z\rangle \quad (31)$$

and

$$\begin{aligned} & \frac{2}{N_a} \sum_{q^0} \left[\hat{N}_c(q^0) \tan^{-1} \hat{S}(q) - (4t=U) \sin \hat{K}(q^0) \right] \\ & \sum_{q^0} \hat{N}_s(q^0) \tan^{-1} \frac{1}{2} \hat{S}(q) - \hat{S}(q^0) \quad |j_z; S_z\rangle = q |j_z; S_z\rangle : \end{aligned} \quad (32)$$

These equations fully define the rapidity operators in terms of the pseudomomentum distribution operators (19) and (25).

Simple generalizations of these equations hold for other multicomponent quantum liquids: the many-pseudoparticle terms of the Hamiltonian can be expressed, exclusively, in terms of the rapidity operators. Moreover, the BA solutions always provide: (a) the expressions of the Hamiltonian in terms of the rapidity operators; and (b) operator equations which define (implicitly) the expressions of the rapidity operators in terms of the pseudomomentum distribution operators (19).

The forms of the eigenfunction equations (20) and (29) show that the operator equations (31) and (32) are equivalent to the BA equations presented previously [15,20,21]. Formally, these are obtained from Eqs. (31) and (32) by replacing the operators by the corresponding eigenvalues and projecting onto the various $|j_z; S_z\rangle$ states. As in the case of Eqs. (31) and (32), these equations are general and apply to all eigenstates. Their solution gives the rapidity-Landau functionals in terms of the pseudomomentum distributions (19). Insertion into these functionals of the pseudomomentum distributions of a given Hamiltonian eigenvalue $|j_z; S_z\rangle$ leads to the corresponding rapidity eigenvalue of the RHS of Eq. (29).

Unsurprisingly, it is difficult to solve the BA operator equations (31)–(32) directly and to obtain the explicit expression for the rapidity operators in terms of the pseudomomentum distribution operators (19). In contrast, it is easier to calculate their normal-ordered expression in terms of the normal-ordered operators (25). In the ensuing section, we introduce the pseudoparticle perturbation theory which leads to the normal-ordered expressions for the Hamiltonian and rapidity operators.

IV . PSEUDOPARTICLE PERTURBATION THEORY

In the pseudoparticle basis the normal-ordered rapidity operators $:\hat{R}^{\pm}(q):$ contain an infinite number of terms, as we shall demonstrate below. The first of these terms is linear in the pseudomomentum distribution operator $:\hat{N}^{\pm}(q):$ (25), whereas the remaining terms consist of products of two, three, ..., until infinity, of these operators. The number of $:\hat{N}^{\pm}(q):$ operators which appears in these products equals the order of the scattering in the corresponding rapidity term.

A remarkable property is that in the pseudoparticle basis the seemingly "non-perturbative" quantum liquids become perturbative: while excitations associated with adding or removing of one electron have zero lifetime and decay into collective pseudoparticle excitations, all low-energy Hamiltonian eigenstates of the system with one electron more or less are pseudoparticle-pseudohole states of the form (11) or (18). Furthermore, while the two-electron forward scattering amplitudes and vertices diverge, the two-pseudoparticle functions (given by Eq. (42) below) and the corresponding two-pseudoparticle forward-scattering amplitudes, which were calculated in Ref. [15], are finite. By "perturbative" we also mean here the following: since at each point of parameter space (canonical ensemble) the excited low-energy eigenstates are of form (18) and correspond to quantum-number configurations involving a density of excited pseudoparticles relative to the ground-state configuration (11), (24), the expectation values of the Hamiltonian in these states are functions of the density of excited pseudoparticles. This density is given by

$$n_{\text{ex}} = \sum_{\mathbf{q}} n_{\text{ex}}(\mathbf{q}); \quad (33)$$

where

$$n_{\text{ex}} = \frac{1}{N_a} \sum_{\mathbf{q}} [1 - N^0(\mathbf{q})] N^{\pm}(\mathbf{q}) \quad (34)$$

defines the density of excited pseudoparticles (and pseudoholes) associated with the Hamiltonian eigenstate $|j_z; S_z\rangle$.

When all the densities n_{ex} are small, we can expand the expectation values in these densities. The perturbative character of the quantum liquid rests on the fact that the evaluation of the expectation values up to the n^{th} order in the densities (33) requires considering only the corresponding operator terms of scattering orders less than or equal to n . This follows from the linearity of the density of excited pseudoparticles, which are the elementary "particles" of the quantum liquid, in $N(q) = \langle j_z; S_z j | \hat{N}(q) | j_z; S_z \rangle$ and from the form of (34). The perturbative character of the quantum liquid implies, for example, that, to second order in the density of excited pseudoparticles, the energy involves only one- and two-pseudoparticle Hamiltonian terms [18,23], as in the case of the quasiparticle terms of a Fermi-liquid energy functional [36,37].

We note parenthetically that Eqs. (33) and (34) do not apply to the excitations (A) studied in [18] and in II. In the case of these excitations, which are associated with changes in the numbers N , the densities (33) and (34) are replaced by the density of "removed" or "added" pseudoparticles. In this case the change in the integer or half-integer character of the quantum numbers must be taken into account.

The eigenvalue equations (26) imply that the problem of using the rapidity Eqs. (31) and (32) to derive the expression of the operators $\hat{R}(q)$ in terms of the operators $\hat{N}(q)$ is equivalent to the problem of evaluating the corresponding expansion of the rapidity eigenvalues $R(q)$ in terms of the pseudoparticle deviations $N(q)$ (28). This last problem, which leads to the Landau-liquid expansions, was studied in Ref. [21] for the case of the Hubbard chain. Furthermore, we emphasize that it is the perturbative character of the pseudoparticle basis which justifies the validity of these Landau-liquid deviation expansions [15,19,20,21].

Based on the connection between the two problems we can derive the expressions for the rapidity operators $\hat{R}(q)$. This corresponds to expanding the expressions of the operators $\hat{R}(q)$ in terms of increasing pseudoparticle scattering order. It is convenient to define these expressions through the operators $\hat{Q}(q)$. These are related to the rapidity operators as follows

$$:\hat{R}^0(q) := R^0(q) + :\hat{Q}^0(q) : - R^0(q); \quad (35)$$

where $R^0(q)$ is the ground-state eigenvalue of $\hat{R}^0(q)$, i.e.

$$\hat{R}^0(q) |j; z; S_z\rangle = R^0(q) |j; z; S_z\rangle. \quad (36)$$

The operators $:\hat{Q}^0(q) :$ contain the same information as the rapidity operators, and involve, exclusively, the two-pseudoparticle phase shifts $\theta_0(q; q^0)$ defined in Ref. [21]. Following the related studies of Refs. [15,20,21], in Appendix A we introduce Eq. (35) in the BA equations (31) and (32) and expand in the scattering order. This leads to

$$:\hat{Q}^0(q) := \sum_{i=1}^X \hat{Q}^{(i)}(q); \quad (37)$$

where i gives the scattering order of the operator term $\hat{Q}^{(i)}(q)$. For example, for the first-order term we find

$$\hat{Q}^{(1)}(q) = \frac{2}{N_a} \sum_{q^0, -q^0}^X \theta_0(q; q^0) : \hat{N}_0(q^0) :. \quad (38)$$

We emphasize that while the expressions for the phase shifts $\theta_0(q; q^0)$ are specific to each model because they involve the spectral parameters [21], the form of the operator term $\hat{Q}^{(1)}(q)$ (38) is universal and refers to the multicomponent quantum liquids discussed in Secs. I and V.

All the remaining higher-order operator terms of expression (37), $\hat{Q}^{(i)}(q)$, can be obtained from the rapidity equations (31) and (32). For simplicity, we provide only the expression of the first-order operators (38) and of the second-order operators, which are given in Eqs. (A 20)–(A 23) of Appendix A. In that Appendix we evaluate the expressions of these operators and use the Hamiltonian expression (30) in terms of the rapidity operators to derive the following expression for the normal-ordered Hamiltonian :

$$:\hat{H} := \sum_{i=1}^X \hat{H}^{(i)}; \quad (39)$$

where, to second pseudoparticle scattering order

$$\begin{aligned}\hat{H}^{(1)} &= \sum_{q,j} v(q) : \hat{N}^{(j)}(q) : ; \\ \hat{H}^{(2)} &= \frac{1}{N_a} \sum_{q,j} \sum_{q^0,j^0} \frac{1}{2} f_{jj^0}(q;q^0) : \hat{N}^{(j)}(q) : : \hat{N}^{(j^0)}(q^0) : : \end{aligned} \quad (40)$$

Here (40) are the Hamiltonian terms which are relevant at low energy [18,23]. Furthermore, it is shown in Refs. [18,23] that at low energy and small momentum the only relevant term is the non-interacting term $\hat{H}^{(1)}$. This property justifies the Landau-liquid character of these systems and plays a key role in the symmetries of the critical point [18,23].

The form of the normal-ordered Hamiltonian (39)–(40) is universal for the integrable multicomponent quantum liquids. On the other hand, the expressions for the pseudoparticle bands $v(q)$ involve the spectral parameters and are specific to each model. For the case of the Hubbard chain they are defined in Ref. [20]. The c and s pseudoparticle bands are shown in Figs. 7 and 8, respectively, of that reference.

All $\hat{Q}^{(i)}(q)$ terms of the RHS of Eq. (37) are such that both the f functions of the RHS of Eq. (40) and all the remaining higher order coefficients associated with the operators $\hat{H}^{(i)}$ of order $i > 1$ have universal forms in terms of the two-pseudoparticle phase shifts and pseudomomentum derivatives of the bands and coefficients of order $< i$. This follows from the fact that the S-matrix for i -pseudoparticle scattering factorizes into two-pseudoparticle scattering matrices, as in the case of the usual BA S-matrix [28,38,39,40]. For example, although the second-order term $\hat{H}^{(2)}$ of Eq. (40) involves an integral over the second-order function $\hat{Q}^{(2)}(q)$ (see Eq. (A 24) of Appendix A), this function is such that $\hat{H}^{(2)}$ can be written exclusively in terms of the first-order functions (38),

$$\hat{H}^{(2)} = \sum_{q,j} v(q) \hat{Q}^{(1)}(q) : \hat{N}^{(j)}(q) : + \frac{N_a}{2} \sum_{j=1}^X \frac{v}{2} \hat{Q}^{(1)}(jq_F)^2 ; \quad (41)$$

as shown in Appendix A, and the four (or, in the general case, X) "Landau" f functions, $f_{jj^0}(q;q^0)$, have universal forms which read

$$\begin{aligned} f_{jj^0}(q;q^0) &= 2 v(q) v(q^0) + 2 v(q^0) v(q) \\ &+ \sum_{j=1}^X \sum_{j^0=1}^X 2 v(q) v(q^0) f_{jj^0}(jq_F;q^0) ; \end{aligned} \quad (42)$$

where the pseudoparticle group velocities are given by

$$v(q) = \frac{d \epsilon(q)}{dq}; \quad (43)$$

and depend on the spectral parameters. In particular, the velocities

$$v = v(q_F); \quad (44)$$

play a determining role at the critical point, representing the "light" velocities which appear in the conformal-invariant expressions [13,18,23]. In the case of the Hubbard chain, the velocities (44) are plotted in Fig. 9 of Ref. [20].

We note that the Hamiltonian term $\hat{H}^{(1)}$ in (40) has, from the point of view of the pseudoparticle basis, a non-interacting character. However, in the electron basis, $\hat{H}^{(1)}$ is of interacting character, as revealed by the U dependence of the bands $\epsilon(q)$ [20]. Furthermore, as we mentioned above, both the two-pseudoparticle f functions (42) and forward-scattering amplitudes [15] are finite. This is in contrast to the non-perturbative electronic basis, where the two-electron forward-scattering vertices and amplitudes diverge.

Obviously, Eqs. (27) and (28) imply that the operator expressions (35) and (37)–(40) are fully equivalent to the corresponding Landau expansions already studied in Refs. [15,20,21] for the case of the Hubbard chain. Thus we have achieved one of the aims of the present paper, which is to use the pseudoparticle operator algebra to justify the validity of the Landau-liquid properties of the quantum problem already studied in these earlier papers.

We emphasize that in the present $U(1) \times U(1)$ sector of parameter space and at energy scales smaller than the gaps for the non-LWS and LWS II, Eqs. (30) and (39)–(40) refer to the expression of the full quantum-liquid Hamiltonian. In the electronic basis this is given by Eq. (1). Our operator representation leads, in a natural way, to the low-energy spectrum studied in Refs. [13,14,15,20,21]. In particular, the study of Refs. [14,15,20,21] shows that the low-energy of the Hamiltonian (39)–(40) is, in many points, similar to that of a Fermi liquid, with the pseudoparticles playing the role of the quasiparticles and the colors of the spin projections. For example, the static susceptibilities can be calculated

as in a Fermi liquid [21], and involve, exclusively, the velocities (44) and the expressions of the two-pseudoparticle phase shifts at the pseudo-Fermi points. Also the low-temperature thermodynamics can be studied as in a Fermi liquid, being described by pseudoparticle Fermi-Dirac distributions of the form [20]

$$N(q) = \frac{1}{1 + e^{(q) - k_B T}} : \quad (45)$$

As in a Fermi liquid, the low-temperature specific heat is linear in the temperature and involves the static masses $m^* = \frac{q_F}{v}$, where the pseudo-Fermi pseudomomenta and velocities are defined in Eqs. (15) and (44), respectively. The low-frequency and low-momentum dynamical properties can be studied by means of kinetic equations [14,15], again as in a Fermi liquid. This allows the evaluation of the pseudoparticles elementary currents and transport masses. The latter masses, the expressions for which differ from the above expressions for the static masses, define the low-energy dynamical separation which characterizes the multicomponent quantum liquid [15,18,23]. All this refers to the coherent part of the conductivity. The incoherent part is determined by the LWS II and non-LWS multiplets [15,32,33].

In this section we have presented the expression of the normal-ordered Hamiltonian and rapidity operators in the pseudoparticle basis of H_I . This confirms the consistency of the one-dimensional Landau-liquid theory which was shown to refer to this basis. The advantage of using the pseudoparticle basis is that the problem becomes perturbative, i.e. from the point of view of the pseudoparticle interactions it is possible to classify which scatterings are relevant. Consistent with earlier results [18] and our more detailed study in paper II, while at low energy and small momentum all pseudoparticle interactions are irrelevant, at low energy and large momentum only the two-pseudoparticle interactions are relevant. The same happens for low-energy excitations involving changes in the numbers N_i . For simplicity and for further use in paper II, we have presented in this paper only the two first Hamiltonian terms of expression (39). However, Eqs. (30)–(32) contain full information about all the remaining terms of higher-scattering order.

In the companion paper II (see also [18]), the perturbative character of the present class of quantum liquids is used to study the symmetries and algebras at the critical point. These studies confirm that the Hamiltonian (39) is the correct starting point to construct a critical-point Hamiltonian.

V. CONCLUDING REMARKS

In the previous sections we have introduced and studied a new operator algebra [18] describing the low-energy Hamiltonian eigenstates of integrable quantum liquids. Considering explicitly the case of the Hubbard chain in a magnetic field and chemical potential, we have shown that the new operators create and annihilate the pseudoparticles of the one-dimensional Landau liquid [14,15]. Our algebraic approach permits an operator analysis of the BA solutions. Further, it allows us to construct a normal-ordered Hamiltonian which has in the pseudoparticle basis a universal form involving only $k = 0$, forward-scattering pseudoparticle interaction terms. Further, the perturbative character of the pseudoparticle basis implies that the corresponding two-pseudoparticle Landau f functions and forward-scattering amplitudes are finite, in contrast to the non-perturbative electronic representation, in which the two-electron forward-scattering amplitudes and vertices diverge. To clarify further the context and implications of our results, we close with several remarks.

First, concerning the relation to previous work on pseudoparticles [14,15,19,20,21], we note that the pseudoparticle operator algebra motivates and justifies the validity of the Landau-liquid studies. These studies were based on explicit use of eigenenergies and eigenfunctions, rather than the general algebraic approach.

Second, although our explicit calculations were presented for the Hubbard model, which in the general terminology corresponds to a "two-component integrable quantum fluid", our results in fact apply to the class of BA solvable models [8,13,18], which in the largest sector of the parameter space have symmetry $[U(1)]$ [18]. As remarked previously, in this sector, the model has independent branches of gapless elementary excitations. Therefore, when $\gamma > 1$

these models are usually referred as multicomponent integrable quantum liquids [8,13,18]. Note in this sector that r represents the number of independent conserved quantum numbers and also the number of external fields (magnetic field, chemical potential, etc.) which are "conjugate" to these conserved quantum numbers in the statistical mechanics sense. Although some of these systems are described by continuum models [2], most of them are defined in an one-dimensional chain with N_a sites, $j = 1; \dots; N_a$, refer to interacting fermionic particles, and, in particular, to interacting electrons, as in the case of the Hubbard model studied in this paper.

More precisely, the class of solvable quantum liquids to which our results apply is defined by the fact that the dressed-charge matrix [8,13,15,18] is determined exclusively by the R matrix associated with the solution of the Yang-Baxter equation [8,18]. In this regard, we note that our study here, when supplemented by the results in II and Ref. [18], shows that the present operator algebra is consistent with the finite-size-correction results of Frahm and Korepin [13], who found that the complete critical theory of multicomponent integrable systems is a direct product of Virasoro algebras.

Third, the pseudoparticle algebra allows us to see immediately the origin of the "universal" character of these one-dimensional integrable quantum liquids, which can be understood as a straightforward generalization to pseudoparticles of Wilson's renormalization group arguments [31]: the pseudo-Fermi points of the pseudoparticles replace the particle Fermi points, and close to the pseudo-Fermi points only few types of two-pseudoparticle scattering processes are relevant for the low-energy physics.

Fourth, concerning the very important problem of the relation of pseudoparticles to the original electrons, we should make several remarks. As we have stressed throughout, the essential simplifying feature of the pseudoparticle basis is that the quantum problem becomes perturbative there, in the sense that it is possible to identify which pseudoparticle scatterings are relevant for the low-energy physics. For instance, in the new basis the two-pseudoparticle forward-scattering vertices and amplitudes do not diverge and one can define a many-pseudoparticle perturbation theory in which the non-interacting ground state is the

exact ground state of the many-electron problem. This is directly analogous to Landau's Fermi-liquid theory, in which the interacting ground state is used as reference state [36,37]. As is well known, the present class of quantum liquids are not Fermi liquids because the single-particle spectral function has no quasiparticle peak (i.e., $Z_F = 0$), but they do have what was previously called a "Landau-liquid" character [14,15,19,20,21]: in the pseudoparticle basis the low-energy physics is fully controlled by two-pseudoparticle forward scattering, and writing the Hamiltonian and other operators in the pseudoparticle basis allows the use of perturbation-theory techniques, as we have shown in this paper and will exploit in II.

The non-perturbative character of the usual electronic basis { e.g., the divergences of the two-electron forward-scattering vertices and amplitudes } is reflected in the complex and exotic properties of the electron-pseudoparticle operator transformation, which we have studiously avoided in this paper. The construction of this energy, momentum, and parameter dependent nonlinear transformation is in general very involved. For instance, we know that at low energies and large momenta it maps one-pair electron operators onto multipair pseudoparticle operators, and vice versa [22]. Further, both the colors and pseudomomentum q of the pseudoparticles are not, for general values of the parameters, simply related to the usual electronic quantum numbers (i.e., charge, spin, up spin, down spin, and momentum) [29,34]. We shall return to this problem elsewhere [34] and solve it for the generators which transform the exact ground state onto the low-energy and small momentum Hamiltonian eigenstates, which we are able to write both in terms of pseudoparticle and electron operators. Importantly, as we have demonstrated in our calculations here, since the BA solution refers at low energy to the pseudoparticle operator representation, we can study the quantum problem in the corresponding basis without explicit knowledge of the complex electron-pseudoparticle transformation.

Finally, concerning applications of our ideas to other theoretical models and to problems in real materials, we note that although our present analysis refers to a class of one-dimensional integrable quantum liquids only, it has been argued that similar non-Fermi liquid behavior occurs in some sectors of parameter space of two-dimensional interacting quantum liquids

[6,41,42,43,44]. If this is so, techniques similar to ours may apply. Further, it is of considerable interest to determine the extent to which the algebraic structure and pseudoparticle perturbation theory remain valid in systems which are not integrable but which behave as Luttinger (or Landau) liquids (about the relation between Luttinger and Landau liquids see Sec. V III of Ref. [15]); an example of such a system is the one-dimensional extended Hubbard model. In terms of real materials, it is known that the present one-dimensional quantum systems provide useful (albeit idealized) descriptions of the physics of quasi-one-dimensional solids [22,45,46]. At low frequency the pseudoparticles are the transport carriers of the quantum liquid and couple to external fields [14,15]. These couplings determine the exotic instabilities observed in quasi-one-dimensional synthetic metals [22].

Finally, we turn to directions for further research. In addition some specific calculations mentioned above, our immediate aims are to apply the concepts and techniques developed here to establish that the pseudoparticle perturbation theory leads to the correct effective Hamiltonian, which is obtained from the universal expression (39)–(40) of the quantum-liquid normal-ordered Hamiltonian in the pseudoparticle basis, and to study the Virasoro algebras [13,18] of conformal field theory. The initial results of these studies are described in II. The construction of a generalized Landau-liquid theory referring to the Hilbert space spanned by both the LWS I and LWS II is in progress.

ACKNOWLEDGMENTS

This work was supported principally by the C.S.I.C. [Spain] and the University of Illinois. We thank F.H.L. Essler, E.H. Fradkin, F. Guinea, P. Horsch, V.E. Korepin, A.A. Ovchinnikov, N.M.R. Peres, and K. Maki for stimulating discussions. The hospitality and support of the C.S.I.C. and U.I.U.C. are gratefully acknowledged by J.M.P.C. A.H.C.N. thanks CNPq (Brazil) for a scholarship. D.K.C. and A.H.C.N. acknowledge the partial support of this research by the U.S. National Science Foundation under grants NSF-DMR 89-20538 and NSF-DMR 91-22385, respectively.

Following the discussion of Sec. IV, the perturbative character of the system implies the equivalence between expanding in the pseudoparticle scattering order and/or in the pseudomomentum deviations (28). This, together with Eqs. (20) and (26), reveals that the problem of calculating the normal-ordered operator expansions (37) and (39) is equivalent to the Landau-energy functional studies of Refs. [15,20,21]. Since in these papers the full expression of the second-order function $Q^{(2)}(q)$ (which corresponds to the operator $\hat{Q}^{(2)}(q)$) and a number of other functions were not presented, we give in this Appendix a short description of the calculation of the normal-ordered operator expansions (37) and (39). We focus our brief study on the points which were omitted in Refs. [15,20,21].

We start by the evaluation of the first-order and second-order terms of the operator (37). For simplicity, we consider here pseudomomentum deviations and eigenvalues. Equation (26) then allows the straightforward calculation of the corresponding operator expressions.

In the thermodynamic limit, Eqs. (31) and (32) lead to the following equations

$$K(q) - \frac{1}{q_{Fs}^{(-)}} \sum_{q_{Fs}^{(+)}} N_s(q^0) \tan^{-1} S(q^0) - \frac{\sin K(q)}{u} = q \quad (A1)$$

and

$$\begin{aligned} & \frac{1}{q_{Fc}^{(-)}} \left[\sum_{q_{Fc}^{(+)}} N_c(q^0) \tan^{-1} S(q) - \frac{\sin K(q^0)}{u} \right. \\ & \left. + \sum_{q_{Fs}^{(+)}} N_s(q^0) \tan^{-1} \frac{1}{2} (S(q) - S(q^0)) \right] = q; \end{aligned} \quad (A2)$$

where we have replaced the operators by the corresponding eigenvalues and the summations by integrations and $u = U/4t$. The eigenvalue form of Eq. (35) is

$$R(q) = R^0(q + Q(q)) - R^0(q); \quad (A3)$$

where $R(q)$ and $Q(q)$ are the eigenvalues of the operators $\hat{R}(q)$ and $\hat{Q}(q)$, respectively. From Eq. (37) $Q(q)$ can be written as

$$Q(q) = Q^{(1)}(q) + Q^{(2)}(q) + \dots; \quad (A4)$$

where $Q^{(i)}(q)$ is the eigenvalue of the operator $\hat{Q}^{(i)}(q)$. Expanding the $R(q)$ expression (A 3) we find

$$R(q) = \sum_{i=0}^{\infty} R^{(i)}(q); \quad (A 5)$$

(and $R(q) = \sum_{i=1}^{\infty} R^{(i)}(q)$) where the zero-order ground-state functions $R^{(0)}(q)$ are defined by Eqs. (A 1) and (A 2) with the distributions $N(q)$ given by Eq. (24). From the resulting equations we can obtain all derivatives of the ground-state functions $R^{(0)}(q)$ with respect to q . The terms of the RHS of Eq. (A 5) involve these derivatives. For instance, the first-order and second-order terms read

$$R^{(1)}(q) = \frac{dR^{(0)}(q)}{dq} Q^{(1)}(q); \quad (A 6)$$

and

$$R^{(2)}(q) = \frac{dR^{(0)}(q)}{dq} Q^{(2)}(q) + \frac{1}{2} \frac{d^2 R^{(0)}(q)}{dq^2} [Q^{(1)}(q)]^2; \quad (A 7)$$

respectively, and involve the first and second derivatives. Let us use the notation of Eqs. (A 1) and (A 2), i.e.,

$$R_c(q) = K(q) \quad R_s(q) = S(q); \quad (A 8)$$

and

$$R_c^{(i)}(q) = K^{(i)}(q) \quad R_s^{(i)}(q) = S^{(i)}(q); \quad (A 9)$$

From Eqs. (A 1) and (A 2) (with $N(q^0)$ given by the ground-state distribution (24)) we find that the first derivatives $\frac{dR^{(0)}(q)}{dq}$ can be expressed in terms of the functions $R^{(0)}(q)$ as follows

$$\frac{dK^{(0)}(q)}{dq} = \frac{1}{1 + \frac{1}{u} \cos K^{(0)}(q) \frac{R_{q_s^{(+)}}}{q_s^{(-)}}} \frac{1}{dq^0} \frac{1}{1 + [S^{(0)}(q^0) \frac{\sin K^{(0)}(q)}{u}]^2}; \quad (A 10)$$

and

$$\frac{dS^{(0)}(q)}{dq} = \frac{1}{\frac{1}{2} \frac{R_{\mathbf{q}_c^{(+)}}}{\mathbf{q}_c^{(-)}} \frac{dq^0}{1 + \left[\frac{\mathbb{S}^{(0)}(q)}{u} \frac{\sin K^{(0)}(q^0)}{u} \right]^2} + \frac{1}{2} \frac{R_{\mathbf{q}_s^{(+)}}}{\mathbf{q}_s^{(-)}} \frac{dq^0}{1 + \left[\frac{\mathbb{S}^{(0)}(q)}{2} \frac{1}{\mathbb{S}^{(0)}(q^0)} \right]^2}} : \quad (\text{A } 11)$$

From the same equations we find that the second derivatives $\frac{d^2 R^{(0)}(q)}{dq^2}$ can be expressed in terms of the functions $R^{(0)}(q)$ and its first derivatives (A 10) and (A 11) and read

$$\begin{aligned} \frac{d^2 K^{(0)}(q)}{dq^2} = & \left[\frac{dK^{(0)}(q)}{dq} \right]^3 f - \frac{1}{u} \frac{\sin K^{(0)}(q)}{u} \frac{Z_{\mathbf{q}_s^{(+)}}}{\mathbf{q}_s^{(-)}} \frac{dq^0}{1 + \left[\frac{\mathbb{S}^{(0)}(q^0)}{u} \frac{\sin K^{(0)}(q^0)}{u} \right]^2} \\ & - \frac{2}{u} \left[\frac{\cos K^{(0)}(q)}{u} \right]^2 \frac{Z_{\mathbf{q}_s^{(+)}}}{\mathbf{q}_s^{(-)}} \frac{dq^0}{f 1 + \left[\frac{\mathbb{S}^{(0)}(q^0)}{u} \frac{\sin K^{(0)}(q^0)}{u} \right]^2} g; \end{aligned} \quad (\text{A } 12)$$

and

$$\begin{aligned} \frac{d^2 S^{(0)}(q)}{dq^2} = & \left[\frac{dS^{(0)}(q)}{dq} \right]^3 f - \frac{2}{u} \frac{Z_{\mathbf{q}_c^{(+)}}}{\mathbf{q}_c^{(-)}} \frac{dq^0}{f 1 + \left[\frac{\mathbb{S}^{(0)}(q)}{u} \frac{\sin K^{(0)}(q^0)}{u} \right]^2} \frac{\mathbb{S}^{(0)}(q)}{u} \frac{\sin K^{(0)}(q^0)}{u} \\ & + \frac{1}{2} \frac{Z_{\mathbf{q}_s^{(+)}}}{\mathbf{q}_s^{(-)}} \frac{dq^0}{f 1 + \left[\frac{\mathbb{S}^{(0)}(q)}{2} \frac{1}{\mathbb{S}^{(0)}(q^0)} \right]^2} \frac{\mathbb{S}^{(0)}(q)}{u} \frac{\sin K^{(0)}(q^0)}{u} g : \end{aligned} \quad (\text{A } 13)$$

Let us introduce the functions $\mathcal{Q}_c^{(1)}(k)$ and $\mathcal{Q}_s^{(1)}(v)$ such that

$$\mathcal{Q}_c^{(1)}(q) = \mathcal{Q}_c^{(1)}(K^{(0)}(q)); \quad \mathcal{Q}_s^{(1)}(q) = \mathcal{Q}_s^{(1)}(S^{(0)}(q)) : \quad (\text{A } 14)$$

Since we will often consider k and v integrations instead of q integrations, it is useful to define the following limiting values

$$\mathcal{Q}^{(1)} = K^{(0)}(\mathbf{q}_c^{(-)}); \quad \mathcal{B}^{(1)} = S^{(0)}(\mathbf{q}_s^{(-)}); \quad (\text{A } 15)$$

which refer to the pseudo-Fermi points (12)–(14). However, in order to be consistent with the order of the expressions below, these values are to be replaced by the corresponding leading-order terms \mathcal{Q} and \mathcal{B} , respectively, which read

$$\mathcal{Q} = K^{(0)}(\mathbf{q}_c); \quad \mathcal{B} = S^{(0)}(\mathbf{q}_s); \quad (\text{A } 16)$$

where the pseudo-Fermi points \mathbf{q}_c are defined in Eq. (15). Introducing both the distributions

$$N(q) = N^0(q) + N(q); \quad (\text{A } 17)$$

and the first-order functions (A 6) into Eqs. (A 1) and (A 2), we find after expanding to first order that the functions $Q_c^{(1)}(q)$ and $Q_s^{(1)}(q)$ are given by Eq. (A 14) with $\mathcal{Q}_c^{(1)}(k)$ and $\mathcal{Q}_s^{(1)}(v)$ defined by the following system of coupled integralequations

$$\mathcal{Q}_c^{(1)}(k) = \frac{1}{2} \int_{q_s^{(-)}}^{q_s^{(+)}} dq^0 N_s(q^0) \tan^{-1} \left(\frac{S^{(0)}(q^0)}{u} \right) + \frac{1}{2} \int_B dv^0 \frac{\mathcal{Q}_s^{(1)}(v^0)}{1 + [v^0 \frac{\sin k}{u}]^2}; \quad (\text{A } 18)$$

and

$$\begin{aligned} \mathcal{Q}_s^{(1)}(v) = & \frac{1}{2} \int_{q_c^{(-)}}^{q_c^{(+)}} dq^0 N_c(q^0) \tan^{-1} \left(v \frac{\sin K^{(0)}(q^0)}{u} \right) \\ & + \frac{1}{2} \int_{q_s^{(-)}}^{q_s^{(+)}} dq^0 N_s(q^0) \tan^{-1} \left(\frac{1}{2} [v - S_0(q^0)] \right) + \frac{1}{2} \int_Q dk^0 \frac{\cos k^0}{u} \frac{\mathcal{Q}_c^{(1)}(k^0)}{1 + [v \frac{\sin k^0}{u}]^2} \\ & + \frac{1}{2} \int_B dv^0 \frac{\mathcal{Q}_s^{(1)}(v^0)}{1 + [\frac{1}{2}(v - v^0)]^2}; \end{aligned} \quad (\text{A } 19)$$

The use of Eq. (A 18) in Eq. (A 19) allows the introduction of a single integral equation for $\mathcal{Q}_s^{(1)}(v)$. Combining the obtained equations with Eqs. (A 14) and the phase-shift equations (23)–(38) of Ref. [21] leads to expressions (38).

In order to evaluate the second-order functions $Q_c^{(2)}(q)$ and $Q_s^{(2)}(q)$ of the RHS of Eq. (A 4) we introduce the functions (A 6) and (A 7) and distributions (A 17) in Eqs. (A 1) and (A 2). Expanding to second order we find after some algebra

$$Q^{(2)}(q) = Q^{(2;-)}(q) + \frac{1}{2} \frac{d}{dq} [Q^{(1)}(q)]^2; \quad (\text{A } 20)$$

where

$$Q^{(2;-)}(q) = \mathcal{Q}^{(2;-)}(R^{(0)}(q)); \quad (\text{A } 21)$$

and the functions $\mathcal{Q}_c^{(2;-)}(k)$ and $\mathcal{Q}_s^{(2;-)}(v)$ are defined by the following system of coupled integralequations

$$\begin{aligned} \mathcal{Q}_c^{(2;-)}(k) = & \frac{1}{2} \int_{q_s^{(-)}}^{q_s^{(+)}} dq^0 N_s(q^0) \frac{dS^{(0)}(q^0)}{dq^0} \frac{Q_s^{(1)}(q^0)}{1 + [S^{(0)}(q^0) \frac{\sin k}{u}]^2} \\ & + \frac{1}{2} \left[\frac{dS^{(0)}(q)}{dq} \right]_{q=q_s} \times \sum_{j=-1}^{\infty} \frac{[Q_s^{(1)}(jq_s)]^2}{1 + [jB \frac{\sin k}{u}]^2} \\ & + \frac{1}{2} \int_B dv^0 \frac{\mathcal{Q}_s^{(2;-)}(v^0)}{1 + [v^0 \frac{\sin k}{u}]^2}; \end{aligned} \quad (\text{A } 22)$$

and

$$\begin{aligned}
\mathcal{G}_s^{(2; \cdot)}(v) = & \frac{1}{2} \int_{q_c^{(-)}}^{q_c^{(+)}} dq^0 N_c(q^0) \frac{dK^{(0)}(q^0)}{dq^0} \frac{\cos K^{(0)}(q^0)}{u} \frac{Q_c^{(1)}(q^0)}{1 + [v - \frac{\sin K^{(0)}(q^0)}{u}]^2} \\
& + \frac{1}{2} \int_{q_s^{(-)}}^{q_s^{(+)}} dq^0 N_s(q^0) \frac{dS^{(0)}(q^0)}{dq^0} \frac{Q_s^{(1)}(q^0)}{1 + [\frac{1}{2}(v - S_0(q^0))]^2} \\
& + \frac{1}{2} \left[\frac{dK^{(0)}(q)}{dq} \right]_{q=q_c} \times \frac{\cos Q}{u} \sum_{j=1}^{\infty} \frac{Q_c^{(1)}(jq_c)}{1 + [v - \frac{j \sin Q}{u}]^2} \\
& + \frac{1}{4} \left[\frac{dS^{(0)}(q)}{dq} \right]_{q=q_s} \times \sum_{j=1}^{\infty} \frac{Q_s^{(1)}(jq_s)}{1 + [\frac{1}{2}(v - jB)]^2} \\
& + \frac{1}{2} \int_Q dk^0 \frac{\cos k^0}{u} \frac{\mathcal{G}_c^{(2; \cdot)}(k^0)}{1 + [v - \frac{\sin k^0}{u}]^2} \\
& + \frac{1}{2} \int_B dv^0 \frac{\mathcal{G}_s^{(2; \cdot)}(v^0)}{1 + [\frac{1}{2}(v - v^0)]^2}; \tag{A 23}
\end{aligned}$$

respectively. Note that the free terms involve the first-order functions.

In order to derive the first-order and second-order Hamiltonian terms of Eqs. (40) and (41), we again consider eigenvalues and deviations. We introduce in the Hamiltonian expression (30) Eqs. (A 6), (A 7), and (A 17) and expand the obtained expression. Besides simpler terms, the first-order expression includes the term $\int_Q dk [2t \sin k] \mathcal{G}_c^{(1)}(k)$. After some algebra we find the first-order term of Eq. (40) with the pseudoparticle bands given by expressions (19)–(21) of Ref. [20]. Expanding (30) to second-order leads to the following energy

$$\begin{aligned}
E_2 = & \frac{N_a}{2} \int_{q_c^{(-)}}^{q_c^{(+)}} dq^0 N_c(q^0) \frac{dK^{(0)}(q^0)}{dq^0} [2t \sin K^{(0)}(q^0)] Q_c^{(1)}(q^0) \\
& + \frac{1}{2} \left[\frac{dK^{(0)}(q)}{dq} \right]_{q=q_c} [t \sin Q] \sum_{j=1}^{\infty} Q_c^{(1)}(jq_c) + \int_Q dk [2t \sin k] \mathcal{G}_c^{(2; \cdot)}(k) g; \tag{A 24}
\end{aligned}$$

such that $\hat{H}^{(2)} |j_z; S_z\rangle = E_2 |j_z; S_z\rangle$. Inserting the suitable functions in the RHS of Eq. (A 24), performing some integrations by using symmetry properties of the integral equations (A 22) and (A 23), and replacing deviations by pseudomomentum normal-ordered operators (25) we find after some algebra expression (41). This can be rewritten in terms of the f functions (42) as given in the RHS of Eq. (40).

Permanent address, Department of Physics, University of Evora, Apartado 94, P-7001 Evora codex, Portugal.

REFERENCES

- ¹ This ansatz was introduced for the case of the single-component ($\ell = 1$) isotropic Heisenberg chain by H. A. Bethe, Z. Phys. 71, 205 (1931).
- ² For one of the first generalizations of the Bethe ansatz to multicomponent ($\ell = 2$) systems see C. N. Yang, Phys. Rev. Lett. 19, 1312 (1967).
- ³ For a classical review of the Bethe ansatz and integrability in field theory and statistical mechanics see H. B. Thacker, Rev. Mod. Phys. 53, 253 (1981).
- ⁴ Elliott H. Lieb and F. Y. Wu, Phys. Rev. Lett. 20, 1445 (1968).
- ⁵ For a modern and comprehensive discussion of these issues, see V. E. Korepin, N. M. Bogoliubov, and A. G. Izergin, Quantum Inverse Scattering Method and Correlation Functions (Cambridge University Press, 1993).
- ⁶ Philip W. Anderson, Phys. Rev. Lett. 64, 1839 (1990); 65 2306 (1990); P. W. Anderson and Y. Ren, in High Temperature Superconductivity, edited by K. S. Bedell, D. E. Meltzer, D. Pines, and J. R. Schrieffer (Addison-Wesley, Reading, MA, 1990).
- ⁷ F. D. M. Haldane, J. Phys. C 14, 2585 (1981).
- ⁸ A. G. Izergin, V. E. Korepin, and N. Yu Reshetikhin, J. Phys. A: Math. Gen. 22, 2615 (1989).
- ⁹ F. W. Woytovich, J. Phys. A 22, 4243 (1989).
- ¹⁰ Alberto Parola and Sandro Sorella, Phys. Rev. Lett. 64, 1831 (1990).
- ¹¹ H. J. Schulz, Phys. Rev. Lett. 64, 2831 (1990).
- ¹² Yong Ren and P. W. Anderson, Cargèse Lectures, 1990 (unpublished).
- ¹³ Holger Frahm and V. E. Korepin, Phys. Rev. B 42, 10553 (1990); 43, 5653 (1991).
- ¹⁴ J. M. P. Carmelb and P. Horsch, Phys. Rev. Lett. 68, 871 (1992).

- ¹⁵ J.M .P .Carm elb, P .H orsch, and A .A .O vchinnikov, Phys.Rev.B 46, 14 728 (1992).
- ¹⁶ C .N .Yang, Phys.Rev.Lett. 63, 2144 (1989).
- ¹⁷ Fabian H .L .Essler, V ladim ir E .K orepin, and K areljan Schoutens, Phys.Rev.Lett. 67, 3848 (1991); Nucl.Phys.B 372, 559 (1992).
- ¹⁸ J.M .P .Carm elb and A .H .Castro Neto, Phys.Rev.Lett. 70, 1904 (1993).
- ¹⁹ J.Carm elb and A .A .O vchinnikov, J.Phys.: Condens.M atter 3, 757 (1991).
- ²⁰ J.Carm elb, P .H orsch, P .A .Bares, and A .A .O vchinnikov, Phys.Rev.B 44, 9967 (1991).
- ²¹ J.M .P .Carm elb, P .H orsch, and A .A .O vchinnikov, Phys.Rev.B 45, 7899 (1992).
- ²² J.M .P .Carm elb, P .H orsch, D .K .Campbell, and A .H .Castro Neto, Phys.Rev.B 48, 4200 (1993).
- ²³ J.M .P .Carm elb, A .H .Castro Neto, and D .K .Campbell preprint (1993).
- ²⁴ O .J .Heilm ann and E .H .Lieb, Ann.N .Y .Acad.Sci. 172, 583 (1971).
- ²⁵ E .H .Lieb, Phys.Rev.Lett. 62, 1201 (1989).
- ²⁶ C .N .Yang and S .C .Zhang, M od.Phys.Lett.B 4, 759 (1990).
- ²⁷ Stellan O stlund, Phys.Rev.Lett. 69, 1695 (1992).
- ²⁸ Fabian H .L .Essler and V ladim ir E .K orepin, preprint ITP -SB -93-40 (1993) (to be published); preprint ITP -SB -93-45 and BONN -HE -93-39 (1993).
- ²⁹ J.M .P .Carm elb, D .K .Campbell, A .H .Castro Neto, and N .M .R .Peres, preprint (1994).
- ³⁰ J.M .P .Carm elb and N .M .R .Peres, preprint (1994).
- ³¹ W alter M etzner and Carlo D iCastro, Phys.Rev.B 47, 16107 (1993).
- ³² J.M .P .Carm elb and N .M .R .Peres, unpublished (1994).

- ³³ J.M .P.Carmelo, P.Horsch, and N.M .R .Peres, unpublished (1994).
- ³⁴ J.M .P.Carmelo, A.H .Castro Neto, D.K .Campbell, and N.M .R .Peres, preprint (1994).
- ³⁵ C.N .Yang and C.P .Yang, J.Math.Phys.10, 1115 (1969).
- ³⁶ D.Pines and P.Nozières, in The Theory of Quantum Liquids, (Addison-Wesley, Redwood City, 1966 and 1989), Vol. I.
- ³⁷ Gordon Baym and Christopher J.Pethick, in Landau Fermi-Liquid Theory Concepts and Applications, (John Wiley & Sons, New York, 1991).
- ³⁸ V.E.Korepin, Teor.Mat.Fiz.41, 169 (1979) [Theor.Math.Phys.41, 953 (1980)].
- ³⁹ Alexander B.Zamolodchikov and Alexey B.Zamolodchikov, Nucl.Phys.B 133, 525 (1978).
- ⁴⁰ N.Andrei and J.H.Lowenstein, Phys.Lett.91B, 401 (1980).
- ⁴¹ C.M.Varma, P.B.Littlewood, S.Schmitt-Rink, E.Abrahams, and A.E.Ruckenstein, Phys.Rev.Lett.63, 1996 (1989).
- ⁴² F.Guinea, E.Louis, and J.A.Verges, Phys.Rev.B 45, 4752 (1992).
- ⁴³ Fabian H.L.Essler, Vladimir E.Korepin, and Kareljan Schoutens, Phys.Rev.Lett.68, 2960 (1992).
- ⁴⁴ Hua Chen and Daniel Mattis, preprint University of Utah (1992).
- ⁴⁵ J.P.Pouget, S.K.Khanna, F.Denoyer, R.Comes, A.F.Garito, and A.J.Heeger, Phys.Rev.Lett.37, 437 (1976).
- ⁴⁶ C.S.Jacobsen, Ib Johannsen, and K.Bechgaard, Phys.Rev.Lett.53, 194 (1984).