

MULTIFRACTAL ANALYSIS OF BROADLY DISTRIBUTED OBSERVABLES AT CRITICALITY

MARTIN JANSSEN

Institut für Theoretische Physik, Universität zu Köln, Zùlpicher Str. 77, 50937 Köln, Germany

Received

ABSTRACT

The multifractal analysis of disorder induced localization-delocalization transitions is reviewed. Scaling properties of this transition are generic for multi parameter coherent systems which show broadly distributed observables at criticality. The multifractal analysis of local measures is extended to more general observables including scaling variables such as the conductance in the localization problem. The relation of multifractal dimensions to critical exponents such as the order parameter exponent and the correlation length exponent is investigated. We discuss a number of scaling relations between spectra of critical exponents, showing that all of the critical exponents necessary to characterize the critical phenomenon can be obtained within the generalized multifractal analysis. Furthermore we show how bounds for the correlation length exponent are obtained by the typical order parameter exponent ν_0 and make contact between the multifractal analysis and the finite size scaling approach in 2-d by relying on conformal mapping arguments.

1. Introduction

Anderson localization^{1,2} has gathered interest for over three decades. Electrons in disordered conductors can undergo a transition to an insulating state. At zero temperature the diffusion constant is a function of the Fermi energy ϵ_F and takes a finite value in the conducting phase while it vanishes in the insulating phase which is reached by crossing a certain value $\epsilon_F = E_c$ called the critical energy. This transition is due to a disorder induced localization-delocalization transition of the electrons wave functions. Localization occurs for strong enough disorder because of quantum interference effects brought about by the randomness of the disorder which is assumed to be static (quenched disorder).²

The modeling of disorder induced localization-delocalization transitions (LD transitions) refers to independent electrons of effective mass m moving in a random one-particle potential $V(\mathbf{r})$, i.e. the Hamiltonian is $H = \frac{p^2}{2m} + V(\mathbf{r})$ where \mathbf{r} is

the kinetic momentum which may include magnetic fields. The potential is characterized by its mean value $\langle V \rangle$ and a finite range correlation function $\langle V(\mathbf{r})V(\mathbf{r}^0) \rangle$. A corresponding tight-binding version with on-site random energies and nearest-neighbor hopping is referred to as Anderson model.

Transport properties of electrons are related to two-particle Green functions which are able to describe the transition probability from point \mathbf{r} to point \mathbf{r}^0 at a given energy.³ Here the notion of an n -particle Green function is introduced for the disorder average of a product of n Green functions involving retarded, G^+ , and advanced, G^- , Green functions. These are defined as the matrix elements of the resolvent operators corresponding to H , i.e. $G^\pm(E) = (E - H \pm i)^{-1}$ for $\pm = +, -$.

For quasi-1-d systems (characterized by their length being much larger than their width) rigorous results are available⁴, saying that all states become localized as the length grows infinite. For localized states, the two-particle Green function shows exponential decay, $\langle G^+(\mathbf{E}; \mathbf{r}; \mathbf{r}^0) \rangle \propto \exp(-|\mathbf{r} - \mathbf{r}^0|/\xi(E))$, from which a localization length $\xi(E)$ can be identified.

The lower critical dimension of the LD transition is believed to be $d = 2$.^{5,6} While in the absence of strong magnetic fields or spin-orbit scattering most indications^{5,7} tell that all states are localized in the thermodynamic limit of 2-dimensional systems (though the localization length can be quite large), 2-d systems with strong magnetic fields undergo LD transitions which are believed to be responsible for the occurrence of the quantum Hall effect^{8,9} (QHE) and are referred to as quantum Hall systems (QHS). There is now also striking numerical evidence that 2-d systems with strong spin-orbit scattering show an LD transition.^{10;11;12;13}

In 3-dimensional disordered conductors the LD transition is believed to occur more generally⁵ which has been confirmed numerically.^{14;15;16;17}

Most studies on the LD transition (e.g. Refs. 14;15;17) were focused on the determination of the critical value of the Fermi energy, E_c , (or of the disorder strength) and on the critical exponent of the localization length. Since the late 80's, however, it became clear that the wave functions at criticality are multifractal measures leading to a whole spectrum of critical exponents.^{17;18;19;20;21;22} Nevertheless, only rare connection between these new exponents and the localization length exponent was made.^{17;20;23}

In this article, we wish to emphasize that the LD transition can be viewed as a prototype for a class of critical phenomena characterized by broadly distributed observables at criticality. For this class the multifractal analysis provides a framework to investigate all of the critical exponents. Though, in the following, we often will refer to the LD transition for the sake of concreteness, we hope to make recognition of the generalities possible (a related approach has been put forward by Ludwig²⁴ and by Ludwig and Duplantier²⁵).

The multifractal analysis is a scaling approach relying on the principle of "absence of length scales". Systems without characteristic intrinsic length scales (defining the interrelations of the systems constituents) obey homogeneity laws with respect to rescaling. Let A be an operator or a complex valued function defined for

values x describing a length scale of descriptive nature (e.g. wavelength, system size). Absence of length scales means that A shows a typical homogeneity law

$$A(s^{-1}x) = s^{-\alpha} A(x) \quad (1)$$

where α is called the homogeneity exponent and s is a real number. In other words: the absence of length scales is reflected by the property that a rescaling of x can be compensated by a rescaling of the observable A . For real valued functions $A(x)$ the solution of the homogeneity equation (1) is a power law: $A(x) \sim x^{-\alpha}$. Now, assume that $A^{[q]}(x)$ is a functional of powers q of those observables which are involved in the definition of $A(x)$, then the simplest situation that may appear is that $\alpha(q)$, defined by

$$A^{[q]}(s^{-1}x) = s^{-\alpha(q)} A^{[q]}(x); \quad (2)$$

is a linear function of q ; the operators simply add their exponents. If $\alpha(q)$ shows a significant deviation from linearity we say that the scaling behavior of $A(x)$ is anomalous. Multifractality will turn out to be a generic case of anomalous scaling behavior.

The article is organized as follows. In Sec. 2 several aspects of critical phenomena, especially the notion of order parameters, correlation lengths and critical exponents, are reviewed and the LD transition is put into this perspective. It turns out that the LD transition shows unusual features, the most important of which is the occurrence of a dependence of the local susceptibility on the system size. In Sec. 3 we review the multifractal analysis of measures and the notion of $f(\cdot)$ spectra describing the scaling of broad distributions for local probabilities. Results for the wave function (defining a local measure) of QHS are discussed in detail. Generalizing to positive box-observables in Sec. 4 will allow us to relate the critical exponent of the correlation length, ν , to the multifractal exponents α_0 (the typical scaling exponent of box-probabilities) and X , the normalization exponent of a scaling variable. Furthermore, we discuss bounds for ν in terms of multifractal exponents. In Sec. 5 we focus on correlations in local box-observables. This topic is touched already in previous sections, but is then treated more systematically. As a result we find scaling relations which determine all of the correlation exponents by the original $f(\cdot)$ -spectrum and the corresponding normalization exponents. By relying on conformal mapping arguments we conjecture scaling relations to determine the multifractal spectra via finite size scaling methods in 2-d. Sec. 6 contains our conclusions.

2. Aspects of Critical Phenomena

2.1. Critical Exponents

A defining feature of a critical phenomenon is the divergence of the correlation length ξ_c , defined by the spatial decay of the statistical correlations of a local field, $\langle \phi(0) \phi(r) \rangle \sim \exp(-r/\xi_c)$, has to diverge at the critical point T_c . In the sequel of this

article we refer to the following nomenclature for critical phenomena. The physical states are described by state coordinates (we will refer to them by one symbol T) and by order parameters (m). For infinite system sizes the state coordinates exhibit a critical point T_c where the order parameters vanish with a power law in $t = |T - T_c|$

$$m = 0 \quad (T < T_c) ; \quad m \propto t^\beta \quad (T > T_c) : \quad (3)$$

There exists a generating function $F(t;h)$ where h is the conjugate field with respect to the order parameter $m = \frac{\partial F}{\partial h} (h \neq 0)$. In analogy to equilibrium systems we call F the free energy per unit volume. Higher derivatives of F also show power law behavior for $h = 0, t \neq 0$. For example

$$\chi = \frac{\partial^2 F}{\partial h^2} (h \neq 0) \propto t^{-\gamma} ; \quad c = \frac{\partial^2 F}{\partial t^2} (h \neq 0) \propto t^{-\alpha} : \quad (4)$$

The exponents γ and α (the latter may be negative) are called critical exponents of the susceptibility, χ , and specific heat, c , respectively.

To have a concept of spatial correlations we assume that the order parameter and its conjugate are local fields $m(\mathbf{r}), h(\mathbf{r})$ in a d -dimensional system of volume V for which a generating functional

$$Z(h(\mathbf{r});t) = \int \mathcal{D}[\varphi(\mathbf{r})] \exp \left\{ -H[\varphi(\mathbf{r});h(\mathbf{r});t] \right\} \quad (5)$$

exists where H is commonly called Hamiltonian although it does not have to be hermitean. The functional $F(h;t)$ is given as $F(h;t) = -\frac{1}{V} \ln Z(h;t)$ for the volume $V \rightarrow \infty$. While the mean value m of the order parameter is given by the first moment of the generating functional, $m = h'(\mathbf{r})$, correlation functions are given by higher order cumulants

$$h'(\mathbf{r}_1) \dots h'(\mathbf{r}_n) i_c = \frac{1}{n!} \frac{\delta^n \ln Z}{\delta h(\mathbf{r}_1) \dots \delta h(\mathbf{r}_n)} (h \neq 0) : \quad (6)$$

which are related to moments in a combinatoric way, for example: $h'(\mathbf{r}) i_c = h'(\mathbf{r})$ and

$$h'(\mathbf{r}) h'(\mathbf{r}^0) i_c = h'(\mathbf{r}) h'(\mathbf{r}^0) i - h'(\mathbf{r}) i h'(\mathbf{r}^0) i = : (h'(\mathbf{r}) h'(\mathbf{r}^0)) : \quad (7)$$

$R(r)$ is called local susceptibility, since the generating formalism allows to conclude $\int d^d r R(r) = 1$. The correlation length ξ_c can be identified from $R(r) \sim \exp(-r/\xi_c)$ and is assumed to diverge with a power law approaching $t = 0$,

$$\xi_c(t) \propto t^{-\nu} ; \quad (8)$$

where ν is called critical exponent of the correlation length.

To illustrate the analogies of the LD transition with critical phenomena as outlined here we refer to the fact³ that the LD transition is characterized by the analytic behavior of the disorder averaged two-particle Green function,

$$G^2(\mathbf{r} - \mathbf{r}^0) = \overline{\langle \mathbf{r} \mathbf{r}^\dagger + \mathbf{r}^0 \mathbf{r}^0 \rangle} \langle \mathbf{r}^0 \mathbf{r}^\dagger \mathbf{r} \rangle ; \quad (9)$$

with respect to the infinitesimal parameter in $G = (E - H - i)^{-1}$ which distinguishes retarded and advanced Green functions and controls the long time averages of transport quantities such as the diffusion constant $D(E)$,

$$D(E) = \lim_{t \rightarrow 0^+} \frac{1}{2t} \int_0^t \int_0^t \langle \mathbf{r} \mathbf{r}^T \rangle dt dt' \quad (10)$$

where $\rho(E) = \frac{1}{2\pi i} \langle \mathbf{r} \mathbf{r}^T \rangle (E - G^+(E))^{-1}$ is the average density of states. Thus, it seems natural that a critical phenomenon description of the LD transition may start from a generating function for disorder averaged Green functions. For our purposes we do not need to go into technical details and leave the explicit form of the corresponding Hamiltonian open. We only like to refer to some of the main features of such theories (for details see e.g. Refs. 26; 27) which can be summarized as follows.

The degrees of freedom are matrix fields Q with a block structure $Q^{pp'}$ where $p, p' = \pm$ correspond to retarded (+) and advanced (-) Green functions, respectively. The source field is the infinitesimal parameter and the corresponding source term in the Hamiltonian is $(Q^{++} - Q)$ leading to

$$\frac{\ln Z}{N} = \int d\mathbf{r} \int d\mathbf{r}' \langle \mathbf{r} \mathbf{r}'^T \rangle \quad (11)$$

which is proportional to the average density of states $\rho(E)$. Thus, the density of states in disordered electron systems seems to be a candidate for an order parameter of the LD transition. Unfortunately, the density of states is a smooth function of energy^{3; 28} unable to reflect the LD transition which is characterized by the vanishing of the diffusion constant related to the susceptibility via Eqs. (9, 10). Before coming back to this peculiarity of the LD transition let us proceed in listing common knowledge about scaling ideas applied to critical phenomena.

2.2. Scaling Relations

The crucial assumption of any scaling approach to critical phenomena (see e.g. Ref. 29) is that the critical exponents ν , β , and γ have their origin in the divergence of one relevant length scale, the correlation length ξ_c . Since thermodynamic quantities are generated by the free energy we make the homogeneity assumption that the free energy, in the vicinity of the critical point, can be written as

$$F(t; h) = at^2 f(h/t) + f_r \quad (12)$$

where f is a regular function of the argument h/t and f_r stands for regular terms which do not show scaling behavior for $h \rightarrow 0$, $t \rightarrow 0$. By differentiating F , the following scaling relations

$$2 + \nu = 2 \quad ; \quad 2 = 2 \quad (13)$$

can be concluded. The scaling exponent of the correlation length ξ_c which relates the divergence of ξ_c with the vanishing of t is related to these exponents by the

following scaling argument. Consider a finite system of volume L^d for which the free energy per unit volume is

$$F_L(t;h) = \frac{\ln Z_L(t;h)}{L^d} : \quad (14)$$

A change of L in $Z_L(t_0;h_0)$ to $L^0 = L$ for fixed values $t_0;h_0$ is expected to be compensated by an appropriate change in $t;h$ such that the partition sum remains unchanged, i.e.

$$Z_L(t_0;h_0) = Z_{L^0}(t^0;h^0) : \quad (15)$$

This procedure defines implicitly functions $t^0(L;L^0);h^0(L;L^0)$ with $t(L;L) = t_0$, $h(L;L) = h_0$. Consequently,

$$F_L(t_0;h_0) = F_{L^0}(t^0;h^0) : \quad (16)$$

Choosing the value $L = \xi_c(t_0)$ we see $F_{\xi_c(t_0)}(t_0;h_0) = F_{L^0}(t^0;h^0)_{\xi_c(t_0)}$ and find a scaling relation

$$2 = d : \quad (17)$$

We can cast the scaling relations Eq. (13) in a form, where only ν , β , and γ appear

$$2 + \nu = 2 = d : \quad (18)$$

This means that only two of these four exponents are independent.

We introduced the correlation length with the help of the local susceptibility. At the critical point, where χ_c diverges, the correlation function can show a homogeneity law

$$\langle r \rangle \sim r^{-z} : \quad (19)$$

Owing to the general relation between the global susceptibility and the correlation function $\chi = \int d^d r \langle r \rangle$, and by introducing the correlation length $\xi_c(t)$ as a cut-off length in the integral over $\langle r \rangle$, which diverges at $t = 0$, we find another scaling relation which is independent of the homogeneity assumptions for the free energy

$$= (d - z) : \quad (20)$$

Together with the scaling relations of Eq. (18) this tells that only two of the five exponents ν , β , γ , and z are independent.

We now come to a crucial point for what follows on the multifractal analysis of broadly distributed observables at criticality. Imagine that in addition to the distance r the finite system size L is in a regime of absence of length scales. Assume that the function $C(r;L) \equiv h'(\epsilon)'(\epsilon^0)_L$ shows a homogeneity law with respect to both lengths, r and L ,

$$C(r;L) \sim r^{-z} L^{-y} : \quad (21)$$

Note, that in the literature on critical phenomena the exponent z is often written in a form $z = d - 2 + \dots$. We will avoid this notation since it could lead to confusions in the context of the LD transition where an exponent ν is used sometimes in a different context (see the discussion below).

The exponent y describes the system length dependence and $z \in \mathbb{R}$ describes the distance dependence. That such a situation will indeed appear in the LD transition problem will be discussed in Sec. 2.4 and in more detail in Secs. 3.3; 5.2. A similar reasoning now shows that

$$= (d - (y + z)) ; y + z = z \quad (22)$$

where the last relation follows from consistency with Eq. (20). Notice that identifying the exponent in $z = d - 2 +$ as the correlation exponent means that such is not determined by and alone, but requires the additional knowledge of y .

2.3. ϕ -Functions

A method to test scaling assumptions and calculate critical exponents is the so-called renormalization group. The definition of the renormalization group is not unique, but can in general be described as a transformation acting on the Hamiltonian H of the system where a length scale serves as transformation parameter. Instead of introducing a finite system size, one can also work in the thermodynamic limit considering the system to be defined on a lattice. Changing the lattice constant a_0 to $L_b = ba_0$, and changing the coordinates $t; h$ in an appropriate way, one tries to keep $\ln Z$ fixed. After many iterations of this procedure (which defines the renormalization group) one ends up with a new Hamiltonian the structure of which should not have changed except for the change in the coordinates and that the fields are now defined on a new lattice. That such a procedure indeed works makes the model renormalizable. Otherwise one had to introduce a number (in the worst case an infinite number) of new terms and coordinates into the new Hamiltonian in order to keep $\ln Z$ fixed. Thus, the application of the renormalization group to the free energy density leads to

$$F_0 = F(t; h) = b^{-d} F(t^0; h^0) \quad (23)$$

where $t^0 = b^{y_t} t$, $h^0 = b^{y_h} h$ in the critical regime and y_h, y_t are scaling exponents. Choosing the factor b such that $t^0 = \text{const.}$, we arrive again at the scaling relations of Eqs. (13, 18). By considering the system size L being the scaling parameter, an extension of renormalization group ideas can be made, based on the following assumption. Far away from the critical regime (where power law scaling holds) there still exist scaling variables $\phi(L)$ which fulfill a restrictive functional equation:

$$\phi(bL) = f(\phi(L); b) : \quad (24)$$

For simplicity we consider the case of only one scaling variable here. This means that the value of ϕ for system size bL only depends on the value of ϕ for system size L and the scaling factor b where f is called a scaling function. This assumption is very strong since, in general, ϕ will depend on many microscopic parameters. The idea behind this assumption is that most of these parameters will become unimportant for describing the flow of the scaling variable with increasing system size, and that the scaling function f is the only information needed to determine

this flow. The existence of a scaling function f is equivalent to the existence of a β -function

$$(\ln \beta) := \frac{d \ln \beta(L)}{d \ln L} = -\frac{1}{b} \frac{\partial f}{\partial b} - \frac{\partial f}{\partial \epsilon} \quad (b=1) \quad (25)$$

which is indeed a function of $\ln L$ alone. Provided the β -function is smooth, the flow is given by the solution of the differential equation

$$(\ln \beta) = \frac{d \ln \beta}{d \ln L} : \quad (26)$$

Eq. (26) is called a renormalization group equation. The regime where β can be linearized around a fixed point of the flow ($(\ln \beta) = 0$) is called critical regime,

$$(\ln \beta) = \beta^0 (\ln L - \ln L_0) : \quad (27)$$

Here β^0 is the slope of the β -function at L_0 . Starting from a system of size $L = L_0$, with $\epsilon = \epsilon_0$ chosen close to ϵ_c , and turning on the renormalization flow until the system reaches a size ξ_c marking the bound of validity of Eq. (27) (which is hereby by definition the correlation length), one finds

$$\frac{\xi_c}{L_0} = \frac{\ln L_0 - \ln \xi_c}{\ln L_0 - \ln L_0} : \quad (28)$$

The critical regime is narrow, we expand $(\ln \beta)$ and β^0 around ϵ_c and get for the correlation length

$$\xi_c = L_0 \frac{1 - \beta^0(\epsilon_c)}{\beta^0(\epsilon_c) - \beta^0(\epsilon)} : \quad (29)$$

Consider the situation where the width of the critical regime $\epsilon_c - \epsilon_0$ is triggered by the parameter t (leaving L_0 and ϵ_0 unchanged), the divergence

$$\xi_c / L_0 \sim 1 / t \quad (30)$$

follows, where the exponent is given by the inverse slope of the β -function at the fixed point, i.e.

$$\nu = 1 / \beta^0(\epsilon_c) : \quad (31)$$

2.4. Unusual Features of the LD Transition

It is now time to put the LD transition into the perspective of the critical phenomena terminology. A common feature of field theoretic approaches to the LD transition is that disorder averaged n -particle Green functions are generated by a field theoretic generating functional in the sense of Sec. 2.1. As already mentioned, the density of states which is a one-particle Green function appears as the formal order parameter and the conjugate field to the density of states is the infinitesimal parameter of the Green function. The density of states does not show the LD transition, but is a smooth function of the energy which, in the problem of the LD transition, is

the analog of the temperature in equilibrium phase transitions. Consequently, the "critical exponent" of this formal order parameter is zero

$$\beta = 0 : \quad (32)$$

The LD transition cannot be described by the formal order parameter, Eq. (11), of the generating functional. The LD transition can only be obtained by analyzing quantities which are related to the two-particle Green function such as the diffusion constant $D(E)$.

Furthermore, the scaling relations, Eqs. (18,20), tell that the correlation exponent of the local susceptibility (a two-particle Green function) vanishes, i.e.

$$\alpha = 0 : \quad (33)$$

This result seems to rule out a power law behavior of the two-particle Green function (e.g. the density correlator). However, Chalker and Daniel³⁰ demonstrated, by numerical calculations, the density correlator to show a power law behavior at the LD transition of a quantum Hall system, $\propto r^{-0.38}$. In addition, by relying on an exact inequality obtained by Chalker³¹ one has analytical evidence that the density correlator at the critical point cannot be described by a vanishing distance exponent. This phenomenon is often referred to as anomalous diffusion. Fortunately, from our discussion of correlation exponents in Sec. 2.2 where we distinguished between different types of distance exponents z and \bar{z} we can fairly conclude that the value of 0.38 does not correspond to \bar{z} , but corresponds to the distance exponent z in a regime where the correlation function shows power law also with respect to another length scale which can be interpreted as a system size L being much smaller than the correlation length ξ_c , i.e.

$$z = 0.38 \neq \bar{z} = 0 : \quad (34)$$

The phenomenon of anomalous diffusion and the inability of the density of states to describe the LD transition are two of the main unusual features of the LD transition. By Eq. (34) it is indicated that both are not independent of each other.

Another unusual feature of the LD transition which, at first sight, seems to be independent of the previous ones was observed by Altshuler et al.^{32,33} They reexamined the phenomenological scaling theory for the LD transition of Abrahams et al.³³ In this scaling theory the conductance $g(L)$ of a cube with linear dimension L is considered to be a scaling variable in the sense of Sec. 2.3, i.e. g obeys a differential equation in terms of a β -function

$$\frac{d \ln g}{d \ln L} = \beta(g) \quad (35)$$

which is a unique function of $\ln g$. As shown by Altshuler et al.^{32,33} the conductance of a mesoscopic system highly depends on the individual properties of a given system (e.g. on the concrete disorder potential). As a result an ensemble of different systems

reflects a broad distribution in g which cannot be characterized by the mean value of g alone. Consequently, the mean value of the conductance g_i is not a suitable scaling variable. Instead, in any scaling approach to the LD transition one has to consider the whole distribution function of the conductance. However, it may be possible to apply the ideas of the renormalization group to certain parameters of this distribution function. Following Shapiro³⁴ we refer to an s -parameter scaling theory for the LD transition if s parameters $g_{rel,1,...,s}$ obeying scaling equations

$$\frac{d \ln g_{rel}}{d \ln L} = (fg_{rel,k} g_{k=1,...,s}) \quad (36)$$

are required to determine the distribution function of a scaling variable g . If, at least in the vicinity of the transition point ($fg_{rel,k} g_{k=1,...,s} = 0$), only one relevant length scale ξ_c and one type of scaling variable $g_{rel}(L)$ exists, then the critical exponent ν of the correlation length ξ_c is defined by

$$g_{rel}(L) \sim g_{rel} / L^{1/\nu} \quad \text{for } L \gg \xi_c \quad (37)$$

and $1/\nu$ is given by the slope of the corresponding \ln -function at g_{rel} . A context where this concept will serve to be fruitful is provided by the generalized multifractal analysis to be discussed in Sec. 4. There we will also see that anomalous diffusion and broad distributions of physical quantities are deeply connected.

3. Multifractal Analysis of Measures

After having discussed some aspects of critical phenomena, where the absence of length scales is reflected by the existence of critical exponents, the multifractal analysis of measures is reviewed in this section.^{35;36;37;38} This analysis is appropriate to describe self-similar local observables which can be interpreted as measures. It can also be viewed as an extension of the fractal dimension approach to self-similar geometric objects invented by Mandelbrot.³⁹

3.1. Scaling of Moments

To introduce the notion of the multifractal analysis and for the sake of concreteness we consider a quantum Hall system at criticality. For a finite two dimensional quantum Hall system of linear size L with double periodic boundary conditions we study the electrons wave function ψ , the modulus of which defines a normalized measure. The probability for an electron to be found in a box of linear size L_b is given by the box-probability

$$P(L_b) := \int_{\text{box}} d^2 r |\psi|^2 : \quad (38)$$

Covering the system by a mesh of $N(L_b; L)$ boxes the fractal dimension D of the wave functions support is defined by $N(L_b; L) \sim L_b^{-D}$ where $N(L_b; L)$ is the number of boxes with non-vanishing box-probability for a given ratio $L_b = L$. Since the wave

function is never exactly zero in any box, $N(\epsilon)$ equals the total number of boxes $N(L_b; L)$ and, consequently, the fractal dimension is the Euclidean dimension of the system, i.e. $D = d = 2$. At the level of the fractal dimension D the wave function shows no interesting fractal behavior. Let us now focus our attention on the scaling behavior of the box-probability. The normalization condition $\sum_i^{N(\epsilon)} P_i(L_b) = 1$ yields the scaling behavior for the average

$$\langle P(L_b) \rangle_L / L^{-D} : \quad (39)$$

Here the average of quantities A_i (corresponding to the box numbers i) is defined by $\langle A \rangle_L = \frac{1}{N(\epsilon)} \sum_i^{N(\epsilon)} A_i$. Let us consider higher moments of the box-probability $P(L_b)$ with respect to this averaging procedure. The general assumption underlying the multifractal analysis is that for a finite interval of values of the moments $\langle P^q(L_b) \rangle_L$ show power law behavior indicating the absence of length scales in the system, i.e.

$$\langle P^q(L_b) \rangle_L / L^{-D + (q)} \quad (40)$$

where (q) does not depend on L . In quantum Hall systems the assumption about the absence of length scales means that on the one hand the correlation length ξ_c of the LD transition has to be much larger than the lengths $L_b; L$. On the other hand microscopic length scales l , such as the cyclotron radius r_c , have to be much smaller than $L_b; L$. In summary the condition for the multifractal analysis to be useful is

$$l \ll L_b < L \ll \xi_c : \quad (41)$$

For finite systems the states which are candidates for the multifractal analysis are states with localization lengths being much larger than the system size. This corresponds to the critical states of the LD transition in finite systems. In the thermodynamic limit $L \rightarrow \infty$ such states can only be found at the critical energy E_c . In this limit ξ_c goes to zero and the function (q) can be defined uniquely for the critical state by

$$(q) = \lim_{L \rightarrow \infty} \frac{\ln \langle P^q(L_b) \rangle_L}{\ln L} \quad D : \quad (42)$$

For finite systems one can give an estimate for (q) by considering the slope in a plot of $\ln \langle P^q(L_b) \rangle_L$ versus $\ln L$ over an interval of values L which fulfill the above mentioned constraints. Now we want to discuss how multifractality is reflected by the properties of the function (q) . Note that by the construction of $\langle P^q(L_b) \rangle_L$ we have found an observable in the sense of Eq. (2). The function (q) corresponds to the homogeneity exponents (q) and multifractality of the wave function means that

(q) is a non-linear function of q . According to the normalization condition, $(1) = 0$; $(0) = D$ and (q) can thus be parametrized by the generalized dimensions $D(q)$, defined by

$$(q) = : D(q) - (q - 1); \quad (43)$$

and the fractal dimension $D = D(0)$. Consequently, a single-fractal wave function is characterized completely by the fractal dimension D , i.e. $D(q) = D$. $D(q)$ being not a constant function distinguishes multifractals from single-fractals.

The reason why the multifractal analysis is a powerful method to analyse scaling behavior comes from the fact that one can make very general statements about the analytic behavior of the $\phi(q)$ function and the $D(q)$ function, respectively. Assuming smoothness of both functions for q being real numbers one can derive the following results:^{38;40}

The function $\phi(q)$ is a monotonically increasing function with negative curvature, i.e.

$$\frac{d\phi(q)}{dq} > 0; \quad \frac{d^2\phi(q)}{dq^2} < 0: \quad (44)$$

The generalized dimensions $D(q)$ are positive, monotonically decreasing and bounded by $D_{-1} \leq D(q) \leq D_1$, i.e.

$$\frac{dD(q)}{dq} < 0; \quad 0 \leq D_1 \leq D(q) \leq D_{-1}: \quad (45)$$

The function $\phi(q)$ has asymptotically constant slopes given by D_1

Here a sketch of the proves of these statements is given (for details see Refs. 38;40). The main part of the derivations relies on the homogeneity, reflected by power laws, already for finite ratio b , and on the normalization condition which keeps the box probabilities less or equal to unity. First, the monotonicity of $\phi(q)$ is a simple consequence of $P_i \leq 1$ and, thus, $P_i^q < P_i^{q^0}$ for $q > q^0$. A much stronger statement is the monotonicity of $D(q)$ which can be shown by writing $D(q)$ as

$$D(q) = \frac{1}{q} \frac{\ln \left(\sum_i P_i^q \right)}{\ln \left(\sum_i P_i \right)} = \ln \left(\frac{\sum_i P_i^q}{\sum_i P_i} \right)^{\frac{1}{q-1}}; \quad (\ln)^{-1} \quad (46)$$

with the help of a generalized Hölder inequality,

$$\left(\sum_i P_i (P_i)^r \right)^{\frac{1}{r}} \leq \left(\sum_i P_i (P_i)^{r^0} \right)^{\frac{1}{r^0}} \quad \text{for } r > r^0: \quad (47)$$

Notice that for distinct values of q and q^0 the equality $D(q) = D(q^0)$ only holds in the case of a constant function $D(q) = D$. For each ratio b there exist maximum and minimum values for the box-probabilities. Now raising the box-probabilities to positive powers $q \neq 1$ means that in the sum $\sum_i P_i^q$ only the maximum values contribute significantly yielding the dimension $D_1 > 0$ by

$$P_{\max}(L_b) / L_b^{D_1}: \quad (48)$$

A similar consideration for $q \neq -1$ shows that the minimum values of the box-probabilities dominate in the sum $\sum_i P_i^q$ yielding the dimension D_{-1} by

$$P_{\min}(L_b) / L_b^{D_{-1}}: \quad (49)$$

Thus, the $\phi(q)$ function has asymptotic slopes given by D_1 which shows that the limits of $D(q)$ for $q \rightarrow \pm 1$ exist and are just given by D_1 . Notice that we referred to finite ratios guaranteeing the existence of maximum and minimum values. Extrapolating from small but finite values of q to the thermodynamic limit $q \rightarrow 0$ becomes possible due to scaling, i.e. the absence of a length scale in the system. To complete the proof we have to show that $\phi(q)$ has negative curvature. Due to the smoothness assumption

$$\phi(q) = \frac{d}{dq} \ln Z(q); \quad \phi'(q) = \frac{d}{dq} \phi(q) \quad (50)$$

exist and are continuous functions of q . The definition of $\phi(q)$ yields

$$\begin{aligned} \phi(q) &= \frac{1}{Z(q)} \sum_i \frac{\ln P_i}{\ln} P_i^q > 0 \\ \phi'(q) &= \frac{\sum_i \frac{\ln P_i}{\ln} P_i^q}{Z(q)} - \frac{\phi(q)^2}{Z(q)} \ln \\ &= \ln \frac{\sum_i \frac{\ln P_i}{\ln} P_i^q}{Z(q)} - \phi(q)^2 \end{aligned} \quad (51)$$

where

$$Z(q) = \sum_i P_i^q \quad (52)$$

is called the partition sum of the multifractal. The equality in the relation $\phi'(q) = 0$ only holds if $\phi(q) = D$, i.e. in the single-fractal situation. From Eq. (50) one can see that the objects characterizing multifractals are constructed in a similar way to thermodynamic quantities where the partition sum $Z(q)$ is replaced by the canonical partition sum $Z(\beta; V)$ in Boltzmann statistics. Indeed, one can write down for each of the quantities like $q; \ln$; $\phi(q) ::$ its thermodynamic counterpart (see e.g. Ref. 41). Keeping this analogy between multifractals and thermodynamics in mind one can imagine that a violation of the smoothness condition for $\phi(q)$ could be viewed as an intrinsic phase transition in each multifractal wave function. Such kind of phase transition (which has not been observed in systems showing LD transitions up to now) should not be confused with the LD transition itself.

3.2. Distributions

The $\phi(q)$ function describes the scaling behavior of moments, i.e. of $\langle P^q \rangle$. We wish to describe the whole distribution function $\rho(P)$ that corresponds to these moments. For a function $F(P)$ the corresponding average value reads

$$\langle F(P) \rangle_L = \int_0^1 dP \rho(P) F(P) \quad (53)$$

Since we are interested in the scaling behavior of the distribution function itself we change the variable P to $\ln P$ by defining

$$\tilde{P}(\ln P) d \ln P = P(P; \ell) dP : \quad (54)$$

Now, average values can be calculated with the help of $\tilde{P}(\ln P)$ via

$$\langle F(P; \ell) \rangle_L = \int_0^{\infty} d \ln P \tilde{P}(\ln P) F(\ln P) : \quad (55)$$

Absence of length scales forces the distribution function \tilde{P} to show power-law scaling with respect to ℓ . Since, in the thermodynamic limit $L \rightarrow \infty$, the values $\langle F \rangle_L$ are uniquely defined one can apply the method of steepest descent when calculating the average for $F = P^q$. This leads to the following conclusion

There exists a positive function, the $f(q)$ -spectrum of the multifractal, characterizing the scaling behavior of the distribution function \tilde{P} , and is related to $\langle P^q \rangle_L$ by a Legendre transformation.

$$\ln \langle P^q \rangle_L = f(q) / q : \quad (56)$$

Here the function $f(q)$ is the same as introduced in Eq. (50).

Using the analogy between multifractality and thermodynamics one can check that $f(q)$ corresponds to the microcanonical statistics and resembles the entropy as a function of energy. Let's focus on the meaning of the function $f(q)$ in the context of multifractality. This function describes the scaling behavior of the whole distribution function of box-probabilities in the absence of length scales. In the single-fractal case the function $f(q)$ shrinks to the point $(0; D)$ in an $(q; f(q))$ diagram. Consequently, the distribution function \tilde{P} is singular, and the distribution function with respect to the variable P , $P(P; \ell)$, is a narrow distribution on all length scales. In the single-fractal case $P(P; \ell)$ is typically a gaussian distribution and can be characterized by a few cumulants.

In the multifractal case there appears a new and unexpected behavior of the distribution function $P(P; \ell)$ which we want to describe in the following. Due to general properties of Legendre transformations we have

$f(q)$ is a positive, single-humped function of negative curvature on a finite interval of q values:

$$D_1 < q < D_2 ; 0 < f(q) < D ; f(0) = D \quad (57)$$

where $q_0 = (q = 0) > D$.

$f(\cdot)$ terminates at the points $(D_1; 0)$ with infinite slopes and has slope $f'(1) = 1$ at the point $(1; 1)$ where $\tau = 1$, as well as $f(1) = D_1$.

These statements can most easily be derived by writing the functions $\tau(q)$, $\tau(q)$ and $f(q)$, defined by $f(\tau(q)) = \tau(q)$, as⁴²

$$\begin{aligned}\tau(q) &= \frac{1}{\ln} \ln Z(q; \cdot) \\ \tau(q) &= \frac{1}{\ln} \sum_i P_i^q \ln P_i \\ f(q) &= \frac{1}{\ln} \sum_i P_i^q \ln P_i : \end{aligned} \quad (58)$$

Here the q -dependent normalized quantity

$$P_i(q; L_b) = \frac{P_i^q(L_b)}{\sum_i P_i^q(L_b)} \quad (59)$$

is a generalization of the original box-probability P_i and is called the q -microscope since it increases the large box-probabilities for positive values of q and increases the small box-probabilities for negative values of q , respectively. The second of these equations (58) is only a reparametrization of Eq. (50) and by Legendre transforming $\tau(q)$ one can easily check that $f(q)$ is correctly described by the third equation of (58). We mention that numerical calculations of $f(\cdot)$ -spectra are of higher precision when using Eqs. (58) compared to numerical Legendre transformation of $\tau(q)$.

Let us summarize the instruments for describing the multifractal scaling behavior of normalized box-probabilities: the $\tau(q)$ function describes the scaling behavior of moments and $f(\cdot)$ describes the scaling behavior of the corresponding distribution function. Both of them are related by Legendre transformation. The functions $D(q)$ and $\tau(q)$ can also serve to describe the multifractal nature of box-probabilities but have, for our purposes, less appealing interpretations.

Motivated by the single-humped shape of the $f(\cdot)$ -spectrum we try to give a physical interpretation of multifractality, at least in the context of the LD transition. It is illuminating to approximate the $f(\cdot)$ -spectrum by a parabola and to see to which kind of distribution function the parabolic approximation (PA) corresponds. The PA, which fulfills most of the desired constraints, is determined by one parameter, τ_0 , besides the geometric fractal dimension D ,^{20;37} i.e.

$$f(\tau) = D - \frac{(\tau - \tau_0)^2}{4(\tau_0 - D)} : \quad (60)$$

This corresponds to a log-normal distribution of the box-probabilities

$$P(\tau; \cdot) / \exp \left(-\frac{(\tau - \tau_0)^2}{4(\tau_0 - D)} \right) \ln : \quad (61)$$

A log-normal distribution is the prototype of a broad distribution which cannot be characterized by a few cumulants. For example, the n -th cumulant grows exponentially with n , $\propto e^{n^2}$. Scaling requires that the distribution is broad on all length scales and with the normalization condition on the box-probabilities the form given by Eq. (61) can be viewed as the paradigm of a distribution function which is broad on all length scales. The physical interpretation of such a behavior is the following.

In phase coherent disordered conductors the actual value of a local box-quantity, like $P_i(L_b)$, depends on a large number of conditions simultaneously. This can be simulated by writing $P_i(L_b)$ as a product of a large number of independent random factors,

$$P_i(L_b) = p_0 \cdot p_1 \cdot p_2 \cdots; \quad (62)$$

and the central-limit theorem tells that one can expect $P_i(L_b)$ to be log-normal distributed. The scaling condition (absence of length scales) requires the distribution function to exhibit power law behavior. Thus, we conclude

$$\sim \left(\frac{\ln P}{\ln} \right)^{-D} \quad f(\ln P = \ln) \quad (63)$$

with $f(\cdot)$ being almost parabolic. For completeness we list the PA for $D(q)$ and (q)

$$D(q) = D_0 - q(q - D_0); \quad (q) = q_0 - 2q(q - D_0); \quad (64)$$

However, the parabolic approximation can never be exact since it violates some of the constraints we have derived for $f(\cdot)$; mainly positivity and the boundary conditions at $(D_0, 0)$. PA can only serve as a good approximation in the vicinity of the most probable value q_0 for the scaling exponents. The breakdown of the PA can be indicated by either the values $q = q_0 - \frac{D_0}{2} \frac{D(q - D_0)}{D_0}$ (where $f(\cdot)$ vanishes in the PA), or by $q_+ = \frac{q_0}{2} \frac{D_0}{(q - D_0)}$ (where (q) loses monotonicity in the PA). They give rough estimates of D_0 .

The most probable value of the box-probability corresponding to q_0 is given by the typical value of $P(L_b)$,

$$P_{\text{typ}}(L_b) = \exp \left(\frac{f(\ln P(L_b))}{\ln} \right) \quad (65)$$

which is a geometric type of mean value.

Finally, we present results of the multifractal analysis for wave functions in the critical regime of a finite quantum Hall system.²⁰ In Fig. 1 the squared amplitude of a wave function is shown where increasing darkness corresponds to higher probability to find an electron at the corresponding point in the system. The picture demonstrates the self-similar structure of the wave functions and the tremendous amount of fluctuation of local probabilities, even within one given state. In Fig. 2 $f(\cdot)$ spectrum for this wave function is graphically depicted, showing that the box probabilities are almost log-normal distributed and show power law behavior. The most interesting values of scaling exponents are

$$q_0 = 2.3 \pm 0.07; D_1 = 0.95 \pm 0.1; D_2 = 3.7 \pm 0.1; \quad (66)$$

Fig. 1. Squared amplitude of a multifractal wave function in the critical regime of a quantum Hall system. Increasing darkness corresponds to higher probability of finding an electron (on a logarithmic scale).

Fig. 2. The $f(\cdot)$ spectrum of a multifractal wave function at the center of the lowest Landau level.

Fig. 3. Comparison of $f(\nu)$ -spectra for three different wave functions in the critical regime of a quantum Hall system. The dots correspond to the $f(\nu)$ -spectrum and other curves correspond to the parabolic approximations for the $f(\nu)$ -spectra of this and two distinct states.

The fractal dimension is, of course, $D = 2$.

Striking is the observation that the $f(\nu)$ function seems to be universal for all of the critical states. This can be seen in Fig. 3 where the $f(\nu)$ -spectrum of one state is shown together with the parabolic approximations of this and two distinct states chosen in the critical regime. The coincidence is remarkable. Calculations of $f(\nu)$ -spectra, carried out by Pook²⁰ for the same model system, but with different realizations of the random potential and different system sizes, confirm this result. Further evidence was provided by calculations of Huckestein et al.^{22,43} for a tight-binding version of the quantum Hall system which give the same values for the quantities ν_0, D_{-1} within the error bars. This means that the LD transition of quantum Hall systems (at least for the lowest Landau level where all of the numerics has been performed) is uniquely characterized by those critical numbers.

3.3. Inverse Participation Numbers

In this subsection we describe how disorder-averaged Green functions are involved in the multifractal analysis via inverse participation numbers. Thereby two applications will be addressed: the first concerns the calculation of the critical exponent of the correlation (localization) length in a quantum Hall system. The second concerns the anomalous diffusion found by Chalker et al.^{30,44} as mentioned in Sec. 2.4.

The role of fractality for the LD transition was already observed by Aoki⁴⁵ in

1983 by looking at the inverse participation number defined by

$$P = \frac{1}{Z} \int d^d r j(\mathbf{r})^4 \quad (67)$$

where $\int d^d r$ denotes a d -dimensional system with linear dimension L . He gave an argument that the wave function $\psi(\mathbf{r})$ of electrons at the LD transition point shows fractal structure. If the wave function is uniformly distributed, $P \propto L^{-d}$, the participation ratio $p = (P L^d)^{-1}$ is constant. In the localized regime $P \propto L^{-d}$ and p vanishes in the thermodynamic limit. At the transition point where the wave function is extended the participation ratio still vanishes in the thermodynamic limit. Consequently, P scales with a power $d < d$. By a box-counting method Aoki⁴⁶ calculated the fractal dimension of wave functions of a QHS at the transition point to be approximately 1.5. Although his method was not free of systematic errors, plots of the electron density $j(\mathbf{r})^2$ indicated self-similarity.

Castellani et al.¹⁹ considered the $d = 2 + \epsilon$ expansion of Wegner's non-linear model⁴⁷ for the generalized inverse participation numbers

$$P^{[q]} = \frac{1}{Z} \int d^d r j(\mathbf{r})^{2q} / L^{-d(q)} \quad (68)$$

and concluded that the wave functions at the transition point show multifractal behavior since $d(q)$ is not equal to $d(q=1)$ but defines a set of fractal dimensions $d(q)$:

$$d(q) = d(q)(q=1) : \quad (69)$$

To show the relation between $d(q)$ and $d(q)$ we consider $P^{[q]}$ calculated in a tight binding model with lattice constant b ,

$$P^{[q]} = \frac{1}{X} \sum_i j(i)^{2q} = Z(q; b=L) : \quad (70)$$

If both quantities b and L are in a regime of self-similarity the identity of $d(q)$ and $d(q)$ follows immediately. However, they are different in the continuum limit ($b \rightarrow 0$) for fixed L since self-similarity breaks down once b is smaller than the microscopic length scales l . On the other hand, in the continuum limit, we can consider $Z(q; b=L)$ to be an approximation for $P^{[q]}$ which becomes more and more accurate as L goes to infinity. If self-similarity is preserved up to length scales $L \gg 1$, $d(q) \approx d(q)$ can be concluded. The assumption of self-similarity for $L \gg 1$ is fulfilled at the critical point of the LD transition where the only relevant length scale, the localization length ξ , diverges. Thus, at the critical point of the LD transition

$$d(q) = d(q) : \quad (71)$$

On the other hand the correlation length ξ_c is defined as an upper bound for power law behavior and $d(q)$ can be determined from the scaling behavior of $P^{[q]}$ with ξ_c :

$$P^{[q]} \propto \xi_c^{-d(q)} : \quad (72)$$

The critical exponent ν_c of the correlation length ξ_c is defined by

$$\xi_c \propto |E - E_c|^{-\nu_c} \quad (73)$$

where E_c is the difference of a critical observable (e.g. the energy) from its critical value. In analogy to this, Wegner⁴⁷ introduced critical exponents $\nu(q)$ for the generalized inverse participation numbers

$$P^{[q]} \propto |E - E_c|^{-\nu(q)} \quad (74)$$

Thus, we conclude a scaling relation between multifractal scaling numbers $\nu(q)$ and critical exponents ν_c and ν_2 ,^{17;48}

$$\nu(q) = \nu_c + \nu_2(q) \quad (75)$$

and $\nu(q)$ depends on q in a non-trivial way if the wave functions are multifractals. Still, Eq. (75) doesn't yield ν_c as a function of only multifractal scaling numbers. Nevertheless, we can see what was overlooked by Hikami⁴⁹ when calculating the critical exponent for a QHS (restricted to one Landau level) to be 1.9 ± 0.2 . What he did calculate was the exponent ν_2 . He found $\nu_2 = 3.8 \pm 0.4$. The result for ν_c given above follows if $P^{[2]}$ is assumed to scale with the localization length as ξ^2 , ignoring that multifractality leads to the anomalous scaling behavior $P^{[2]} \propto \xi^{D(2)}$. Taking multifractality into account, $D(2) = 1.62 \pm 0.02$,⁵⁰ we get

$$\nu_c = 2.4 \pm 0.3 \quad (76)$$

This is in agreement with the high accuracy result, $\nu_c = 2.34 \pm 0.04$, of Huckestein and Kramer⁵¹ obtained by a finite size scaling technique.

As a further application to the QHE we show that multifractality relates the exponent ν of anomalous diffusion to $D(2)$. We have already anticipated that the spectrum of multifractal dimensions has universal features for states in the vicinity of the LD transition since we omitted to distinguish between inverse participation numbers of individual states and their ensemble average. In fact, as already discussed, different wave functions of several systems show, in the critical regime, the same $f(q)$ -spectra within the error bars.

The observation of anomalous diffusion (see Sec. 2.4) means that the averaged two-particle Green function

$$\overline{G^+(r; E)} = \overline{\frac{1}{N} \sum_j \langle j | G(E - H + i0^+) | j \rangle} \quad (77)$$

behaves as

$$\overline{G^+(r; E)} \propto r^{-(d+2)} \quad (78)$$

This power law, with a value $\nu = 0.38 \pm 0.04$, was found⁵⁰ in quantum Hall systems ($d=2$) for length scales r in the regime of multifractality, i.e. $l \ll r \ll \xi$. Replace the ensemble average of the inverse participation ratio (69) of wave functions ψ_i with

$$P^{[q]}(E) = \frac{1}{N} \sum_i \frac{|\psi_i(E)|^{2q}}{|\psi_i(E)|^2} \quad (79)$$

where ϵ_i (i) corresponds to energy eigenvalue ϵ_i . The ensemble averaged inverse participation number $P(\epsilon)$ is now defined with respect to energy. It can be calculated from the Green function⁴⁷

$$P(\epsilon) = \lim_{L \rightarrow 0} \frac{1}{L^d} \overline{|\mathcal{G}^+(0; \epsilon)|^2} \quad (80)$$

where $\rho(\epsilon) = \lim_{L \rightarrow 0} \frac{1}{L^d} \int^R d^d r \overline{|\mathcal{G}^+(r; \epsilon)|^2}$ is the density of states. The expected scale independence of this quantity is excellently confirmed by numerical results²⁰. Thus, with the ansatz

$$\overline{|\mathcal{G}^+(r; \epsilon)|^2} = (\epsilon; L) r^{-z} \quad (81)$$

one concludes

$$P(\epsilon) \sim (\epsilon; L) / L^{z-d}; \quad (82)$$

and consequently

$$d + d - 2 = z = d - D(2); \quad (83)$$

More precisely, in the case of multifractal behavior one expects the Green function to have a spectrum of exponents and the one occurring in Eq. (83) should be interpreted as the dominating one. In the regime of multifractality, besides the distance r , a second length scale, the system size L , appears in the correlation function $\overline{|\mathcal{G}^+|^2}$ (Eq. (81)). This was anticipated in Sec. 2.4 when discussing the difference between the distance exponents z and \mathbf{z} . Using the terminology of Sec. 2.2 we identify the exponents y and \mathbf{z} to be

$$y = \mathbf{z} = D(2) - d; \quad \mathbf{z} = y + z = 0; \quad (84)$$

In a single-fractal situation, $z = 0$ and $y = \mathbf{z}$. Thus, in systems without multifractal structure there is no need for introducing two exponents when discussing correlation functions in finite systems. The numerical value⁵⁰ $D(2) = d(2) = 1.62 \pm 0.02$ is in accordance with our considerations, and anomalous diffusion can be interpreted as being a direct consequence of the multifractal nature of wave functions at the critical point of the LD transition. This was already conjectured in Ref. 30. There is no contradiction between a finite value of \mathbf{z} , as interpreted via the distance exponent z instead of \mathbf{z} , and the vanishing of the exponent corresponding to the formal order parameter of the LD transition. The physical reason for the phenomena of anomalous diffusion and of multifractality is the quantum coherence of highly disordered systems.

4. Generalized Multifractal Analysis

So far, only measures, i.e. box-probabilities P , have been considered. To get into contact with usual critical exponents we introduce a generalized multifractal analysis for non-normalized observables.²⁰ A related approach in the context of field theory was put forward by A. Ludwig.²⁴

Consider a physical observable Q which can be defined for systems of any linear extension L . For simplicity, we assume the system to be a box of volume $(L_b)^d$ where d is the euclidean dimension of the system. Thus, for any of such boxes we have a box-observable $Q_i(L_b)$ similar to the box-probability in the foregoing sections. Examples are the magnetic moment or the two-terminal conductance of a d -dimensional cube with volume $(L_b)^d$.

We are going to study the statistical and scaling properties of box-observables. We restrict to positive values for Q ,

$$Q(L_b) \geq 0; \quad (85)$$

to avoid cancellations in averaging procedures and to apply the results of the foregoing sections. At first glance, this seems to be a strong restriction. However, still a large class of observables can be studied which is specified by the following consideration. Let a physical observable $A(L_b)$, measured on a box, be an element of some observable algebra, then the scaling behavior of $A(L_b)$ may show up in a positive scalar factor Q ,

$$\begin{aligned} A(s \cdot x) &= s \cdot A(x) \\ A(x) &= Q(x) \cdot \hat{A}(x) \\ Q(x) &\propto x \end{aligned} \quad (86)$$

where the algebra element \hat{A} does not show significant scaling behavior, or it's average might even vanish due to symmetry. The following generalized multifractal analysis applies to such observables. With this restriction in mind we proceed as follows.

4.1. Normalization Exponents

To study the statistics of box-observables one takes a large number N of equivalently prepared systems and records the values of the box observable. This leads to the set $\{Q_i(L_b)\}_{i=1, \dots, N}$. The mean value of a function of Q is then calculated as

$$\langle F(Q(L_b)) \rangle_N = \frac{1}{N} \sum_{i=1}^N F(Q_i(L_b)); \quad (87)$$

A corresponding smooth distribution function $F(Q; L_b; N)$ is assumed to describe average values (provided N is large enough),

$$\langle F(Q(L_b)) \rangle_N = \int_0^\infty dQ F(Q) F(Q; L_b; N); \quad (88)$$

In the absence of intrinsic length scales, i.e. compared with L_b intrinsic length scales are either much less or much larger than L_b , one expects power law scaling to occur in averages. To make contact to the multifractal analysis of measures at

an early state of the investigations let us introduce a symbolic system size L by reparametrizing the number N as

$$L := L_b^d N^{1-d} \quad (89)$$

Let $N = N$ be the number of non-vanishing values for $Q_i(L_b)$, define the normalization of Q by

$$\|Q(L_b; L)\| := \sum_i^{X^N} Q_i(L_b) \quad (90)$$

and, by keeping L fixed in the averaging procedure, identify the normalization exponent $X^{[Q]}$ in the regime of power law scaling via

$$\|Q(L_b; L)\| / L_b^{X^{[Q]}} : \quad (91)$$

$X^{[Q]}$ can be any real number. The crucial step is the construction of a normalized box-observable $P(Q; L_b; L)$ associated with Q :

$$P_i(Q; L_b; L) = \frac{Q_i(L_b)}{\|Q(L_b; L)\|} ; \quad \sum_i^{X^N} P_i(Q; L_b; L) = 1 : \quad (92)$$

$P_i(Q; L_b; L)$ is normalized with respect to the symbolic system of linear size L but, in general, not additive,

$$P(Q; L; L) \neq \sum_i^{X^N} P_i(Q; L_b; L) : \quad (93)$$

In the usual multifractal analysis one imagines the box-observables to be measures for which the normalization condition is very natural. Nevertheless, when proving the general features of the functions q and $f(\cdot)$ the normalization condition was essential, but no use was made of additivity. By fixing L we are able to attach to the positive, normalized box-observable $P(Q; L_b; L)$ the multifractal analysis by defining $^{[Q]}(q)$, $^{[Q]}(q)$ and $f^{[Q]}(^{[Q]})$ with the help of the partition sum,

$$Z(Q; q;) := \sum_i^{X^N} P_i^q(Q; L_b; L) ; \quad (94)$$

in direct analogy to Eqs. (58). For these functions all the general features explained in Sec. 3 follow if the ratio $= L_b = L$ is chosen small enough, and the observables $Q_i(L_b)$ are in the scaling regime. The results of Sec. 3 can be translated to the following statements

For positive box-observables $Q(L_b)$ the scaling behavior of its distribution function can be described by the corresponding normalization exponent $X^{[Q]}$ and a single-humped function $f^{[Q]}(^{[Q]})$

$$\sim \frac{\ln Q}{\ln} ; L_b \rightarrow L_b^D h\left(\frac{\ln Q}{\ln}\right) ; h\left(\frac{\ln Q}{\ln}\right) = f^{[Q]}(^{[Q]}) \quad (95)$$

with

$$\frac{\ln Q}{\ln} = \mathbb{Q} + X \mathbb{Q} : \quad (96)$$

The scaling relations for the moments and the typical value,

$$Q_{\text{typ}}(L_b) = \exp f \ln Q(L_b) ; \quad (97)$$

read

$$\begin{aligned} \langle P^Q(Q; L_b; L) \rangle & / L_b^{D + \mathbb{Q}(q)} \\ \langle Q^q(L_b) \rangle & / L_b^{x(q)} ; x(q) = D + \mathbb{Q}(q) + q X \mathbb{Q} \\ Q_{\text{typ}}(L_b) & / L_b^{\mathbb{Q} + X \mathbb{Q}} \end{aligned} \quad (98)$$

The statements of Sec. 3 about properties of ρ , μ , f such as monotonicity, curvature, Legendre transform, etc. do also apply for \mathbb{Q} ; \mathbb{Q} and $f \mathbb{Q}$. In general one expects $D = d$, unless a systematic scaling dependence of $N(N)$ appears.

4.2. Local observables

In case that Q is a local additive observable, i.e. for each subdivision of a starting system of linear size L into N boxes of linear size L_b , $Q(L_b)$ is an observable and additivity holds,

$$Q(L) = \sum_{i=1}^N Q_i(L_b); \quad (99)$$

the generalized multifractal analysis can also be applied to the investigation of scaling properties with respect to the system size L which is then a true system size. We focus on the situation where L_b and L are in a regime of power law scaling and vary both lengths independently. We thus have to be aware of an additional normalization exponent $Y \mathbb{Q}$ with respect to L which is, in general, different from $X \mathbb{Q}$,

$$\langle Q(L_b; L) \rangle / L_b^{X \mathbb{Q}} L^{Y \mathbb{Q}} : \quad (100)$$

The box-probabilities studied in Sec. 3 fall, of course, into the class of local observables with trivial normalization exponents $X = Y = 0$. For local observables one finds, besides the scaling relations of Eqs. (95, 98) with respect to L_b , (by varying L and taking the normalization exponent $Y \mathbb{Q}$ of Eq. (100) into account)

$$\langle Q^q(L_b) \rangle_L / L_b^{x(q)} L^{-y(q)} ; Q_{\text{typ}}(L_b; L) / L_b^{\mathbb{Q} + X \mathbb{Q}} L^{\mathbb{Q} + Y \mathbb{Q}} \quad (101)$$

with

$$x(q) = D + \mathbb{Q}(q) + q X \mathbb{Q} ; y(q) = D + \mathbb{Q}(q) + q Y \mathbb{Q} : \quad (102)$$

As an illustration let's discuss two examples. A trivial example is the box-observable $Q = P L_b^X L^{-Y}$ where P is a box-probability with multifractal functions ρ , μ and

f and $X; Y$ are real numbers. Then, $[\mathbb{Q}] = \mathbb{Q}$, $[\mathbb{Q}] = \mathbb{Q}$, $f^{[\mathbb{Q}]} = f$, and $X^{[\mathbb{Q}]} = X$, $Y^{[\mathbb{Q}]} = Y$. As another example consider the box-observable $Q = P^m$ where P is a box-probability and m some real number. All the relevant scaling information is already contained in the (q) function of P . Thus, it is easy to check that

$$\begin{aligned} X^{[P^m]} &= Y^{[P^m]} = (m) \\ [P^m](q) &= m (q - m) [P^m] \\ f^{[P^m]}([P^m](q)) &= (qm) (qm) (qm) : \end{aligned} \quad (103)$$

The last of these three equations yields, due to the Legendre transform property of $f(\cdot)$ with respect to (q) ,

$$f^{[\mathbb{Q}]}([\mathbb{Q}]) = f(qm) : \quad (104)$$

It is tempting to expect from this examples that $f(\cdot)$ -spectra of different observables, defined for one ensemble of equivalently prepared systems, are either identical, if only the normalization is distinct, or are related in a simple manner, if powers of elementary observables are involved.

4.3. Typical Scaling Variables

In the vicinity of a phase transition one expects a universal behavior of the scaling exponents. They may depend on general properties of the systems dynamic, however not on microscopic details. We have already observed that different wave functions in the LD transition regime showed the same $f(\cdot)$ -spectrum. A further indication for universality of $f(\cdot)$ -spectra was provided by the observation of Hukestein and Schweitzer⁴³ that the local equilibrium current density as well as the local magnetization⁵² show the same $f(\cdot)$ -spectrum as the wave functions in a quantum Hall system. However, universality of scaling variables in the LD transition (which are non equilibrium properties) is even more interesting.

Recall the phenomenological scaling theory of Abraham et al.⁵ for the LD transition (see Sec. 2.4) where the conductance g is assumed to be a scaling variable on all length scales. As we already mentioned in Sec. 2.4 the conductance of mesoscopic systems has a broad distribution which cannot be described in terms of the mean value of g . Consequently, one has to consider the distribution function $(g; L)$ which can be characterized by relevant parameters g_{rel} for which ϕ -functions exist. If g is chosen for the $f(\cdot)$ analysis a candidate for g_{rel} is obviously given by the typical value $g_{typ}(L) = \exp f \ln g(L) i g$

$$g_{typ}(L) = g_{yp} / L^{[g]_0 + X[g]} \quad (105)$$

and by comparing Eqs. (37,98) one concludes

$$1 = [g]_0 + X[g] : \quad (106)$$

The universality of ϕ suggests the $f(\cdot)$ -spectra of normalized scaling observables being universal, too. Notice that Eq. (106) provides a method to calculate the critical

Fig. 4. Thouless numbers for 6 realizations of a quantum Hall system with linear size $120r_c$ restricted to the lowest Landau level. ω_c is the cyclotron frequency. Energy windows for data evaluation as described in the text are indicated by vertical lines (after Ref. 23).

exponent with only the help of multifractal critical numbers. This gives the possibility for calculating by a different method than the finite size scaling method.^{14;15} In addition, if both methods coincide the multifractal analysis demonstrates that typical observables are relevant for scaling; a consequence of the fluctuating self similar structure underlying multifractal objects.

In order to check the predictions of the multifractal analysis Fastenrath et al.²³ calculated numerically Thouless numbers for the QHS and applied the multifractal analysis to the data. The Thouless numbers are defined by the ratio of the energy shift due to a change of boundary conditions (e.g. periodic \rightarrow antiperiodic) with respect to the mean level spacing. They can be thought of as being a transport quantity showing the same qualitative (and scaling) behavior of a (dimensionless) dc conductance.⁵³

In Fig. 4 data for the Thouless numbers denoted by g are shown which are calculated for the lowest Landau level of 6 realizations of a QHS with linear size $120r_c$. One can see the critical regime outside of which the states are localized and have zero dc conductance. In the critical regime there are large fluctuations and the corresponding histogram of Fig. 5 demonstrates the broadness of the distribution function. To apply the multifractal analysis, Thouless numbers $g(L)$ for a large number of different systems with varying system sizes from $30r_c$ to $120r_c$ were calculated. As the box observable Q the modulus of the deviation $Q(L) := |g(L) - g_{typ}|$ was chosen where g_{typ} was indeed size independent. Fig. 6 shows that the corresponding $f(\cdot)$ -spectrum of the Thouless numbers is very similar to that of the

Fig. 5. Histogram of Thouless numbers calculated for the lowest Landau level of one quantum Hall system.

box-probabilities.²⁰ Data have been taken from a narrow window in the critical region as indicated in Fig. 4. The authors find

$$g_0^{[1]} = 2.22 \pm 0.05; X^{[1]} = 1.75 \pm 0.05; \nu = 2.2 \pm 0.3: \quad (107)$$

They also calculated ν directly from $j \ln g_{\text{typ}} - \ln g_{\text{typ}} j / L^1 =$ and found

$$\nu = 2.3 \pm 0.2: \quad (108)$$

In either case they used $g_{\text{typ}} = 0.27 \pm 0.03$, determined at the band center. The data for ν are in agreement with the high precision value of Huckestein and Kramer, obtained by finite size scaling calculations.⁵¹

The multifractal analysis of Fastenrath et al.²³ confirms the one-parameter scaling theory of the LD transition in terms of the typical conductance and demonstrates the broadness of the conductance distribution at criticality.

4.4. Bounds for the Correlation Length Exponent

In this section we show how the multifractal analysis provides a lower bound for the critical exponent ν . We refer to the LD transition for concreteness but the results generally are reflected by the fact that only a few assumptions such as one-parameter scaling are needed. To obtain upper bounds for ν more restrictive assumptions are needed.

That ν has a lower bound given by

$$\nu > \frac{2}{d} \quad (109)$$

Fig. 6. $f(\epsilon)$ spectra of normalized (right) and unnormalized (left) Thouless numbers. The displacement of the extrema is just the normalization exponent X . The broken line to the right is a parabolic approximation (after Ref. 23).

where d is the Euclidean space dimensionality is known since the work of Chayes et al.⁵⁴ A trivial upper bound for ν is given by

$$\nu < 1 \quad (110)$$

otherwise the definition of the exponent would be meaningless (e.g. in the Kosterlitz-Thouless transition).⁵⁵ We give arguments which improve both bounds. Our arguments rely on the assumption that one parameter scaling holds true in the vicinity of the LD transition point, on the analytic properties of $\chi(q)$ functions and on the universality hypothesis.

In a field theoretical statistical model for a critical phenomenon as outlined in Sec. 2 a scaling operator ϕ couples to the scaling field t (in the LD transition problem t corresponds to $\beta(E_c) = t$) in the Hamiltonian

$$H = t \int d^d r S(\mathbf{r}) : \quad (111)$$

Here $S(\mathbf{r})$ is a local scaling operator of the field theoretical statistical model. The scaling dimensions $\gamma(n)$ corresponding to the scaling operators

$$[n] = \int d^d r S^n(\mathbf{r}) \quad (112)$$

are defined with respect to the renormalization group action (see e.g. Ref. 56). Renormalizability of the theory then means that only a finite number of these

exponents are positive (relevant) and most of them are negative (irrelevant). Here we identify them via the finite size scaling properties of the average

$$\langle Z^* \rangle = \frac{1}{L} \int_0^L d\mathbf{r} S^n(\mathbf{r}) / L^{(n)} \quad (113)$$

where the statistical model is defined on a finite system of linear size L . For a relevant scaling operator the integral diverges for $L \rightarrow 1$ and consequently the scaling fields $t^{[n]}$ in $t^{[n]} - t^{[n]}$ have to be zero, i.e. the critical point is reached. One-parameter scaling means that there is exactly one relevant operator and thus

$$(n) < 0 \text{ for } n \geq 2 : \quad (114)$$

If by accident or symmetry properties the expectation value in Eq. (113) vanishes we assume that there exists a corresponding positive scalar observable Q which has the same scaling behavior as $S(\mathbf{r})$, in the spirit of Eq. (86). If this assumption holds true, then the general analytic properties of (q) functions tell that

$$\begin{aligned} (n) &= \gamma^{[1]}(n) + X^{[1]} - n \\ \frac{d(n)}{dn} &= \gamma^{[1]}(n) + X^{[1]} \\ \frac{d^2(n)}{dn^2} &= 0 : \end{aligned} \quad (115)$$

Consequently the scaling dimensions have positive curvature. Since the exponent is given by $\gamma(1) = 1 =$ (for consistency with the divergence of the correlation length) we have

$$X^{[1]} = 1 = \quad (116)$$

In order to have (n) monotonically decreasing guaranteeing renormalizability the lower bound

$$> 1 = D_1^{[1]} \quad (117)$$

has to be respected. Furthermore, the one-parameter scaling condition Eq. (114) requires

$$> \frac{2}{D^{[1]}(2)} : \quad (118)$$

This resembles and improves the Chayes et al. criterion Eq. (109). The observation of universality in $f(\cdot)$ spectra for different observables suggests $D^{[1]}(q) = D(q)$. Relying on the universality hypothesis for $f(\cdot)$ the multifractal analysis of the wave function in the LD transition problem already allows to obtain a lower bound for γ , $\gamma > 2 = D(2)$.

In Sec. 4.3 we have found that it is the typical conductance g_{typ} rather than the average conductance g_i which serves as a one-parameter scaling variable for the LD transition and scales like

$$g_{\text{typ}}(L) = g_{\text{typ}} / L^{1=} : \quad (119)$$

The corresponding β -function has an unstable fixed point. The exponents $x^{[g]}(q)$ describing the scaling of moments $\langle g(L) \rangle \sim q_{yp}^f / L^{x^{[g]}(q)}$ are given by

$$x^{[g]}(q) = D + \frac{[g]}{0}(q) + qX^{[g]} : \quad (120)$$

The analytic properties of $x^{[g]}$ are summarized as follows.

$$\begin{aligned} \frac{dx^{[g]}(q)}{dq} &= \frac{[g]}{0}(q) + X^{[g]} \\ \frac{d^2 x^{[g]}(q)}{dq^2} &= \frac{d}{dq} \frac{[g]}{0}(q) < 0 : \end{aligned} \quad (121)$$

Since $x^{[g]}(0) = 0$ and $1 = \frac{[g]}{0} + X^{[g]} > 0$ the exponents $x^{[g]}(q)$ are positive for small positive values of q . For large enough values of q they will become negative if $\frac{d}{dq} \frac{[g]}{0} + X^{[g]} < 0$ which means that high moments of $\langle g(L) \rangle \sim q_{yp}^f$ may decrease when approaching the transition point. This is not in contradiction to renormalizability and does not yield a rigorous upper bound for β . However, if at least the first moment has positive scaling exponent, $x^{[g]}(1) = D + X^{[g]} > 0$, then an upper bound for β can be concluded

$$\beta < \frac{1}{\frac{[g]}{0} - D} : \quad (122)$$

Note, that this upper bound relies on the additional, though intuitive, assumption about a positive scaling exponent for the mean deviation $\langle g(L) \rangle \sim q_{yp}^f$.

We give three examples for which the validity of the bounds can be checked relying on the universality of $f(\beta)$ spectra.

1. For the QHS the values for β , $D(q)$ and $\frac{[g]}{0}$ reported above are compatible with the bounds,

$$1.3 < \beta = 2.3 < 3.3 : \quad (123)$$

2. A similar conclusion holds for the 2-d spin-orbit model (the calculation of $\beta = 2.75$ was done by finite size scaling methods¹¹ and $\frac{[g]}{0} = 2.175$ was calculated by the multifractal analysis⁵⁷),

$$1.2 < \beta = 2.75 < 6 \quad (124)$$

3. The one-loop results of Wegner's non linear ϕ -model¹⁷ in $d = 2 + \epsilon$ leads to a parabolic $f(\beta)$ -spectrum as shown in Ref. 19. The results are

$$\beta = 1 - \frac{\epsilon}{2} ; \frac{[g]}{0} = d + \frac{\epsilon}{2} ; D(2) = d - 2 + \frac{1}{2} \quad (125)$$

Thus, the one loop result meets the upper bound and the lower bound is valid up to $\beta < 2.3$. Higher loop orders⁴⁷ lead to $f(\beta)$ spectra which violate the condition of constant curvature if taken seriously for arbitrary values of ϵ . This indicates that the higher orders are improvements only for very small values of ϵ (as often

happens in asymptotic expansions). The applicability of the loop expansion is an open question.

5. Correlations in Multifractals

In this section we ask for correlations in positive, local observables as introduced in Sec. 4.2. For example, a local field $\phi(\mathbf{r})$ gives rise to local box-observables

$$Q_i(L_b) = \frac{1}{Z} \sum_{\text{box}(i)} \phi(\mathbf{r}_j) \phi(\mathbf{r}_j) : \quad (126)$$

We are interested in the scaling properties of

$$M^{[q]}(\mathbf{r}; L_b; L) = \overline{Q_i^q(L_b) Q_{i+s}^q(L_b)}_L \quad (127)$$

where the average is over all pairs of boxes with fixed distance $r = sL_b$.

5.1. Scaling Relations

Usually, in critical phenomena one studies correlations for infinite system size (and L_b being microscopic) as a function of r alone. Here we consider a regime, where both r and L are able to show scaling behavior. We thus start with an ansatz

$$M^{[q]}(\mathbf{r}; L_b; L) \sim L_b^{x_2(q)} L^{-y_2(q)} r^{-z(q)} \quad (128)$$

and try to find relations of the exponents $x_2(q)$, $y_2(q)$, $z(q)$ to the previous ones $x(q)$ and $y(q)$.⁵⁷ A typical heuristic scaling argument to fix scaling relations relies on considering limiting cases where scaling will be already violated. However, since the limit is reached continuously and scaling exponents are unique one can conclude scaling relations due to consistency. We thus proceed by considering the limiting cases for $M^{[q]}$: (i) $r = L_b$ and (ii) $r = L$. In case (i) we expect the asymptotics

$$M^{[q]} \sim \overline{Q^{2q}(L_b)}_L \sim L_b^{x(2q)} L^{-y(2q)} : \quad (129)$$

In case (ii) we expect the asymptotics

$$M^{[q]} \sim \overline{Q^q(L_b)} \overline{Q^q(L_b)}_L \sim h Q^q(L_b) L_L^2 \sim L_b^{2x(q)} L^{-2y(q)} \quad (130)$$

where $\overline{Q^q(L_b)}$ is the value for $Q^q(L_b)$ already averaged over one system of size L . Comparison with Eq. (128) yields the scaling relations

$$\begin{aligned} y_2(q) &= y(2q) \\ x_2(q) &= x(2q) \\ z(q) &= 2y(q) - y(2q) = 2x(q) - x(2q) \end{aligned} \quad (131)$$

which coincide with those of a more sophisticated derivation by Cates and Deutsch⁵⁸ in the case of Q being a local measure. Notice that the normalization exponents X

⁵⁷In the following we suppress the superscript $[q]$ in exponents.

and Y cancel in $z(q)$ leading to $z(q) = D + 2 - (q) - (2q)$. The analytic properties of $z(q)$, $x_2(q)$ and $y_2(q)$ are summarized as follows. $z(q)$ vanishes at $q = 0$ where it has a minimum. It is monotonically increasing (decreasing) for $q > 0$ ($q < 0$), and its slope vanishes for $q = 1$. $x_2(q)$ and $y_2(q)$ have negative curvature. Their monotonicity depend on the sign of $(2q) + X$ and of $(2q) + Y$, respectively. Since (q) is monotonically decreasing they are monotonically increasing provided $D_1 > X$; Y . Notice that $x_2(q)$ and $y_2(q)$ depend on the normalization exponents and may become negative for positive values of q .

The fact that $M^{[q]}$ can depend on the system size L is unusual compared to ordinary critical phenomena where the correlation function, the susceptibility $\chi(r)$, is expected to depend on the distance r only, provided the thermodynamic limit is reached and the correlation length is infinite at the critical point T_c . We like to comment on this in the following.

In Sec. 3.2 we discussed an interpretation of multifractality of the box-probability in the context of the LD transition: Multifractality reflects broadness of the box-probabilities distribution function on all length scales which is due to the dependence of the box-probability in each box on a large number of conditions, simultaneously. More general, we call a situation, where a local box observable depends on a large number of conditions for the entire system of linear size L , simultaneously, a situation of "many parameter (MP) coherence". In the context of the LD transition coherence at zero temperatures is due to quantum mechanical phase coherence of the electrons wave function and disorder introduces a huge number of parameters, e.g. the position of point-scatterers. It may happen that MP coherence is valid only up to a certain scale $L < \hat{L}$, defined implicitly by $M^{[q]}$ being independent of L for $L > \hat{L}$. We call \hat{L} a MP coherence length. If such crossover in $M^{[q]}$ exists, two situations have to be distinguished. First, \hat{L} introduces a cut-off for correlations. For example, the correlation length is a MP coherence length of this kind. Alternatively, \hat{L} does not introduce a cut-off and correlations still show homogeneity exponents for distances $r > \hat{L}$. An example for this kind of MP coherence length in the LD transition problem will be discussed below. In the latter situation the following scaling behavior of $M^{[q]}$ is expected to occur for $r \ll \hat{L} \ll L$,

$$M^{[q]}(r; L; \hat{L}) / \hat{L}^{-y_2(q)} r^{-z(q)} ; \quad (132)$$

and for $L \ll r \ll \hat{L}$,

$$M^{[q]}(r; L; \hat{L}) / r^{-z(q)} ; \quad (133)$$

respectively. Thus, for $r \gg \hat{L}$ the usual behavior is recovered. However, for this situation the multifractality on scales less than \hat{L} is still reflected by the q -dependence of $z(q)$ the scaling relation of which can be concluded by a similar reasoning as that leading to Eqs. (131)

$$z(q) = y_2(q) + z(q) = 2y(q) : \quad (134)$$

The analytic properties of $z(q)$ are such: $z(q)$ has negative curvature and its monotonicity properties depend on the sign of $(q) + Y$. It can thus happen, that $z(q)$

is negative for a wide range of q values. A system which is never MP coherent, i.e. \hat{L} is microscopic, the single-fractal situation applies and, then, $z(q) = 0$ and $\bar{z}(q) = 2\gamma(q) = \gamma(2q)$.

5.2. Application to LD Transitions

In the LD transition problem the density of states of an individual system is a suitable candidate for a local box-observable. It is defined by

$$\rho(E; \mathbf{r}) := \frac{1}{N} \sum_{j=1}^N \langle \mathbf{r} | \delta(E - H_j) | \mathbf{r} \rangle \quad (135)$$

where ψ_j are eigenstates of H with respect to energy E . One peculiar feature of the LD transition is that the average density of states, $\bar{\rho}(E) = \frac{1}{L^d} \text{Tr}(\rho(E))$, is not an order parameter and is L -independent. The corresponding box observable is

$$Q_i(L_b; E) := \frac{1}{L_b^d} \int_{\text{box}(i)} d^d \mathbf{r} \rho(E; \mathbf{r}) \quad (136)$$

The scale-independence of $\bar{\rho}(E)$ determines the normalization exponents to be

$$X = 0; Y = D : \quad (137)$$

The fact that the average of $\rho(E)$ is not an order parameter ($\beta = 0$) is equivalent to $x(1) = 0$. However, the typical value of Q_{typ} can serve as an order parameter, because

$$Q_{\text{typ}} / L^{-(D+D)} \quad (138)$$

and consequently, by L approaching ∞ , we have

$$Q_{\text{typ}} / L^{-(D+D)} : \quad (139)$$

Thus, in contrast to the mean value \bar{Q} , Q_{typ} is able to indicate the LD transition and we have reason to call

$$Q_{\text{typ}} \approx (0 + Y) = (0 + D) \quad (140)$$

the typical order parameter exponent.

The exponents $z(q)$ do not depend on the normalization exponents and are given as $z(q) = D + 2 - (q)$ whereas the exponents $\bar{z}(q)$ do depend on $Y = D$,

$$\bar{z}(q) = 2(q - 1) + (q - D) \quad (141)$$

have negative curvature, are positive for $0 < q < 1$, vanish at $q = 0; 1$ and are negative elsewhere. The function $\bar{z}(q)$ is, due to Eq. (135), the same as for the wave function itself.

For $q = 1$ we can compare with the result of Chalker and Danieli³⁰ for the scaling of the density correlator at the LD transition of a quantum Hall system,

$$\langle \rho(E + i\epsilon; \mathbf{r}) \rho(E - i\epsilon; \mathbf{r}') \rangle \quad (142)$$

where l sets the scale of a MP coherence length $L_l = (l)^{-1/2}$. L_l can be interpreted as the linear system size for a system with mean level spacing of about l .³⁰ Their result is: for $r \ll L_l$ correlations scale like $r^{0.38}$ whereas for $r \gg L_l$ correlations behave like r^0 . The first of these results was already discussed in Secs. 2,4 and can be recovered from Eq. (131)

$$z = z(1) = D \quad D(2) = 0.38 : \quad (143)$$

The second result is consistent with the interpretation of z being the distance exponent in a regime $r \ll L_l$, i.e.

$$z = z(1) = 0 : \quad (144)$$

A similar observation was made in Ref. 12 for the LD transition in 2-d system with spin-orbit coupling, though the results were not conclusive concerning quantitative results for $z(1)$. Furthermore, the scaling of the density correlator with respect to L_l fulfills the scaling relation of Eq. (131) which means here

$$y_2(1) = y(2) = z = 0.38 : \quad (145)$$

This came out in the work of Chalker et al. by studying the combined variable $L_l k$ where k is the inverse wavelength in the Fourier-transform of the density correlator and probes long distances in the limit $k \rightarrow 0$. Instead of interpreting l via L_l one can (more directly) look at the density correlator of Eq. (142) as describing the density correlations in energy and, owing to Fourier transformation for the limit $l \rightarrow 0$, the long time correlations of wave packets. Huckestein and Schweitzer⁵⁰ checked the latter point of view by explicitly calculating the long time correlations of wave packets in the critical regime of a quantum Hall system and found excellent agreement, i.e. $z = 0.38 \pm 0.02$.

Unfortunately, no results for values $q \neq 1$ for $z(q)$; $y_2(q)$ and $z(q)$ are available up to now. Thus, the picture developed here needs further tests.

5.3. Conformal Mapping in 2-d

We apply conformal mapping arguments to find a relation between finite size scaling (FSS) methods relying on strip-like systems and the multifractal analysis for square-like systems in 2-d. To begin with some ideas about conformal mapping are reviewed (for a satisfactory treatment see e.g. Ref. 59).

For critical correlation functions $A(r) \sim r^{-\alpha}$ the following assumption seems plausible. Scale invariance, as reflected by Eq. (1), should hold also for local transformations which preserves angles but may change scales locally in the sense that correlations of one system with a certain geometry are mapped onto those correlations of a similar system the geometry of which is determined by the transformation. Such transformations are called conformal mappings. The corresponding Jacobian J has to fulfill

$$\frac{v \cdot w}{(v^2 w^2)^{1/2}} = \frac{Jv \cdot Jw}{((Jv)^2 (Jw)^2)^{1/2}} \quad (146)$$

where $\mathbf{v}; \mathbf{w}$ are vectors of the tangent space at a given point. In 2-d any holomorphic function $f(z = x + iy)$ is a conformal mapping. Especially, the complex logarithm provides a conformal mapping of the entire plane onto an infinite strip with periodic boundary conditions. Let's introduce Cartesian $(x; y)$ and polar coordinates $(r; \theta)$ on the plane by $x + iy = re^{i\theta}$. Then the conformal mapping $F(x + iy) = \frac{M}{2} \ln(x + iy) = u + iv$ maps a ring area, denoted by C , with inner radius $l=L$ and outer radius L onto a strip of width M with length $L^0 = \frac{M}{2} \ln L$, centered at $u = 0$ and fulfilling periodic boundary conditions in v -direction. For a correlation function on the plane $A(r) / r^z$ which behaves regular for $r \neq 0$ we can express the exponent by an integral expression in coordinate free terms by (cf. Ref. 60)

$$z = \frac{1}{2} \int_C \star d(\ln A), \quad (147)$$

where the two-form $\star = d \wedge d(\ln A)$, \star is the Hodge-operator corresponding to the Euclidean metric, and $l=L$ is assumed to be small enough to give vanishing contribution to \int_C by applying Stokes' theorem. By lifting \star with the conformal mapping F , and assuming exponential decay of the corresponding correlation function along the strip, $A(u; v) \sim \exp(-|u|/M)$, one arrives at the result

$$\frac{M}{2} = \frac{1}{z} : \quad (148)$$

We wish to apply this result to correlation functions of multifractal correlations $M^{[q]}$ with L_b being microscopic. However, in the regime $r \sim \hat{L}; L$ conformal mapping arguments of the kind presented here can never apply since a second length scale besides the distance r appears. We thus have to focus on the regime $r \sim \hat{L}$ where $M^{[q]}$ behaves as

$$M^{[q]}(r) \sim r^{-z(q)} : \quad (149)$$

At the critical point of the LD transition, where power law in the plane is valid, the decay lengths $\lambda_c^{[q]}(M)$ of the corresponding correlations in the strip-geometry vary linearly with M . This fact is exploited in the FSS analysis of the LD transition (see e.g. Ref. 14). Thus, the FSS variable $\lambda_c^{[q]}(M) \sim 2^{-[q]}(M) = M^{z_c^{[q]}}$ becomes a constant $\lambda_c^{[q]}$ at the critical point.

Making the hypothesis, that the conformal mapping result Eq. (148) applies to the multifractal correlations in the regime described by Eq. (149) one would conclude

$$\lambda_c^{[q]} = 2 = (-z(q)) : \quad (150)$$

However, in the situation of the LD transition $z(q)$ takes negative values for $q > 1$ which is counterintuitive and suggests exponential growth of correlations in the strip geometry. Furthermore, the exponents $z(q)$ and the decay lengths $\lambda_c^{[q]}$ are attached to averages of powers of the correlation function, though the precise statement on

²The factor of 2 is due to convention.

the existence of the decay length in the strip-geometry relies on the average of the logarithms of the correlations (see e.g. Ref. 14). Thus, we leave the validity of Eq. (150) as an open question and turn over to the typical values defined by the average of logarithms. In FSS calculations it is the average of the logarithm of the correlation function which is calculated by knowing that this quantity is self-averaging. Therefore, we denote results of such FSS calculations by ζ_c^{typ} and a conjecture about the relation to multifractality (relying on conformal mapping arguments) reads

$$\zeta_c^{\text{typ}} = 2 = (\zeta_{\text{typ}}) ; \zeta_{\text{typ}} = 2(\zeta_0 + Y) \quad (151)$$

where the expression for ζ_{typ} is a consequence of Eqs. (134,101).

In the context of the LD transition we found $Y = D$ for the local observable being the density of states. This observable involves the square amplitude of wave functions and localization lengths are usually defined with respect to the modulus of the wave function. This explains the conventional factor of 2 in the definition of (M) .

Up to now, only data for two universality classes are known where one can check the prediction of Eq. (151): (i) quantum Hall systems and (ii) 2-d Anderson model with spin-orbit scattering. Taking into account that present days calculations, lacking larger systems, are able to produce results for ζ_c^{typ} and ζ_0 with a precision hardly exceeding 5%, the results are not in conflict with Eq. (151): (i) $\zeta_c^{\text{typ}} = 1.14^{61}$ and $1 = (\zeta_0 - 2) = 1.06^{43}$ (ii) $\zeta_c^{\text{typ}} = 1.88^{62}$ and $1 = (\zeta_0 - 2) = 1.82^{57}$.

6. Conclusions

After reminding some aspects of critical phenomena in Sec. 2 we described the multifractal analysis of broadly distributed observables in the critical regime of a critical phenomenon. Broadness of the distribution on length scales much less than the correlation length ζ_c is determined by the single-humped $f(\cdot)$ spectrum of normalized box-observables. The parabolic approximation for $f(\cdot)$ corresponds to a log-normal distribution. Scaling of the q -th moment is given by the function $\Phi(q)$ defining generalized dimensions $D(q) = \Phi(q)/(q-1)$ ($D = D(0)$ being the geometric fractal dimension) and by the normalization exponents $X; Y$. The function $\Phi(q)$ is monotonically increasing and has negative curvature. It is related to $f(\cdot)$ by Legendre transformation (Secs. 3-4). Since the distribution is broad it is not possible to characterize it by the mean value. Instead, the typical value defined as the geometric mean is a self-averaging quantity which scales with exponent $\zeta_0 + X$ where ζ_0 is the maximum position of $f(\cdot)$, $f(\zeta_0) = D$. We interpreted the appearance of broad distributions being due to many parameter coherence, saying that local observables depend on a large number of conditions for the entire system, simultaneously.

Furthermore, correlation functions of broadly distributed observables at criticality show scaling dependence with respect to the distance between local observables and with respect to the linear system size L or, equivalently, with respect to a length \hat{L} indicating the range of many parameter coherence. For distances r exceeding the

length \hat{L} correlations will only depend on r (Sec. 5). Although compatible with known results, this scenario needs further investigations.

Scaling exponents of correlations are related to (q) , X , Y by scaling relations (Eqs. (131,134)). Therefore, a complete characterization of critical exponents in terms of multifractal spectra is possible.

We applied the multifractal analysis to the localization-delocalization transition induced by disorder (LD transition) leading to the following conclusions. The formal order parameter, the average density of states, does not show the LD transition which means the order parameter exponent vanishes, $\beta = 0$. Consequently, usual scaling relations tell that correlations show a distance exponent $z = 0$. In contrast to the average density of states, the typical value is able to reflect the LD transition and the corresponding typical order parameter exponent and distance exponent are $\beta_{\text{typ}} = (\nu - D)$ and $\nu_{\text{yp}} = 2(\nu - D)$, respectively (Sec. 5). Here ν is the critical exponent of the localization length. By choosing the typical conductance as a scaling variable an expression for ν in terms of only multifractal exponents can be given, $\nu = (\nu_0 + X^{[q]})^{-1}$. The multifractal analysis yields lower as well as upper bounds for ν in terms of $D(2)$ and ν_0 , respectively (Eqs. (118, 122)). The exponent z describing the distance exponent in the regime $r \ll L; \hat{L}$ is given by $z = d - D(2) \notin \mathbb{Z}$ which is excellently confirmed for the LD transition in quantum Hall systems (Sec. 5).

With the help of conformal mapping arguments in $d = 2$ we suggested a relation between the critical value of the (typical) renormalized localization length ν_c of strip-like systems (being the scaling variable in finite size scaling calculations) and the exponent ν_0 (Eq. (151)). This relation is in agreement with presently available data. A corresponding relation between moments (Eq. (150)) needs further numerical investigations. Such relation is highly desirable since it would allow to demonstrate complete equivalence between the multifractal analysis and the finite size scaling approach.

Finally, we comment on the question of how to determine the critical point T_c by the multifractal analysis. For finite system sizes L the critical regime of values T around T_c , characterized by $L \ll \nu_c$, is finite. Within this regime the $f(q)$ spectra can be calculated by determining linear regimes in log-log plots (see Eqs. (58)). These values will slightly differ from the universal values at T_c . The difference (finite size corrections) will scale, e.g.,⁶³ $\nu_0(T) - \nu_0(T_c) \propto |T - T_c|^{-1}$. Thus, in principle, one can determine T_c and $1/\nu$ from finite size corrections.¹⁷ There have been speculations^{64;65} to determine T_c , in the context of the LD transition, by requiring a certain fractal dimension to coincide with the lower critical dimension d_1 which is believed to be $d_1 = 2$. Since it is known that there is a spectrum of fractal dimensions, such criteria are not evident. To the authors knowledge the only motivation for conjecturing $D(1) = d_1$ as a criterium to $x T_c$ comes from the one-loop result of Wegner's non-linear σ -model in $d = 2 + \epsilon$ ^{19;47} where $D(q) = 2 + \frac{q}{2}$. However, this approximation is expected to deviate from exact values as $\epsilon \rightarrow 1$. Thus, in order to determine T_c by the multifractal analysis one has to establish

universality in multifractal exponents rather than to establish certain values for these exponents.

Acknowledgements

This work was supported by the Deutsche Forschungsgemeinschaft. I thank the Department of Theoretical Physics in Oxford, U.K., for the kind hospitality during a visit in which most part of this work was performed. I gratefully acknowledge helpful discussions with J. Chalker, U. Fastenrath, J. Hajdu, B. Huckestein, W. Pook, L. Schweitzer and M. Zimbauer.

References

1. P.W. Anderson, Phys. Rev. 109, (1958) 1492.
2. P.A. Lee, T.V. Ramakrishnan Rev. Mod. Phys. 57, (1985) 287.
3. A.J.M. McKane, M. Stone, Ann. Phys. 131, (1981) 36.
4. R. Johnston, H. Kunz, J. Phys. C 16, (1983) 3913.
5. E. Abrahams, P.W. Anderson, D.C. Licciardello, T.V. Ramakrishnan, Phys. Rev. Lett. 42, (1979).
6. J.T. Chalker, M. Bernhardt, Phys. Rev. Lett. 70, (1993) 982.
7. U. Fastenrath, Solid State Commun. 76, (1990) 855.
8. K. von Klitzing, G. Dorda, M. Pepper, Phys. Rev. Lett. 45, (1980) 494.
9. R. Prange, S. Girvin (eds.), The Quantum Hall Effect, (Springer, New York, 1990).
10. S.N. Evangelou, T. Ziman, J. Phys. C 20, (1987) L235.
11. U. Fastenrath, G. Adams, R. Bundschuh, T. Hermes, B. Raab, I. Schlosser, T. Wehner, T. Wichmann, Physica A 172, (1991) 302.
12. J.T. Chalker, G. J. Danielli, S.N. Evangelou, I.H. Noh, J. Phys.: Condens. Matter 5, (1993) 485.
13. U. Fastenrath, Physica A 189, (1992) 27.
14. J.L. Pichard, G. Samà, J. Phys. C 14, (1981) L127.
15. A. Maackinnon, B. Kramer Phys. Rev. Lett. 47, (1981) 1546.
16. B. Kramer, K. Broderix, A. Maackinnon, M. Schreiber, Physica A 167, (1990) 163.
17. J. Bauer, T.M. Chang, J.L. Skinner, Phys. Rev. B 42, (1990) 8121.
18. M. Schreiber, H. Grunbach, Phys. Rev. Lett. 67 (1991) 607.
19. C. Castellani, L. Peliti, J. Phys. A 19, (1986) L429.
20. W. Pook, M. Janen, Z. Phys. B 82, (1991) 295; W. Pook, PhD thesis, Universität zu Köln, 1991.
21. B. Huckestein, L. Schweitzer, in High Magnetic Fields in Semiconductor Physics III: Proceedings of the International Conference, Würzburg 1990, ed. G. Landwehr (Springer Series in Solid State Sciences 101, 1992) p. 84.
22. B. Huckestein, B. Kramer, L. Schweitzer, Surface Science 263, (1992) 125.
23. U. Fastenrath, M. Janen, W. Pook, Physica A 191, (1992) 40.
24. A.W. W. Ludwig, Nucl. Phys. B 330, (1990) 639.
25. B. Duplantier, A.W. W. Ludwig, Phys. Rev. Lett. 66, (1991) 247.
26. K.B. Efetov, Adv. Phys. 32, (1983) 53.
27. A.M.M. Pruisken, in Ref. 9, p. 119.
28. F. Wegner, Z. Phys. B 51, (1983) 279.
29. H.E. Stanley, Phase Transitions and Critical Phenomena (Clarendon, Oxford, 1971).
30. J.T. Chalker, G. J. Danielli, Phys. Rev. Lett. 61, (1988) 593.

31. J. T. Chalker, *J. Phys. C* **20**, (1987) L493.
32. B. L. Altshuler, V. E. Kravtsov, I. V. Lerner, *Sov. Phys. JETP* **64**, (1986) 1352.
33. B. L. Altshuler, V. E. Kravtsov, I. V. Lerner *Phys. Lett. A* **134**, (1989) 488.
34. B. Shapiro, *Phys. Mag. B* **56**, (1987) 1031.
35. B. B. Mandelbrot, *The Fractal Geometry of Nature* (W. H. Freeman, San Francisco, 1983).
36. U. Frisch, G. Parisi, in *Turbulence and Predictability of Geophysical Fluid Dynamics and Climate Dynamics*, Proceedings of the International School of Physics \Enrico Fermi\ LXXXV III, Varena, ed. M. Ghil (North Holland, New York, 1985).
37. P. Meakin, A. Coniglio, H. E. Stanley, T. A. Witten, *Phys. Rev. A* **34**, (1986) 3325; M. E. Cates, T. A. Witten, *Phys. Rev. A* **35**, (1987) 1809.
38. T. C. Halsey, M. H. Jensen, L. P. Kadano, I. Procaccia, B. I. Shraiman *Phys. Rev. A* **33** (1986) 1141.
39. B. B. Mandelbrot, *J. Fluid Mech.* **62**, (1974) 331.
40. H. G. E. Hentschel, I. Procaccia, *Physica D* **8**, (1983) 435.
41. T. Tel, *Z. Naturforsch.* **43a**, (1988) 1154.
42. A. Chhabra, R. V. Jensen, *Phys. Rev. Lett.* **62**, (1989) 1327.
43. B. Huckestein, L. Schweitzer, *Helv. Phys. Acta* **65**, (1992) 317.
44. J. T. Chalker, *J. Phys. C* **21**, (1988) L119.
45. H. Aoki, *J. Phys. C* **16**, (1983) L205.
46. H. Aoki, *Phys. Rev. B* **33**, (1986) 7310.
47. F. Wegner, *Z. Phys. B* **36**, (1980) 209; F. Wegner, *Nucl. Phys. B* **316**, (1989) 663.
48. G. Paladin, A. Vulpiani, *Phys. Rep.* **156**, (1987) 147.
49. S. Hikami, *Prog. Theor. Phys.* **76**, (1986) 1210.
50. B. Huckestein, L. Schweitzer, *Phys. Rev. Lett.* **72** (1994) 713.
51. B. Huckestein, B. Kramer, *Phys. Rev. Lett.* **64**, (1990) 1437.
52. L. Schweitzer, 1992 (private communication).
53. D. J. Thouless, *Phys. Rep.* **13C**, (1974) 93.
54. J. T. Chayes, L. Chayes, D. S. Fisher, T. Spencer, *Phys. Rev. Lett.* **57**, (1986) 299.
55. J. M. Kosterlitz, D. J. Thouless, *J. Phys. C* **6**, (1973) 1181.
56. F. Wegner, in *Phase Transitions and Critical Phenomena*, Vol. **6**, eds. C. Domb, M. S. Green, (Academic, London, 1976).
57. L. Schweitzer, 1993 (private communication).
58. M. E. Cates, J. M. Deutsch, *Phys. Rev. A* **35**, (1987) 4907.
59. J. L. Cardy, *J. Phys. A* **17**, (1984) L385.
60. J. L. Pichard, G. Samra, *J. Phys. C* **14**, (1981) L617.
61. B. Huckestein, 1993 (preprint).
62. U. Fastenrath, 1992 (unpublished).
63. B. Huckestein, L. Schweitzer, *Physica A* **191**, (1992) 406.
64. M. H. Cohen, E. N. Economou, C. M. Soukoulis, *Phys. Rev. Lett.* **51**, (1983) 1202.
65. M. Schreiber, *Phys. Rev. B* **31**, (1985) 6146.