

Tricritical behavior of the frustrated XY antiferromagnet

M. L. Plumer, A. M. Ailhot, and A. Caille

Centre de Recherche en Physique du Solide et Département de Physique
Université de Sherbrooke, Sherbrooke, Québec, Canada J1K 2R1

(May 1994)

Extensive histogram Monte-Carlo simulations of the XY antiferromagnet on a stacked triangular lattice reveal exponent estimates which strongly favor a scenario of mean-field tricritical behavior for the spin-order transition. The corresponding chiral-order transition occurs at the same temperature but appears to be decoupled from the spin-order. These results are relevant to a wide class of frustrated systems with planar-type order and serve to resolve a long-standing controversy regarding their criticality.

75.40 Mg, 75.40 Cx, 75.10 Hk

The nature of phase transitions in frustrated systems which can be mapped onto spin models has been studied extensively over the past twenty years by means of renormalization-group (RG) methods and Monte-Carlo (MC) simulations [1]. The triangular antiferromagnet serves as the simplest example of geometry-induced frustration, which in this case gives rise to the non-collinear 120° spin structure. This type of magnetic order can also be described as a helically polarized spin density. The appropriate Landau-Ginzburg-Wilson Hamiltonian is usually taken as

$$H = rS^\dagger S + rS^\dagger S + rS^\dagger S + U_1(S^\dagger S)^2 + U_2 jS^\dagger S^2 j$$

where S is a complex vector and there are two fourth-order terms as a consequence of frustration. One reason for the plethora of studies of this Hamiltonian is that in addition to describing helical spin systems, it is also relevant to the dipole phase of superfluid ^3He , Josephson-junction arrays in a transverse magnetic field, as well as the fully frustrated bipartite lattice (see [1] and [2] for references). Although some of the earlier RG studies suggested a continuous transition within standard universality classes, others found evidence for a first-order transition. In view of these results, the tricritical behavior suggested by the histogram MC simulations of the XY stacked triangular antiferromagnet (STAF) reported here may not be too surprising.

Interest in the criticality of frustrated spin systems has recently been enhanced due to the suggestion by Kawamura of new "chiral" universality classes associated with the XY and Heisenberg STAF's [2,3]. This claim is partially supported by arguments which demonstrate that the symmetry of the order parameter V in the XY case involves a discrete two-fold chiral degeneracy as well as that of the two-dimensional rotation group, so that $V =$

$Z_2 \times S_1$ [4]. If this is indeed the relevant symmetry, and if the transition is neither first-order nor tricritical, then the universality class should be different from the standard ones. The strongest support for the existence of new universality classes comes from Kawamura's MC simulations. These were of the conventional type using rather large lattices $L_x \times L_y$ with $L = 18-60$ but with a possibly modest number of Monte Carlo steps per site (MCS), 6-20 runs with 2×10^4 MCS each. Critical exponents were estimated by the conventional "data collapsing" method, which necessitates a simultaneous estimation of the critical temperature. In the Heisenberg case, the reported values are $\nu = 0.24(8)$; $\beta = 0.30(2)$; $\gamma = 1.17(7)$ and $\alpha = 0.59(2)$. These results have recently been corroborated by three different groups using the more accurate finite-size scaling based on histogram MC data [5-7]. In the XY case, the exponent values reported by Kawamura are $\nu = 0.34(6)$; $\beta = 0.253(10)$; $\gamma = 1.13(5)$ and $\alpha = 0.54(2)$. Both of these sets of exponents are quite different from those of any standard class in 3D. However, they are somewhat suggestive of mean-field tricriticality (apart from the values for ν), where $\nu = \frac{1}{2}$; $\beta = \frac{1}{4}$; $\gamma = 1$ and $\alpha = \frac{1}{2}$. Note that it is only Kawamura's estimate for ν in the XY case which coincides with these values within error.

Unique sets of exponents were also found for the chiral-ity order, which Kawamura suggests occurs at the same critical temperature. The coincidence, or not, of the two critical temperatures has received much attention in corresponding 2D frustrated systems [4,8]. Most authors appear to support the notion that they are the same.

In contrast with the suggestion of new chiral universality, the conclusion of a study of the non-linear sigma model in $2 + d$ dimensions by Azaria et al. [9] is that the criticality of such frustrated systems, at least in the Heisenberg case, is nonuniversal. Depending on unspecified system parameters, one can have either standard $O(4)$ criticality, a first-order transition, or mean-field tricritical behavior. It is natural to extend these arguments to the XY case and speculate either a first-order or tricritical transition [10]. We note that the nature of the chiral transition was not addressed by these authors.

This already confusing situation was recently further exacerbated by the results of Zumbach's local-potential-approximation treatment of the RG [11]. This work emphasizes the possibility of "almost second-order phase transitions" for frustrated systems, where there can be a set of effective critical exponents (also see Refs. [2] and

[12]). At least in the Heisenberg case, the distinction between this possibility and that of a new chiral universality class may never be satisfactorily resolved by MC simulations.

The finite-size scaling of thermodynamic functions evaluated from histogram MC data has demonstrated the ability to yield highly accurate critical-exponent estimates for unfrustrated systems [13,14]. In addition, this procedure when combined with the cumulant crossing method gives an independent and accurate estimate of critical temperatures [13]. This latter feature is particularly useful for frustrated systems in view of the possibility that spin and chiral degrees of freedom order at different, but nearby, temperatures.

It is hoped that the results of MC simulations presented here will provide convincing evidence that the XY STAF exhibits mean-field tricritical behavior.

Near-neighbor antiferromagnetic exchange coupling in the basal plane, $J_z = 1$, and ferromagnetic coupling along the c-axis, $J_k = -1$, were used [15]. The Metropolis MC algorithm was employed in combination with the histogram technique on lattices with $L = 12$ -33 and runs using $1 \cdot 10^6$ MCS for the smaller lattices and $1.2 \cdot 10^6$ MCS for the larger lattices, after discarding the initial $2 \cdot 10^5 - 5 \cdot 10^5$ MCS for thermalization. Averaging was then made over 6 (smaller L) to 17 (larger L) runs. For the largest lattices, this gives a respectable $20.4 \cdot 10^6$ MCS for averaging. The advantage of performing many runs is that errors can be estimated (approximately) by taking the standard deviation. The present work represents one of the very few reports of finite-size scaling of MC data which includes error bars. All histograms were generated at Kawamura's estimate of the critical temperature, $T_c = 1.458$.

The correlation time for the spin order parameter was estimated [16] to be about 620 MCS at $L = 24$ and $T = 1.458$. With the assumption $\tau \propto L^2$, it can thus be expected that averaging was performed using roughly 500 independent configurations [17] in a single run for our largest lattice size. Although τ decreases sharply away from T_c , it remains rather large. At $T = 1.440$, for example, τ was found to be approximately 200 MCS. This result implies that averaging was made using not more than about 8 independent configurations in a single run for Kawamura's simulations at this temperature with $L = 60$.

Results of applying the cumulant-crossing method [13] to estimate the critical temperatures associated with both spin and chiral orderings are presented in Fig. 1. The points represent the temperatures at which the order-parameter cumulant $U_m(T)$ at L^0 crosses the cumulant at $L = 12$ or $L = 15$. There is considerable scatter in the data and care must be taken to use only results with L sufficiently large to be in the asymptotic region where a linear extrapolation is justified [13]. In the case of the spin order, this appears to be for $\ln^{-1}(L^0=L)$

< 1.5 but somewhat larger in the case of chiral order. As with the Heisenberg model, finite-size effects appear much less pronounced for the chiral degrees of freedom [7]. These results suggest that the two types of order occur at the same temperature, $T_c = 1.4584(6)$. The possibility that there are two very close but distinct ordering temperatures can never be ruled-out based on finite-size simulations.

The possibility that the transition is weakly first order was also examined. No evidence for a double-peak structure was found in the energy histograms, consistent with a continuous transition. In addition, the fourth-order energy cumulant U_4 evaluated at T_c yielded a result extrapolated to $L \rightarrow \infty$ of $U_4 = 0.666652(20)$, consistent with a value $\frac{2}{3}$ expected for a continuous transition [18]. A somewhat smaller value occurs in the case of a weak first-order transition [15]. We note, however, that other continuous transitions which occur in this model under the influence of an applied magnetic field have values of U_4 closer to $\frac{2}{3}$ than found in the present case [15]. Finally, the assumption of volume-dependent scaling of various thermodynamic quantities did not yield a straight-line fit, even for the data at large L .

Finite-size scaling results at T_c for the specific heat C , spin order parameter M , susceptibility (as defined in Kawamura's work [3,7,13]), and the first logarithmic derivative of the order parameter $V_1 = \partial \ln(M)/\partial K$ (where $K = T^{-1}$) [14] are shown in Figs. 2-4. Exponent ratios were estimated by performing \ln - \ln plots and also by assuming a scaling dependence $F = aL^x$ for a function of interest (plus a constant term in the case of the specific heat). Except for the specific heat where errors are very large, the two methods gave essentially the same results only if the smaller lattices $L = 12$ and $L = 15$ were excluded. In order to estimate errors due to the uncertainty in T_c , identical scaling was also performed at $T = 1.4579$ and $T = 1.4590$. The resulting exponent ratios, along with those associated with chiral order, are presented in Table I. All results correspond to fits performed on data for $L = 18$ -33, except in the case of the specific heat where $L = 15$ data was also included to reduce the error (otherwise, the result is $\gamma = 0.47(20)$). The given errors represent the robustness of the fitting procedure and do not account for error bars on the figures. Results for the exponents β and γ estimated from the second logarithmic derivative V_2 [7,14] are 0.51(1) and 0.55(1), respectively.

For ease of comparison, the results of Kawamura's work are also included in the table. Note that in order to obtain best-fit exponents by the data-collapsing method, Kawamura used two different values of T_c : 1.458 for spin order and 1.459 for chiral order (within the range of our estimate for T_c). Within errors, however, he concludes that the two transitions were the same.

From the results in Table I, it can be seen that β ; γ ; γ/β ; and γ/β are the most sensitive to the choice of T_c . Vari-

ation in the exponents due to the error bars on the figures was about the same. With these considerations, our best estimate of the exponents and their associated errors at $T_c = 1.4584(6)$ are given by: $\nu = 0.46(10)$; $\nu_\perp = 0.24(2)$; $\nu_\parallel = 1.03(4)$; $\nu_\perp = 0.50(1)$ for spin order, and $\nu = 0.38(2)$; $\nu_\perp = 0.90(9)$; $\nu_\parallel = 0.55(1)$ for chiral order. These results for the spin order strongly suggest that the transition is mean-field tricritical.

A possible interpretation for this behavior can be found in our recent results of applying an in-plane magnetic field [15]. This work reveals that T_c is a multicritical point, with one phase having the symmetry of the three-state Potts model and consequently a weak first-order transition. The effect of the field is to generate a term third-order in S in the free energy, of the form $m S^3$, where m is the $q = 0$ Fourier component of the spin density induced by the field. Since the system is frustrated in the triangular plane, one might expect that short-range order along the c -axis is already well developed at temperatures near T_c , in the present case corresponding to the spin component m . At zero applied field, coupling between this short-range order and S could generate a third-order term which influences the critical behavior. For other types of systems, there may be similar coupling to other Fourier components [19].

It is also noteworthy that even though our simulations on the XY model were made with about a factor of ten more MCS than in the Heisenberg system [7], larger fluctuations were observed in the present case for the spin order. This can be observed by comparing the cumulant-crossing data of Fig. 1 with the corresponding results of Ref. [7]. Larger fluctuations are expected if the transition is at or near a tricritical point.

Our results for the chiral order are more difficult to interpret. The values for the exponents are not too different from those of Kawamura (when the errors are accounted for), and do not correspond to any known universality class (also see Ref. [8]). The estimates for ν are in agreement but are significantly different from our value of ν for the spin order. This may indicate that the chiral order has a distinct correlation length and that its criticality is decoupled from the spin order [8,10]. Another possibility [20] currently under investigation is that chirality is associated with one of many possible symmetry-breaking fields, in a manner analogous to the infinite set of crossover exponents previously determined for the XY model [21].

In conclusion, these results of extensive histogram MC simulations of the stacked triangular XY antiferromagnet strongly support the scenario of tricritical behavior associated with the spin order, in agreement with Azzaria et al. and in contrast with the proposal of new XY-chiral universality by Kawamura. (Recent MC simulations do support, however, the existence of a new Heisenberg-chiral universality class [5(7)].) This resolves a long-standing controversy in the literature, and is relevant to a wide

class of frustrated systems. Further work is necessary to fully understand the nature of the chiral ordering transition which appears to be decoupled from the spin order.

ACKNOWLEDGMENTS

We thank P. Azzaria, H. Kawamura and G. Zumbach for useful discussions and the Service de l'Informatique for the partial use of 10 RISC 6000 Workstations. This work was supported by NSERC of Canada and FCAR du Québec.

Present address: INRS-EAU, 2800 rue Einstein, C.P. 7500, Sainte-Foy, Québec, G1V 4C7 Canada.

- [1] Magnetic Systems with Competing Interactions, edited by H.T. Diep (World Scientific, in press).
- [2] H. Kawamura, Phys. Rev. B 38, 4916 (1988); 42, 2610 (E) (1990); J. Phys. Soc. Jpn. 55, 2157 (1986).
- [3] H. Kawamura, J. Phys. Soc. Jpn. 54, 3220 (1985); 55, 2095 (1986); 56, 474 (1987); 61, 1299 (1992).
- [4] D.H. Lee et al., Phys. Rev. Lett. 52, 433 (1984); Phys. Rev. B 33, 450 (1986).
- [5] T. Bhattacharya, A. Billoire, R. Lacaze, and Th. Jolicœur, J. Phys. I (France) 4, 122 (1994).
- [6] D. Loison and H.T. Diep, unpublished.
- [7] A. Mailhot, M.L. Plumer, and A. Caille, Phys. Rev. B (submitted).
- [8] M. Gabay et al., Phys. Rev. B 40, 264 (1989); E. G. Ranto and M.P. Nightingale, Phys. Rev. B 48, 7438 (1993); G. Ramirez-Santiago and J.V. Jose, Phys. Rev. Lett. 68, 1224 (1992); Phys. Rev. B 49 (1994).
- [9] P. Azzaria, B. Delamotte, and Th. Jolicœur, Phys. Rev. Lett. 64, 3175 (1990); P. Azzaria, B. Delamotte, F. Delduc, and Th. Jolicœur, Nucl. Phys. B 408, 485 (1993).
- [10] P. Azzaria, private communication.
- [11] G. Zumbach, Phys. Rev. Lett. 71, 2421 (1993); Nucl. Phys. B 413, 754, 771 (1994).
- [12] P. Peczak and D.P. Landau, Phys. Rev. B 39, 11932 (1989).
- [13] P. Peczak, A.M. Ferrenberg, and D.P. Landau, Phys. Rev. B 43, 6087 (1991).
- [14] A.M. Ferrenberg and D.P. Landau, Phys. Rev. B 44, 5081 (1991), C. Holm and W. Janke, Phys. Rev. B 48, 936 (1993); K. Chen, A.M. Ferrenberg, and D.P. Landau, Phys. Rev. B 48, 3249 (1993).
- [15] M.L. Plumer, A. Mailhot, and A. Caille, Phys. Rev. B 48, 3840 (1993); 49, June 1 (1994).
- [16] S. Tang and D.P. Landau, Phys. Rev. B 36, 567 (1987).
- [17] A. Ferrenberg, D.P. Landau, and K. Binder, J. Stat. Phys. 63, 867 (1991).
- [18] M.S.S. Challa, D.P. Landau, and K. Binder, Phys. Rev. B 34, 1841 (1986).
- [19] O. Heinonen and R.G. Petschek, Phys. Rev. B 40, 9052

(1989); M L. Plumer et al, Phys. Rev. B 47, 14312 (1993).

[20] G. Zum bach, private communication.

[21] A. Aharony et al, Phys. Rev. Lett. 57, 1012 (1986); B. Fourcade and A. M. S. Tremblay, Phys. Rev. B 39, 6819 (1989).

TABLE I. Variation of exponents with assumed critical temperature, along with Kawamura's results.

T_c	1.4579	1.4584	1.4590	Kawamura
	0.47 (8)	0.46 (10)	0.39 (13)	0.34 (6)
	0.23 (1)	0.24 (1)	0.26 (1)	0.25 (1)
	1.07 (2)	1.03 (2)	0.99 (2)	1.13 (5)
	0.51 (1)	0.50 (1)	0.50 (1)	0.54 (2)
	0.36 (1)	0.38 (1)	0.40 (1)	0.45 (2)
	0.96 (2)	0.90 (2)	0.81 (2)	0.77 (5)
	0.56 (1)	0.55 (1)	0.54 (1)	0.55 (2)

FIG. 1. Results of applying the cumulant-crossing method (see text) to estimate the critical temperatures associated with the spin and chiral orderings where $b = L^0 = L$.

FIG. 2. Finite-size scaling of the specific heat data for $L = 12 - 33$. Data at $L=12$ is excluded from the fit. Error bars are estimated from the standard deviation found in the MC runs.

FIG. 3. Finite-size scaling of the order parameter. Data at $L=12$ and 15 are excluded from the fit. Error bars are estimated from the standard deviation found in the MC runs.

FIG. 4. Finite-size scaling of the susceptibility as well as the logarithmic derivative of the order parameter V_1 (see text), as in Fig. 2.