

Simulations of Solid-Solid Models of Spreading of Viscous Droplets

O. Venäläinen¹, T. Ala-Nissila^{1,2}, and K. Kaski¹

¹Tampere University of Technology
Department of Electrical Engineering
P.O. Box 692, FIN-33101 Tampere, Finland

²University of Helsinki
Research Institute for Theoretical Physics
P.O. Box 9 (Siltavuorenpenger 20 C)
FIN-00014 University of Helsinki, Finland
and

Brown University
Department of Physics, Box 1843
Providence, R.I. 02912, U.S.A.

May 26, 1994

Abstract

We have studied the dynamics of spreading of viscous non-volatile fluids on surfaces by Monte Carlo simulations of solid-on-solid (SOS) models. We have concentrated on the complete wetting regime, with surface diffusion barriers neglected for simplicity. First, we have performed simulations for the standard SOS model. Formation of a single precursor layer, and a density profile with a spherical cap shaped center surrounded by Gaussian tails can be reproduced with this model. Dynamical layering, however, only occurs with a very strongly attractive van der Waals type of substrate potential. To more realistically describe the spreading of viscous liquid droplets, we introduce a modified SOS model. In the new model, tendency for dynamical layering and the effect of the surface potential are in part embedded into the dynamics of the model. This allows a relatively simple description of the spreading under different conditions, with a temperature like parameter which strongly influences the droplet morphologies. Both rounded droplet shapes and dynamical layering can easily be reproduced with the model. Furthermore, the precursor width increases proportional to the square root of time, in accordance with experimental observations.

PACS numbers: 68.10.Gw, 05.70.Ln, 61.20.Ja.

Keywords: Droplet Spreading, Dynamics of Wetting, Solid-on-Solid Model

1 Introduction

Surface processes such as wetting [1], coating and adhesion depend strongly on phenomena occurring in the first few layers on top of the substrate. Recent progress in optical techniques [2-8] has focussed attention to the dynamics of spreading of tiny, viscous nonvolatile droplets on solid surfaces. With ellipsometry, measurements can now be done on molecular scales. They have revealed fascinating phenomena previously unaccounted for in macroscopic droplets, such as dynamical layering and diffusive spreading of the precursor layer [2, 4, 7, 8].

In the experiments, the most common liquids have been tetrakis (2-ethylhexoxy) silane, squalane, and polydimethylsiloxane (PDMS). It has been shown [2] that the thickness profiles of tetrakis and PDMS droplets on a silicon wafer exhibit strikingly different shapes under spreading. On a "low energy" surface (with a relatively low solid-vacuum surface tension [1]) tetrakis produces clear dynamical layering (Fig. 1(a)), i.e. stepped shapes of exactly one molecular layer in thickness. In contrast, on the same substrate the spreading of PDMS proceeds by a fast evolving precursor of about one molecular layer in thickness (Fig. 1(b)).

On the other hand, the same liquid on different surfaces has been shown to form distinct morphologies [4]. On a "high energy" surface (with a relatively high surface tension and adsorption energy [1]) PDMS develops dynamical layering, too, in analogy to tetrakis. On a low energy surface the formation of a single precursor layer with a central cap is observed [4]. Other experiments show [6, 8] that PDMS droplets form stepped, dynamically layered profiles on two different silicon wafers (Fig. 1(c)). In addition, a recent experiment

of PDM S spreading on a silver substrate shows [5] the formation of a profile with spherical cap shaped center and Gaussian tails at late times (Fig. 1(d)).

On the theoretical side progress has been more moderate. The continuum hydrodynamic approach [1, 9, 10, 11], while being able to qualitatively explain the precursor film and its eventual diffusive ($\propto t^{1/2}$) growth [11] is simply not valid for the measurements on scale of Angstroms. The first theoretical model to successfully account for dynamical layering was given by Abraham et al. [12]. They used the two dimensional horizontal solid-on-solid (SOS) model, which describes coarse grained layers of fluid emanating from a droplet "reservoir". Dynamical layering was observed to occur with an attractive van der Waals type of surface potential in the complete wetting regime [12, 13]. The model also predicts diffusive behaviour for the bulk layers of the droplet on a high energy surface, but linear time dependence for the precursor layer.

A different approach was recently taken by de Gennes and Cazabat [14], who proposed an analytic model of an incompressible, stratified droplet. The stepped morphology of the spreading droplet is postulated a priori, and spreading is assumed to proceed by a longitudinal flow of the molecular layers, and a transverse permeation flow between them. The model introduces an effective Hamaker constant, which has been fitted [6] to experimental data on stepped PDM S droplets.

On a more microscopic level, molecular dynamics simulations have been performed [15, 16, 17]. Yang et al. [16] studied droplets of Lennard-Jones (LJ) particles and dimers on corrugated surfaces, surrounded by a vapor phase. They observed terraced droplet shapes in the complete wetting regime, but

obtained logarithmically slow growth for the layers. In contrast, Nieminen et al. [17] were able to qualitatively observe a crossover from "adiabatic" to "diffusive" spreading for LJ droplets on a smooth surface. To account for the chainlike molecular structure of some of the liquids, Nieminen and Ala-Nissila [18] recently carried out a systematic study of droplets of LJ solvent particles and flexible oligomers on smooth surfaces. By varying the strength of the surface potential, they were able to qualitatively conclude that the actual molecular structure of the liquid can play an important role in the formation of the density profiles, even beneath the entanglement regime [19]. Namely, relatively weaker substrate interactions and longer chains favour more rounded droplet shapes, due to the mixing of the molecular layers under spreading. On the other hand, a very strong surface attraction promotes layer separation and leads towards stepped shapes. In the latter case, the spreading process qualitatively proceeds in analogy to the ideas of Ref. [14], with layers of fluid "leaking" at the edges of the steps. They also observed a crossover from "adiabatic" to "diffusive" behaviour of the precursor film and developed a scaling form for its time dependence. These results are in good qualitative agreement with experiments; however, droplets consisting of just a few thousand molecules were simulated, while experimental droplets are microscopic in the vertical direction only.

In the present work we have taken to approach the problem of droplet spreading on a more coarse-grained level, in order to further clarify the role of the effective viscosity of the liquid, and the surface potential in the droplet morphologies. We have carried out systematic Monte Carlo simulations with two different SSS models to study the dynamics of the spreading process. In these models, the droplet is assumed to consist of effective, coarse-grained

blocks of molecules [12]. The size of these blocks can be highly anisotropic between the vertical and horizontal directions, depending on the details of the interactions. All the simulations have been done in the complete wetting regime in the canonical ensemble, where the final state of the droplet is a surface monolayer. No evaporation from the surface is allowed, and in this work the surface diffusion barriers have been neglected for simplicity.

The first model that we have studied is the standard vertical SOS model [20, 21]. Besides computing density profiles of the droplets, we have also followed the time evolution of the width of the precursor $l_m(t)$, and the droplet height. Our results demonstrate that with a strongly attractive short range substrate potential, only a single precursor layer evolves, with the center of the droplet assuming a spherical cap shape rather than a Gaussian profile. At later times, a surface monolayer forms. These results are analogous to the observations in Ref. [21]. In addition, formation of a spherical cap with Gaussian tails occurs with relatively weak attractive short range substrate potentials. To obtain stratified droplet shapes corresponding to dynamical layering, however, an extremely strong long range van der Waals type of surface potential must be used. In addition, the behaviour of the precursor l_m was not observed to follow the correct dynamical behaviour. This is due to the somewhat unrealistic dynamics of the model.

To better take into account the fluid-like nature of the spreading droplets, we have developed a modified SOS model in which the tendency for dynamical layering and the effect of the surface potential have been partially embedded into the dynamics of the model, in analogy with some of the ideas presented by de Gennes and Cazabat [14]. In its simplest form, the model enables a description of the spreading process with a single temperature like parameter.

When varied, this parameter strongly influences the droplet morphologies and reproduces both rounded and stepped droplet shapes. In particular, dynamical layering can be reproduced without explicitly including a long range van der Waals substrate attraction. When such an attraction is added the range of layering is further enhanced, as expected. Our quantitative results also show that the precursor width increases proportional to the square root of time. A preliminary account of these results has been published in Ref. [22].

2 Standard Solid-on-Solid Model of Spreading Dynamics

2.1 Definition of the Model

The standard solid-on-solid (SOS) model used in our simulations is defined by the following Hamiltonian [23]:

$$H(\{h_i\}) = \frac{\chi}{2} \sum_{i,j} \min(h_i, h_j) + \sum_{i=2}^N h_i + \sum_i V(h_i); \quad (1)$$

where in the first term $\chi < 0$ is the nearest neighbour interaction parameter and the summation goes over the four nearest neighbours of the column i whose height is h_i . The heights, which measure the average concentration of particles and obey a vertical SOS restriction [20] are integers. The second term on the right hand side of Eq. (1) describes the vertical interaction between the effective molecules. $V(h_i)$ denotes the strength of an attractive, height dependent substrate potential at site i . $V(h_i)$ is defined as

$$V(h_i) = \frac{A}{h_i^3} + B h_i h_{i+1}; \quad (2)$$

where $A < 0$ is an effective Hamaker constant, and $B < 0$ describes the interaction between the substrate and the fluid.

In the model, the effective particles follow standard Monte Carlo (MC) Metropolis dynamics with transition probability

$$P(h_i \rightarrow h_i^0) = \min[1; e^{-H/k_B T} g]; \quad (3)$$

where H is the energy difference between the final (h_i^0) and initial (h_i) states (with the SOS restriction obeyed). Time is measured in units of MC steps per site (MC S/s). The initial configuration of the droplet is usually chosen to be a three-dimensional cube, which after a transient time assumes its characteristic shape. During the spreading stage, we calculate the density profile of the droplet (in analogy to the experiments), the time dependence of the precursor width $r(t)$ and the height of the droplet. All the simulations have been done without allowing evaporation from the surface; thus the final state is a surface monolayer. In this work the activation energy for surface diffusion of isolated particles is neglected, and thus the model may not be applicable to some high energy surfaces.

2.2 Results for the Standard Model

First, we study the spreading dynamics without the van der Waals potential by setting $A = 0$ in Eq. (2). In this case, the results are not very sensitive to the droplet size, and in most cases a $11 \times 11 \times 10$ droplet was used, where ten is the initial height. We also $\chi = 0.05$ in this section. With a relatively strong surface attraction, only a single precursor film is formed, with a rounded central cap. At the edge of the film the migration of the effective molecules causes the formation of a monolayer of separate particles.

This happens typically in the range $\beta = j = 2.0 - 4.0$, and $j k_B T = j = 0.6 - 1.0$. These results are in complete agreement with the qualitative observations of Ref. [21]. Additionally, we find that the time development of the precursor \ln approximately follows $r(t) \sim t^{0.14}$, in contrast to $t^{1/2}$. If we then lower the surface attraction and set $B = 0$, and $j k_B T = j = 0.6$, the precursor \ln vanishes and the droplets assume a rounded, spherical shape with Gaussian tails as shown in Fig. 2.

Next, we study the influence of the long range van der Waals type potential. For a moderately large A , the results are virtually indistinguishable from those above. Namely, for $A = B = 0.15$, a single precursor \ln forms while for $A = B = 1$ the \ln disappears ($j k_B T = j = 0.6$). However, for a very large surface attraction $A = B = 1.0$, the droplets show indications of separation of layers. At $j k_B T = j = 1.0$ shown in Fig. 3(a), a small central cap is formed at submonolayer heights. When temperature is further lowered to 0.6, three separate layers can be observed (Fig. 3(b)). We have calculated $r(t) \sim t$ for the lower temperature case, and find ~ 0.17 for both $A = B = 0.15$ and 1.0 . We also find that at very late times, the height of the submonolayer \ln decreases approximately as t^{-2} . This describes the decrease of the almost two dimensional island, which leaks into a surface monolayer. This monolayer which spreads diffusively consists of all the particles, which are two or more lattice sites apart from the cluster. Thus the precursor \ln width decreases and the areal density of the monolayer correspondingly increases. We note that due to the lack of surface diffusion activation energy, the case of larger surface attraction enhances the rate of spreading, which is opposite to what happens on high energy surfaces [4, 18].

From the results presented above, it is clear that the standard SOS model

can produce dynamical layering only for extremely large values of the long range surface attraction in the model. In fact, although standard SOS models have been very useful in describing solid surfaces and their roughening [20], and even growth of surface layers [23], it is clear that they are less realistic for fluid spreading. For example, the diffusion length of the effective particles is not well defined, due to the fact that very large height differences between nearest neighbour are allowed. Second, nonmonotonic variations in the density profiles are possible, although not significant for the parameter range used in the present work. Due to these limitations, in the next section we shall introduce a modified SOS model which better accounts for the fluid-like nature of the spreading droplets.

3 Modified Solid-on-Solid Model of Spreading Dynamics

3.1 Definition of the Model

The basic idea behind the modified SOS model is to more realistically describe the viscous flow of the droplets. We have done this by modifying the dynamics of the standard SOS model defined by Eqs. (1)–(2) in the following way. First, the effective molecules are not allowed to climb over height barriers, or jump down over terrace edges. For example, two individual particles located initially at different sites on the surface can never form a column of height $h_i = 2$. Second, when the difference between the neighbouring initial and final sites is $(h_i - h_i^0) \geq 2$, a transient non-SOS excitation (a hole or vacancy) can be created into the initial column by a particle which jumps to the neighbouring position with a probability

$$P = \min_i (h_i - h_j - 1) \theta(h_i - h_j - 1) e^{-H_t/k_B T} g \quad (4);$$

where $\theta(x)$ is the Heavyside step function. The energy difference H_t is calculated for this transient state. The hole left behind is immediately filled by the column of particles above it, which are all lowered by a unit step. This means that $H_t > H$ (as calculated for the final state). We note that the effective dynamics used here does not satisfy microscopic reversibility at all times.

The motivation for the unusual dynamics comes from the physics of dissipative spreading of viscous fluids on attractive surfaces. Microscopic calculations have revealed [18] that at least in the layered state, the droplets spread by the viscous flow of rather well separated layers, which "leak" on the surface at the terrace edges. Furthermore, the analytic model for stratified droplets by de Gennes and Cazabat [14] assumes, that the horizontal layers of fluid flow with friction, and that there is a narrow region of permeation flow at the terrace edges. In our model, this is roughly described by the creation of the transient hole at the step edges. Since this is temperature controlled (cf. Eq. (4)), we expect that the model be able to describe droplet spreading under a variety of different conditions.

3.2 Results

First, we present results for $A = B = \gamma = 0$, which corresponds to an infinite temperature. The effective dynamics in the model renders even this simplest case nontrivial. As seen in Fig. 4, at early and intermediate stages of spreading the droplet assumes a non-Gaussian, spherical shape with Gaussian tails surrounding the center of the droplet. At early times the spreading

is relatively fast, but slows down at later times as the droplet density profile approaches a Gaussian shape at submonolayer heights. In contrast, we note that an infinite temperature for the standard SOS model always leads to Gaussian shaped profiles. At late times, the width of the precursor film approaches the expected $t^{1/2}$ behaviour.

Next, we set $A = B = 0$, $\beta = 0.05$ and let the temperature ratio $k_B T =$ vary. For the lowest temperature of 0.6, dynamical layering of the density profiles is immediately evident in Fig. 5(a). For larger droplets, up to five or six layers can easily be detected (Fig. 5(b)). In this case the droplet maintains a rather compact shape and the surface monolayer formation is relatively slow. At later stages (not shown in Fig. 5) a flat profile is formed with a tiny central cap at the center. This cap disappears at approximately after 6000 MCS/s. The precursor width follows $t^{0.50(3)}$. We also studied the spreading of an initially ridge-shaped droplet of size $120 \times 11 \times 10$, where the precursor width follows $t^{0.75}$ rather than $t^{1/2}$. In this case the spreading is highly anisotropic and rather different from the three-dimensional droplet.

When temperature is raised, the layer formation weakens and droplets become more spherical. For $k_B T = 0.8$ (Fig. 6) and 1.0, the precursor film grows as $t^{0.49(7)}$ and $t^{0.47(6)}$, respectively. The surface monolayer formation due to enhanced thermal fluctuations is faster, and for the higher temperature the whole precursor film tends to break down. In Fig. 7 we show the time development of the droplet height at the three different temperatures. The fast spreading rate, and consequently the enhanced edge flow leading to more rounded shapes at higher temperatures is clearly reflected in the results.

Next we study the spreading with a long range van der Waals type of surface

potential with $\beta A = \beta B = 1$. In this case simulations have been performed for $B = 0.025$, $B = 0.05$, $B = 0.07$, and $B = 0.2$. The temperature was set to $k_B T = 0.6$. In our model with modified dynamics, the inclusion of the van der Waals energy enhances the tendency for dynamical layering, as demonstrated in Fig. 8. Furthermore, the results indicate clearly that with $B = 0.2$ the precursor film breaks down, giving for the time development of the precursor film width the behaviour $t^{0.3}$. With the other choices, however, the time dependence of the precursor width follows $t^{1/2}$ rather accurately. For $B = 0.2$ spreading becomes unrealistically fast due to the lack of the surface diffusion activation barriers, which have been set to zero. As mentioned earlier, we do not expect the model to be applicable to high energy surfaces if the diffusion barriers control the dynamics.

4 Discussion of the Results and Comparison with Experiments

Next, we analyze the density profiles and make comparisons with relevant experiments [2-8], whenever possible. Due to the coarse-grained nature of the SSS models quantitative comparisons may not be meaningful, however. The aim here in part has been to elucidate the physical processes behind different droplet morphologies, and thus we will also discuss our results in connection to other coarse-grained models [12, 13, 14, 21], and microscopic simulations [16, 17, 18].

First, we discuss results for the standard SSS model. The model qualitatively reproduces single precursor film formation with a relatively strong attractive short range substrate potential, while with a weaker potential, rounded droplet shapes with Gaussian tails appear. This is in accord with

the notion that increased surface attraction should enhance the precursor ℓ_m [4, 5, 12, 13]. The precursor ℓ_m was also qualitatively observed in the simulations of Ref. [21]. Also, with the inclusion of a long range van der Waals potential, the droplets exhibited tendency towards dynamical layering. This is in accordance with the horizontal SSS model results of Refs. [12, 13], and qualitatively with the microscopic picture as well [17, 18]. Although the standard SSS model can reproduce density profiles resembling experimentally observed ones (Fig. 3(a) [3] and Fig. 3(b) [2, 6]) the strength of the long range potential must be considered unrealistically large as compared with the particle-particle interactions on the surface level. Most importantly, the time dependence for the width of the precursor ℓ_m does not follow the expected $t^{1/2}$ behavior, which reflects the unrealistic nature of the dynamics in the model.

In contrast, the modified SSS model yields results which seem much more realistic. At low temperatures, stepped droplet shapes are obtained (Figs. 5(a) and (b)). On the other hand, when particle motion from one layer to another is more likely, rounded droplet shapes occur (Figs. 4 and 6). This is in qualitative accord with conclusions from coarse-grained models [12, 13, 14] and microscopic calculations [17, 18]. Furthermore, the inclusion of a van der Waals potential further enhances the range of layering, as expected. The results are also in qualitative agreement with experiments [4, 5, 6] where increasing the viscosity of the liquid enhances layering, and layering is also more prominent on high energy surfaces. Most importantly, the model reproduces the expected $t^{1/2}$ dependence for the width of the precursor layer. However, it should be noted again that since the model neglects diffusion barriers, our results may not be applicable for all cases.

The remarkable feature of the modified SOS model is that by just varying a single temperature like parameter, droplet morphologies from rounded to stepped shapes can be easily reproduced. This can be understood based on the physical analogy behind the modified dynamics [14]. High effective temperatures in the model correspond to enhanced flow at the edges, and reduced viscosity and surface interactions allowing transient hole formation with relative ease. On the other hand, lowering the temperature suppresses this hole formation which is analogous to suppressing interlayer flow in the phenomenological model of de Gennes and Cazabat [14]. In fact, our model gives further support to the physical ideas behind their model regarding dynamical layering.

5 Summary and Conclusions

To summarize, in this work we have tried to unravel physical processes behind different morphologies observed in the spreading of tiny liquid droplets on surfaces. The present approach of using coarse-grained descriptions of the fluid is complementary to other approaches, and can indeed give a lot of physical insight to the problem. In particular, by introducing a new, more realistic coarse-grained lattice SOS model of fluid spreading we have demonstrated how the interplay between the internal viscosity of the fluid and the nature of the surface attraction can produce a number of different droplet shapes, in qualitative agreement with experiments. Our results also give further support to ideas behind the analytic models, in particular that of Ref. [14] regarding the spreading of terraced, stratified droplets. They are also in agreement with microscopic studies of Refs. [17, 18]. However, the experimental situation remains complicated and should not be oversimplified

{ there exist very few attempts to systematically study the effects of surface attraction, microscopic structure of the liquid, finite size effects and other factors, which can play an important role in spreading dynamics. We hope that the present work inspires more systematic studies in these directions.

Acknowledgements: We wish to thank R. Dickman, S. Herminghaus, and J. A. Nieminen for useful discussions, and the Academy of Finland for financial support. We also wish to thank P. Leiderer for sending his data on the density profiles of PDM S spreading on silver.

Figure captions

Fig. 1. (a) Experimental thickness profiles of tetrakis on a "low energy" silicon surface [2]. Dynamical layering occurs at late times. (b) Thickness profiles of PDM S on the same surface (the "Mexican hat" shape) [2]. For even smaller droplets, the central cap becomes more distinct and a Gaussian shape is assumed at latest times [3]. (c) Thickness profiles of PDM S on a silicon wafer covered by a grafted layer of trimethyl groups. This surface behaves as a "low energy" surface [8], with strong dynamical layering occurring. (d) Thickness profiles of PDM S on a silver surface after 50, 80, 150, and 360 minutes [5]. The submonolayer profiles can be fitted by a spherical cap, with Gaussian convolution at the edges.

Fig. 2. Density profile for the standard SOS model, with $A = 0$, $B = 1$, and $j_B T = j = 0.6$. The droplet assumes a rounded profile, with no precursor (cf. Fig. 1(d)). The initial height of the droplet is 41 units, and the result is an average over 50 runs after 25 000 MCS/s. The solid line indicates a Gaussian fit.

Fig. 3. Spreading droplet profiles for the standard SOS model with $A = B = 1$, and (a) $j_B T = j = 1$. At submonolayer heights, a central cap forms. The curves correspond to 800, 1200, 1400 and 1600 MCS/s, and are averages over 500 runs. The initial size of the droplet is $11 \times 11 \times 10$. (b) Results for $j_B T = j = 0.6$. Dynamical layering tends to occur, with three layers visible. The curves correspond to 1600, 2200, and 3800 MCS/s, with averages over 300{400 runs.

Fig. 4. The density profile of a spreading droplet for the modified SOS model at an infinite temperature, after 650 MCS/s. A distinctly non-Gaussian, spherical shape occurs above monolayer heights, as evidenced by a Gaussian fit shown with a solid line. The initial height of the droplet is 41 units, and the profile is an average over 500 runs.

Fig. 5. Density profiles for the modified SOS model at $k_B T = \epsilon = 0.6$. Dynamical layering of the profiles occurs after an initial transient. (a) Results for a $11 \times 11 \times 10$ droplet, with curves corresponding to 1800, 2800, and 3800 MCS/s. (b) Results for a $11 \times 11 \times 40$ droplet, with curves at 5800, 6800, and 8000 MCS/s. All profiles are averages over 500 runs.

Fig. 6. Density profiles for the modified SOS model at $k_B T = \epsilon = 0.8$. A shoulder and a central cap are clearly seen, with a spherical cap shaped center and a Gaussian foot at late times. The curves correspond to 2000, 2200, and 2600 MCS/s. Results are averages over 500 runs.

Fig. 7. The height of the droplets for the modified SOS model at the three different temperatures studied, as a function of time. Filled squares correspond to $k_B T = \epsilon = 0.6$, open circles to 0.8, and filled circles to 1.0. Breaks in the curves correspond to separate layers. While the spreading rate strongly increases as a function of temperature, the dynamical layering considerably weakens and droplets become more rounded.

Fig. 8. Density profiles for the modified SOS model, with $\beta_A = \beta_B = 1$, $k_B T = \epsilon = 0.6$, and $B = 0.2$. A stronger van der Waals potential clearly enhances dynamical layering (cf. Fig. 5). See text for details.

References

- [1] P. G. de Gennes, *Rev. Mod. Phys.* 57, 827 (1985).
- [2] F. Heslot, N. Fraysse, and A. M. Cazabat, *Nature* 338, 640 (1989).
- [3] F. Heslot, A. M. Cazabat, and P. Levinson, *Phys. Rev. Lett.* 62, 1286 (1989).
- [4] F. Heslot, A. M. Cazabat, P. Levinson, and N. Fraysse, *Phys. Rev. Lett.* 65, 599 (1990).
- [5] U. Albrecht, A. Otto, and P. Leiderer, *Phys. Rev. Lett.* 68, 3192 (1992); *Surface Sci.* 283, 383 (1993).
- [6] M. P. Valignat, N. Fraysse, A. M. Cazabat, and F. Heslot, *Langmuir* 9, 601 (1993).
- [7] N. Fraysse, M. P. Valignat, A. M. Cazabat, F. Heslot, and P. Levinson, *J. Coll. Interface Sci.* 158, 27 (1993).
- [8] J. DeConinck, N. Fraysse, M. P. Valignat, and A. M. Cazabat, *Langmuir* 9, 1906 (1993).
- [9] L. Leger and J. F. Joanny, *Rep. Prog. Phys.* 55, 431 (1992).
- [10] L. H. Tanner, *J. Phys. D* 12, 1473 (1979).
- [11] J. F. Joanny and P. G. de Gennes, *J. Physique* 47, 121 (1986).
- [12] D. B. Abraham, P. Collet, J. de Coninck, and F. Dunlop, *Phys. Rev. Lett.* 65, 195 (1990); *J. Stat. Phys.* 61, 509 (1990).
- [13] D. B. Abraham, J. Heinio, and K. Kaski, *J. Phys. A* 24, L309 (1991); J. Heinio, K. Kaski, and D. B. Abraham, *Phys. Rev. B* 45, 4409 (1992); E. Cheng and C. Ebner, *Phys. Rev. B* 45, 12056 (1992).
- [14] P. G. de Gennes and A. M. Cazabat, *C. R. Acad. Sci. Paris* 310, Ser. II, 1601 (1990).
- [15] J. Hautman and M. L. Klein, *Phys. Rev. Lett.* 67, 1763 (1991).

- [16] J. X. Yang, J. Koplik, and J. R. Banavar, Phys. Rev. Lett. 67, 3539 (1991); Phys. Rev. A 46, 7738 (1992).
- [17] J. A. Nieminen, D. B. Abraham, M. Karttunen, and K. Kaski, Phys. Rev. Lett. 69, 124 (1992); J. A. Nieminen, A. Lukkarinen, K. Kaski, and D. B. Abraham, unpublished (1993).
- [18] J. A. Nieminen and T. Ala-Nissila, Europhys. Letters 25, 593 (1994); University of Helsinki preprint HU-TFT-93-62 (to appear in Phys. Rev. E (1994)).
- [19] P. G. de Gennes, Scaling Concepts in Polymer Physics (Cornell University Press, Ithaca, N.Y. 1979); M. Doi and S. F. Edwards, The Theory of Polymer Dynamics (Oxford University Press, Oxford, 1986).
- [20] H. J. Leamy, G. H. Gilmer and K. A. Jackson, in Surface Physics of Materials, vol. I, ed. J. M. Blakely (Academic Press, New York 1975).
- [21] J. Heinio, K. Kaski and D. B. Abraham, Phys. Scripta T 38, 28 (1991).
- [22] O. Venalainen, T. Ala-Nissila, and K. Kaski, Europhys. Letters 25, 125 (1994).
- [23] O. Venalainen, J. Heinio, and K. Kaski, Phys. Scripta T 38, 66 (1991); O. Venalainen, Phys. Scripta 48, 487 (1993).