

The Fractional Quantum Hall Effect on a Lattice.

F.F. Assaad⁽¹⁾ and S. Biskamp⁽²⁾

⁽¹⁾ ISSP, The University of Tokyo, Roppongi, Minato-ku, Tokyo 106, Japan.

⁽²⁾ Physikalisches Institut, Universität Würzburg 97074 Würzburg, FRG.

Abstract

Starting from the Hofstadter butterfly, we define lattice versions of Landau levels as well as a continuum limit which ensures that they scale to continuum Landau levels. By including a next-neighbor repulsive interaction and projecting onto the lowest lattice Landau level, we show that incompressible ground states exist at filling fractions, $\nu = 1/3; 2/5$ and $3/7$. Already for values of $l_0 = a^2$ where $l_0(a)$ is the magnetic length (lattice constant), the lattice version of the $\nu = 1/3$ state reproduces with nearly perfect accuracy the continuum Laughlin state. The numerical data strongly suggests that at odd filling fractions of the lowest lattice Landau level, the lattice constant is an irrelevant length scale. We find a new relation between the hierarchy of incompressible states and the self-similar structure of the Hofstadter butterfly.

There has been a considerable amount of numerical work on the fractional quantum Hall effect [1] (FQHE) for continuum models on planar or spherical geometries. The question we wish to address in this paper is: can one observe the FQHE on a lattice and if so, how does it compare to continuum results? Since on the lattice rotational invariance is broken, the pseudopotential [2] approach to the FQHE is not applicable and there is a priori no reason to expect that ground states at certain filling fractions of Hofstadter [3] bands are characterized by a specific power law decay of the density-density correlation $g(r)$ for small values of r . On the other hand there is an efficient way to construct incompressible quantum liquids (IQL) states which implicitly assumes the existence of Laughlin states [4] on a lattice: the parton approach [5,6]. The Hamiltonian which has parton wave functions as mean-field solutions can be derived from first principles only on a lattice [5]. Hence, this approach requires that properties of the Laughlin states survive discretization of the plane. We show that even for small lattice sizes there exist incompressible states at odd denominator filling fractions of the lowest Hofstadter level if it has the Landau level degeneracy. These states reproduce with remarkable accuracy the relevant properties of the Laughlin states.

Our starting point are fully spin-polarized electrons on a square lattice in a magnetic field:

$$H_0 = \sum_{\langle i,j \rangle} t_{ij}^X c_i^y c_j^y \exp \left[\frac{2}{\phi_0} \int_i^j \tilde{A} d\mathbf{l} \right] \quad (1)$$

Here, the sum runs over next-neighbors, c_i^y creates a spinless electron on site i , $n_i = c_i^y c_i^y$ and in the Landau gauge $\tilde{A}(\mathbf{x}) = B(x_2; 0; 0)$. The square lattice lies in the $\mathbf{e}_1, \mathbf{e}_2$ plane and ϕ_0 is the flux quantum. In order to put the system on a torus, we impose the boundary conditions: $c_{i+\mathbf{L}}^y = \exp \left[\frac{2i}{\phi_0} \int_{\mathbf{L}} \tilde{A} \right] c_i^y$. Here, \mathbf{L} is a vector with norm equal to the linear size of the square lattice oriented along the \mathbf{e}_1 or \mathbf{e}_2 direction, and $\tilde{A}(\mathbf{x} + \mathbf{L}) = \tilde{A}(\mathbf{x}) + \tilde{\Phi}_{\mathbf{L}}(\mathbf{x})$. To obtain a single-valued wave function the total flux traversing the lattice has to be an integer multiple of the flux quantum [7]. The spectrum of the above Hamiltonian yields the Hofstadter butterfly [3]. To make contact with the corresponding continuum Hamiltonian, we define the field operators $c_i^y = a^{-R} \int d\mathbf{r} (\mathbf{r} - \mathbf{R}_i)^{-y}(\mathbf{r})$. Expanding the phase factors in

equation (1) and the delta-function up to second order in the lattice constant, one obtains up to a constant term : $H_0 = \frac{\hbar^2}{2m} \int d\mathbf{r} \nabla^2 \psi(\mathbf{r}) + \frac{e}{c} \mathbf{A}(\mathbf{r}) \cdot \nabla \psi(\mathbf{r})$. Setting $\hbar^2/2m = 1$ where m denotes a constant mass, yields the desired continuum limit. The continuum Hamiltonian has an energy spectrum described by Landau levels: $E_n = \hbar \omega_c (n + 1/2)$ where $\omega_c = eB/mc$. Each Landau level has a macroscopic degeneracy given by $N = L^2 B / \phi_0$ and the magnetic length is given by $l_0 = \sqrt{\hbar / 2eB}$.

To achieve such a situation on the lattice, we set

$$\frac{p}{q} = \frac{B a^2}{\phi_0} = \frac{1}{mN} \text{ for a } N \times N \text{ lattice:} \quad (2)$$

Since the number of flux quanta traversing the lattice, $N = N/m$, has to be an integer, N has to be divisible by m . Before arguing why this choice of $p=q$ is suitable, let us note that in order to obtain a well defined continuum limit for a fixed value of m , we have to set B or equivalently l_0 to a constant. This yields: $a = \sqrt{\phi_0 / N}$ as well as $L = \sqrt{\phi_0 N}$. Hence, our continuum limit corresponds to a combined continuum ($a \rightarrow 0$) and thermodynamic ($L \rightarrow \infty$) limit. Since we want to keep the filling fraction $\nu = N_p/N$ constant during the scaling to the combined continuum and thermodynamic limit, $N_p = N^2 \nu = N$. Here, N_p denotes the number of particles on the lattice.

For the above choice of $p=q$ and in the Landau gauge, the boundary conditions are: $c_{1+N a_1} = c_1$ and $c_{1+N a_2} = \exp(-2i\pi) c_1$. Here, $\mathbf{r} = a(\mathbf{x}; n)$. We define the partial Fourier transform $c_k^{(n)} = \frac{1}{N} \sum_{\mathbf{x}} e^{ik \cdot \mathbf{x}} c_{a(\mathbf{x}; n)}$, $k = 2\pi n_k / N$ where $n_k = 1 :: N$ to write the Hamiltonian (1) as:

$$H_0 = \sum_k \sum_n \left(2t \cos \frac{2\pi n}{mN} + k \cdot \frac{y_k^{(n)}}{k} - \frac{y_k^{(n)}}{k} \frac{y_{k+n}^{(n+1)}}{k} \right) + h.c. \quad (3)$$

$\frac{y_k^{(n+N)}}{k} = \frac{y_{k+k}^{(n)}}{k+k} \text{ where } k = 2\pi m :$

The boundary condition in the n index of the operators mixes m k -sectors. Hence, the resulting Hamiltonian is block diagonal and contains $N = N/m$ blocks, each of dimension $mN \times mN$. We label the blocks with the quantum number k_0 , and denote the $mN \times mN$ matrices by $A(k_0)$. For this special choice of $p=q$ (2), one may show that the matrices

$A(k_0)$ are equivalent so that the spectrum of $A(k_0)$ is independent of k_0 . Therefore, each energy level of the Hamiltonian (3) is at least N -fold degenerate. Numerically, one may see that the eigenvalues of one $A(p)$ matrix are up to accidental degeneracies at zero energy, non-degenerate. The spectrum of the Hamiltonian (3) is thus described by N m , N -fold degenerate energy levels. We identify those energy levels to lattice Landau levels (LL). To proceed further, we have to diagonalize numerically the matrices $A(k_0)$ so as to obtain the single particle wave functions spanning a given LL. For more general values of $p=q=\frac{n}{m}N$, the Hamiltonian again splits into N/m block diagonal matrices of dimension $mN \times mN$. For $n > 1$ these matrices are not equivalent, and the spectrum is better described as mN , N/m -fold degenerate bands. One may see numerically that as the lattice size is increased, n low-lying adjacent bands merge very rapidly together to form a Landau level [8].

Having defined LL, one may now compare their properties with those of the continuum Landau levels. Consider the magnetic length which is given in the continuum by: $l_b = \frac{2}{\phi_0} \sqrt{\frac{\hbar}{eB}}$. For our choice of $p=q$ the former equation is equivalent to $a=l_b = \frac{1}{\sqrt{2}} \sqrt{\frac{\hbar}{mN}}$. In order to obtain an estimate of the magnetic length on the lattice, one notes that for a filled lowest Landau level (i.e. $\nu=1$) [9]:

$$\frac{N_p}{N} \langle n(n+1) \rangle = 1 + \exp\left(-\frac{r^2}{2\lambda_0^2}\right) \quad (4)$$

In order to obtain the magnetic length on a finite lattice, we have fitted our lattice-data for the density-density correlations (lhs of the above equation) to the continuum form (rhs of the above equation). We consider two directions: $\mathbf{r} = \mathbf{r}(1;1)$ and $\mathbf{r} = \mathbf{r}(1;0)$. For both directions, and already for small lattice sizes (i.e. $N = 12$) the data fits the above form extremely well for all considered values of m [10]. Figure 1 plots the lattice magnetic-length for both considered directions (symbols) as well as its value predicted by the continuum equations (solid line). As may be seen, the agreement, already for small lattice sizes is very good. It is to be noted that the rotational invariance present in the continuum is violated by the presence of the square lattice. However, this symmetry is restored since as may be seen the discrepancy between the two lattice values of l_b vanishes rapidly. To

understand this striking result we have considered the continuum eigenfunctions of H_0 and the corrections due to finite lattice constant and volume. We find that the decay of the eigenfunctions remains gaussian in any order of perturbation theory. Similarly, one may check that the cyclotron frequency scales rapidly to its continuum value. Hence, the above defined LL, reproduce very accurately the physics of the continuum Landau levels already on small lattice sizes.

We show that IQ L states exist on the lattice by considering a next-neighbor density-density interaction:

$$H_I = \frac{V}{2} \sum_{\langle i,j \rangle} :n_i n_j: + \frac{V a^4}{2} \int dx : (\psi)^\dagger \psi (\psi)^\dagger \psi : \quad (5)$$

Here, $:O:$ denotes the normal ordering of the operator O . A similar Hamiltonian was considered in reference [11] in the context of anyon superconductivity. For this continuum interaction (rhs of the above equation), it has been shown that Laughlin wave function is exact [12]. Therefore, we should ultimately reproduce the physics of the Laughlin wave function.

In order to carry out the calculations, we project the Hamiltonian onto the lowest LL. This projection is justified in the limit $B \rightarrow 1$. The resulting Hamiltonian is an effective one dimensional Hamiltonian which one may solve numerically by using a standard implementation of the Lanczos method. The projected Hamiltonian has the same symmetries (conservation of the one-dimensional total momentum) as the Hamiltonian considered by Yoshioka et al. [13]. The scaling of the hopping matrix element, t , and interaction strength, V , follow from the continuum limit: $V = V_0 N^2$ and $t = t_0 N$ where V_0 and t_0 are constants independent of the lattice size. The former follows from the fact that $a^{-1} = \frac{p}{N}$.

For the numerical calculations we chose $m = 1$ so that $N = N$. This choice gives the smallest magnetic length, and hence lattice effects should be maximized. The following refers to this special case. Our numerical analysis is based on the calculation of the quantity

$E_0(N_p=N) - E_0(N_p=N-1) \sim N^{-1}$ where $E_0(N_p=N)$ is the ground state energy of the projected Hamiltonian for an N -site chain with N_p particles and $E_0(N_p=N-1)$ is the single particle

energy of the first LL. The filling fraction is given by $\nu = N_p/N$. Since the first LL is dispersionless $E_0(N_p=N)$ is independent of t . Hence only interaction effects are taken into account. We search for incompressible states by computing the energy gap to adding a full electron,

$$E_p(\nu) = E_0\left(\frac{N_p+1}{N}\right) - 2E_0\left(\frac{N_p}{N}\right) + E_0\left(\frac{N_p-1}{N}\right) \quad (6)$$

as well as the energy gap to adding one flux quantum,

$$E_o(\nu) = E_0\left(\frac{N_p}{N-1}\right) - 2E_0\left(\frac{N_p}{N}\right) + E_0\left(\frac{N_p}{N+1}\right) : \quad (7)$$

Figure 2, plots a summary of our results for various filling fractions and as a function of the lattice size. Several points may be made: i) The gap to adding a full electron shows more fluctuations as a function of lattice size than the gap to adding a flux quantum.

$E_o(\nu = 2/5)$ and $E_o(\nu = 1/3)$ show a very smooth behavior as a function of lattice size. This provides convincing data for the occurrence of incompressible states at $\nu = 1/3$ and $\nu = 2/5$ on the lattice. The order of the gaps to adding a flux quantum is correct:

$E_o(\nu = 1/3) > E_o(\nu = 2/5) > E_o(\nu = 3/7)$ [14]. Note that we obtained for particle excitations: $E_p(\nu = 1/3) > E_p(\nu = 3/7) > E_p(\nu = 2/5)$ (data not presented). Finally, due to particle hole symmetry within the lowest LL identical gaps at $\nu = 2/3$, $\nu = 3/5$, and $\nu = 4/7$ were observed. ii) At $\nu = 1/3$, and in the spirit of the Laughlin wave function, removing one flux quantum is equivalent to adding a quasiparticle. Since a full electron is composed of three quasiparticles, one expects if the interaction between quasiparticles is small:

$E_o(\nu = 1/3) \approx E_p(\nu = 1/3) = 3$. This is confirmed in figure 2. However, it appears that $3E_o(\nu = 1/3)$ is slightly smaller than $E_p(\nu = 1/3)$ which support the existence of an attractive interaction between the quasiparticles. Hence, the quasiparticle picture seems to be justified also on the lattice. iii) The gap at $\nu = 1/2$ shows very large fluctuations as a function of system size. From our numerical data, we clearly may not obtain reliable information on the nature (incompressible or fermi liquid [15]) of the $\nu = 1/2$ ground state for the considered short range interactions.

Figure 3 plots the density-density correlation functions (4) for the $\nu = 1/3$ incompressible state on a 24×24 lattice. The lattice results (solid circles) are compared to the prediction of the Laughlin wave function [16] (solid line). For this lattice size, $l_0 = a/2$ and it is remarkable to see that there is already nearly perfect agreement between the lattice and continuum results. Figure 3 also plots the quasiparticle $g(r)$ which is obtained by removing a flux quantum from the $\nu = 1/3$ state. $g(r)$ for the quasiparticle state is also a smooth function at small distances, and as expected decreases for $r \rightarrow 0$ with a smaller power law than the $\nu = 1/3$ state.

The numerical data show explicitly that the Hamiltonian $H_0 + H_I$ is scale invariant at the numerically accessible odd denominator filling fractions. The lattice constant is an irrelevant length scale, and the characteristic correlation functions scale with l_0 , the only length scale involved. Hence, the choice of the interaction (5) is a renormalization group fixed point.

Mean field theories for partially filled LL at $\nu = \frac{1}{2m+1}$ can be obtained by decomposing the electron operators into an odd number of fermionic partons with charges $\frac{e}{2m+1}$, $\psi_r = Q_m \psi_{r-1}$, subjected to the constraints $\psi_r^\dagger \psi_r = \psi_{r-1}^\dagger \psi_{r-1} ; 8 ; [5]$. If each species fills up the LL with $N = \frac{N}{2m+1}$ the mean field decoupling of the product of parton operators is stable if the interaction between the electrons is repulsive and short ranged [5]. The parton wave functions have $2m+1$ components when we put them onto a torus. On one hand, this implies that the product wave function is periodic and hence is a candidate for an incompressible electron state [6]. On the other hand the constraints on the parton currents induce fractional statistics [5]. The important point is that in our approach the correct multivaluedness of anyons on a torus [18] is automatically provided. The hierarchy of IQH states, which we partially observe on the lattice, can be understood analogously to the continuum situation. The continual condensation of localized quasiparticle excitations into Laughlin states relies at the mean field level on the self-similarity of the Hofstadter spectrum. This property assures that the levels at smaller magnetic field strengths, which are populated by quasiparticle excitations, have the same overall structure. The continuum one particle spectrum is trivially self-similar, since in this case the energies depend linearly

on the magnetic field strength. By the introduction of the lattice the hierarchy of IQ L states is unfolded.

To conclude, we have considered tight-binding spinless electrons on a square lattice in a strong magnetic field and submitted to a next-neighbor repulsive interaction. We defined lattice Landau levels (LL) as well as a continuum limit which insures that they scale to continuum Landau levels. By projecting onto the lowest LL and already on lattices where $l_0 = a^2$ the FQHE at odd filling fractions was observed with remarkable precision. Those observations open a new region, the continuum edge of the Hofstadter spectrum, where the FQHE may be observed and studied.

We are very indebted to A. Muramatsu who suggested this approach to the FQHE. We would like to thank W. Hanke, M. Imada, C. Kubert, K. Kusakabe, F. V. Kusmartsev and S. Meixner for motivating discussions. F.F.A. would like to thank the DFG for financial support under the grant number Ha 1537/6-2 as well as the Japanese Ministry of Education and S.B. the financial support of the BMFT under the grant number 03-HA 3W UE.

Figure captions

Fig. 1 The magnetic length as a function of the lattice size. An estimate of l_b on the lattice (symbols) is obtained by fitting the lattice value of the density-density correlation functions to their continuum value (4) in the case of a filled first Landau level [10]. We consider two lattice directions, $\mathbf{r} = \mathbf{r}(1;0)$ (downward triangles) and $\mathbf{r} = \mathbf{r}(1;1)$ (circles) as well as $m = 1;3$ and 5 . The solid lines correspond to the continuum value: $l_b = L = \frac{p}{m} \frac{p}{2N}$ where N corresponds to the linear size of the lattice.

Fig. 2 Gaps to adding a full electron $\epsilon_p(\mathbf{r})$ (6) as well as a flux quanta $\epsilon_0(\mathbf{r})$ (7) for various fillings and as a function of the lattice size. The data refers to the special case $m = 1$ (see equation (2)).

Fig. 3 Density-density correlation functions for the $\nu = 1/3$ state on a 24×24 lattice (solid circles). The open circles plot $g(r)$ for the quasiparticle on a 23×23 lattice. The solid line corresponds to $g(r)$ as obtained from the Laughlin wave function at $\nu = 1/3$ [16]. The data refers to the special case $m = 1$ (see equation (2)).

REFERENCES

- [1] The Quantum Hall Effect, edited by R.E. Prange and S.M. Girvin, Springer Verlag, 1990.
- [2] F.D.M. Haldane, Phys. Rev. Lett. 51, 605, (1983).
- [3] D.R. Hofstadter, Phys. Rev. B 14, 2239, (1976).
- [4] R.B. Laughlin, Phys. Rev. Lett. 50, 1395, (1983).
- [5] X.G. Wen, Phys. Rev. Lett. 66, 802, (1991).
- [6] J.K. Jain, Phys. Rev. B 40 8079, (1989)
- [7] E. Fradkin, Field Theories of Condensed Matter Systems, Addison-Wesley Publishing Company, 1991. Chapter 9, page 256.
- [8] For the special case $n=m = 2=5$ and $N = 10$ the $N = 4$ lowest single particle energy levels are degenerate up to 10^{-12} .
- [9] B. Jancovici, Phys. Rev. Lett. 46, 386, (1981).
- [10] The sum of square errors per data point for an $N = 15$ lattice, $m = 5$ and for both considered lattice directions was of the order of 10^{-4} . This corresponds to the worst case.
- [11] G.S. Canright and S.M. Girvin, Int. J. of Mod. Phys. B 3, 1943, (1989).
- [12] S.A. Trugman and S. Kivelson, Phys. Rev B 31, 5280, (1985).
- [13] D. Yoshioka, B.I. Halperin and P.A. Lee, Phys. Rev. Lett. 50, 1219, (1983). D. Yoshioka, Phys. Rev B 29, 6833, (1984).
- [14] N. d'Ambrosio and R.H. Morf, Phys. Rev. B 40 6108, (1989).
- [15] B.I. Halperin, P.A. Lee, N. Read, Phys. Rev. B 47, 7321, 1993.

[16] $g(x)$ for the Laughlin wave function at $\nu = 1/3$ was obtained by using the analogy to the two-dimensional one-component plasma. (See reference [17].) Our results were obtained through the use of classical Monte-Carlo methods for 256 particles.

[17] J.M. Caillol, D. Levesque, J.J. Weiss and J.P. Hansen, J. of Statistical Mechanics, 28, 325, (1982). D. Levesque, J.J. Weiss and A.H. MacDonald, Phys. Rev. B 30, 1056, (1984).

[18] T. Einarsson, Phys. Rev. Lett. 64, 1995, (1990)




Figure 1




Figure 2




Figure 3