

OUTP-9419P
cond-mat/9409011

Polymer Collapse on Fluctuating Random Surfaces

Simon Dalley

Department of Physics, Theoretical Physics
Oxford University, Oxford OX1 3NP, U.K.

Abstract

The conformations of interacting linear polymers on a dynamical planar random lattice are studied using a random two-matrix model. An exact expression for the partition function of self-avoiding chains subject to attractive contact interactions of relative strength $1=c$, $0 < c < 1$, is found as a function of chain length L and c . The number of configurations $L^a e^{bL}$ as $L \rightarrow \infty$ is determined, showing that a chain undergoes a third-order collapse transition at $c = \frac{1}{2}$; the universal scaling laws are found.

1 Introduction

A long polystyrene chain molecule in a solvent undergoes a coil-to-globule transition [1] at a finite temperature $T = T_c$ due to attractive polymer-polymer interactions. Many approximate theoretical techniques have been developed to describe this and other processes of macromolecular folding [2]. In this letter exact results for this problem of polymers, modeled by self-avoiding chains with contact interactions, will be given on the ensemble of two-dimensional simplicial lattices with random fluctuations of the local intrinsic curvature which appear in the study of two-dimensional quantum gravity [3, 4]. This ensemble represents an annealed average over a certain class of fractals. The advantage is that statistical mechanics problems are often exactly solvable in this case using random matrix models [4], which yield expressions for the thermodynamic and some of the geometrical quantities of interest. Moreover it is intuitively plausible, and found to be the case in all of the many solved examples, that the phase structure of such statistical systems is the same as on a fixed lattice but with weaker transitions when one averages over fluctuating lattices. Although a simple structureless polymer will be studied here, the same techniques could be extended to more complicated polymer types.

A two-dimensional simplicial lattice is obtained by gluing, pairwise along edges, A squares each of edge length a , so as to form a surface with spherical topology say. A polymer chain of L links walks isotropically on the edges of squares, cannot cross itself (excluded volume), but can occupy a given edge any number of times (figure 1); the latter represent contacts. The partition function is a sum over all possible simplicial lattices and configurations of a polymer on such lattices with a weight for contacts. The β -temperature is found exactly in this model and represents a third-order collapse transition. One is therefore led to predict that on a fixed two-dimensional lattice (e.g. honeycomb), with the rules of contact given in Fig.1, the transition will be first or second order.¹ The universal exponent a , defined by the number of configurations as $L \rightarrow \infty$, $L^a e^{bL}$, changes from $a = 5/2$ at $T > T_c$ to $a = 3/2$ at

¹ Some previous exact results [5] on fixed lattices have used a different definition of contact, forbidding entirely configurations 'a' and 'b' of Figure 1; a contact occurs when two non-adjacent parts of the polymer come within one lattice spacing. This could make a difference in the collapsed regime since the polymer can be much more compact if multiple occupation of an edge is allowed (type 'a' in Figure 1). Such multiple occupation necessarily occurs, for example, in the high-temperature expansion of the lattice Ornstein model with action $S[\phi]$

$T = \infty$. For $T < \infty$ it was not possible to determine a. The universal behaviour around $T = \infty$ is described in terms of a reduced temperature $T = L^{-1/6}$.

2 Matrix Model.

The self-avoiding walk on fluctuating random surfaces was described using a one-matrix model by Duplantier and Kostov [6]. In order to include an attractive interaction whenever contacts occur consider a two-matrix model with hermitian $N \times N$ matrices M and N ,

$$Z = \int dM dN \exp \left[-N \text{Tr} \left(\frac{1}{2} M^2 + \frac{1}{2} N^2 - c \frac{g(1 - c^2)^2}{4} M^4 + (1 - c^2)^L N^{2L} \right) \right] : \quad (1)$$

As usual, the Feynman diagram representation of the expansion in g generates graphs dual to simplicial lattices and as $N \rightarrow 1$ those of genus zero dominate. c is a chemical potential conjugate to the number of polymers n , each of length L , i.e. the surfaces are built from 4 vertices dual to squares while the polymers from M matrices. The last term in the action $\text{Tr} V(M; N)$ represents a hole of length $2L$ in the surface with one point on the boundary marked; one can visualize the polymer by sewing up the hole starting at the marked point. Each possible hole and each possible location of the marked point give a different polymer configuration. The polymer is connected into the surfaces by c so that c represents a polymer-lattice Boltzmann weight, while the polymer-polymer (contact) weight c^2 has been normalised to one. Thus $0 < c < 1$ sets the temperature scale, with $c = 1$ being infinite temperature.

$$Z_{N \rightarrow 1} = \sum_{w, n, A} c^w n! g^A H(A; w; n; L) \quad (2)$$

where the sum is over genus zero surfaces and H is the number of configurations with n polymers, each of length L with w polymer-contacts, on all surfaces of area A . Since H grows as const^A there exists a critical $g = g_c$ at which large area surfaces become significant and a universal continuum limit can be established [4].

The matrix model (1) can be solved in the large N limit by the method of orthogonal polynomials [7]. The polynomials $P_i(x) = x^i + \dots$ $Q_j(x) = x^j + \dots$ are such that

$$\int_1^{Z+1} dx d e^{V(x)} P_i(x) Q_j(x) = h_{ij} \quad (3)$$

and the free energy is $\log Z = \text{const} \sum_{i=1}^N (N - n) \log f_i$, where $f_i = h_i/h_{i-1}$ and the constant is g -independent. P and Q satisfy recursion relations

$$P_i(\epsilon) = P_{i+1} + r_i P_{i-1} + s_i P_{i-3} + \dots \quad (4)$$

$$Q_j(\epsilon) = Q_{j+1} + q_j Q_{j-1} + t_j Q_{j-3} + \dots \quad (5)$$

That for P terminates after $L + 1$ terms. From identities such as

$$\sum_i \frac{d}{d\epsilon} e^{V(\epsilon)} P_{i-1}^0 Q_i = 0 \quad (6)$$

one can find a closed set of equations for r, s, q, t and f which can be solved for f . In the large N limit one can define a continuous variable $x = i/N$ and functions $f_i = N f(x); r_i = N r(x);$ etc. and these equations are explicitly:

$$cr = f \sum_i \frac{P_i}{f} [i; i-1] \quad (7)$$

$$cq = f - 3g(1 - c^2)^2 fr \quad (8)$$

$$ct = -g(1 - c^2)^2 f^3 \quad (9)$$

$$x = -cf + q \sum_i \frac{P_i}{f} [i-1; i]; \quad (10)$$

where

$$[i; j] = 2L(1 - c^2)^L \sum_i \frac{d}{d\epsilon} e^{V(\epsilon)} \frac{P_i}{h_i} - 2L - 1 \frac{Q_j}{h_j} : \quad (11)$$

Substituting eqs. (7) and (8) into eq. (10) gives an equation (the 'string equation' in quantum gravity) for f . It will be evaluated shortly for large N .

The critical point for continuum surfaces is identified by setting $\epsilon = 0$ (no polymers) and finding the singularity in the solution for $f(1)$ of the quadratic equation (10) as g is varied, yielding $g_c = \frac{1}{12}$ and $f_c = 2c/(1 - c^2)$. Defining $g = g_c(1 - \epsilon^2)$, where ϵ is the lattice spacing, ϵ is the renormalised 2D 'cosmological constant' conjugate to renormalised surface area $A = A/\epsilon^2$. In the limit $\epsilon \rightarrow 0$, ϵ couples to universal continuum surfaces (finite) and surfaces finite in lattice units ($A = 0$). The partition function will have a part regular in ϵ , usually representing the latter surfaces, and a non-analytic part representing the former. It is

appropriate to define scaling variables $x = 1 - z^{-2}$ and $f(x) = f_c(1 - u(z))$ in which case, in the limit $\epsilon \rightarrow 0$,

$$u^2 = \epsilon + z \quad (12)$$

from eq.(10) at $\epsilon = 0$ and

$$\log Z = N^{-2/5} \int_0^{Z^{-1/5}} dz zu + \text{regular} \quad (13)$$

3 Polymer Collapse.

First the case of a single polymer will be addressed, when eq.(12) is appropriate and one studies the expectation value of $\text{Tr}(z^L)$ in this theory, i.e. Z expanded to first order in ϵ ,

$$Z = Z_0 + Z_1 + \dots,$$

$$\begin{aligned} Z_1 &= \int \mathcal{D}D \mathcal{D}d (1 - c^2)^L \text{Tr}(z^{2L}) e^{-N \text{Tr}(V(\epsilon=0))} \\ &= (1 - c^2)^L \int \prod_{i=1}^N d d e^{-V(\epsilon=0)} \prod_{i=1}^P \frac{P_i}{h_i} z^{2L} \prod_{i=1}^Q \frac{Q_i}{h_i} : \end{aligned} \quad (14)$$

Thus, if $Z_1 = \sum_{i=1}^P Z_1^0$, from eqs.(5) (3) (8) and (9) one finds

$$\begin{aligned} Z_1^0(L; c; u; \epsilon) &= (1 - c^2)^L (2L)! \epsilon^L \sum_{p=0}^{[L/2]} \frac{(c - 3gf)^L - 2p (gf)^p c^{3p-2L}}{(L-2p)! (p+L)! p!} \\ \frac{gf}{c} &= \frac{1 - c^2}{6} (1 - u + O(\epsilon^2)) \end{aligned} \quad (15)$$

($[L/2]$ indicates greatest integer $L/2$). In order that the polymer be an extended object on the surfaces of finite A one must scale its length as $L = L\epsilon$ in which case

$$Z_1^0 = G(L; c) e^{-(c)u + O(\epsilon)} + O(\text{const } O(\epsilon^{-1})) \quad (16)$$

where $G \sim L^a e^{b(c)L}$. Despite many attempts the author has not managed to derive the asymptotic form (16) for all c . It is generally expected however that the number of configurations has the form of G as $L \rightarrow \infty$, which numerical evaluation of (15) confirms. Figure 2 plots (c) obtained from

$$u_L = \log \frac{Z_1^0(L; c; 0)}{Z_1^0(L; c; u_L)} \quad (17)$$

for large but finite L | corrections to the right hand side above are $O(\epsilon^2)$.

$G = \lim_{L \rightarrow \infty} Z_1^0(L; c; 0)$ is a kind of wavefunction renormalisation which represents the effects of surfaces of zero renormalised area, $A = O(\epsilon^2)$; it does not depend upon u or ϵ . On the other hand u represents an induced free energy per unit length of the polymer due to fluctuations of the finite A surfaces; it is non-analytic in ϵ from eq.(12). The behaviour of $G(c)$ thus governs the interesting geometrical properties of the polymer. In particular, $G(c)$ seems to show a transition to $G(c) = 0$ at $c = 0.4$, which will be verified shortly. Physically, as c is reduced the polymer favours polymer-polymer rather than polymer-solvent bonds and so the induced free energy falls. Below a certain $c_c = 0.4$ the polymer collapses entirely and is only connected into the surface by a small number of polymer-solvent bonds. This is the coil-to-globule transition on fluctuating random surfaces. The polymer-polymer contacts also induce free energy per unit length but at a lower order in ϵ , and it will be necessary to choose a different scaling law for L in order to see this contribution near $c = c_c$.

In order to find c_c exactly, determine the universal exponent a , and the details of the phase transition, it is useful to rewrite Z_1^0 as

$$Z_1^0 = (1 - c^2)^L \left[\frac{dp}{2} p \frac{1}{f} e^{-ip} + p \frac{q}{f} e^{-ip} + \frac{t}{f^{3/2}} e^{3ip} \right]^{2L} \quad (18)$$

As the continuum limit is approached it is expedient to rescale $p \rightarrow p \epsilon^d$ for some exponent d . Expanding in ϵ one has

$$= \frac{1}{2c} [a_1 + a_2 i p^d + a_3 u + a_4 p^{2-2d} + a_5 i u p^{d+1} + a_6 u^{2-2} + a_7 \epsilon^2 + \dots] \quad (19)$$

Consider first the high temperature $c \rightarrow 1$ limit. It is known from the $c = 1$ solution [6] that $d = 1/2$, so this ansatz will be used in the high temperature (coil) phase. Writing $c = 1 - \epsilon$ for small ϵ , the coefficients a_i can be found as power series in ϵ ; in particular $a_2 = O(\epsilon)$ and one finds from (18)

$$Z_1^0 = \frac{1}{4} [8 + O(\epsilon)] \exp \left[u L \left(1 - \frac{4}{3} \epsilon + O(\epsilon^2) \right) \right] \quad (20)$$

in the continuum limit $\epsilon \rightarrow 0$. This is of the form (16) with a that correctly matches onto

the answer of Fig. 2. The partition function itself is then

$$Z_1 / N^{5/2} \frac{\text{const}^L}{L^{3/2}} p^{-} + \frac{1}{L} e^{-p^{-}} \quad (21)$$

using eq.(12). Large area $A \rightarrow 1$ corresponds to $\beta \rightarrow 0$, in which case the universal power exponent is $a = 5/2$. If c is reduced by a finite amount from one, a_2 is finite and the resultant oscillatory behaviour means that it is no longer possible to neglect the higher orders in p as $\beta \rightarrow 0$. Nevertheless one expects the form (21) to persist down to $c = c_c$.

Fortunately there is one other value of c where a_2 vanishes, given by $c = \frac{p}{2} - 1$; a_3 also vanishes at this point indicating that this is the collapse temperature. A non-trivial scaling solution is obtained with the scaling laws $c = \frac{p}{2} - 1 + \frac{1}{L^3}$, $L = L^2$, and $d = 1$ (note that polymer length scales as area now). In this case one derives from (18)

$$Z_1 / \text{const}^L N^{3/2} \frac{e^{Z_1}}{L^{3/2}} dy e^{-3yL^2} \frac{1}{L^{3/2}} \quad (22)$$

in the continuum limit, where $a = 3/2$ at $\beta = 0$. As the reduced temperature $\beta \rightarrow +1$ one recovers the form (21) with $\beta \rightarrow 3$ matching in the high temperature phase (g2). At $\beta = 0$, Z_1 , while now analytic in β , in this case seems to represent finite A surfaces. $A = 0$ surfaces $| A = 0$ (2) more precisely $|$ are characterised by the fact that a finite number of derivatives with respect to β can remove their contribution, since each derivative brings down a factor of A . But $Z_1(\beta = 0)$ cannot be removed by differentiating. It can be interpreted as a surface dense with polymer whose area is proportional to the polymer length; for small L differentiating does remove its contribution. It is tempting to extrapolate into the collapsed phase $\beta \rightarrow 1$ also, but the above scaling laws do not seem to be consistent since Z_1 becomes a saddle point integral with the result $Z_1 \sim (L^3=1) \exp(-L)$. This cannot be correct because Z_1 should be monotonically decreasing with β .

If polymers are present with fugacity one must go back to eq.(12) and modify it accordingly. From eq.(8) and eq.(10) one has

$$x = cf + \frac{f}{c} \frac{3gf^2}{c^2} + \frac{6gf^{3/2}}{c^2} [i; i-1] \frac{p}{2} \frac{1}{f} [i-1; i] \quad (23)$$

In order that the new terms contribute to the scaling part of the equation, but do not disturb the critical condition for continuum surfaces, $g_c = \frac{1}{12}$ to leading order in β , one must tune

! 0 appropriately. For $c > \frac{p-2}{2} - 1$ one finds at order u^2 in eq.(23)

$$0 = \frac{1}{2} + z - u^2 - \frac{p-1}{2} e^{-1/u} : \quad (24)$$

The wavefunction renormalisation G of Z_1 has been absorbed in \bar{g} and a renormalised fugacity defined, of dimension $5=2$. When $\bar{g} \neq 0$ the singularity at $g = g_c$ ($\bar{g} = 0$) is now shifted by an $O(u^2)$ term, $\bar{g} = \bar{g}_c$ say, which therefore couples to renormalised area A . \bar{g}_c represents the free energy of the polymer system per unit A on finite surfaces. The positive root of eq.(24) is the correct one since $u \rightarrow \frac{p-2}{2} + z$ as $\bar{g} \rightarrow 0$. A singularity in u appears when $2u = \frac{1}{2} p^{3/2} e^{-1/u}$. Then $u(\bar{g})$ is singular at $\bar{g} = \bar{g}_c$. In particular at $c = \frac{p-2}{2} - 1 + \epsilon$, $\epsilon > 0$, one finds $u(\bar{g}_c) = \frac{1}{2} p^{3/2} e^{-1/u} + O(\epsilon^2)$ and $\bar{g}_c = \frac{p-1}{2} u^3 + O(\epsilon^4)$. For $\epsilon < 0$, $\bar{g} = 0 = \bar{g}_c$, so that $\partial^3 \bar{g}_c / \partial c^3$ is discontinuous implying that the collapse transition is third order. The numerical factor is not expected to be universal but the p and l dependence should be. Finally, it would be interesting to study the possibility of inter-polymer aggregation in the low temperature phase.

Acknowledgements: I have benefited from valuable discussions with J.Cardy, I.Kostov, and especially S.Flesia. Part of the work was done while visiting Service de Physique Theorique de Saclay and the Aspen Center for Physics.

References

- [1] G Swislow, S-T Sun, INishio, and T Tanaka, Phys. Rev. Lett. 44 (1980) 796.
- [2] P G .de Gennes, Scaling concepts in Polymer Physics, Cornell University Press, Ithaca, NY, (1979).
- [3] J Ambjorn, B Durhuus, and J Frohlich, Nucl. Phys. B 257 (1985) 433;
- [4] F David, Nucl. Phys. B 257 45 (1985); V A Kazakov, IK Kostov, and A A Migdal, Phys. Lett. B 157 (1985) 295.
- [5] B Duplantier and H Saleur, Phys. Rev. Lett. 59 (1987) 539; Nucl. Phys. B 290 (1987) 291; D Dhar and J Vannimenus, J. Phys. A 20 (1987) 199.
- [6] B Duplantier and IKostov, Nucl. Phys. B 340 (1990) 491.
- [7] D Bessis, C Itzykson, and J-B Zuber, Adv. in Appl. Math. 1 (1980) 109; M L Mehta, Commun. Math. Phys. 79 (1981) 327.

FIGURE CAPTIONS

Figure 1. { A portion of a two-dimensional simplicial lattice made of square simplices (broken lines) with the polymer chain shown as a continuous line. This figure shows one contact interaction on edge a' (b' is not counted as a contact).

Figure 2. { The induced free energy per unit length u estimated from eqs.(15) (17) at $L = 150$. Broken lines are the leading perturbative results around $c = 1$ and $c = \frac{P}{2} - 1$ for $L = 1$.