

# Generalized Laughlin Wave Functions

## Including the Effect of Coulomb Interaction

O. J. Kwon

*Department of Physics, Pai-Chai University, Taejon 302-162, Korea*

and

*Department of Physics, Hanyang University, Seoul 133-791, Korea*

Bum-Hoon Lee and Sang-Jin Sin

*Department of Physics, Hanyang University, Seoul 133-791, Korea*

### Abstract

We present improved wave functions for the ground state, Laughlin quasihole and quasiparticle excitations of the fractional quantum Hall effect. These depend explicitly on the effective strength of Coulomb interaction and reproduce Laughlin's original result in the limit of no Coulomb interaction.

PACS numbers: 73.40.Hm, 72.20.My

The fractional quantum Hall effect (FQHE) has attracted much interest since it was discovered in the 2D system of electrons subject to a high perpendicular magnetic field [1]. Laughlin first proposed a liquid-type ground state wave function (GSWF) [2]. His wave functions for the ground state, quasihole and quasiparticle excitations are independent of the effective strength of the inter-electron Coulomb interaction. However it has been observed in experiments that the energy gap is greatly reduced as the effective strength of Coulomb interaction increases [3].

After Laughlin's seminal work, most of the discussions on the fractional quantum Hall effect using a microscopic trial wave function has been confined to the strong magnetic field limit, at which the effect of Landau level mixing can be ignored. The only exception is the recent work by Price et. al. [4] on the spherical geometry. Their trial wave function includes a term which is similar to the pseudopotential proposed by Ceperley [5] to include the effect of Coulomb interaction on the Wigner crystallization of the fermion one-component plasma.

We present in this Letter a ground state wave function depending explicitly on the Coulomb interaction, which is derived in a plausible manner from a Chern-Simons gauge field theory in the plane geometry. The modifying term is different from the term in Ref. [5]. Once a ground state wave function is given and if it is nondegenerate, wave functions for Laughlin quasihole and quasiparticle excitations can be written down directly following the Laughlin's argument of adiabatic flux insertion. To support our reasoning, we also have done numerical calculations. We have compared numerically the Laughlin ground state wave function and our trial wave function in the case of up to five particles. This calculation shows the superiority of our trial wave function. This positive result provides a motivation for further numerical efforts.

Let's start with a non-relativistic Chern-Simons gauge field theory. The Hamiltonian is

$$H = \int d^2x \left( \frac{1}{2m} |D_j \psi|^2 + \frac{\alpha}{m} (\psi^\dagger \psi)^2 + V_c(\psi) \right), \quad (0.1)$$

where  $j = 1, 2$ ,  $m$  is the effective mass of charge carrier (electron or hole),  $iD_j = i\partial_j + a_j - qA_j$  and  $V_c(\psi)$  is the Coulomb interaction term given by

$$V_c(\psi) = \frac{q^2}{2\epsilon} \int d^2y (\psi^\dagger \psi(x) - \rho) \frac{1}{|x - y|} (\psi^\dagger \psi(y) - \rho), \quad (0.2)$$

where  $q$  is the charge of the particle,  $\epsilon$  is the dielectric constant,  $\rho$  is the particle density of uniform neutralizing background charges, and  $\psi^\dagger \psi$  is the particle density of charge carriers whose mean value is  $\rho$ . We use the unit such that  $c = \hbar = 1$ . In two spatial dimensions a spin polarized fermion can be traded with a composite of a hard-core boson and odd integer number of flux quanta [6]. We realize hard-core bosons by bosons interacting each other with a repulsive  $\delta$ -function type interaction, which gives rise to the term  $(\psi^\dagger \psi)^2$ . The interaction between the attached fluxes can be described by the Chern-Simons gauge field  $a_j$  satisfying the Gauss constraint equation  $(1/2\alpha)\epsilon_{ij}\partial_i a_j = \psi^\dagger \psi$  where  $\epsilon_{12} = -\epsilon_{21} = 1$ . Here  $\alpha$  is the odd integer  $n$  times  $\pi$ , which is called the statistics parameter [7]. If  $B$  is the strength of external magnetic field, the gauge potential  $A_j = -(B/2)\epsilon_{jk}x^k$  in the symmetric gauge. Terms in the Hamiltonian are understood to be normally ordered. We will not specify the operator orderings here since the effects to the ground state wave function are irrelevant for our purpose. We emphasize that our model is not the one conventionally used in the effective field theory of the fractional quantum Hall effect [8]. In the effective theory the hard-core interaction term comes from the Coulomb interaction in the microscopic theory and thus there is no explicit Coulomb interaction term in the effective theory [9]. However in our derivation the hard-core interaction term turns out to have no relevance to the ground state wave function as shown by Haldane [10]. Hence we will adopt the second quantized version of a microscopic theory of Ezawa et.al., in which the Coulomb interaction term appears explicitly [11]. Later we will take the particle limit to obtain the many-body quantum mechanical wave function.

We are interested in the ground state wave function describing FQHE at the filling factor  $\nu = 1/n$ , therefore  $\rho = eB/2\alpha$ . It has also been known that a kind of Meissner effect occurs in the fractional quantum Hall liquid system [12], which, in this formulation, can be stated as  $a_i - eA_i = \delta a_i = 0$ . It is easy to show that above two conditions are unique constant solution satisfying the equations of motion coming from the Hamiltonians with as well as

without the Coulomb interaction term [11].

We now take an ansatz for the boson field operator that is known to be good for the strongly correlated cases [13]

$$\psi = e^{i\theta(x)} \sqrt{\rho + \eta(x)}, \quad (0.3)$$

where  $\theta(x)$  is the phase operator and  $\eta(x)$  is the density fluctuation operator. Solving the Gauss constraint equation, we get

$$\delta a_i(x) = -\frac{\alpha}{\pi} \epsilon_{ij} \int d^2 y \frac{x^j - y^j}{(x - y)^2} \eta(y). \quad (0.4)$$

Note that the statistics parameter  $\alpha$  determines the strength of statistical interaction ( $\delta a_i(x) \delta a_i(y) \sim \alpha^2 \eta(x) \ln |x - y| \eta(y)$ ), and therefore considering fermion case means that the interaction is strong [14]. This justifies our choice of ansatz.

We substitute the ansatz in the Hamiltonian. Since fluctuations are very small in the liquid we keep terms only up to quadratic order of  $\eta(x)$  and  $\theta(x)$ . This is just the static version of harmonic approximation used in Ref. [12]. Then the resulting quadratic Hamiltonian is

$$\begin{aligned} H = \int d^2 x \bigg[ & \frac{\rho}{2m} \partial_j \theta(x) \partial_j \theta(x) + \frac{1}{8m\rho} \partial_j \eta(x) \partial_j \eta(x) \\ & - \frac{\rho\alpha}{m\pi} \int d^2 y \eta(x) \ln |x - y| \eta(y) + \frac{\alpha}{m} \eta(x) \eta(y) \\ & + \frac{e^2}{2\epsilon} \int d^2 y \eta(x) \frac{1}{|x - y|} \eta(y) \bigg]. \end{aligned} \quad (0.5)$$

In the quadratic Hamiltonian the first three, the fourth and the last terms come from the kinetic term, the hard-core interaction term and the Coulomb interaction term, respectively.

We next Fourier transform the quadratic Hamiltonian. The transformation rule is

$$\begin{aligned} f(k) &= \frac{1}{2\pi} \int d^2 x e^{ikx} f(x), \\ f(x) &= \sum_{\mathbf{k} \neq 0}^{k_{max}} \frac{1}{2\pi} e^{-ikx} f(k). \end{aligned} \quad (0.6)$$

We here introduce the cut-off of momentum  $k_{max}(= \sqrt{8\pi\rho})$  to preserve the number of degrees of freedom given by the particle number in the original many body problem. We also exclude

$\mathbf{k} = 0$  case in the summation, since this mode can be absorbed in the condensate, the constant part of  $\psi^\dagger\psi$ . The Fourier transformed Hamiltonian is then given by

$$H = \sum_{\mathbf{k} \neq 0}^{k_{max}} \left[ \frac{\rho}{2m} k^2 \theta(-k) \theta(k) + \left( \frac{1}{8m\rho} k^2 + \frac{\alpha}{m} + \frac{\pi e^2}{\epsilon} \frac{1}{k} + \frac{2\rho\alpha^2}{m} \frac{1}{k^2} \right) \eta(-k) \eta(k) \right]. \quad (0.7)$$

We note that a collective field theory, in which Fourier transforms of the density operator are regarded as collective fields, gives the same quadratic Hamiltonian in the large- $N$  ( $N$  is the number of particles) expansion [15].

In the non-relativistic quantum field theory the field operators  $\psi(x)$  and  $\psi^\dagger(x)$  satisfy the following commutation relations:

$$\begin{aligned} [\psi(x), \psi(y)] &= [\psi^\dagger(x), \psi^\dagger(y)] = 0, \\ [\psi(x), \psi^\dagger(y)] &= \delta(x - y). \end{aligned} \quad (0.8)$$

We remember that  $\psi(x)$  and  $\psi^\dagger(x)$  have been traded by  $\eta(x)$  and  $\theta(x)$  through the ansatz. Since we consider here the case where  $\eta(x)$  is much smaller than  $\rho$  we may get an approximate representation  $\psi(x) \simeq e^{i\theta(x)}(1 + \eta/2)\sqrt{\rho}$ . From these, the following commutation relations of  $\eta(x)$  and  $\theta(x)$  are obtained

$$\begin{aligned} [\theta(x), \theta(y)] &= [\eta(x), \eta(y)] = 0, \\ [\theta(x), \eta(y)] &= -i\delta(x - y). \end{aligned} \quad (0.9)$$

From these commutation relations we may write down directly commutation relations of  $\eta(k)$  and  $\theta(k)$

$$\begin{aligned} [\theta(k), \theta(k')] &= [\eta(k), \eta(k')] = 0, \\ [\eta(k), \theta(k')] &= i\delta_{k+k'}. \end{aligned} \quad (0.10)$$

Due to these commutation relations the Fourier transformed Hamiltonian can be interpreted as the Hamiltonian of a system of  $2N$  uncoupled oscillators with the following mass  $M_k$  and natural frequency  $\omega_k$  for each mode

$$M_k = m/(\rho k^2),$$

$$\omega_k = \sqrt{\omega_c^2 + 2\pi \frac{\rho}{m} \frac{e^2}{\epsilon} k + \frac{2\rho\alpha}{m^2} k^2 + \frac{1}{4m^2} k^4}, \quad (0.11)$$

where  $\omega_c (= eB/m)$  is the cyclotron frequency.

Recall that a simple harmonic oscillator with Hamiltonian  $h = p^2/2M + (1/2)M\omega^2 x^2$  and commutation relations  $[x, x] = 0 = [p, p]$ ,  $[x, p] = i$ , has the GSWF  $\Psi_0 \propto \exp(-1/2 M\omega x^2)$ .

Using this harmonic oscillator analogy [12] we reach at the GSWF  $\Psi_C$  of our system

$$\Psi_C \propto \exp \left[ - \sum_{\mathbf{k} \neq 0}^{k_{max}} \frac{1}{2} M_k \omega_k \eta(k) \eta(-k) \right]$$

$$\simeq \exp \left[ - \sum_{\mathbf{k} \neq 0}^{k_{max}} \left( \alpha \frac{1}{k^2} + \frac{\pi \lambda l_B}{2} \frac{1}{k} + \left( \frac{\alpha l_B^2}{2} - \frac{\pi^2}{8\alpha} l_B^2 \right) \right) \eta(k) \eta(-k) \right], \quad (0.12)$$

where  $l_B (\equiv 1/\sqrt{eB})$  is the magnetic length and  $\lambda (\equiv (e^2/\epsilon l_B)/\omega_c)$  is the so-called Landau-level mixing parameter measuring the effective strength of the Coulomb interaction. It is very easy to rewrite  $\omega_k$  as  $\omega_k = \omega_c \sqrt{1 + 2\nu\sqrt{\nu}\lambda p + 4\nu p^2 + 4\nu^2 p^4}$  with  $p \equiv k/k_{max}$ . Since  $p \leq 1$  the approximation  $\omega_k \simeq \omega_c [1 + \nu\sqrt{\nu}\lambda p + O(p^2)]$  converges as long as  $\lambda \leq (\nu\sqrt{\nu})^{-1}$ . Even for the massive hole case,  $\lambda$  is  $3 \sim 5$  [3,16] and thus this approximation is good even for  $\nu = 1/3$  case. We emphasize that  $p^M$  ( $M \geq 2$ ) terms will give only contact terms of no physical importance.

We now Fourier transform  $\Psi_C$  back to the coordinate space.

$$\Psi_C = \exp \left[ \begin{aligned} & \frac{\alpha}{2\pi} \int d^2x d^2y \eta(x) \ln |x - y| \eta(y) \\ & - \frac{\lambda l_B}{4} \int d^2x d^2y \eta(x) \frac{1}{|x - y|} \eta(y) \\ & - \left( \frac{\alpha l_B^2}{2} - \frac{\pi^2}{8\alpha} l_B^2 \right) \int d^2x \eta(x) \eta(y) \end{aligned} \right]. \quad (0.13)$$

At this stage the GSWF is expressed in terms of fields. To obtain the many-body quantum mechanical GSWF, we take the particle limit

$$\eta(x) \rightarrow \sum_{a=1}^N \delta(x - x_a) - \rho. \quad (0.14)$$

From Eqs. (13) and (14) we get

$$\Psi_C \propto \prod_{a < b} |z_a - z_b|^{1/\nu} \exp \left[ -\frac{1}{4} \sum_a |z_a|^2 - \frac{\lambda}{2} \sum_{a < b} \frac{1}{|z_a - z_b|} \right], \quad (0.15)$$

where  $z_a \equiv x_a/l_B$ . We neglected the contact terms ( $\sim \delta(z_a - z_b)$  and its derivatives) as well as the terms coming from the finite size effect, which break the translational symmetry of the original Hamiltonian.

The  $\Psi_C$  is the wave function of charged boson. After (singular) gauge transformation to get the wave function of the original spin polarized fermions the final form can be written as

$$\Psi_C = \Psi_L \times \exp \left[ -\frac{\lambda}{2} \sum_{a < b} \frac{1}{|z_a - z_b|} \right], \quad (0.16)$$

where  $\Psi_L$  denotes the Laughlin wave function. This is the main result of our work.

We now calculate energies for  $\Psi_L$  and  $\Psi_C$  for  $N = 5$ . If we do not consider the particle-background and background-background Coulomb interactions, the Hamiltonian is

$$H = \frac{N}{2} + 2 \sum_a^N \phi_a^\dagger \phi_a + \lambda \sum_{a < b} \frac{1}{|z_{ab}|}, \quad (0.17)$$

where  $\phi = \partial_z + \bar{z}/4$  and  $\phi^\dagger = -\partial_{\bar{z}} + z/4$ . Here we take  $\omega_c = 1$ .  $(H - N/2)\Psi_C = H_C\Psi_C$  can be obtained analytically

$$\begin{aligned} H_C = & \frac{5}{4} \lambda \sum_{a < b}^N \frac{1}{|z_{ab}|} - \sum_{a < b}^N \left( n - \frac{1}{2} \right) \frac{\lambda}{|z_{ab}|^3} - \frac{\lambda^2}{4} \sum_{a < b}^N \frac{1}{|z_{ab}|^4} \\ & - \frac{\lambda^2}{8} \sum_{a \neq b \neq c}^N \frac{\text{Re}(z_{ab} \bar{z}_{ac})}{|z_{ab} z_{ac}|^3} - \frac{n\lambda}{2} \sum_a^N \sum_{b \neq a}^N \sum_{c \neq a, b}^N \frac{z_{ab}}{|z_{ab}|^3} \frac{1}{z_{ac}}. \end{aligned} \quad (0.18)$$

Energies for  $\Psi_L$  and  $\Psi_C$ , which are defined by  $\int d^2z \Psi_L^* \Psi_L (\sum_{a < b} \lambda/|z_{ab}|) / \int d^2z \Psi_L^* \Psi_L$  and  $\int d^2z \Psi_C^* \Psi_C H_C / \int d^2z \Psi_C^* \Psi_C$ , are calculated for  $N = 5$ ,  $\lambda = 2/3$  case using Monte Carlo method, and are given by 2.05 and 1.75, respectively. That is, the energy is lowered by about 15%. This positive result from the five-body calculation provides a motivation for more numerical efforts. There is another support for the would-be superiority of our wave function. The total energy per particle is given in terms of the radial distribution function  $g(|z|)$ . Laughlin argued that the main reason why the fractional quantum Hall liquid is variationally superior to the Wigner crystal at  $\nu = 1/3$  is the fact that  $g(|z|)$  for the liquid

is pushed out further from the origin than  $g(|z|)$  for the crystal [17]. It is evident that the modifying term in our trial wave function further push out  $g(|z|)$  from the origin. Thus we expect our wave function will be variationally superior to the Laughlin wave function.

Based on our improved ground state wave function, we can directly write down the wave functions  $\Psi_C^{-z_0}$  and  $\Psi_C^{+z_0}$  for the Laughlin quasihole and quasiparticle excitations following the Laughlin's argument [2,17,18] of adiabatic flux insertion

$$\begin{aligned}\Psi_C^{-z_0} &= \exp \left[ -\frac{1}{4} \sum_a |z_a|^2 - \frac{\lambda}{2} \sum_{a<b} \frac{1}{|z_a - z_b|} \right] \prod_a (z_a - z_0) \prod_{a<b} (z_a - z_b)^m, \\ \Psi_C^{+z_0} &= \exp \left[ -\frac{1}{4} \sum_a |z_a|^2 - \frac{\lambda}{2} \sum_{a<b} \frac{1}{|z_a - z_b|} \right] \prod_a \left( 2 \frac{\partial}{\partial z_a} - z_0^* \right) \prod_{a<b} (z_a - z_b)^m.\end{aligned}\quad (0.19)$$

One might try to derive these wave functions using the mean field method used above from a vortex field theory obtained via the Hubbard-Stratonovich transformation [11,12]. However this is not so promising, since the linear approximation (or the harmonic approximation) is not good for the intrinsically nonlinear object like a vortex and moreover the vortex size is comparable with the magnetic length, which is a typical length of the system.

The appearance of the FQHE at  $\nu = 1/7$  has been controversial. Goldman et. al's experiment [19] and Jain's classification [21] insist on the appearance. However, Lam and Girvin's classic work [19] and the most updated numerical calculation [4,22] show that the critical filling factor  $\nu_c$  for the liquid-solid transition is larger than  $1/7$ . In Ref. [4], they used a trial wave function including the Landau level mixing effect, which however has a form different from ours in (16). It is interesting to see if our improved wavefunction gives the ground state energy lower than that of Ref. [4] making  $\nu_c$  smaller. In case  $\nu_c$  is less than  $1/7$  with our modified wave function, then the appearance of the FQHE at  $\nu = 1/7$  could be explained theoretically.

As discussed in the beginning of this paper, the Laughlin's wave functions for the ground state, quasihole and quasiparticle excitations are independent of  $\lambda$  and therefore cannot explain the experimentally observed reduction of energy gap between the ground state and the Laughlin quasiexciton as  $\lambda$  increases. However, our wave functions in (16) and (19) are explicitly dependent on  $\lambda$  and therefore there is a chance to explain the reduction of the



energy gap.

In conclusion we propose systematically derived improved trial wave functions for the ground state, and Laughlin quasihole and quasiparticle excitations. These include the Landau level mixing effect explicitly. Since the validity of our wavefunction will shed new light on the appearance of the fractional quantum Hall effect at  $\nu = 1/7$  [19] and the reduction of energy gap as the effective strength of Coulomb interaction increases [3,20], it will be very interesting to continue the numerical study.

This work was supported in part by the KOSEF and by the Korean Ministry of Education.

## REFERENCES

- [1] D. C. Tsui, H. L. Stormer and A. C. Gossard, Phys. Rev. Lett. **48**, 1559 (1982).
- [2] R. B. Laughlin, Phys. Rev. Lett. **50**, 1395 (1983); Surf. Sci. **142**, 163 (1984).
- [3] M. B Santos, Y. W. Suen, M. Shayegan, Y. P. Li, L. W. Engel, and D. C. Tsui, Phys. Rev. Lett. **68**, 1188 (1992). They measured activation energy  $\Delta \simeq 4K$  for a 2D electron system, but  $\Delta \simeq 0.4K$  for a 2D hole system with roughly the same density and quality as the 2D electron system. In these systems amounts of the effect of impurity and finite thickness of the layer are roughly the same. The drastic reduction therefore comes from the increment of the effective strength of Coulomb interaction.
- [4] R. Price, P. M. Platzman and Song He, Phys. Rev. Lett. **70**, 339 (1993).
- [5] D. Ceperley, Phys. Rev. B **18**, 3126 (1978).
- [6] D. Sen, Nucl. Phys. **B360**, 397 (1991).
- [7] J. M. Leinass and J. Myrheim, Nuovo Cimento **37B**, 1 (1977); F. Wilczek, Phys. Rev. Lett. **49**, 957 (1982); F. Wilczek and A. Zee, Phys. Rev. Lett. **51**, 2250 (1983).
- [8] S. M. Girvin, Chap. 10 in *The Fractional Quantum Hall Effect*, edited by R. E. Prange and S. M. Girvin (Springer-Verlag, New York, 1986); S. M. Girvin and A. H. MacDonald, Phys. Rev. Lett. **58**, 1252 (1987).
- [9] S. C. Zhang, T. H. Hansson and S. Kivelson, Phys. Rev. Lett. **62**, 82 (1989).
- [10] F. D. M. Haldane, Phys. Rev. Lett. **51**, 605 (1983); F. D. M. Haldane and E. H. Rezayi, Phys. Rev. Lett. **54**, 237 (1985).
- [11] Z. F. Ezawa, M. Hotta and A. Iwazaki, Phys. Rev. B **46**, 7765 (1992).
- [12] S. C. Zhang, Int. J. Mod. Phys. B **6**, 25 (1992) and references therein.
- [13] B. Sakita, *Quantum Theory of Many Variable System and Fields* (World Scientific,

- Singapore, 1985).
- [14] For a near-boson case ( $\alpha \simeq 0$ ) the ansatz  $\psi = \sqrt{\rho} + \eta$  ( $\eta$  is complex) is more appropriate. Using this ansatz and keeping terms up to quadratic order of  $\eta(x)$  we may get the GSWF (the Jastrow wave function) for charged superfluid,  $\Psi_s^\nu = \prod_{a < b} (z_a - z_b)^{1/\nu}$  (see Ref. 11).
  - [15] Hiro-Oka, H. Minakata, and Hisakazu, TMUP-HEL-9306, Jun 1993; B. Sakita, D-N. Sheng, and Z-B. Su, Phys. Rev. B **44**, 11510 (1991).
  - [16] T. Ando, J. Phys. Soc. Jpn. **54**, 1528 (1985).
  - [17] R. B. Laughlin, Chap. 7 in *The Quantum Hall Effect*, edited by R. E. Prange and S. M. Girvin (Springer-Verlag, New York, 1990).
  - [18] B. I. Halperin, Helv. Phys. Acta **56**, 75 (1983). In this paper there appears a different form of  $\Psi_L^{+z_0}$  motivated by a classical statistical-mechanics problem.
  - [19] V. J. Goldman, M. Shayegan, and D. C. Tsui, Phys. Rev. Lett. **61**, 881 (1988); P. K. Lam and S. M. Girvin, Phys. Rev. B **33**, 473 (1984).
  - [20] D. Yoshioka, J. Phys. Soc. Jpn. **53**, 3740 (1984); **55**, 885 (1986).
  - [21] Fig. 1 in J. K. Jain, Phys. Rev. Lett. **63**, 199 (1989).
  - [22] X. Zhu and S. G. Louie, Phys. Rev. Lett. **70**, 335 (1993).