Level Statistics and Localization for Two Interacting Particles in a Random Potential

Dietmar Weinmann and Jean-Louis Pichard

CEA, Service de Physique de l'Etat Condensé, Centre d'Etudes de Saclay,

91191 Gif-sur-Yvette Cedex, France

We consider two particles with a local interaction U in a random potential at a scale L_1 (the one particle localization length). A simplified description is provided by a Gaussian matrix ensemble with a preferential basis. We define the symmetry breaking parameter $\mu \propto U^{-2}$ associated to the statistical invariance under change of basis. We show that the Wigner-Dyson rigidity of the energy levels is maintained up to an energy E_{μ} . We find that $E_{\mu} \propto 1/\sqrt{\mu}$ when Γ (the inverse lifetime of the states of the preferential basis) is smaller than Δ_2 (the level spacing), and $E_{\mu} \propto 1/\mu$ when $\Gamma > \Delta_2$. This implies that the two-particle localization length L_2 first increases as |U| before eventually behaving as U^2 .

PACS numbers: 72.15, 73.20

For a single particle diffusing in a disordered system of size L smaller than the one particle localization length L_1 , there are two characteristic energies: the Thouless energy $E_{\rm c} = \hbar D/L^2$ and the level spacing $\Delta_1 \approx B_1/L^d$ $(B_1, D \text{ and } d \text{ are the band width, the diffusion constant})$ and the system dimension, respectively). If one writes the distribution of energy levels as a Gibbs factor of a fictituous Coulomb gas, the corresponding pairwise interaction for levels with separation $\epsilon < E_{\rm c}$ coincides [1] with the logarithmic repulsion characteristic of the matrix ensembles which are statistically invariant under change of basis, e. g. the Gaussian Orthogonal Ensemble (GOE). For $\epsilon > E_{\rm c}$, the level repulsion vanishes more or less quickly, depending on the system dimension. The dimensionless conductance g_1 is given by E_c/Δ_1 . This ratio is the single relevant parameter in the scaling theory of localization. In quasi-one dimension, the size where $g_1 \approx 1$ defines L_1 . In three dimensions, the mobility edge is characterized by $g_1 \approx g_c$ where g_c is of order 1.

We shall generalize those concepts to two particles with a local (repulsive or attractive) interaction. This two interacting particle (TIP) problem has received a particular attention since Shepelyansky [2] pointed out that certain TIP states may extend over a scale L_2 much larger than L_1 . Shepelyansky's original reasoning consists in mapping the problem for $L \gg L_1$ onto a random band matrix model with a superimposed diagonal matrix (SBRM-model). Imry [3] used later the Thouless scaling block picture to arrive at precisely the same results as Shepelyansky. The smearing due to the interaction of the energy levels within L_1 was estimated using Fermi's golden rule, yielding $L_2 \propto U^2$. This delocalization effect has been confirmed by transfer matrix studies [4,5], and unambiguously illustrated from numerical studies [6] of rings threaded by an AB-flux. However, in one dimension, for system sizes which can be numerically investigated, one obtains [5] $L_2 \propto |U|$ contrary to Fermi's golden rule, and a disorder dependence [4] $L_2 \propto L_1^{\alpha}$ with $\alpha \approx 1.5 - 1.7$ and not 2, as predicted by Shepelyansky

and Imry.

To understand those contradictory results, we study the TIP energy level statistics at a scale L_1 in order to identify the energy which plays the role of $E_{\rm c}$ in this case, and to determine its dependence on U. For the original TIP-problem, we assume a tight-binding model [2] on a d-dimensional lattice $(L_1^d \text{ sites } p \text{ where the random poten-})$ tial is taken with a box distribution of width 2W). The nearest neighbor hopping term takes a constant value V = 1 and U is the on-site interaction. Assuming two electrons with opposite spins, we consider only the symmetric states. The TIP-Hamiltonian [2] can be written in the basis of the $N = L_1^d (L_1^d + 1)/2$ (symmetrized) products of one particle states $|AB\rangle$. We denote by R_{pA} the value on site p of the one particle eigenstate with energy ϵ_A . In this basis, the diagonal terms are dominated by one particle contributions $\epsilon_A + \epsilon_B$ and the interaction Hamiltonian yields a full matrix (for $L \leq L_1$) with entries $U \cdot Q_{ABA'B'} = U \sum_{p} R_{pA} R_{pB} R_{pA'} R_{pB'}$. The magnitude of those terms is of order $U/L_1^{3d/2}$ with a random sign which does not preserve the sign of the interaction.

Before considering the TIP-Hamiltonian, it is instructive to discuss a simplified matrix model where the correlations between matrix elements are neglected: an ensemble of real symmetric matrices G with independent entries, characterized by Gaussian distributions with variances $\langle G_{ii}^2 \rangle \approx B_1^2/3 \ (B_1 = 4Vd + 2W)$ and $\langle G_{ij}^2 \rangle \approx$ U^2/L_1^{3d} for the diagonal and off-diagonal terms, respectively. The averages are set to zero, which neglects a shift of the diagonal terms by an amount $U \cdot Q_{ABAB} \approx U/L_1^d$ assumed to be much smaller than B_1 . These shifts preserve the sign of U and for large U, eventually split the energy band into two parts. For the sake of simplicity, we ignore them, restricting us to small U and to a Gaussian matrix with preferential basis (GMPB)-model which has been used previously [7] to study the GOE to Poisson crossover for the level statistics, and to define a maximum entropy model [8] where the range of the level interaction depends on a parameter. When $\langle G_{ii}^2 \rangle$ is very large



FIG. 1. The strength function for a TIP-Hamiltonian (5 by 5 lattice in the metallic regime W = 2, V = 1). Diamonds, squares and triangles are for U=0.2, 0.6 and 1.0, respectively. Lines are Breit–Wigner distribution functions, fitted to the numerical data. The inset shows how Γ depends on U. The line represents $\Gamma = U^2/22$.

compared to $\langle G_{ij}^2 \rangle$, one has indeed a strongly preferential basis and it is convenient to re-order the diagonal terms such that $G_{11} < G_{22} < \ldots < G_{NN}$. Those G_{ii} may be considered as the positions of the energy levels in the zeroth approximation, when the interaction with the other states is neglected. The small coupling terms G_{ij} spread those basis states over Γ/Δ_2 neighbors. $\Delta_2 \approx \sqrt{2\pi \langle G_{ii}^2 \rangle} / N$ is the level spacing and Γ can be estimated using Fermi's golden rule: $\Gamma \approx 2\pi < G_{ii}^2 >$ $/\Delta_2$. Diagonalizing G by an orthogonal transformation $O(G = OG_dO^t)$, where G_d is a diagonal matrix with real entries E_{α}), we consider the strength function introduced by Wigner $\rho_{\rm W}(E,p) = \sum_{\alpha=1}^{N} O_{p\alpha}^2 \delta(E-E_{\alpha})$. If one averages over the ensemble, one gets for the eigenvector amplitudes $\langle O_{p\alpha}^2 \rangle = \Delta_2^{-1} \langle \rho_W(E_\alpha, p) \rangle$. For $L \gg L_1$, $\rho_{\rm W}(E,n)$ has been found [9–11] in agreement with the Breit-Wigner form

$$<\rho_{\rm W}(E,n)>=\frac{\Gamma}{2\pi[(E-G_{nn})^2+\Gamma^2/4]}$$
. (1)

We have checked that this holds too for $L \leq L_1$, and we show in Fig. 1 that this Breit-Wigner form of the eigenstates characterizes also the original TIP-Hamiltonian, once the shifts $U \cdot Q_{nn}$ of the quasi-energies H_{nn} are taken into account. This shows us that a basis state $|n\rangle \equiv |AB\rangle$ (i. e. an eigenstate of the TIP-Hamiltonian for U = 0) becomes delocalized over Γ/Δ_2 of its neighbors (i.e. over the basis states $|n'\rangle$ where $H_{n'n'}$ is close to $H_{nn} = \epsilon_A + \epsilon_B$), with a Lorentzian shape centered in $H_{nn} + U \cdot Q_{nn}$. Γ plays the role of a localization length in the preferential basis, and is given by Fermi's golden rule for small values of U ($U \leq 2$ in Fig. 1).

Having understood how the eigenstates are delocalized by the interaction over the preferential basis (see also Refs. [9,12]), we focus our attention on the energy levels. We introduce a symmetry breaking parameter μ in the probability density

$$\rho(G) \propto \exp\left(-\sum_{i=1}^{N} \frac{G_{ii}^2}{2\sigma^2} - (1+\mu) \sum_{i$$

with $\sigma^2 \approx B_1^2/3$ and $\sigma^2/(2(1 + \mu)) \approx U^2/L_1^{3d}$. When $\mu = 0$, one recovers the GOE-ensemble with $\rho_{\text{GOE}}(G) \propto \exp(-tr(G^2)/2\sigma^2)$. When $\mu \neq 0$, there is a factor $\rho_{\mu}(G)$ which removes the statistical invariance under change of basis. Expressed [8] in eigenvalue-eigenvector coordinates, it reads

$$\rho_{\mu}(G) \propto \prod_{\alpha < \beta}^{N} \exp\left(-\frac{\mu}{2\sigma^2} (E_{\alpha} - E_{\beta})^2 \sum_{p} O_{p\alpha}^2 O_{p\beta}^2\right) .$$
(3)

The question is to understand how this additional factor, after integration over the matrices O (distributed with Haar measure $\mu(dO)$ over the orthogonal group) can destroy the logarithmic level repulsion coming from the measure $\mu(dG) = \prod_{\alpha < \beta}^{N} |E_{\alpha} - E_{\beta}| \prod_{\alpha}^{N} dE_{\alpha}\mu(dO)$. This will allow us to identify the characteristic scale E_{μ} below which one recovers the GOE rigidity, and above which the levels become uncorrelated. Two cases have to be considered:

(i) $\Gamma < \Delta_2$. The G_{ij} are so small that one can just consider the coupling between two nearest neighbor diagonal entries: i. e. a 2 × 2 matrix which can be diagonalized by a rotation of an angle θ . One finds [8] $\int d\theta \rho_{\mu}(G) = f(x) = \exp(-x) \cdot I_0(x)$ where $x = \mu \cdot \epsilon^2/(8\sigma^2)$, ϵ denoting the separation of the two coupled levels. For x < 1, $f(x) \approx 1$ and decreases as $1/\sqrt{x}$ for x > 1. This gives

$$\frac{E_{\mu}}{\Delta_2} = \frac{\sqrt{8\sigma^2/\mu}}{\Delta_2} \propto \frac{N}{\sqrt{\mu}} \,. \tag{4}$$

For $\epsilon < E_{\mu}$, one has the GOE statistics, while for $\epsilon > E_{\mu}$, the levels are uncorrelated.

(ii) $\Gamma > \Delta_2$. Many neighboring G_{ii} are coupled by the off-diagonal terms. First, we consider the case where $\epsilon = |E_{\alpha} - E_{\beta}| < \Gamma$: i. e. the case where the two corresponding eigenvectors have a strong overlap. Assuming that the eigenvectors $|O_{\alpha}\rangle$ have non-zero coordinates of order $O_{n\alpha}^2 \approx \Delta_2/\Gamma$ over Γ/Δ_2 neighboring basis states only, one gets $\sum_{p=1}^{N} O_{p\alpha}^2 O_{p\beta}^2 \approx \Delta_2/\Gamma$, and $\exp\left(-\mu\epsilon^2\Delta_2/(2\sigma^2\Gamma)\right) \approx 1$, independent on $\epsilon(<\Gamma)$. Writing $O = \exp A$, with A a real antisymmetric matrix $(\mu(dO) = \prod_{\alpha < \beta} dA_{\alpha\beta})$, one can see that the small fluctuations of the $A_{n\alpha}$ around their typical values will not yield a correction to the GOE level repulsion. This means that there is no coupling between eigenvalues and eigenvectors as far as $\epsilon < \Gamma \equiv E_{\mu}$ with

$$\frac{E_{\mu}}{\Delta_2} \propto \frac{N^2}{\mu} \tag{5}$$



FIG. 2. Σ_2 for the GMPB-ensemble (N = 500). Diamonds, squares and triangles are for $\mu = 30000, 5000$ and 1000, respectively. The inset shows how E_{μ} depends on μ . The triangles give the energy where Σ_2 is 20 percent above the GOE value. The solid and the dotted line represent $E_{\mu}/\Delta_2 = 0.039N^2/\mu$ and $E_{\mu}/\Delta_2 = 0.19N/\sqrt{\mu}$, respectively. The squares result from a fit $\Sigma_2 = (E/E_{\mu})^{\alpha}$ valid for $E \gg E_{\mu}$.

now, instead of $N/\sqrt{\mu}$ previously. When $|\epsilon| > \Gamma$, the eigenvectors do not overlap and the levels should become uncorrelated. In Ref. [8], it was noted that if $O_{p\alpha} \approx \delta_{p,\alpha} + A_{p\alpha}$ where $A_{p\alpha} \ll 1$, $\sum_{p=1}^{N} O_{p\alpha}^2 O_{p\beta}^2 \approx 2A_{\alpha\beta}^2$, which gives a $1/|\epsilon|$ factor, after integration over $A_{\alpha\beta}$. This level attraction exactly compensates the level repulsion due to $\mu(dG)$. Qualitatively, one can adapt this reasoning to produce the requested level attraction, after integration of the exact form of the level interaction as a function of ϵ , taking into account the Breit-Wigner form of the eigenvectors, is postponed to a further study.

We have carried out a numerical study of the GMPBensemble (N = 500) as a function of μ , to illustrate the two regimes. The number variance $\Sigma_2(E)$ (variance of the number of levels in an energy interval E) is shown in Fig. 2. For small energy intervals, $\Sigma_2(E)$ coincides with the GOE-logarithmic increase observed when $\mu = 0$. For larger energy intervals, $\Sigma_2(E)$ can be fitted by $(E/E_{\mu})^{\alpha(\mu)}$, which gives a first method for calculating E_{μ} . A second method consists in calculating the energy interval where Σ_2 is above the GOE-curve by a certain threshold (e. g. 20 percent). Note that those methods give a non zero E_{μ} (depending on the chosen threshold) even for uncorrelated levels, which has been subtracted from the data. The inset of Fig. 2 confirms that the two methods are in agreement and exhibits the predicted crossover for E_{μ} when $\Gamma \approx \Delta_2$ ($\mu \approx N^2$), from a N^2/μ dependence (small μ) towards a $N/\sqrt{\mu}$ -dependence. The μ -dependence of the exponent α (see inset in Fig. 4) depends on the exact form of the pairwise level repulsion.

We now turn our attention to the TIP-Hamiltonian in two dimensions. The corresponding Σ_2 is shown in Fig. 3, for different U. We have obtained the same curves



FIG. 3. Σ_2 for the symmetric states of a 2*d* TIP-Hamiltonian (ring with 10x10 sites, W = V = 1). The ring is threaded by a magnetic flux $\Phi = \Phi_0/4$. Diamonds, squares and triangles are for U=0.25, 0.75 and 2.0, respectively. The inset shows how E_U depends on U. The data are obtained as described in the caption of Fig. 2. The dotted line (solid line) corresponds to $E_U/\Delta_2 = 1.35U/V (E_U/\Delta_2 = 2.1(U/V)^2)$.

for U = 1 and U = -1, and consider in more detail repulsive interactions. We took rings containing 10×10 sites threaded by a magnetic flux, so that the level statistics should have a GUE behavior for energy intervals $E < E_{\rm U} \equiv E_{\mu}$, with $(1 + \mu)^{-1} \approx 6U^2/(L_1^{3d}B_1^2)$. Except this trivial change from orthogonal to unitary symmetry, the similarity with the curves calculated for the GMPB-model is very striking, if one disregards large energy intervals and the U-dependence of the exponent α (see inset of Fig. 4). The first difference can be explained from the fact that in the TIP Hamiltonian, the one particle level rigidity cannot be ignored when $E > \Delta_1$ (one has Δ_1/Δ_2 superimposed GUE series when U = 0, correlations which are neglected in the GMPB-model. Another difference results from the shifts $U \cdot Q_{nn}$ of the diagonal terms which become important when U is large. The crossover value $U_{\rm c} \approx \sqrt{2}(8Vd + 4W)/(\sqrt{\pi}L^{d/2})$ between the two regimes is of order 1, for the considered parameters. When $U > U_c$, we observe the U^2 -behavior of E_U .

A similar study in one dimension is very instructive. When W = V = 1, we have $L_1 \approx 25$, which gives again $U_c \approx 1$. As expected, one can see in Fig. 4 that $E_U \propto |U|$ when $|U| < U_c$, but when $U > U_c$, the splitting of the energy band occurs, and E_U decreases. Note that one recovers a Poisson statistics when U is very large (for d = 2, there is only a saturation of E_U). For d = 1, this means that one can couple only two basis states within $L_1 \approx 25$, with a small enough value of U to justify the simplified GMPB ensemble. The observation of the U^2 behavior of E_U requires larger values of L_1 in d = 1 than considered in the numerical studies [4–6].



FIG. 4. $E_{\rm U}$ for a 1*d* TIP-Hamiltonian (W = V = 1and $L = L_1 = 25$ sites). The data are obtained as described in the caption of 2. The dotted line corresponds to $E_{\rm U}/\Delta_2 = 1.2U/V$. The inset gives the exponent α occurring in $\Sigma_2 \propto E^{\alpha}$ at large *E*. Squares are for the 1*d* TIP Hamiltonian, plotted as a function of $U\sqrt{L}/V$. Diamonds represent α for the GMPB ensemble (see Fig. 2), as a function of $N/\sqrt{\mu}$.

We now follow the argument developed by Imry [3] to estimate the localization length L_2 . First, we consider a series of building blocks of size L_1^d . Γ_U is the smearing of the TIP levels of one of the blocks, due to the interactioninduced coupling with the neighboring block. For such a quasi-1*d* wire, the dimensionless conductance at scale L_1 is given by

$$g_2(L_1) \equiv \frac{L_2}{L_1} \approx \frac{1}{2} + A \frac{\Gamma_{\rm U}}{\Delta_2}.$$
 (6)

The factor 1/2 gives [5] the right limit when $U \to 0$ and A is a constant. Obviously, one should have $\Gamma_U \equiv E_U$. When $\Gamma_U > \Delta_2$, Γ_U is given by Fermi's golden rule, the case considered in Ref. [3] and we only discuss the case $\Gamma_U < \Delta_2$, where $\Gamma_U \approx \sqrt{U^2/L_1^{3d}}$. Physically, this means that U is so small that it couples only a single TIP state in one of the blocks to another TIP state in the next block, giving rise to "Rabi oscillations" between those two coupled states. The inverse life time is no longer given by the square of the coupling term, as in Fermi's golden rule, but by its absolute value. In addition, we have shown that this inverse life time gives the scale below which one has a GOE spectral rigidity, extending the known results for non-interacting particles to the TIP problem.

We conclude by discussing a few implications for TIP localization. For d = 1 and $U < U_c$ ($U_c \approx 1$ when $W \approx V \approx 1$, see Fig. 4), one gets (L_2/L_1) $\approx 1/2 + A(|U|/B_1)\sqrt{L_1}$ which is in agreement with the dependence on U observed in the numerical studies [5]. The conjecture proposed in Ref. [5] gives L_1 instead of $\sqrt{L_1}$. As noted in Ref. [4], the distribution of the $Q_{ABA'B'}$ is far from being Gaussian, which can matter as far as the description of the L_1 -dependence by the GMPB-model is concerned. However $L_2 \propto L_1^{3/2}$ is close to the behavior observed in Ref [4]. The *U*-dependence is not affected by this consideration and is correctly described by the GMPB-model.

If one considers two quasi-particles above a Fermi sea, one should replace [3] in Eq. (6) Δ_2 by $\Delta_2(E) \approx \Delta_1^2/E$ where E is the total excitation energy. One immediately obtains that the quasi-particle conductance $g_2^q(E, L_1)$ is of order of $g_2(L_1)$ when $E \approx B_1$, which gives $L_2^q(E \approx B_1) = L_2$, in agreement with Ref. [13]. Similarly, in three dimensions, Imry's relation $(E_{m2} \approx (B_1^2/|U|)E_{m1}^{\nu d/2})$ between the one quasi-particle mobility edge E_{m1} and the two quasi-particle mobility edge E_{m2} does not change when $U < U_c$ (ν denotes the critical exponent associated with L_1).

In summary, we have shown that the basic concepts developed for non interacting particles can be naturally extended to M = 2 interacting particles, after the changes $E_{\rm C} \rightarrow E_{\mu}$ and $\Delta_1 \rightarrow \Delta_2$. A similar conclusion has been obtained from a non linear σ model description of the TIP-Hamiltonian [14], when $L > L_1$. Moreover, our approach can be easily extended to an arbitrary number Mof particles.

We gratefully acknowledge a useful discussion with Boris Shapiro and Dima Shepelyansky. This work was supported by the European HCM program (D. W.).

- R. Jalabert, J.-L. Pichard and C.W.J. Beenakker, *Europhys. Lett.* 24, 1 (1993).
- [2] D. L. Shepelyansky, Phys. Rev. Lett. 73, 2067 (1994).
- [3] Y. Imry, Europhys. Lett **30**, 405 (1995).
- [4] K. Frahm, A. Müller-Groeling, J.-L. Pichard and D. Weinmann, *Europhys. Lett.* **31**, 405 (1995).
- [5] F. von Oppen, T. Wettig and J. Müller, *Phys. Rev. Lett.* 76, 491 (1996).
- [6] D. Weinmann, A. Müller-Groeling, J.-L. Pichard and K. Frahm, *Phys. Rev. Lett.* **75**, 1598 (1995).
- [7] G. Lenz and F. Haake, *Phys. Rev. Lett.* 67, 1 (1991).
- [8] J.-L. Pichard and B. Shapiro, J. Phys. 1 France 4, 623 (1994).
- [9] P. Jacquod and D. L. Shepelyansky, *Phys. Rev. Lett.* 75, 3501 (1995).
- [10] K. Frahm and A. Müller-Groeling, *Europhys. Lett.* **32**, 385 (1995).
- [11] Y. V. Fyodorov and A. D. Mirlin, *Phys. Rev.* B 52, R11580 (1995).
- [12] V. V. Flambaum et al., Phys. Rev. A 50, 267 (1994).
- [13] F. von Oppen and T. Wettig, *Europhys. Lett.* **32**, 741 (1995).
- [14] K. Frahm, A. Müller-Groeling and J.-L. Pichard, to appear in *Phys. Rev. Lett.* (1996).