Landau level bosonization of a 2D electron gas

H.W estfahl Jr¹, A H.C astro N eto² and A D.C aldeira¹ ¹Instituto de F sica G leb W ataghin U niversidade E stadual de C am pinas 13083–970, C am pinas, SP, B razil ²D epartm ent of P hysics U niversity of C alifornia R iverside, CA, 92521, U SA (A ugust 21, 2021)

In this work we introduce a bosonization scheme for the low energy excitations of a 2D interacting electron gas in the presence of an uniform magnetic eld under conditions where a large integral num ber of Landau levels are lled. We give an explicit construction for the electron operator in terms of the bosons. We show that the elementary neutral excitations, known as the magnetic excitons or magnetoplasm a modes, can be described within a bosonic language and that it provides a quadratic bosonic Ham iltonian for the interacting electron system which can be easily diagonalized.

PACS: 73.40.-c,73.40.Hm, 73.20 M f, 71.27.+ a

Bosonization of Ferm i elds has been an important non-perturbative tool for nding the solutions to problems involving many interacting ferm ions. The pioneering attempts by Bloch¹ on sound waves in dense 1D systems, which were later extended and given an intuitive foundation by Tomonaga², called attention to the conceptual and operational simplicity that results if all excitations of a ferm ionic system can be described within a bosonic language^{3 {5}. Now adays the 1D bosonization is a mature eld⁶. Bosonization in higher dimensions was pioneered by Luther⁷ and extended by Haldane a few years ago⁸. More recently this approach has been applied in the context of strongly correlated system swhere the possibility of non-conventional ground states seeks for a non-perturbative m ethod of study^{9 {11}.

In the context of correlated system s, one of the most interesting phenom enon in condensed matter physics is the quantum Halle $\operatorname{ect}^{12 \{14}$ either in its integer (IQHE) or fractional (FQHE) regimes. Despite the fact that there is no therm odynam ic distinction between this two phenom ena¹³, they have quite di erent m icroscopic origins. The IQHE can be understood from the properties of non-interacting electrons whereas the understanding of the FQHE necessarily requires a treatm ent of the electron-electron interaction. Consequently, almost all the e orts to study correlations in a 2D electron gas in the presence of a magnetic eld, B, were directed toward the study of the strong magnetic eld case where the FQHE takes place. In this case the Landau level spacing, the cyclotron energy $h!_c = heB = (m c)$, can be m uch grater than the typical Goulom b energy, which is of the order of $e^2 = 1$ (where 1 = 1ch=(eB) is the magnetic lenght), and therefore the system can be described only in terms of the rst Landau level and the mixing with the other Landau levels can be neglected or taken into account perturbatively. In a weak magnetic eld¹⁵ this is not the case and the typical C oulom b energy can exceed the cyclotron energy resulting in a mixing between the Landau levels. Som e attem pts were made to develop a

m ethod for studying the mixing caused by these correlation e ects starting from a Ferm i Liquid theory^{16;17} or perturbative G reen's function m ethod^{15;18;19}.

In this paper we start from a Landau level description of the system and introduce a non-perturbative bosonization scheme for the 2D interacting ferm ion gas subject to a perpendicular uniform magnetic eld under the condition that there is a large integer number of lled Landau levels. In this case, the excitations are electron-hole pairs offen called magnetic excitons and as we will show can be described within a bosonic language.

Recent experiments in tunneling between parallel 2D electron gases in applied magnetic $elds^{20}$ have revealed the existence of a gap in the tunneling density of states. In this case it is established that correlations have major hole in the tunneling conductance and we expect that the method presented here can be applied to such problem s where the correlations between the electrons in a magnetic eld are so important.

W e start with the Ham iltonian of a spinless ferm ion gas in a plane perpendicular to an uniform magnetic eld

$$H_{o} = \frac{1}{2m}^{Z} d^{2}r^{y}(r) p = \frac{e}{c}A(r)^{2}(r);$$

where (r) is the ferm ion eld operator (for the time being we do not discuss the spin of the electron).

Rewriting (r) in a Landau level basis (symmetric gauge) the Hamiltonian H $_0$ becomes diagonal

$$H_{o} = h!_{c} (n + 1=2)C_{n,m}^{+}C_{n,m};$$

where $c_{n,m}^{\dagger}$ is the ferm ion operator that creates a particle in a Landau level n with the guiding center at a distance 2m + 1' from the origin of the coordinate system. Therefore the m degeneracy is related to the location of the position of the guiding center of the cyclotron orbit.

The non-interacting ground state is generated by uniform ly lling Landau levels (integer) of each guiding center labeled by the quantum num ber m

$$jGi = {\begin{array}{*{20}c} \frac{N}{Y} & Y^{1} \\ m = 0 & n = 0 \end{array}} c^{+}_{n,m} jDi;$$

where $N = \frac{S}{2^{-v^2}}$ is the number of ux quanta. The neutral excitations in the non-interacting system can be generated by creating a hole in a Landau level p 1 at the guiding center m and an electron in a level n + p

1 at the guiding center m 0 .

We note, however, that it is not possible to de ne eigenstates of the quiding center \hat{R}_{0i} for the cyclotronic motion of a single charged particle because of the noncom mutativity of the components of the position operators. Nevertheless we can de ne a vector $\hat{R}_0 = \hat{R}_{0e} + \hat{R}_{0h}$, related to the center of mass of the guiding centers of the electron and the hole excited in the system, whose com ponents x_0 and y_0 commutes with each other and therefore have simultaneous eigenstates. It is also possible to de ne a momentum operator \tilde{P}_0 , canonically conjugated to \mathbb{R}_0 , which is related to the momentum of the center of m ass of the guiding centers. The eigenstates of this operator can be generated by a superposition of the above described neutral excitations in di erent guiding centers and they form a complete basis in the space of the neutral excitations known as magnetic excitons¹⁸. In operator form they can be described as,

$$e_{n,p}^{V}(k) jGi = e^{\frac{(k')^{2}}{4}} \int_{m^{0}\pi = 0}^{X^{1}} G_{m^{0}\pi}(k) C_{n+p,m^{0}}^{+} C_{p,m} jGi;$$

where $k = k_x + ik_y$, = $j_k j$ and $G_{m^0,m}$ is given by¹⁴

$$G_{n+pp}(k) = \frac{p!}{(n+p)!} \frac{ik}{p!} L_p^n \frac{2}{2}$$

with L_p^n being generalized Laguerre polynom ials.

The density uctuations of the system can be expressed in terms of superpositions $a_n^y\left(k\right)$ of these magnetic excitons

$$(k) \qquad {}_{0} = {\begin{array}{*{20}c} X^{k} \\ & a_{n}^{y} (k) + a_{n} (k) ; \\ & n = 1 \end{array}}$$

where

$$a_{n}^{y}(\mathbf{k}) = e^{\frac{(y)^{2}}{4}} \sum_{p=0}^{k} G_{n+p,p}(\mathbf{k}) e_{n,p}^{y}(\mathbf{k})$$
(1)

can be interpreted as magnetic plasmons and $_0 = \frac{1}{2^{-\sqrt{2}}}$ is the electronic average density in the system .

In order to bosonize this problem we proceed exactly like in the Luttinger model and enlarge the H ilbert space of the ferm ionic system by including led energy eigenstates of negative values of the quantum number n. This construction is the analogue of the sea of in nite negative energy of the Luttinger model (the Tom onaga construction can also be obtained in an analogous way by making the Hilbert space nite²¹). This vacuum does not concem us because we are studying the bw energy excitation spectrum of the system. It is clear from this construction that the bosons only make a faithful representation of the electronic system if >> 1. In the same way the bosonization of the electron gas is only possible when the system is very dense (large Ferm im om entum). By doing that one can show that the bosonic operators de ned in (1) obey the exact com mutation relations,

h i
a_n(k);
$$a_{n^0}^{y}(k^0) = a_{n;n^0 k;k^0} M G_n(k^{\circ})$$

 $[a_n(k);a_{n^0}(k^0)] = 0;$

where

$$G_n(x) = nJ_n^2(kR_c);$$

 J_n is a Bessel function of the rst kind and $R_c = \frac{p}{2}$ is the size radius of the last occupied \orbit". We can proceed further and de ne the canonical bosonic operators

$$b_n(k) = \frac{1}{N - G_n(k')} a_n(k)$$

and recover the standard bosonic com mutation relations.

It is easy to show that in term softhese bosonic degrees of freedom the free H am iltonian can be rew ritten, apart from a background energy, as

$$H_{o} = h!_{c} nb_{n}^{y}(k)b_{n}(k)$$
(2)
$$K_{k} n = 1$$

which describes the energy of an assembly of noninteracting bosons.

So far we have succeeded in expressing the neutral excitations of the system of electrons in term s of bosons. In order to complete our scheme one needs to express the electron operator in terms of bosons. This is done by transforming from the momentum representation which is the natural representation for bosons to the jnm i representation which is the appropriate one for ferm ions in a magnetic eld. This should be contrasted with the case of the electron gaswhere the momentum is a good quantum num ber for the electron and the particle-hole excitation. It means that we must construct an operator that creates an electron in a Landau level n and guiding centerm as a function of the density uctuations within this guiding center. This transform ation involves a linear superposition of the magnetic excitons at di erent momentum states. This is done by considering the operator

$$a_{n,m}^{y} = \frac{1}{p_{m}} \sum_{p=1}^{X^{k}} c_{n+p,m}^{y} c_{p,m}$$
$$= \frac{Z}{2} \frac{d^{2}k}{2} e^{-\frac{2}{4}} G_{m,m} (k) b_{n}^{y} (k):$$

Dierently from the operator $b_n^y(k)$ that creates density uctuations with a de nite momentum k, the operator $a_{n,m}^y$ generates density uctuations on a de nite

guiding center. This operator also obeys exact bosonic commutation relations. This is completely analogous to Luther's^{7;8} construction of the Fermi liquid in 2D and 3D as many 1D fermi liquids, one for each direction normal to the Fermi surface. But contrary to that construction here we do not need to \sm ear" the operators close to the Fermi surface. The discrete nature of the Landau states does this sm earing naturally. Here we have also reduced the 2D problem of electrons in a magnetic eld to N 1D chiral problem s, one for each guiding center labeled by m.

In order to complete the bosonic description of the ferm ion operator it will be necessary to de ne an operator that creates a ferm ion in a \phase" state associated with this circular motion. This is done by rewriting the original ferm ion operator as,

$$c_{m}$$
 () = $\frac{1}{\frac{p}{2}} \int_{n=1}^{X} e^{in} c_{n,m}$;

where we have used the enlarged H ilbert space and it is easy to show that

$$fc_{m}^{y} \circ ({}^{0}); c_{m} ()g = {}_{m} \circ_{m} ({}^{0})$$
(3)

$$f_{C_m \circ} ({}^{0}); c_m ()g = 0:$$
 (4)

T he operator c_m^y () creates an electron about the guiding center given by the quantum numberm and at the phase

of the cyclotronic motion. This is the operator that can be properly bosonized. Indeed the M and elstam form $^{4;5}$ of the ferm ion operator is given by

$$c_{m}^{y}() = \frac{1}{2}e^{i} e^{i \frac{y}{m}()}U_{m}e^{i m}();$$

with

m () =
$$\frac{N_m}{2}$$
 i $\lim_{\substack{n \ge 0 \\ n = 1}} e^{\frac{n}{2}} \frac{e^{in}}{p} a_{n,m}$

and

$$U_{m} = e^{s=0} e^{i m}$$

where the Herm itian operators

$$N_{m} = \begin{array}{c} X^{t} \\ c_{p,m}^{V} c_{p,m} < c_{p,m}^{V} c_{p,m} > \\ \end{array}$$

and m are canonically conjugated ($[N_m; m_0] = i_m o_m$) such that the unitary operators U_m anticom mute.

In this representation, it can be shown that the ferm ionic operators obey the desired anticom mutation relations (3) and (4) and it recovers the long time asymptotic behavior of the non-interacting propagator between two di erent phase states which is given by

$$K (;t) = \langle Q_{m_0}^{V}(;t) Q_{m_1}(0;0) \rangle$$
$$= \frac{e^{i}}{1 - e^{i(-1)e^{i}}}$$

and can be related to the space-time propagator by writing the operator (r) in terms of the c_m ($)^{21}$. That completes the construction of the model. Now we have an operator relation between electrons in a magnetic eld and bosons.

O ur next step is to include the interaction between the electrons within this fram ework. In an uniform neutralizing background the total H am iltonian can be written as

H = H
$$_{\circ}$$
 + H $_{I}$

where H $_{\circ}$ is given by (2) and

$$H_{I} = \frac{1}{2} X V(k) \int_{k}^{k} p_{n'} \frac{1}{G_{n'}(k')G_{n'}(k')} \int_{k}^{k} \frac{1}{h'_{n'}(k')} \int_{k}^{k} \frac{1}{h'$$

is a density-density interaction between the fermions which is written in bosonic language. V (k) is the Fourier transform of the interacting potential.

In order to diagonalize H we will adopt a generalized B ogoliubov transform ation¹⁰ with real coe cients $_{n1}$ (k) and $\#_{n1}$ (k) which is given by

$$b_{n}(k) = \int_{1}^{X} b_{n}(k) + \#_{n1}(k) + \#_{n1}(k$$

where $_{1}(k)$ are the new bosonic operators. This canonical transform ation yields

$$H = h!_{c} \qquad \begin{array}{c} X & X^{l} \\ H = h!_{c} & _{1}(k) & _{1}^{y}(k) & _{1}(k) \\ & _{k} & _{l=1} \end{array}$$

where the new eigenfrequencies $\ _1$ (k) are determined by the equation

$$l = \frac{V(k)}{h!_{c}^{2}} \prod_{n=1}^{X} \frac{G_{n}(k')n}{(1(k)^{2} n^{2})};$$

The above equation has the same form as the one obtained for the norm alm odes in the integer quantum H all e ect in the RPA approximation for a C oulom b potential, V () = $2 e^{2} = {}^{18}$. Moreover, the same equation is obtained in the fractional quantum H all e ect for com - posite ferm ions¹⁹. Now we have shown that as far as the bosonization construction is valid, the equation which de-

nes the eigenvalues holds for any value of the m agnetic eld. This is a consequence of the fact that bosonization ful lls the W ard identities. Thus, exactly like in the Ferm i gas, the lim it of validity of bosonization goes beyond w hat is expected.

As a last comment we would like to mention that the method developed in this letter can be applied to the fractional Hall e ect if we use the Chem-Sim ons theory to map the original electrons into the composite fermions. In this case, as it is well established, the fractional quantum Hall e ect is mapped into the integer quantum Hall e ect of composite fermions. The condition that >> 1

becomes equivalent to say that we are boking at llings close to $1=2^{22}$. The nature of the ground state of this system has recently been the source of much controversy and we believe that our methods can provide non-perturbative answers for the structure of the ground state of this problem 21 .

In conclusion, we have proposed a new method of bosonization for 2D ferm ions subject to an uniform magnetic eld by directly working in a Landau basis. We propose an analogue of the Luttinger model which can be bosonized and provides an operator relation between electrons and bosons in the limit of >> 1. This construction is essentially non-perturbative and provides new insights into the problem of electrons interacting in the presence of magnetic elds. We show that bosons are collective excitations of the system which are related with a collection of N 1D Luttinger liquids associated with each quiding center. W e also obtain the dispersion of the collective m odes of the system and discuss the applicability of the method to the integer and fractional quantum Halle ects. W e expect our method will enlarge the understanding of the physics of strongly correlated system s in the presence of magnetic elds.

H W .and A O C.kindly acknow ledge respectively the full and partial supports from the Conselho Nacional de Desenvolvim ento Cieni co e Tecnologico (CNPq). We are also grateful to E H.Fradkin, F D M.Haldane and D.V.K hveshchenko for very useful discussions.

- ¹ F.B bch, Z.P hys. 81, 363 (1933).
- 2 S.Tom onaga, Prog.Theor.Phys. (K yoto), 5, 544, (1950).
- ³ W. Thirring, Ann. Phys. (N. Y.) 3, 91 (1958); JM Luttinger, JM ath Phys. 4, 1154, (1963); D. M attis and E. Lieb, J. M ath. Phys. 6, 304 (1965); A. Luther, Phys. Rev. B 14, 2153 (1975); S. Coleman, Phys. Rev. D. 11, 2088 (1975).
- ⁴ S.M andelstam , Phys.Rev.D 11 ,3026 (1975).
- ⁵ F D M .Haldane, J.Phys.C 14 , 2585 (1981).
- ⁶ For a recent review on this subject see, J. Voit, Rep Prog Phys. 58, 977 (1995) and references therein.
- ⁷ A Luther, PhysRev B, 19, 320 (1979).
- ⁸ F D M .Haldane, Varenna Lectures (1992) and Helv.Phys. Acta. 65, 152 (1992).
- ⁹ A. Houghton and B. Marston, Phys. Rev. B 48, 7790 (1993); H.J. Kwon, A. Houghton and B. Marston, Phys. Rev. B 52, 8002 (1995).
- ¹⁰ A H. Castro N eto and Eduardo Fradkin, Phys. Rev. Lett. 72, 1393 (1993); Phys. Rev. B, 51, 4048 (1995).
- ¹¹ P. Kopietz and G E. Castilla, PhysRevLett. 76, 4777 (1996); PK opietz, J. Herm isson and K. Schonhammer, PhysRevB 52, 10877 (1995).
- ¹² R E. Prange and S M. G irvin eds., The Quantum HallEffect, 2nd ed. (Springer-Verlag, New York, 1990).
- ¹³ A Karlhede, SA. Kivelson, and SL. Sondhi, The Quantum Hall E ect: The article, Lectures pres. at the 9th Jerusalem W inter School on Theor. Phys. (1992) and ref-

erences therein.

- ¹⁴ A H. M acD onald, The Quantum HallE ectA Perspective, K huwer A cadem ic D ordrecht, 1989.
- ¹⁵ A H.M acD onald, Phys. Rev. B 30, 4392 (1984).
- ¹⁶ IL. A leiner and L.I. G lazman, PhysRev.B, 52, 11296, (1995).
- $^{17}\ {\rm L}$ S.Levitov and A V.Shytov, cond.m at./9507058.
- ¹⁸ C.Kallin and B.I.Halperin, Phys.Rev B, 30, 5655, (1984).
- ¹⁹ A Lopez and E Fradkin, Phys. Rev. B 44, 5246 (1991); Phys. Rev. Lett. 69, 2126 (1992); Nucl. Phys. B 33 C, 67 (1993).
- ²⁰ JP Eisenstein, LN. Pfei er and KW. West, Phys. Rev. Lett. 74, 1419 (1995).
- ²¹ H W estfahl Jr., A H. Castro Neto and A D. Caldeira, in preparation.
- ²² B. I. Halperin, P. A. Lee and N. Read, Phys. Rev.B 47, 7312 (1993).