

On the mass transport by a Burgers velocity field

Paolo Muratore Ginanneschi

January 20, 2022

NBI, Blegdamsvej 17 DK-2100, Copenhagen, Denmark

Abstract

The mass transport by a Burgers velocity field is investigated in the framework of the theory of stochastic processes. Much attention is devoted to the limit of vanishing viscosity (inviscid limit) describing the "adhesion model" for the early stage of the evolution of the Universe. In particular the mathematical foundations for the ansatz currently used in the literature to compute the mass distribution in the inviscid limit are provided.

PACS: 05.20, 05.40, 47.20, 47.15

1 Introduction

It was proposed by Zeldovich [1], that a possible model for the description of large scale dynamics of the mass distribution, in the early stage of evolution of the Universe, is provided by:

$$\begin{aligned} \partial_t v + v \cdot \nabla v &= 0; & v(x; t=0) &= v_0(x) \\ \partial_t \rho + \operatorname{div}(\rho v) &= 0; & \rho(x; t=0) &= \rho_0(x) \end{aligned} \quad (1)$$

Here, the mass density is driven by a rotation free velocity field, solution of the Riemann equation.

As it is well known, the Riemann equation with a gradient initial condition has multi-stream solutions [2],[3]. This phenomenon is easily understood in the Lagrangian picture of the flow: the point-mass particles evolving along the characteristics, collide after finite time. The study of the multi-stream solutions requires a very subtle analysis of the caustics formed after a finite time by the Lagrangian solutions of the Riemann equation [4].

One way to avoid multistreaming is to introduce the "adhesion model" (see [3], [5], [6], [7] with references therein).

$$\begin{aligned} \partial_t v + v \cdot \nabla v &= -v; & v(x; t=0) &= v_0(x) \\ \partial_t \rho + \operatorname{div}(\rho v) &= 0; & \rho(x; t=0) &= \rho_0(x) \end{aligned} \quad (2)$$

where the velocity field is now governed by the Burgers equation [8].

The introduction of a small diffusion term is expected to have a smoothing effect only in the regions where shocks are about to occur: the limit of vanishing viscosity of the Burgers equation, the inviscid limit, selects one solution for the Riemann equation.

The main disadvantage of the adhesion model (2) is the loss of a unique Lagrangian picture for both the velocity field and the mass density. The consequences are not only conceptual but also practical, since for (2) the solution of the Burgers equation does not provide immediately a definite algorithm which solves the continuity equation. The problem is therefore moved to the mass density evolution.

In the one dimensional case, because of the strong topological constraint, there is general agreement on the idea that for times long enough that shocks appear, an initial uniform density field evolves into a singular distribution which describes the formation of point-masses on a background of a smooth, diluted, density field. The point-masses are situated at the shock positions of the solution to the Burgers equation and may be referred to as macro-particles. The mass of a macro-particle is equal to the integral of the initial mass density extended over the interval of initial positions which fall into the given shock at some time prior to that under consideration.

In the two dimensional case the situation seems to be more subtle. Here we have the appearance of ruled surfaces, ribbons, where tangent planes touch the graph of the convex hull at a segment instead of a point and of triangles, corresponding to tangent planes with triple contact. In the Eulerian plane to these regions are respectively associated shock-lines and shock nodes.

A further source of complication is that a physically reasonable [9] initial condition for the Zeldovich model (1) is to assume the velocity potential, at time $t = 0$, given by a generalized Brownian motion in the space variables.

According to the adhesion model (2), in order to compute the mass density distribution at any time t , first, we should integrate the rotation free Burgers equation and then use the resulting velocity $v = v(x; t)$ to solve the ordinary differential equation (ODE)

$$\dot{x}_t = v(x_t; t); \quad x_0 = a \quad (3)$$

as a function of time and of the initial condition. Then the limit of vanishing viscosity should be taken to define the inviscid (weak) limit of every mass averaged smooth observable.

It is clear that such a program in the realistic three dimensional case is very resource demanding for its numerical implementation. Therefore the question naturally arises whether a regularized Lagrangian picture exists which simultaneously solves both the velocity field and the mass density equation.

In [3], Vergassola et al. proposed a numerical algorithm (the VDFN algorithm from now) based on the generalized Legendre transform of the potential field $(x; t; \cdot = 0)$. The underlying ansatz is that the values of x , performing the transform, define a solution of (3).

In recent works [6], [7] it has been proposed that a consistent regularization of (1) with a simple numerical implementation, can be found by introducing a small viscosity term also in the right hand side of the continuity equation.

In the present paper it is shown that a Lagrangian picture is uniquely associated to the Burgers equation. The characteristics of the Burgers equation are the realizations of the solution of a stochastic differential equation (SDE) describing a backward diffusion.

In such a context, the mass transport along the characteristics of the Burgers equation is described by a backward Fokker-Planck equation, whose solution is related to a difficult inverse problem when the given boundary conditions are the initial velocity potential and the initial mass density distribution. (section 2).

On the other hand the Lagrangian approach shows that the choice of the Burgers equation in order to regularize the Riemann equation, is not the natural one and it is "exact" only in absence of shocks. In section 3 the natural regularization is introduced. It is shown that it provides a simple algorithm in the inviscid limit for the solution of both the velocity field and the mass density. The limitation of such a procedure is that it imposes, in order to be exact, some restrictions on the initial conditions of the velocity field.

In section 4 the approaches previously introduced in the literature ([6], [7]) are reviewed in light of the results provided by stochastic calculus. In particular it is shown that the VDFN algorithm corresponds to an "effective" mass transport along the trajectories of the backward diffusion underlying the Burgers equation.

Furthermore the pair of PDEs specified by the Burgers equation together with the "effective" mass transport equation, has the Zeldovich model as the weak limit for vanishing viscosity for any initial data. The price to pay is in the artificial nature of the procedure.

The results of the present analysis suggest that both from the physical and the algorithmic point of view, the more natural way to introduce adhesion models is that one based on conservation laws [10].

2 Lagrangian picture for the Burgers equation

Let us consider the backward stochastic differential equation [11], [12], [13], [14]:

$$\begin{aligned} d_s!_s &= -v(!_s; t-s)ds + \sqrt{\frac{p}{2}} dw_s \quad s \leq t \\ !_0 &= x \end{aligned} \quad (4)$$

The Itô differential of the drift field along the flow is:

$$\begin{aligned} d_s v(!_s; t-s) &= \\ &= \int \left[\partial_t v(!_s; t-s) - v(!_s; t-s) \cdot \nabla v(!_s; t-s) + \frac{1}{2} \Delta v(!_s; t-s) \right] ds + \\ &+ \sqrt{\frac{p}{2}} dw_s \cdot \nabla v(!_s; t-s) \end{aligned} \quad (5)$$

where dw_t denotes, as usual, the stochastic differential of Brownian motion.

If $v(x; t)$ satisfies the Burgers equation, the equality holds:

$$v(x; t) = \langle v(!_t^{x;0}; 0) \rangle \quad (6)$$

where average $\langle \cdot \rangle$ is taken over the realizations of the diffusion defined by (4).

The physical meaning is clear: the Burgers equation defines a velocity field constant on the average over the random trajectories of (4). The conservation law allows a straightforward integration of the rotation free Burgers equation.

The basic object to be considered [15] is the transition probability defined by:

$$p_{(\cdot)}(y; s | x; t) = \langle !_0 = x, !_s = y \rangle \quad s \leq t \quad (7)$$

The transition probability (7) satisfies [15] in the variables $(y; s)$ the backward Fokker-Planck equation:

$$\partial_s p_{(\cdot)} + \text{div}(v p_{(\cdot)}) + \frac{1}{2} \Delta p_{(\cdot)} = 0 \quad (8)$$

with the final condition:

$$\lim_{s \rightarrow t} p_{(\cdot)}(y; s | x; t) = \delta(y - x) \quad (9)$$

Let us introduce the auxiliary stochastic process described by the SDE:

$$\begin{aligned} dz_s &= \sqrt{\frac{p}{2}} dw_s \\ z_0 &= x \end{aligned} \quad (10)$$

The transition probability density for this process at times s is easily found to be Gaussian with expectation value x and variance $2(p-s)$.

We can transform (7) into an average over the realizations of z_t :

$$p_{()}(y;s|x;t) = \langle (z_0 - x) (z_{t-s} - y) \frac{dP_{|t-s}}{dP_{z_{t-s}}}(z_{t-s}) \rangle \quad (11)$$

The Jacobian of this transformation is supplied by the famous Girsanov formula (see for example [15] or [16]):

$$\frac{dP_{|t-s}}{dP_{z_{t-s}}}(z_{t-s}) = \exp \left[\int_0^{z_{t-s}} \frac{r(z_u;t-u)}{2} dw_u - \int_0^{z_{t-s}} \frac{k r(z_u;t-u)^2}{4} du \right] \quad (12)$$

where the relation $v = r$ has been used. Furthermore along the trajectories z_s the Ito differential of the velocity potential becomes:

$$d_s(z_s;t-s) = \left[\partial_{t-s}(z_s;t-s) + \frac{1}{2} (z_s;t-s) \right] ds + \frac{1}{2} dw_s - r(z_s;t-s) \quad (13)$$

If we use (13) to eliminate the stochastic integral in (12) and we impose the normalization condition

$$\int_1^{z_1} dy p_{()}(y;s|x;t) = 1 \quad (14)$$

we find the velocity potential to be

$$(x;t) = -2 \ln \left(\frac{1}{4t} \right)^{\frac{D}{2}} \int_1^{z_1} dy \exp \left[\frac{(x-y)^2}{4t} - \frac{V(y)}{2} \right] \quad (15)$$

and

$$p_{()}(y;s|x;t) = \frac{\exp \left[\frac{(y-x)^2}{4(t-s)} - \frac{V(y)}{2} \right]}{\int_1^{z_1} dz \exp \left[\frac{(z-x)^2}{4(t-s)} - \frac{V(z)}{2} \right]} \quad (16)$$

It is easy to verify that (16) is a well defined Markovian transition probability for any $s < t$.

In the physical literature is more common to deal with stochastic calculus in the Stratonovich representation which has the advantage to preserve the ordinary rules of differential calculus [17].

If we express the stochastic Ito integral, appearing in (12), in terms of the corresponding Stratonovich's [18], then what we have done is nothing else than proving that the transition probability (7) is given by a Feynman path integral with Lagrangian:

$$L(x;\underline{x};s) = \frac{k\underline{x} + r(x;t-s)k^2}{4} - \frac{1}{2} V(x;t-s) \quad (17)$$

From (17) the derivation of (16) is then trivial.

The solution of the Burgers equation at arbitrary time takes the form :

$$v(x;t) = \int_1^{z_1} dy r V(y) p_{()}(y;0|x;t) \quad (18)$$

The knowledge of (7) solves also the problem of the forward evolution of a passive scalar driven by the Burgers equation. More explicitly if we consider:

$$\begin{aligned} \partial_t C + v \cdot \nabla C &= C + E \\ C(\mathbf{x}; 0) &= C_0(\mathbf{x}) \end{aligned} \quad (19)$$

where $E = E(\mathbf{x}; t)$ is an external forcing, then we have

$$C(\mathbf{x}; t) = \int_0^t \int_1^{\infty} dy C_0(y) p_{(\cdot)}(y; 0 | \mathbf{x}; t) + \int_0^t \int_1^{\infty} dy du E(y; u) p_{(\cdot)}(y; u | \mathbf{x}; t) \quad (20)$$

As a function of $(y; s)$ the transition probability (7) describes the mass transport along the characteristics (4) of the Burgers equation.

The equation with final condition

$$\partial_s + \text{div}(v \cdot) + \dots = 0; \quad (\mathbf{x}; t = T) = \rho_T(\mathbf{x}) \quad (21)$$

is readily solved for any $t \leq T$ by:

$$\rho(\mathbf{x}; t) = \int_1^{\infty} dz p_{(\cdot)}(\mathbf{x}; t | z; T) \rho_T(z) \quad (22)$$

On the other hand the boundary conditions for the Zeldovich model provide the initial mass distribution. In order to solve the mass density evolution in the interval $[0; T]$, we have, first, to solve the inverse problem:

$$\rho_0(\mathbf{x}) \exp\left[\frac{V(\mathbf{x})}{2}\right] = \int_1^{\infty} \frac{dz}{(4 - T)^{\frac{D}{2}}} \exp\left[\frac{(\mathbf{x} - z)^2}{4 - T} + \frac{(z; T)}{2}\right] \rho(z; T) \quad (23)$$

In principle, once we have solved equation (23), it would be possible to specify the mass density evolution in the interval $[0; T]$. The fact that the interval is finite, is not, from this point of view, a limitation since $\rho(\mathbf{x}; T)$ could be consistently used as a new initial condition at time $t = T$ to iterate the procedure, as far as the integral in (23) is convergent.

It is worth to note [13], [14], that the problem (23) can be reformulated by looking for the conditions that insure the existence of the forward stochastic process specified for any $t \in [0; T]$ by the probability density, solution of (21) with $\rho(\mathbf{x}; T)$, the unknown final condition to be determined.

The crucial point is that, in general, the transition probability density (7) does not have an inverse: because of the non-reversible nature of the diffusion dynamics, it satisfies only a semi-group property [15].

On the other hand, it is interesting to understand the physical meaning for the occurrence of the inverse problem (23), when our starting point is to find a coherent Lagrangian regularization for the Zeldovich model.

3 The forward diffusion

As it is clear from the above discussion, Burgers equation is equivalent to the SDE :

$$\begin{aligned} d_s !_s &= v(!_s; t) \frac{p}{2} ds + \frac{p}{2} dw_s; & !_0 &= x \\ d_s v(!_s; t-s) &= \frac{p}{2} dw_s - r v(!_s; t-s); & v(x; 0) &= v_0(x) \end{aligned} \quad (24)$$

The limit of vanishing viscosity of (24) is

$$\begin{aligned} d_s !_s &= v(!_s; t-s) ds; & !_0 &= x \\ d_s v(!_s; t-s) &= 0; & v(x; 0) &= v_0(x) \end{aligned} \quad (25)$$

On the other hand, the characteristic equation of the Zeldovich model (1) is

$$\begin{aligned} d_t t &= v(t; t) dt; & t_0 &= y \\ d_t v(t; t) &= 0 & v(a; 0) &= v_0(y) \end{aligned} \quad (26)$$

If $y_t^{x;0}$ is a solution of (26) in $[0; t]$ then $!_s^{x;0} = \frac{x; t}{t-s}$ is a well defined solution of (25) only provided that the solution for (26) exists and is unique in $[0; t]$.

If the latter condition is satisfied, (25), allows to solve the inverse problem implicit in the Riemann equation or in the equations for any passive scalar conserved along the flow :

$$d_t v\left(\frac{y; 0}{t}; t\right) = 0 \quad ! \quad v\left(\frac{y; 0}{t}; t\right) = v(y; 0) \quad (27)$$

Namely, if v is well defined for all $s \in [0; t]$ the Riemann equation is equivalent to

$$\partial_s v(x; t-s) - v(x; t-s) - r v(x; t-s) = 0; \quad v(x; 0) = v_0(x) \quad (28)$$

The straightforward consequence is the well known fact [2] that for any t such that no collision occurs between the Lagrangian particles, we have:

$$\begin{aligned} v(x; t) &= v_0\left(\frac{y; 0}{t}; t\right) \\ \frac{y; 0}{t} &= x - v_0(y) t \end{aligned} \quad (29)$$

It is worth to note that (29) and the steps leading to it are exactly the deterministic counterpart of (6) and of the procedure providing the integration of the Burgers equation in the particular case of rotation free initial conditions.

Beside the velocity field, the Zeldovich model requires the mass density field. It is clear already from the deterministic case that the knowledge of $!_t^{x;0}$, alone, does not provide the solution of the continuity equation:

$$\partial_t + \text{div}(v) = 0; \quad (\rho; t=0) = \rho_0(x) \quad (30)$$

The only expression consistent with (30) we can construct, is

$$\rho(y; t-s) = \int_{-1}^1 d^D x \, \rho(y - !_t^{x;0}) \rho(x; t) \quad (31)$$

where $(x;t)$ is a given solution of the continuity equation for $t \leq s$. Again the deterministic case is a mirror of the situation we faced studying the mass transport by the Burgers equation, i.e. equation (23).

The choice to regularize the Riemann equation by resorting to the Burgers equation is equivalent to select a solution for the Riemann's, when shocks occur, by extending the time reversal conjugation between the exact equation for the characteristics (26) and the deterministic limit of (24). This procedure is of clear advantage when dealing with the velocity field alone, but the price to pay consists in the difficulties which arise when turning to the problem of the mass evolution.

The alternative approach is to try to regularize the Riemann equation by looking to the stochastic generalization of its Lagrangian picture.

The SDE to be considered is then:

$$\begin{aligned} d_t t &= v(t;t) dt + \frac{\sigma}{2} dw_t; & v_0 &= y \\ d_t v(t;t) &= \frac{\sigma}{2} dw_t - r v(t;t); & v(x;0) &= v_0(x) \end{aligned} \quad (32)$$

Hence, the system of PDEs implied by (32) is

$$\begin{aligned} \partial_t v + v \cdot \nabla v + \frac{\sigma^2}{2} \Delta v &= 0; & v(x;t=0) &= r V(x) \\ \partial_t + \text{div}(v \cdot) &= 0; & (x;t=0) &= v_0(x) \end{aligned}$$

For a rotation free initial velocity field, the integration of the pair of equations (33) is achieved in terms of the transition probability density defined by (32), following the same steps leading to (16).

The result is, for any $s \leq t$:

$$p_{(+)}(x;t|y;s) = \frac{\exp\left[-\frac{(x-y)^2}{4(t-s)} - \frac{(x;t)}{2}\right]}{\int_1^{\infty} dz \exp\left[-\frac{(z-y)^2}{4(t-s)} - \frac{(z;t)}{2}\right]} \quad (33)$$

where the velocity potential is fixed by the normalization condition and is equal to

$$(x;t) = 2 \log \left[\frac{1}{4(T-t)} \right]^{\frac{D}{2}} \int_1^{\infty} dz \exp\left[-\frac{(x-z)^2}{4(T-t)} - \frac{(z;T)}{2}\right] \quad (34)$$

Therefore, in order to achieve the solution in the interval $[0;T]$ we need to solve the inverse problem:

$$\exp\left[\frac{V(x)}{2}\right] = \int_1^{\infty} \frac{dz}{(4(T-t))^{\frac{D}{2}}} \exp\left[-\frac{(x-z)^2}{4(T-t)} - \frac{(z;T)}{2}\right] \quad (35)$$

Equation (35) is the natural stochastic generalization of the inverse problem (27). Once $(x;T)$ is known, it can also be assumed as the new initial condition at time T to iterate the procedure.

In the limit of vanishing viscosity, equation (35) takes the form:

$$V(x) = \sup_z \left[(z;T) - \frac{(z-x)^2}{2T} \right] \quad (36)$$

The latter expression is basically a (generalized) Legendre transform [3]:

$$T[V](x) = \frac{(x)^2}{2} = \sup_z [T(z;T) - \frac{z^2}{2} + xz] \quad (37)$$

It is tempting to solve (36) as:

$$V^0(x;T) = \inf_z [V(z) + \frac{(z-x)^2}{2T}] \quad (38)$$

This solution is not exact for a general initial condition: due to the convexity properties of the Legendre transform, when we substitute $(x;T)$ back in (36) what we find is not $V(x)$ but its convex hull. Only for this latter the transform (36) is involutive. Therefore, by assuming $(x;T) = V^0(x;T)$ we implicitly adopt the consistent initial condition.

If we accept, despite of the above restrictions, the pair (33) as a regularization for the Zeldovich model then we are provided a simple algorithm for the computation in the inviscid limit of both the velocity field, by means of (34), and the mass distribution, by means of:

$$(x;t) = \int_1^{Z-1} dy p_{(+)}(x;t|y;0) V^0(y) \quad (39)$$

4 Non Lagrangian approaches

In [7] it has been proposed to regularize the Zeldovich model with the pair of PDEs:

$$\begin{aligned} \partial_t v + v \cdot \nabla v &= -\nabla \cdot v; & v(x;t=0) &= \nabla V(x) \\ \partial_t + \text{div}(v) &= 0; & (x;t=0) &= V^0(x) \end{aligned} \quad (40)$$

The meaning of (40) is to assume the velocity field solution of the Burgers equation as an external field in the Langevin equation describing the motion of a point mass particle:

$$dx_t = v(x_t;t) dt + \frac{\sigma}{2} dw_t; \quad x_0 = y \quad (41)$$

Hence, the acceleration felt by the point-mass particle is

$$dv(x_t;t) = (\nabla \cdot v)(x_t;t) dt + \frac{\sigma}{2} dw_t - \nabla v(x;t) \quad (42)$$

It is worth to note that (42) predicts for the case $\sigma = 0$ a mass dynamics different from (32) even in the limit of vanishing viscosity: only in absence of shocks the two dynamics become equivalent.

In the general case the solution of mass distribution resulting from (40) is difficult. The path integral approach clearly shows that it is equivalent to an Euclidean Schrodinger equation in a potential given by the Laplacian of the velocity potential of the Burgers equation. Namely, the transition probability for the mass density in (40) is

$$p(x;t|y;0) = e^{-\frac{(x;t)}{2}} K(x;t;y;0) e^{-\frac{(y;0)}{2}} \quad (43)$$

where K is given by

$$K(x;t;y;0) = \int_{x_0=y}^{x_t=x} D x_u \exp \left[- \int_0^t \left(\frac{x_u^2}{4} + \frac{v_0}{2} (x_u - y) \right) du \right] \quad (44)$$

and it satisfies

$$\partial_t K + \frac{v_0}{2} K = 0 \quad (45)$$

$$\lim_{t \rightarrow 0} K(x;t;y;0) = \delta(x-y) \quad (46)$$

There are two evident cases when the path integral (44) is of practical use. When the Laplacian of the velocity potential does not depend on the trajectory, all the integrations turn out to be Gaussian. This condition is satisfied for:

1. $V(x) = v_0 x$. Then the transition probability is:

$$p(x;t|y;0) = \frac{\exp \left[- \frac{(x - y - v_0 t)^2}{4t} \right]}{(4t)^{D/2}} \quad (47)$$

which, for every smooth enough initial mass distribution, gives for $t \neq 0$:

$$(x;t) = \delta_0(x - v_0 t) \quad (48)$$

in accordance with Galilean covariance.

2. $V(x) = \frac{kx^2}{2}$. By means of (44) we get into

$$p(x;t|y;0) = \left[\frac{1}{2t} \right]^{D/2} \frac{\exp \left[- \frac{1}{4t} \left(x - \frac{v_0}{2} y \right)^2 \right]}{(4t)^{D/2}} \quad (49)$$

In the inviscid limit for smooth enough distribution we obtain:

$$(x;t) = \left(\frac{1}{2t} \right)^{D/2} \delta_0 \left(\frac{x}{2t} \right) \quad (50)$$

The peculiarity of these two examples appears also in the fact that the results do not depend on the viscosity of the driving velocity field. Actually the initial conditions for the velocity potential given above, are such that no shock appears at any time and the Laplacian of the velocity field is zero.

More results can be derived for the case $\eta = 0$. The crucial observation is that the fundamental solution of the mass density problem can be rewritten as

$$\partial_t p + v \cdot \nabla p = - \nabla p \cdot \nabla v$$

$$\lim_{t \rightarrow s} p(x;t|y;s) = \delta(x-y) \quad (51)$$

This means that the transition probability p can be computed as an average over the random trajectories, solutions of (24)

$$p(x;t|y;s) = \langle J(t;s;f|g) \delta \left(\int_s^t \dot{x} - v \right) \rangle \quad (52)$$

where

$$J(t; s; f; g) = \exp \left[\int_0^{t-s} du \, r - v(x_a^{x;0}; t - u) \right] \quad (53)$$

is the Jacobian of the change of variables in the functional integration between the solutions of the backward diffusion (24) and those of (41).

In one dimension this observation leads to a straightforward integration of (51):

$$\langle J(t; s; f; g) (\delta_{t-s}^{x;0} - y) \rangle = \langle (\delta_{t-s}^{x;0} - y) \partial_x \delta_{t-s}^{x;0} \rangle = \partial_x \langle y - \delta_{t-s}^{x;0} \rangle \quad (54)$$

where δ is the step function. It follows:

$$p(x; t | y; s) = \partial_x \int_1^Z dz p_{(1)}(z; s | x; t) \quad (55)$$

One can easily check that (55) verifies (51) and it is a Markovian transition probability.

Unfortunately, the identity in (54) does not hold in more than one dimension.

For the general case, in [7] it has been introduced a "mean field approximation" for (51). Here a different interpretation of the result is proposed.

The mean field theory of [7] is equivalent to substitute to (51) the equation

$$\begin{aligned} \partial_t p + v \cdot r p &= p \cdot p \cdot \langle r \cdot v \rangle \\ \lim_{t \rightarrow s} p(x; t | y; s) &= \delta(x - y) \end{aligned} \quad (56)$$

where the average means

$$\langle r \cdot v \rangle = \int_1^Z d^D x \, r \cdot v(x; t) p(x; t | y; s) \quad (57)$$

Furthermore we need to impose, for every $t > s$, the constraint:

$$\int_1^Z d^D x \, p(x; t | y; t) = 1 \quad (58)$$

The integration of (56) is immediate. Since (57) does not depend on x , the corresponding term can be extracted from the path integral, which is reduced to an average over the realizations of (24):

$$p(x; t | y; s) = e^{\int_s^t du \langle r \cdot v \rangle} p_{(1)}(y; s | x; t) \quad (59)$$

The normalization condition then fixes the value of the prefactor. Finally we get into:

$$p(x; t | y; s) = \frac{\exp \left[\frac{(x - y)^2}{4(t - s)} + \frac{(x; t)}{2} \right]}{\int_1^Z d^D z \exp \left[\frac{(z - y)^2}{4(t - s)} + \frac{(z; t)}{2} \right]} \quad (60)$$

Here the velocity potential of the Burgers equation $(x; t)$ explicitly appears. The effect of the average (57) is to define an "effective" mass transport along the characteristics of the backward diffusion (24). The "effective" theory becomes an exact solution of the

continuity equation in the (weak) inviscid limit: for any smooth observable described by a scalar function $f(\mathbf{x})$, it is readily verified that

$$\lim_{\epsilon \rightarrow 0} \langle f(\mathbf{x}; t) \rangle = \lim_{\epsilon \rightarrow 0} \int_{-\infty}^{\infty} d^D \mathbf{x} d^D \mathbf{y} f(\mathbf{x}) p(\mathbf{x}; t | \mathbf{y}; s; \epsilon) \rho(\mathbf{y}) \quad (61)$$

satisfies in the weak sense the continuity equation and we have,

$$\begin{aligned} \lim_{\epsilon \rightarrow 0} \langle f(\mathbf{x}; t) \rangle &= \lim_{\epsilon \rightarrow 0} \int_{-\infty}^{\infty} d^D \mathbf{y} f[\mathbf{x}(\mathbf{y}; t)] \rho(\mathbf{y}) \\ \mathbf{x}(\mathbf{y}; t) &= \arg \sup_{\mathbf{z}} [(\mathbf{z}; t; \epsilon = 0) - \frac{(\mathbf{z} - \mathbf{y})^2}{2t}] \end{aligned} \quad (62)$$

where $(\mathbf{x}; t; \epsilon = 0)$ is the inviscid limit of velocity potential of the Burgers equation.

The latter result (62) is exactly the ansatz of the VDFN algorithm used in [3], in order to compute the evolution of the mass density field driven by the Burgers equation. It is also worth to note, that in the case of a convex differentiable velocity potential, equation (62) implies free motion for the point-mass particles.

5 Conclusion

The "natural" regularization (33) of the Zeldovich model by means of the introduction of a small viscosity coefficient leads to the inverse problem (35) which is the direct generalization of the one occurring in the deterministic case. The inviscid limit drastically simplifies the situation, although it imposes some restrictions on the initial conditions. Nevertheless, if we neglect such difficulty, the forward diffusion approach (33) provides us a simple algorithm to compute, in the inviscid limit, the velocity field and the mass distribution at any time (equations (34), (38) and (39)). This procedure is exact for convex initial conditions.

To select the solution of Riemann equation by extending at larger times the correspondence with the backward characteristics of the Burgers is of real advantage only if we are interested in the velocity field alone.

The occurrence of inverse problems can be avoided if we associate to the Burgers equation the "effective" equation (56) for the mass transport along the trajectories of the backward diffusion (24). The solutions (15) and (62) of the pair of equations specified by the Burgers together with its "effective" mass transport tend in the limit of vanishing viscosity to a weak solution of the Zeldovich model.

For the mass density field, such solution (equation (62)) was proposed as an ansatz in [3]. The algorithm defined by (62) has the advantage of an easy numerical implementation on a computer like that one provided by the "natural" regularization (33). Furthermore in comparison with the latter it does not impose any restriction on the initial conditions.

The main disadvantage of (62) is its intrinsic non-locality which makes its use artificial from the microscopic point of view.

The interpretation of the transition probability (60), defined by the "effective" theory (56), as an approximate solution of the exact mass transport (51) by a forward diffusion

with external drift field given by the Burgers equation, is certainly correct only before the occurrence of shocks. For larger times the characteristic equation (41) with $\gamma = \gamma_{\text{ext}}$ seems to indicate a different behavior over shock domains. Only in one dimension, heuristic arguments can be provided, [7], to show the equivalence, in the weak inviscid limit, of (60) with the mass transport described by the exact transition probability (55).

Finally it must be remarked that starting from the basic kinetic equations, the Zeldovich equations (1) are not the unique starting point for the construction of adhesion models. Let us consider

$$\partial_t f + p \cdot \nabla_x f = 0 \quad (63)$$

where $f = f(x; p; t)$. The ansatz

$$f(x; p; t) = \delta(x; t) [p - v(x; t)] \quad (64)$$

leads to the equations

$$\begin{aligned} \partial_t v + (v \cdot \nabla) v &= 0 \\ \partial_t + \nabla \cdot (v) &= 0 \end{aligned} \quad (65)$$

By means of the continuity equation, the system can be recasted in the form :

$$\begin{aligned} \partial_t (v_i) + \nabla_j (v_j v_i) &= 0 \\ \partial_t + \nabla_j (v_j) &= 0 \end{aligned} \quad (66)$$

In [10], it is proven that the latter pair of PDEs is equivalent to the Zeldovich model only in absence of shocks or for an uniform initial mass distribution.

The simple meaning in terms of conservation laws of equations (66) allows to achieve an exact algorithm for the solution which turns out to be of simple numerical implementation [10].

Acknowledgments

This work is the result of many discussions with Erik Aurell whom I have to thank for precious explanations and warm hospitality. I also benefited from many explanations and comments by Yuri G. Rykov, Andrei Sobolevsky, Bala Ram Dethlefsen and Gabriele Travaglini Battino Vittorelli.

This work was supported by a 1996 TAO-grant from the European Science Foundation.

The revision of the paper has been done during my stay at the CATS group at Niels Bohr Institute with the support of the TM R grant ERB 4001G T 962476 by the European Commission.

I thank Mogens H. Jensen and all the other cats for useful discussions. During the revision of the paper I also benefited from some comments by Piotr Garbaczewski which I am pleased to acknowledge.

References

- [1] Ya.B. Zeldovich, *Astron. Astrophys.*, 5 (1970), pp. 84-89.
- [2] S. Gurbatov, A. Malakhov and A. Saichev *Nonlinear Random Waves and Turbulence in Nondispersive Media: Waves, Rays and Particles* Manchester U.P., Manchester, 1991.
- [3] M. Vergassola, B. Dubrulle, U. Frisch and A. Noullez, *Astron. Astrophys.*, 289 (1994), pp. 325-356.
- [4] V.I. Arnold Singularities of caustics and wave fronts Kluwer, Dordrecht (1990).
- [5] J.D. Fournier and U. Frisch *J. de Mec. Theor. et Appl.*, 2, (1983) pp. 699-750.
- [6] A.I. Saichev and W.A. Woyczynski *SIAM J. Appl. Math.* Vol 56, No 4, (1996) pp. 1008-1038.
- [7] A.I. Saichev and W.A. Woyczynski (1996) to appear in *Physica D*.
- [8] J.M. Burgers *The Nonlinear Diffusion Equation* D. Reidel, PublCo. (1974).
- [9] Z. She, E. Aurell and U. Frisch, *Comm. Math. Phys.* 148, (1992) pp. 623-641
- [10] E. Weinan, Yu.G. Rykov and Ya.G. Sinai, *Comm. Math. Phys.* (1996) pp. 349-380
- [11] F. Guerra *Phys. Rep.* 77, (1981) pp. 263-312.
- [12] F. Guerra and L.M. Morato, *Phys. Rev. D* 27 No 8, (1983) pp. 1774-1786.
- [13] P. Garbaczewski and G. Kondrat, *Phys. Rev. Lett.* 77 (1996) pp. 2608-2611.
- [14] P. Garbaczewski, G. Kondrat and R. Otkiewicz *Phys. Rev. E* 55 No 2 (1997) pp. 1401-1411.
- [15] I. Karatzas and S.E. Shreve *Brownian Motion and Stochastic Calculus* Springer-Verlag Berlin (1990).
- [16] I.I. Gihman, A.V. Skorohod *The theory of stochastic processes Vol. III*, Springer-Verlag, Berlin (1979).
- [17] L.S. Schulman *Techniques and applications of path integration* John Wiley & Sons (1981)
- [18] L. Bertini, G. Jona-Lasinio and C. Parrinello, *Prog. Theor. Phys. Suppl.* 111 (1993) pp. 83-113.