

# A new family of models with exact ground states connecting smoothly the $S = \frac{1}{2}$ dimer and $S = 1$ Haldane phases of 1D spin chains

A. K. Kolezhuk

Institute of Magnetism, National Academy of Sciences and Ministry of Education of Ukraine, 252142 Kiev, Ukraine  
and Institut für Theoretische Physik, Universität Hannover, 30167 Hannover, Germany

H.-J. Mikeska<sup>y</sup>

Institut für Theoretische Physik, Universität Hannover, 30167 Hannover, Germany  
and Department of Earth and Space Science, University of Osaka, Toyonaka, Osaka 560, Japan  
(November 11, 2021)

We investigate the isotropic two-leg  $S = \frac{1}{2}$  ladder with general bilinear and biquadratic exchange interactions between spins on neighboring rungs, and determine the Hamiltonians which have a matrix product wavefunction as exact ground state. We demonstrate that a smooth change of parameters leads one from the  $S = \frac{1}{2}$  dimer and Majumdar-Ghosh chains to the  $S = 1$  chain with biquadratic exchange. This proves that these model systems are in the same phase.

74.20.Hi, 75.10.Lp, 71.10.+x

Low dimensional quantum antiferromagnets have attracted a large amount of both theoretical and experimental interest in recent years. Theoretically, the results of calculations for model systems in one dimension (1D) clearly show the difference between systems with a gapless spectrum of excitations and power law decay of spin correlations and gapped systems with exponentially decaying correlation functions. The main representatives of these two classes are resp. the  $S = \frac{1}{2}$  chain with isotropic nearest neighbor (NN) exchange interaction [1] and the isotropic  $S = 1$  (Haldane) chain [2]. Gapped elementary excitation spectra are also found in more complicated low-dimensional  $S = \frac{1}{2}$  systems such as chains with sufficiently strong additional next-nearest neighbor (NNN) exchange or with alternating exchange (with noninteracting dimers as the simplest limit) and also isotropic spin ladders with an even number of legs [3].

Many of these models are realized to a high degree of accuracy in simple compounds as demonstrated by the following examples:  $\text{KCuF}_3$  (1D isotropic  $S = \frac{1}{2}$  antiferromagnet [4]),  $\text{CaCuO}_2$  (weakly interacting dimers [5]),  $(\text{VO}_2)\text{P}_2\text{O}_7$  [6] and  $\text{SrCu}_2\text{O}_3$  [7] (two leg spin ladder) and  $\text{Ni}(\text{C}_2\text{H}_8\text{N}_2)_2\text{NO}_2\text{CD}_4$  (NENP, 1D  $S = 1$  antiferromagnet [8]). Although the precise spin Hamiltonians of these substances are not known, the study of the simple theoretical models mentioned above has contributed in an essential way to the understanding of the behaviour of the real materials and these are believed to be in the same phase as the simple models.

In recent years arguments and numerical evidence have been presented that the various models with a gapped ex-

citation spectrum, in particular the  $S = \frac{1}{2}$  dimer chain, the gapped nearest and next nearest neighbor exchange chain, the two-legged  $S = \frac{1}{2}$  spin ladder and the  $S = 1$  (Haldane) chain are all in the same phase [9-14]. In this letter we present the following proof for this conjecture: We describe a class of Hamiltonians with exactly known nondegenerate ground states and show that it is possible, by a continuous change of parameters within this class, to connect smoothly the Hamiltonians of the 1D dimer chain, the Majumdar-Ghosh chain, a generalized spin ladder (which includes biquadratic interactions) and of the  $S = 1$  antiferromagnetic chain with additional biquadratic exchange. Given the relevance of these model systems with exactly known ground states for the traditional model systems and for the real quasi 1D compounds this proves the existence of a single phase for the gapped low dimensional spin systems above. As a byproduct of this investigation, we also present a new set of models of frustrated  $S = \frac{1}{2}$  spin chains (including only bilinear NN and NNN interactions) whose ground states can be found exactly.

Our approach starts from the observation [13] that the exactly known ground states of the Shastry-Sutherland  $S = \frac{1}{2}$  chain [15]

$$H = \sum_n \left( (1 + (-1)^n) S_n S_{n+1} + \frac{1}{2} (1 - (-1)^n) S_n S_{n+2} \right); \quad (1)$$

which includes the Majumdar-Ghosh ( $\alpha = 0$ ) [16] and dimer ( $\alpha = 1$ ) limits as special cases, and the  $S = 1$  chain with biquadratic exchange

$$H = \sum_n S_n S_{n+1} + \alpha (S_n S_{n+1})^2 \quad (2)$$

at  $\alpha = \frac{1}{3}$  (the AKLT model [17] which is believed to be in the same phase as the Haldane chain with  $\alpha = 0$ ) have very similar structure: both ground states can be written as matrix product (MP) wave functions,

$$|\psi\rangle = \sum_i \mathbf{g}_i; \quad \mathbf{g}_i = \sum_{j_1, j_2} \frac{b_{j_1} \mathbf{j}_i + a_{j_2} \mathbf{j}_i}{2a_{j_1} \mathbf{j}_i - b_{j_2} \mathbf{j}_i} : (3)$$

The ground state of the Shastry-Sutherland chain is obtained setting  $a = b = \frac{1}{2}$ , where  $\mathbf{j}_i$  and  $\bar{\mathbf{j}}_i$  are the singlet, resp. triplet formed by two spins  $S = \frac{1}{2}$  on adjacent



$$\begin{aligned}
j_0^{(2)} i &= (3 + u^4)^{-1/2} \frac{P}{3} j_0 s i + u^2 j_0 i = 0 \\
j_1^{(2)} i &= (1 - \frac{P}{2}) (j_1 s i - j_1 t i) \\
j_1^{(3)} i &= (1 + u^2)^{-1/2} (1 - \frac{P}{2}) (j_1 s i + j_1 t i) + u j_1 i = 1 \\
j_2 i &= j_2 i = 2
\end{aligned} \quad (10)$$

Conditions (i) result in 5 equations, corresponding to

$$e_0^{(1,1)} = e_0^{(1,2)} = e_1^{(1,1)} = e_1^{(1,2)} = e_1^{(1,3)} = 0: \quad (11)$$

Thus, in order to lead to M P ground states, the Hamiltonian has to project on the space  $j_0^{(2)} i; j_1^{(2)} i; j_1^{(3)} i; j_2 i$  and condition (ii) results in the inequalities

$$e_0^{(2)} > 0; \quad e_1^{(1)} > 0; \quad e_2 > 0 \quad (12)$$

where  $e_1^{(i,j)}$  ( $i = 1; 2$ ) are the two eigenvalues of the matrix  $\begin{pmatrix} i & j \end{pmatrix}$ ; ( $i, j = 2; 3$ ). The inequalities (12) guarantee that the ground state is nondegenerate (apart from the fact that Eq. (3) describes four ground states with different behaviour at the boundaries of open ladders), since it can be shown by induction with respect to the ladder length that the plaquette states of Eqs. (8,9) do not allow a zero energy state which is different from the M P ground state.

For an explicit discussion of the results we use a somewhat simplified Hamiltonian with less freedom than in the general case: First, we set  $J_0 = J_0^0$ , and from one of the conditions (i) we find that this necessarily means  $J_1 = J_1^0$  (which respects the ladder symmetry) and  $e_1^{(1,2)} = 0$ ; thus the eigenvalues  $e_1^{(i,j)}$  are identical to the  $e_1^{(2,2)}$  and  $e_1^{(3,3)}$ . Further we set  $K = K^0 = K^\infty = 0$  (these combinations are found irrelevant in a more detailed treatment which will be published separately). Then we are left with the 6 coefficients  $C_0, J_0, J_1, J_2, J_2^0, K$  and the conditions (i) take the following form

$$\begin{aligned}
2J_1 - J_2 - J_2^0 + u^2 (4C_0 - 3J_0 + K - 2) &= 0 \\
2(2J_1 + J_2 + J_2^0) - 4C_0 - J_0 - 7K &= 2 \\
&+ u^2 (J_2 + J_2^0 - 2J_1) = 0 \\
2(J_2^0 - J_2) + u (4C_0 - J_0 + 2J_1 - J_2 - J_2^0 - K) &= 0 \\
2J_1 + J_2 + J_2^0 + 3K &= 2 - 4C_0 - J_0 + 2u (J_2 - J_2^0) = 0
\end{aligned} \quad (13)$$

The general solution of these 4 linear equations contains 2 arbitrary constants, when the parameter  $u = b/a$  of the M P wavefunction is fixed. We absorb one of the two constants in the energy scale, denote the remaining one by  $x$  and define

$$u^2 = F$$

to obtain the following family of Hamiltonians with exact M P eigenstate:

$$\begin{aligned}
C_0 &= 3/9 + 2x - 3(1+x)F + 3xF^2 = 32; \\
J_0 &= 3/2 - x - (1+x)F + xF^2 = 4;
\end{aligned}$$

$$\begin{aligned}
2J_1 + J_2 + J_2^0 &= 15 - 9(1+x)F + 7xF^2 = 4; \\
2J_1 - J_2 - J_2^0 &= xF; \\
2J_2 - 2J_2^0 &= F^{1/2} (3(1+x) - 2 - xF); \\
K &= 3(F - 1)(xF - 1) = 4;
\end{aligned} \quad (14)$$

With these definitions the conditions of Eq. (12) that the M P state is a ground state are obtained as follows

$$\begin{aligned}
e_0 &= x(3 + F^2(x)) = 4 > 0 \\
e_1^{(1)} &= (3(1+x) + 2xF(x)) = 8 > 0 \\
e_1^{(2)} &= (1 + F(x))(3(1+x) - 2xF(x)) = 8 > 0 \\
e_2 &= 18 + 8xF^2(x) - 9(1+x)F(x) = 8 > 0
\end{aligned} \quad (15)$$

The eigenvalues  $e_0$  and  $e_1^{(1)}$  are seen to be positive for  $x > 0, F > 0$ . The eigenvalues  $e_1^{(2)}$  and  $e_2$  require a more detailed discussion; it is easily seen that a sufficient (though not necessary) condition for  $e_2 > 0$  is  $J_{\text{eff}} > 0$ , i.e. an effective antiferromagnetic interaction between the two rungs of one plaquette, and the condition on  $e_1^{(2)}$  is equivalent to the requirement  $u(J_2 - J_2^0) > 0$ . It is convenient to discuss the possible Hamiltonians considering  $F$  as function of  $x$ . We use  $F = u^2$  instead of  $u$  keeping in mind that changing the sign  $u \rightarrow -u$  amounts just to interchanging the ladder legs and thus does not bring in any new physics; therefore from now on we assume that  $F^{1/2} > 0$ .

The following illustrative members of this family of Hamiltonians are now easily obtained:

(i) Shastry-Sutherland model:

$$F = 1; \quad x = 3/(2 + \sqrt{3}); \quad 0 < x < 1: \quad (16)$$

Since the M P wave function with  $F = 1$ , i.e.  $a = b$  corresponds to singlets on one type of diagonals [13], this also covers (after a translation of one of the ladder legs) the case  $a = 0$ . The eigenvalue  $e_0$  vanishes for  $x = 0$ , which is to be expected since in the Majumdar-Ghosh limit a second degenerate eigenstate exists (singlets on the alternative bonds).

The solution (16) also applies for partially ferromagnetic interactions,  $x > 1$ , resp.  $(1 - \sqrt{3})/(1 + \sqrt{3}) < 0$ . Although the effective interaction between units on the diagonals is ferromagnetic, the Shastry-Sutherland dimer state is the ground state up to  $x = 9$  (corresponding to  $x = 3$ ), since all  $e$ 's are positive. At the singular point  $x = 3$  the eigenvalue  $e_2$  vanishes and the energy of the dimer ground state coincides with the energy of the fully polarized ferromagnetic configuration. Thus this is the point of the first-order quantum phase transition from dimer to ferromagnetic phase.

(ii) Generalized AKLT model, defined by  $F = 0$ , i.e.  $b = 0$  in the M P wave function (so that only triplets on the rungs occur):

$$F = 0; \quad x \text{ finite} \quad (17)$$

When choosing a convenient scale factor to render  $J_{\text{eff}} = 1$  this corresponds to the following Hamiltonian:

$$\begin{aligned} J_0 &= 4=3 \quad 2x=3; \quad J_1 = J_2 = J_2^0 = 5=6; \\ K &= K^0 = 2=3; \quad K^{\infty} = 0; \quad C_0 = 3=4 + x=3: \end{aligned} \quad (18)$$

This is essentially the Hamiltonian of Ref. [17], but without requiring explicitly the coupling of two spins  $S = \frac{1}{2}$  into a triplet, a result that could be easily obtained directly. The AKLT model in a strict sense is obtained for  $x \rightarrow +1$  and  $x F(x) \rightarrow 0$ .

(iii) For  $x F(x) = 1$  another class of Hamiltonians without biquadratic terms is obtained; the condition  $\alpha > 0$  leads to the restriction  $x = \frac{1}{9}$ . A nontrivial case is  $x = \frac{1}{9}$ , which leads to the presence of only one diagonal exchange interaction, equivalent to a chain with NN and NNN exchange,

$$\begin{aligned} C_0 &= 7=12; \quad J_0 = J_0^0 = 2=3; \quad J_1 = J_1^0 = J_2^0 = 1; \\ J_2 &= K = K^0 = K^{\infty} = 0: \end{aligned} \quad (19)$$

This is a  $S = \frac{1}{2}$  chain with alternating ferro- and antiferromagnetic exchange and ferromagnetic NNN interactions, a frustrated Heisenberg chain. As in (i), the eigenvalue  $\alpha$  vanishes and the energy of the fully polarized ferromagnetic state coincides with that of the MP ground state (3). Thus we have obtained two exact points on the line where the chain with NN and NNN interactions [13] undergoes a first order quantum phase transition to the ferromagnetic state.

(iv) A family of models with exact MP ground states connecting the Majumdar-Ghosh and the AKLT limits is obtained from e.g.

$$F(x) = 1 = (1+x)^2: \quad (20)$$

Since the Majumdar-Ghosh and dimer chains have identical ground states, Eq. (20) explicitly demonstrates the possibility to transform the Hamiltonian of the dimer  $S = \frac{1}{2}$  chain continuously to the Hamiltonian of the AKLT chain without any singularity (or quantum phase transition) in the ground state.

This provides an exact proof that from the point of view of theoretical concepts no essential difference exists between the Haldane gap and the dimer gap and that the gapped  $S = \frac{1}{2}$  models listed in the introduction belong to the same phase as the Haldane  $S = 1$  chain.

The authors gratefully acknowledge financial support from Deutsche Forschungsgemeinschaft (AK) and Volkswagenstiftung (HJM) and the hospitality of the Hannover Institute for Theoretical Physics (AK) and the Department of Earth and Space Science at Osaka University (HJM). AK was also supported by the grant 2.4/27 from the Ukrainian Ministry of Science.

Permanent address: Institute of Magnetism, 36(b) Vernadskii av., 252142 Kiev, Ukraine.

Permanent address: Institut für Theoretische Physik, Universität Hannover, 30167 Hannover, Germany.

- [1] H. A. Bethe, Z. Physik 71, 205 (1931).
- [2] F. D. M. Haldane, Phys. Rev. Lett. 50, 1153 (1983).
- [3] E. Dagotto and T. M. Rice, Science 271, 618 (1996).
- [4] S. E. Nagler, D. A. Tennant, R. A. Cowley, T. G. Perring and S. K. Satija, Phys. Rev. B 44, 12361 (1991).
- [5] A. Zheludov, G. Shirane, Y. Sasago, M. Hase and G. Uchinkura, Phys. Rev. B 53, 11642 (1996).
- [6] R. S. Eccleston, T. Barnes, J. Brody and J. W. Johnson, Phys. Rev. Lett. 73, 2626 (1994).
- [7] M. Azuma, T. Hiroi, M. Takano, K. Ishida and Y. Kitaoka, Phys. Rev. Lett. 73, 3463 (1994).
- [8] J. P. Renard, V. Gadet, L. P. Regnault and M. Verdaguer, J. Magn. Magn. Mat. 90-91, 213 (1990).
- [9] M. Kohmoto and H. Tasaki, Phys. Rev. B 46, 3486 (1992).
- [10] K. Ishida, J. Phys. Soc. Jpn. 60, 1347 (1991); 64, 4896 (1995).
- [11] N. Katoh and M. Imada, J. Phys. Soc. Jpn. 63, 1437 (1995).
- [12] K. Totsuka and M. Suzuki, J. Phys.: Condens. Matter 7, 6079 (1995).
- [13] S. Brehm, H.-J. Mikeska and U. Neugebauer, J. Phys.: Condens. Matter 8, 7161 (1996).
- [14] S. R. White, Phys. Rev. B 53, 52 (1996).
- [15] B. S. Shastry and B. Sutherland, Phys. Rev. Lett. 47, 964 (1981).
- [16] C. K. Majumdar and D. K. Ghosh, J. Math. Phys. 10, 1399 (1969).
- [17] I. A. A. Kennedy, E. Lieb and H. Tasaki, Phys. Rev. Lett. 59, 799 (1987); Commun. Math. Phys. 115, 583 (1988). 2578 (1987).
- [18] M. Fannes, B. Nachtergaele and R. F. Werner, Europhys. Lett. 10, 633 (1989); Commun. Math. Phys. 144, 443 (1992).
- [19] T. Kennedy and H. Tasaki, Phys. Rev. B 45, 304 (1992).
- [20] A. Klumper, A. Schadschneider and J. Zittartz, J. Phys. A 24, L955 (1991); Europhys. Lett. 24, 293 (1993).
- [21] H. Niggemann and J. Zittartz, Z. Physik B 101, 289 (1996).
- [22] Gang Su, Phys. Lett. A 213, 93 (1996).
- [23] K. Totsuka and M. Suzuki, J. Phys.: Condens. Matter 7, 1639 (1995).
- [24] A. K. Kolezhuk, H.-J. Mikeska and Shoji Yamamoto, to appear in Phys. Rev. B 55 (1997).

FIG. 1. Generalized spin- $\frac{1}{2}$  ladder with all possible isotropic exchange interactions between nearest-neighbor rungs. Ovals show spin pairs whose states are involved in the elementary matrix  $g_1$  defined by (3). Couplings are described in the text, see (4).