

The Canonical Flux Quantization and IQHE

F.GHABOUSSI

Department of Physics, University of Konstanz

P.O. Box 5560, D 78434 Konstanz, Germany

E-mail: ghabousi@kaluza.physik.uni-konstanz.de

Abstract

It is shown that the canonical flux quantization, which is described by the uncertainty relation on the phase space of the flux system, can result in the quantization of Hall-measures. Further it is shown that the polarization of this phase space, which is necessary for its quantization, results in the vanishing of longitudinal resistivity and conductivity. The equivalence between this approach and the topological approach to QHE is also discussed.

From the preference of edge currents in IQHE and the restriction of electronic currents in the theory of flux quantization to the boundary regions of superconducting ring, one deduces that there must be a connection between the IQHE and Flux quantization or superconductivity. Hence there is also hints about such a connection in case of FQHE [1]. We discussed such a connection already in a previous paper [2]. Here we will show that one can derive a possible background for the IQHE from the flux quantization.

It is well known that the flux quantization $\oint eA_m dx^m = \int \int eF_{mn} dx^m \wedge dx^n = \Phi = \mathbf{Z}h; m, n = 1, 2$ is an experimentally varified fact, however the conventional theory of flux quantization does not match the canonical quantization. Nevertheless in principle not only the flux quantization but also the QHE ,as quantum theories, should be described by a canonically quantized theory.

We showed recently that the flux quantization can be described as the canonical quantization of $\oint A_m dx^m$ on the $\{A_m, x^m\}$ -phase space by the commutator postulate $e[\hat{A}_m, \hat{x}_m] = -i\hbar$. Thus, there must be an equivalent uncertainty relation that can appear in two dependent forms:

$$\begin{aligned} e\Delta A_m \cdot \Delta x_m &\geq \hbar \\ eB \cdot |\Delta x_m| \cdot |\Delta x_n \epsilon_{mn}| &\geq \hbar, \end{aligned} \tag{1}$$

which are related with each other by the Landau gauge $A_m := Bx_n \epsilon_{mn}$ [3].

Recall that the canonical flux quantization with respect to the $\int \int F_{mn} dx^m \wedge dx^n = eB \int \int dx^m \wedge dx^n = \mathbf{Z}h$ action can also be described by $eB[\hat{x}_m, \hat{x}_n] = -i\hbar \epsilon_{mn}$ commutator which is related with the second form of the uncertainty relation (1) and which is known as the non-commutativity of the relative coordinates of the cyclotron motion of electron [4]. This commutator is related with the above one by the Landau gauge.

If we consider the minimum value of position uncertainty $(\Delta x_m)_{(min)} = l_B$, then we obtain from the second uncertainty relation the defining relation of the magnetic length: $l_B^2 := \frac{\hbar}{eB}$. Thus, one has also the uncertainty equations:

$$e\Delta A_m \cdot l_B = eB \cdot l_B^2 = \hbar \tag{2}$$

Here we will show that the uncertainty relations (1) or (2) can also define the quantization of the Hall- conductivity or resistivity. Recall also that the mentioned commutator $[\hat{x}_m, \hat{x}_n] = -i\ell_B^2 \epsilon_{mn}$ is an example of the non-commutative geometry of configuration space of cyclotron motion, hence there is also a quantization of the Hall-measures according to the non-commutative geometry [5].

We will show also that, if the flux $\Phi := \oint e A_m dx^m = \int \int F_{mn} dx^m \wedge dx^n$ is considered as a canonical action functional on the phase space of the related system, then its variation results in the correct equations of motion of photons. Furthermore, it will be shown that according to the "geometric" quantization of flux, the necessary polarization of the phase space of flux results in the vanishing of longitudinal resistivity and conductivity in QHE [6]. In other words, the canonical quantization of flux should explain both: The quantization of Hall- measure and the vanishing of longitudinal components, in a consistent manner.

To discuss the quantization let us first consider that in the Ohm's equation the Hall resistivity is given as the proportionality factor $\rho_H = \frac{\partial_t A_m}{ne \partial_t x_n} \epsilon_{mn}$ or $\rho_H = \frac{\partial A_m}{ne \partial x_n} \epsilon_{mn}$ which is equivalent to its semi-classical definition $\rho_H := \frac{B}{ne}$, where n is the global density of electrons on the QHE sample. These both definitions are however semi-classical definitions as one expects from the usual (semi-classical) Hall-effect. Whereas in the quantized case, i. e. in QHE, the relation between A_m - and x_m expectation values should be governed either by the expectation values of the above discussed commutators or by the equivalent uncertainty relation (1). In other words, in such a quantum case, one has to do always with dependent variations of A_m and x^m so that: $\partial A_m \geq \Delta A_m$ and $\partial x_m \geq \Delta x_m$ which are related by (1) or (2).

Therefore, in the QHE case the quantized Hall resistivity should be given instead of $\rho_H = \frac{\partial A_m}{ne \partial x_n} \epsilon_{mn}$, consistently, by its quantized version where the differentials ∂A_m and ∂x_m are replaced by the related finite values of ΔA_m and Δx_m uncertainties:

$$\begin{aligned} \rho_H^Q &= \frac{\Delta A_m}{ne \Delta x_n} \epsilon_{mn} , \\ e \Delta A_m \cdot \Delta x_m &= e B \Delta x_m \Delta x_n \geq \hbar , \end{aligned} \tag{3}$$

which means that the uncertainty relation should be fulfilled by the ρ_H^Q relation.

If we replace ΔA_m in the ρ_H^Q relation according to the first uncertainty relation (1), then we arrive at:

$$\rho_H^Q = \epsilon_{mn} \frac{\hbar}{ne^2 \Delta x_m \cdot \Delta x_n} \quad (4)$$

Recall, that the classical and semi-classical relations are given in terms of infinitesimals, e. g. ∂x^m which should tend to zero. Whereas, the quantum relations are in terms of finite quantities, e. g. ΔA_m and Δx^m which are prevented to become zero *in the quantum cases* according to the uncertainty relations.

Now the most closest form of the quantum relation (4) to its differential or local definition: $\rho_H = \frac{\partial A_m}{ne \partial x_n} \epsilon_{mn}$ is achievable for the most minimal possible values of ΔA_m and Δx_n . Nevertheless, this can be achieved, in view of the dependency $A_m = \epsilon_{mn} B \cdot x_n$, also if only $\Delta x_n = \Delta x_m$ is replaced by its most minimal value which is equal to the magnetic length l_B . Because, for a given value of B , the $(\Delta x_n)_{(min)} = l_B$ is the smallest possible length which is available quantum theoretically for the cases under consideration and so it is the smallest variation for x_m under the given quantum conditions. Consistently, also for the ideal case of QHE one has the edge currents which flow within a distance from the edge of sample which is equal to the magnetic length, i. e. $(\Delta x_m)_{(min)} = l_B$ [7]. Moreover, one has for this same ideal case under quantum conditions (QHE) the uncertainty equation (2) where $|\Delta x_m|_{(min)} = |\epsilon_{mn} \Delta x_n|_{(min)} = l_B$. Using this in the relation (4) one obtains the quantized value of the Hall-resistivity:

$$\rho_H^Q = \frac{h}{\nu e^2} \quad , \quad (5)$$

where $\nu := 2\pi n l_B^2$ is as usual the filling factor.

Thus the most closest form of the quantum theoretically allowed definition of ρ_H^Q to its infinitesimal definition $\rho_H = \frac{\partial A_m}{ne \partial x_n} \epsilon_{mn}$, which is also appropriate for the QHE case, is given by the relation (5).

Therefore, the flux quantization on the QHE sample which is described by the uncertainty equation (2) can describe also the quantization of the Hall-resistivity, i. e.: $\rho_H = \frac{\partial A_m}{ne \partial x_n} \epsilon_{mn} \rightarrow \rho_H^Q = \frac{h}{\nu e^2}$ in the above discussed manner [8].

This approach is, despite of its finite character according to Δx^m and ΔA_m , also a topological approach, because on the one hand the area of sample under QHE-conditions is a multiple of the minimal quantum cell area: $2\pi l_B^2$. On the other hand, in the topological approach to QHE [9] the quantized Hall-measures

for the sample are given as the surface integral of the applied magnetic field, which is as the first Chern number on the sample manifold a topological invariant. Now for the applied constant magnetic field B it is obvious that this invariant is given by quantized flux $eB \cdot S = \tilde{N}h$ where S is the total area of the sample and equal to a multiple of the above mentioned quantum cell area. Thus, it is this multiplicity of the total area of sample under quantum conditions, i. e. $S = \tilde{N} \cdot 2\pi l_B^2$ which results in the quantization of the Hall-measures.

Recall that the same area is also involved in the $N = n \cdot S$ [8].

To see the equivalence with the topological approach in a direct way, let us perform the surface integral of the above definition $ne\rho_H = \frac{\partial A_m}{\partial x_n}\epsilon_{mn} := B$ over the area of sample: $\int \int ne\rho_H = \int \int B$. It results in view of the local constancy of ρ_H and for a constant B in the quantized Hall resistivity: $Ne^2\rho_H^Q = \tilde{N}h$ or $\rho_H = \frac{h}{\nu e^2}$ [8].

In summary: the flux quantization defines according to the relation (2) for a given B a quantum measure for the $2 - D$ area which is given by $2\pi l_B^2$, such that the relevant "quantum" topological invariants, e. g. the area, are multiples of it. This is in accordance with the usual quantum cell decomposition of a quantized phase space, which is represented by the uncertainty relation.

Now we show that the flux action functional $S = \oint eA_m dx^m = \int \int F_{mn} dx^m \wedge dx^n$ results in the true equation of motion for the electromagnetic potential, i. e. for the photon field with its true two degrees of freedom.

Herefore, recall first that the mentioned action functional has to be considered on the phase space of the system where its canonical conjugate variables are given by the set $\{A_m, x^m\}$. However, in view of the fact that A_m depends on x^n by $A_m = Bx^n \epsilon_{mn}$, the variation of action δS needs to be considered only with respect to the variation of δx^n , because the variation δA_m is proportional to δx_n . Nevertheless in view of the action preserving canonical transformations on the phase space, x^m as variables on the above phase space are in general functions of x^n , therefore one has to consider $dx^l = \frac{\partial x^l}{\partial x^m} dx^m$. To be precise, let us mention that the situation is the following: In the above phase space the variables $\{A_m, x^m\}$ has to be considered as functions of x^l , whereby the system is constrained by $A_m = Bx^n \epsilon_{mn}$.

Let us consider now the $S = \int \int F_{mn} dx^m \wedge dx^n$ and look for the Euler-Lagrange equations which results from a variation of this action with respect to the variation δx^n . The mentioned Euler-Lagrange equations

$\frac{\partial L}{\partial x^m} = \partial_n \frac{\partial L}{\partial \partial_n x^m}$ results in

$$\partial^n \partial_n A_m - \partial_m \partial^n A_n = 0 , \quad (6)$$

which is the usual vacuum equations of motion for A_m potential, here in two dimensions. If one introduces the Lorentz gauge $\partial^n A_n = 0$, then one obtains the Laplace equation $\partial_n \partial^n A_m = 0$ as the equations of motion of electrodynamics. Recall that the other part of equations of motion in electrodynamics $dF = d^2 A = 0$ are identities.

The same result is also achievable, if one uses the $S = \oint A_m dx^m$ form of the same action functional. Here, one should consider fairly $S = \frac{1}{2} \oint (A_m dx^m - x^m dA_m)$ [10]. Thereafter, one obtains from the above Euler-Lagrange equations, the equations $dA_m = \partial_m A_m$ which is fulfilled only if both sides are equal to zero, whereby $dA_m := \partial_n A_m dx^n \epsilon_{mn}$. Now these together, i. e. $d^\dagger A = \partial_m A_m = 0$ and $dA = dA_m dx^m = \partial_n A_m dx^n \wedge dx^m = 0$, give the same Laplace equation $(d^\dagger d + dd^\dagger)A = \partial_n \partial^n A_m = 0$ as above.

We show further that the necessary polarization in the phase space space of flux quantization, if it is considered for the QHE case, results in the vanishing of the longitudinal resistivity or conductivity [11].

The concept of polarization is an essential tool of the quantization theory [6]. It describes the well known circumstance that the state function, which fulfils the quantum equation of motion, is beyond its time dependency only a function of the half of the phase space variables. This is known usually as the representation of the wave function, so that it should be either in the position or in the momentum representation. Thus, the polarization in the phase space of the canonical system $\{p_m, q^m\}$ is described mathematically by the condition that the wave function Ψ should fulfil either $\frac{\partial \Psi}{\partial p_m} = 0$ or $\frac{\partial \Psi}{\partial q^m} = 0$ [6], [12].

As already discussed the phase space variables of the flux system are $\{A_m, x^m\}$ [11]. Thus the wave function of this system ψ should be, beyond its time dependency, only a function of A_m or of x^m , i. e. either $\psi(A_m, t)$ or $\psi(x^m, t)$. In other words, in the first case it should fulfil the polarization equation:

$$\frac{\partial \psi(A_m, t)}{\partial x^m} = 0 \quad (7)$$

However, in view of the fact that $\psi(A_m(x^n), t)$ does not depend explicitly on x^m , but its A_m -variables can be in general functions of x^m , the following polarization equations should be solved on the integration path, where the action functional is defined:

$$\frac{\partial\psi}{\partial A_m} \frac{\partial A_m}{\partial x^m} + \frac{\partial\psi}{\partial A_n} \frac{\partial A_n}{\partial x^m} \epsilon_{mn} = 0 \quad (8)$$

It is known that in the flux quantization the path of integration in the action functional $\oint A_m dx^m$ must be chosen in a region where the electronic current is absent. This means that the electromagnetic potential *on this path* is a pure gauge potential $\check{A}_m = \partial_m \Phi$ [3]. Therefore, *on* such an integration path the magnetic field strength should vanish $(F_{mn})_{(on)} = F_{mn}(\check{A}_m) = \frac{\partial \check{A}_n}{\partial x^m} \epsilon_{mn} = 0$. This can be understood also in the following way: That because the velocity operators $\hat{V}_m = (\hat{P}_m - e\hat{A}_m)$ which define the current must vanish on this path, hence the field strength defined by the commutator of current operators $(F_{mn})_{(on)} = [\hat{V}_m, \hat{V}_n]$ also should vanish.

Taking this into account the polarization equation (8) reduces to;

$$\frac{\partial\psi}{\partial \check{A}_m} \frac{\partial \check{A}_m}{\partial x^m} = 0 \quad (9)$$

From the semi-classical Ohm's equations $\partial_t \check{A}_m = \rho_L \cdot j_m + \rho_H \cdot j_n \epsilon_{mn}$ in the boundary region, it results that the longitudinal resistivity should be defined by the ratio $\rho_L := \frac{\partial_t \check{A}_m}{ne \partial_t x^m}$ or by $ne \rho_L := \frac{\partial \check{A}_m}{\partial x^m}$. Therefore, in view of the necessary dependency of ψ on \check{A}_m in this polarization direction, the polarization equation (9) is fulfilled only if the longitudinal resistivity vanishes:

$$ne \rho_L = \frac{\partial \check{A}_m}{\partial x^m} = 0 \quad (10)$$

Recall that, if one considers the other possible polarization direction $\frac{\partial\psi}{\partial \check{A}_m} = 0$ in the same phase space, then one arrives in the vanishing of the longitudinal conductivity $\sigma_L := \frac{ne \partial x^m}{\partial \check{A}_m}$. In view of the necessary independency of the quantization from the direction of polarization in a quantized case like QHE [6] [12], both of mentioned longitudinal measures σ_L and ρ_L should vanish.

The absolute vanishing of these longitudinal measures is an indication of an ideal QHE, hence the exact polarization describes an exact quantization; Whereas in real QHE samples, in view of impurities,

these longitudinal measures do not vanish exactly. In an ideal QHE case where the quantization is exact the electronic currents flows within the width of magnetic length on the edge of sample, thus the potential uncertainty in this ideal case should be given by $\Delta A_m = \frac{\hbar}{el_B}$. Whereas in real QHE cases the quantization or polarization is not exact and not only the longitudinal measures do not vanish exactly but also the potential uncertainty should be less than that in the ideal case (see also [3]).

It is also interesting to mention that the vanishing of longitudinal measures or the polarization of the flux/QHE phase space is equivalent to the Lorentz/Coulomb gauge fixing in such a $2 - D$ case.

Thus, the vanishing of longitudinal resistivity or conductivity in QHE is a result of the necessary quantization structure, i. e. of the polarization on the phase space of the system. In this way the vanishing of longitudinal measures in QHE can be understood within the canonical flux quantization of the QH-system.

Footnotes and references

References

- [1] See the contribution by R. B. Laughlin, in : A. Shapere and F. Wilczek (editors): Geometric Phases in Physics (World Scientific, Singapore, 1989); and referencees therein specially the reference by P. Anderson. See Further: V. J. Emery (editor), Correlated Electron Systems, (World Scientific, Singapore 1993).
- [2] F. Ghaboussi, " On the relation between Superconductivity and Quanrum Hall effect", quant-ph/9603013.
- [3] F. ghaboussi, "A new uncertainty relation", cond-mat/ 9611087 v2.
- [4] Landau-Lifschitz, "Quantum Mechanics (Non-Relativistic Theory)" Vol.III (Pergamon Press 1987); H. Aoki: Rep. Prog. Phys. 50, (1987), 655.
- [5] J. Bellissard, " Ordinary quatum Hall effect and non-commutative cohomology". proceedings of localization of disordered systems, Bad Schanau 1986, (Teubner Publication, Leipzig 1988).

If one inserts the commutator for flux quantization $e[\hat{A}_m, \hat{x}_m] = -i\hbar$ in the mentioned commutator $[\hat{x}_m, \hat{x}_n] = -il_B^2 \epsilon_{mn}$ and use $\nu := 2\pi n l_B^2$, then one obtains for $\sigma_H = \frac{ne}{B}$ its quantized values $\sigma_H = \nu \frac{e^2}{h}$ according to the Landau gauge. Therefore, there is a close relation between the non-commutative approach and our flux quantization approach to QHE.

[6] For the theory of geometric quantization and the polarization see: N. Woodhouse, "Geometric Quantization", (Clarendon Press, 1980, 1990) Oxford University.

[7] K. von Klitzing, Physica B 204 (1995) 111-116; R. Knott, W. Dietsche, K. von Klitzing, K. Eberl and K. Ploog, Semicond. Sci. Technol. 10 (1995) 117-126

[8] It is also possible to use the equivalent semi-classical definition of $\rho_H := \frac{B}{ne}$ and to consider $N = n \cdot S$ the total numbers of electrons on the sample. Then, one arrives from $\rho_H := \frac{B}{ne}$ by flux quantization relation $\int \int e F_{mn} dx^m \wedge dx^n = eB \cdot S = \tilde{N}h$ in the quantized relation for the Hall resistivity: $\rho_H^Q = \frac{\tilde{N}h}{Ne^2} = \frac{h}{\nu e^2}$; $N, \tilde{N} \in \mathbf{Z}$.

Now, there are two possibilities for $\nu := \frac{N}{\tilde{N}}$ either $\nu \in \mathbf{Z}$ again or not, in the first case one is in the IQHE situation, whereas in the second case, *for the right fractions*, one is in the FQHE case.

[9] D. J. Thouless, et al., Phys. Rev. Lett., 49, 405 (1982)

[10] In view of the fact that, according to the symplectic structure of the canonical phase space, the symplectic 2-form can be given by $\omega := d(p_m dq^m) = -d(q^m dp_m) = \frac{1}{2}d(p_m dq^m - q^m dp_m)$. Then also the symplectic (closed) 2-form F on the $\{A_m, x^m\}$ -phase space of flux quantization can be given by: $F_{mn} dx^m \wedge dx^n = dA_m \wedge dx^m = d(A_m dx^m) = -d(x^m dA_m) = \frac{1}{2}d(A_m dx^m - x^m dA_m)$.

Thus, the proper form of the closed line integral action should be given by the

$$\frac{1}{2} \oint (A_m dx^m - x^m dA_m).$$

[11] In the first part we used merely the action $\int \int F$. In the polarization part we use the equivalent action $\oint A$ which is predestinated for quantization procedure.

[12] Of course it is possible also to use polarizations which are linear complex combinations of the mentioned two ones. Nevertheless, the quantization should be independent of the polarization and the polarization should be compatible with the constraint of the theory (see Ref. [6]).