

Nonequilibrium phase transition in the kinetic Ising model: Divergences of fluctuations and responses near the transition point

Muktish A Charyya

Department of Physics

Indian Institute of Science, Bangalore-560012, India

and

Condensed Matter Theory Unit

Jawaharlal Nehru Centre for Advanced Scientific Research

Jakkur, Bangalore-560064, India

(January 11, 2022)

The nonequilibrium dynamic phase transition, in the kinetic Ising model in presence of an oscillating magnetic field, has been studied by Monte Carlo simulation. The fluctuation of dynamic order parameter has been studied as a function of temperature near the dynamic transition point. The temperature variation of appropriately defined 'susceptibility' has also been studied near the dynamic transition point. Similarly, fluctuation of energy and appropriately defined 'specific heat' have been studied as a function of temperature near the dynamic transition point. In both the cases, the fluctuations (of dynamic order parameter and energy) and the corresponding responses diverge (as power law fashion) near the dynamic transition point with similar critical behavior (with identical exponent values).

PACS number(s): 05.50.+q

I. INTRODUCTION

The physics of equilibrium phase transition in the Ising model is well understood [1]. However, the mechanism behind the nonequilibrium phase transition is not yet explored rigorously and the basic phenomenology is still undeveloped. It is quite interesting to study how the system behaves if it is driven out of equilibrium. The simplest prototype example is the kinetic Ising model in oscillating magnetic field. In this context, the dynamic response of the Ising system in presence of an oscillating magnetic field has been studied extensively [2-7] in the last few years. The dynamic hysteresis [2-4] and the nonequilibrium dynamic phase transition [5-7] are two important aspects of the dynamic response of the kinetic Ising model in presence of an oscillating magnetic field. The nonequilibrium dynamic phase transition in the kinetic Ising model in presence of an oscillating magnetic field, was first studied by Tome and Oliviera [5]. They solved the mean-field (MF) dynamic equation of motion (for the average magnetization) of the kinetic Ising model in presence of a sinusoidally oscillating magnetic field. By defining the dynamic order parameter as the time averaged magnetization over a full cycle of the oscillating magnetic field, they showed that depending

upon the value of the field amplitude and the temperature, the dynamic order parameter takes nonzero value from a zero value. In the field amplitude and temperature plane there exists a distinct phase boundary separating dynamic ordered (nonzero value of order parameter) and disordered (order parameter vanishes) phase. A tricritical point (TCP), (separating the nature (discontinuous-continuous) of the transition) on the phase boundary line, was also observed by them [5]. However, one may argue that such a mean-field transition is not truly dynamic in origin since it exists even in the quasi-static (or zero frequency) limit. This is because, if the field amplitude is less than the coercive field (at temperature less than the transition temperature without any field), then the response magnetization varies periodically but asymmetrically even in the zero frequency limit; the system remains locked to one well of the free energy and cannot go to the other one, in the absence of fluctuation.

The true dynamic nature of this kind of phase transition (in presence of fluctuation) was first attempted to study by Lo and Pelcovits [6]. They have studied the dynamic phase transition in the kinetic Ising model in presence of an oscillating magnetic field by Monte Carlo (MC) simulation which allows the microscopic fluctuations. Here, the transition disappears in the zero frequency limit; due to the fluctuations, the magnetization tips to the direction of the magnetic field and the dynamic order parameter (time averaged magnetization) vanishes. However, they [6] have not reported any precise phase boundary. A Charyya and Chakrabarti [7] studied the nonequilibrium dynamic phase transition in the kinetic Ising model in presence of oscillating magnetic field by extensive MC simulation. They [7] have also identified that this dynamic phase transition (at a particular nonzero frequency of the oscillating magnetic field) is associated with the breaking of the symmetry of the dynamic hysteresis (m-h) loop. In the dynamically disordered phase (where the value of order parameter vanishes) the corresponding hysteresis loop is symmetric, and loses its symmetry in the ordered phase (giving nonzero value of dynamic order parameter). They [7] also studied the temperature variation of the ac susceptibility components near the dynamic transition point. Their major observation was that the imaginary (real) part

of the ac susceptibility gives a peak (dip) near the dynamic transition point (where the dynamic order parameter vanishes). The important conclusions were: (i) this is a distinct signal of phase transition and (ii) this is an indication of the thermodynamic nature of the phase transition. The Debye relaxation of the dynamic order parameter and the critical slowing down have been studied very recently [11] both by MC simulation and by solving the dynamic MF equation [5] of motion for the average magnetization. The specific-heat singularity [11] near the dynamic transition point is also an indication of the thermodynamic nature of this dynamic phase transition. It is worth mentioning here that the statistical distribution of dynamic order parameter has been studied by Sides et al [8]. The nature of the distribution changes (from bimodal to unimodal) near the dynamic transition point. They have also observed [8] that the fluctuation of the hysteresis loop area becomes considerably large near the dynamic transition point.

In the case of equilibrium phase transitions, the fluctuation-dissipation theorem (FDT) states (due to the applicability of Gibbs formalism) that the mean square fluctuations of some intrinsic physical quantities (say, energy, magnetization etc.) are directly related with some responses (specific heat, susceptibility etc.) of the system. Consequently, near the ferro-param transition point, both the fluctuation of magnetization and the susceptibility show same singular behavior. If it is of power law type, the same singular behavior will be characterised by the same exponent. This is also true for fluctuation of energy and the specific heat. These are the consequences of fluctuation-dissipation theorem [1]. Here, the main motivation is to study the fluctuations and corresponding responses near the dynamic transition temperature.

In this paper, the fluctuations of dynamic order parameter and the energy are studied as a function of temperature near the dynamic transition point. The temperature variations of 'susceptibility' and the 'specific-heat' are also studied near the transition point. The temperature variation of the fluctuation of dynamic order parameter and that of the 'susceptibility' are compared. Similarly, the temperature variation of the fluctuation of energy and that of the 'specific-heat' are compared. The paper is organised as follows: the model and the simulation scheme are discussed in section II, the results are reported in Section III, section IV contains the summary of the work.

II. MODEL AND SIMULATION

The Ising model with nearest neighbor ferromagnetic coupling in presence of a time varying magnetic field can be represented by the Hamiltonian

$$H = \sum_{\langle ij \rangle} J_{ij} s_i^z s_j^z - h(t) \sum_i s_i^z \quad (2.1)$$

Here, $s_i^z (= \pm 1)$ is Ising spin variable, J_{ij} is the interaction strength and $h(t) = h_0 \cos(\omega t)$ represents the oscillating magnetic field, where h_0 and ω are the amplitude and the frequency respectively of the oscillating field. The system is in contact with an isothermal heat bath at temperature T . For simplicity all $J_{ij} (= J > 0)$ are taken equal to unity and the boundary condition is chosen to be periodic. The temperature (T) is measured in the unit of J/k , where k is the Boltzmann constant (here k is taken unity).

A square lattice of linear size $L (= 100)$ has been considered. At any finite temperature T and for a fixed frequency (ω) and amplitude (h_0) of the field, the microscopic dynamics of this system has been studied here by Monte Carlo simulation using Glauber single spin-flip dynamics with a particular choice of the Metropolis rate of single spin-flip [12]. Starting from an initial condition where all spins are up, each lattice site is updated here sequentially and one such full scan over the entire lattice is defined as the unit time step (Monte Carlo step or MCS). The instantaneous magnetization (per site), $m(t) = (1/L^2) \sum_i s_i^z$ has been calculated. From the instantaneous magnetization, the dynamic order parameter $Q = \frac{1}{2} \int_0^{2\pi} m(t) dt$ (time averaged magnetization over a full cycle of the oscillating field) is calculated. Some of the transient loops have been discarded to get the stable value of the dynamical quantities.

III. RESULTS

A. Temperature variations of susceptibility and fluctuation of dynamic order parameter

The fluctuation of the dynamic order parameter is

$$Q^2 = \langle Q^2 \rangle - \langle Q \rangle^2 ;$$

where the $\langle \rangle$ stands for the averaging over various Monte Carlo samples.

The 'susceptibility' is defined as

$$\chi = \frac{d \langle Q \rangle}{dh_0} ;$$

Here, a square lattice of linear size $L (= 100)$ has been considered. $\langle Q^2 \rangle$ and $\langle Q \rangle$ are calculated using MC simulation. The averaging has been done over 100 different (uncorrelated) MC samples.

The temperature variations of fluctuation of Q , i.e., Q^2 and 'susceptibility' have been studied here and both plotted in Fig. 1. From the figure it is observed that both Q^2 and χ diverge near the dynamic transition point (where Q vanishes).

This has been studied for two different values of field amplitude h_0 (Fig. 1a is for $h_0 = 0.2$ and Fig. 1b is for $h_0 = 0.1$). The dynamic transition temperatures $T_d(h_0)$, at which Q and Q^2 diverge, are 1.91 \pm 0.01 for $h_0 = 0.2$

and 2.15 ± 0.01 for $h_0 = 0.1$. These values of $T_d(h_0)$ agree with the phase diagram estimated from vanishing of Q . The $\log_e(\cdot)$ versus $\log_e(T_d - T)$ and $\log_e(Q^2)$ versus $\log_e(T_d - T)$ plots show (insets of Fig. 1) that $\log_e(Q^2)$ and $\log_e(T_d - T)$ are linear. For $h_0 = 0.2$, $\log_e(Q^2)$ versus $\log_e(T_d - T)$ is linear with slope 2.5 (inset of Fig. 1a) and for $h_0 = 0.1$, $\log_e(Q^2)$ versus $\log_e(T_d - T)$ is linear with slope 2.5 (inset of Fig. 1b). Results show that both Q and Q^2 diverge near T_d as a power law with the same exponent, though there is a crossover region (where the effective exponent values are different).

B. Temperature variations of specific-heat and fluctuation of energy

The time averaged (over a full cycle) cooperative energy of the system is

$$E = \frac{1}{L^2} \sum_{i,j} \langle s_i^z s_j^z \rangle = \frac{1}{L^2} \sum_{i,j} \langle s_i^z s_j^z \rangle_A$$

and the fluctuation of the cooperative energy is

$$E^2 = \langle E^2 \rangle - \langle E \rangle^2$$

The 'specific-heat' C [11] is defined as the derivative of the energy (defined above) with respect to the temperature, and

$$C = \frac{d\langle E \rangle}{dT}$$

Here, also a square lattice of linear size L ($= 100$) has been considered. $\langle E^2 \rangle$ and $\langle E \rangle$ are calculated using MC simulation. The averaging has been done over 100 different (uncorrelated) MC samples.

The temperature variation of the 'specific-heat' has been studied [11] and found prominent divergent behavior near the dynamic transition point (where $\langle Q \rangle$ vanishes). The temperature variation of fluctuation of energy, $\langle E^2 \rangle$ has been studied and plotted in Fig. 2. From the figure it is clear that the mean square fluctuation of energy ($\langle E^2 \rangle$) and the specific heat (C) both diverge near the dynamic transition point (where the dynamic order parameter Q vanishes).

This has been studied for two different values of field amplitude h_0 (Fig. 2a is for $h_0 = 0.2$ and Fig. 2b is for $h_0 = 0.1$). Here also (like the earlier case) the specific heat C and $\langle E^2 \rangle$ are observed to diverge at temperatures T_d . The temperatures $T_d(h_0)$, at which C and $\langle E^2 \rangle$ diverge, are 1.91 ± 0.01 for $h_0 = 0.2$ (Fig. 2a) and 2.15 ± 0.01 for $h_0 = 0.1$ (Fig. 2b). These values also agree with the phase diagram estimated from vanishing of Q . The $\log_e(C)$ vs. $\log_e(T_d - T)$ and $\log_e(\langle E^2 \rangle)$ vs. $\log_e(T_d - T)$ plots show (insets of Fig. 2) that C versus $\log_e(T_d - T)$ and $\langle E^2 \rangle$ versus $\log_e(T_d - T)$ are linear. For $h_0 = 0.2$, $\log_e(C)$ versus $\log_e(T_d - T)$ is linear with slope 0.35 (inset of Fig. 2a) and for $h_0 = 0.1$, $\log_e(C)$ versus $\log_e(T_d - T)$ is linear with slope 0.43 (inset of Fig. 2b). Like the earlier

case, here also the results show that both C and $\langle E^2 \rangle$ diverge near T_d as a power law with the same exponent, though there is a crossover region (where the effective exponent values are different).

IV. SUMMARY

The nonequilibrium dynamic phase transition, in the kinetic Ising model in presence of an oscillating magnetic field, is studied by Monte Carlo simulation. Acharrya and Chakrabarti [7] observed that the complex susceptibility components have peaks (or dips) at the dynamic transition point. Sides et al [8] observed that the fluctuation in the hysteresis loop area grows (seems to diverge) near the dynamic transition point. It has been observed [11] that the 'relaxation time' and the appropriately defined 'specific-heat' diverge near the dynamic transition point.

The mean square fluctuation of dynamic order parameter and the 'susceptibility' are studied as a function of temperature, near the dynamic transition point. Both shows the power law variation with respect to the reduced temperature near the dynamic transition point with the same exponent values. Similar, observation has been made for the case of mean square fluctuation of energy and the 'specific-heat'. It appears that although the effective exponent values for the fluctuation and the appropriate linear response differ considerably, away from the dynamic transition point T_d , they eventually converge and give identical value as the temperature interval $|T_d - T|$ decreases and falls within a narrow crossover region. These numerical observations indicate that the fluctuation-dissipation relation [1] holds good in this case of the nonequilibrium phase transition in the kinetic Ising model. However, at this stage there is no analytic support of the FDT in this case.

Finally, it should be mentioned, in this context, that experiments [9] on ultrathin ferromagnetic Fe/Au(001)

ins have been performed to study the frequency dependence of hysteresis loop areas. Recently, attempts have been made [10] to measure the dynamic order parameter Q experimentally, in the same material, by extending their previous study [9]. The dynamic phase transition has been studied from the observed temperature variation of Q . However, the detailed investigation of the dynamic phase transitions by measuring variations of associated response functions (like the ac susceptibility, specific-heat, correlations, relaxations etc) have not yet been performed experimentally.

ACKNOWLEDGMENTS

The author would like to thank B. K. Chakrabarti for critical remarks and for a careful reading of the

manuscript. The JNCASR, Bangalore, is gratefully acknowledged for financial support and for computational facilities.

(triangles). All the data points are plotted in arbitrary units. Here, $\Delta = 2 \pm 0.01$. Corresponding insets show the plots of $\log_e(C)$ (circles) and $\log_e(E^2)$ (triangles) against $\log_e(T_d - T)$. Solid lines represent the linear best-fit in the region very closed to T_d .

Electronic address: muktish@physics.iisc.ernet.in

- [1] H. E. Stanley, Introduction to the Phase Transition and Critical Phenomena, (1971), Oxford University Press, Oxford
- [2] M. Rao, H. R. Krishnamurthy and R. Pandit, Phys. Rev. B 42, 856 (1990).
- [3] P. B. Thomas and D. Dhar, J. Phys. A: Math Gen 26, 3973 (1993).
- [4] S. Sengupta, Y. Marathe and S. Puri, Phys. Rev. B 45, 4251 (1990).
- [5] T. Tome and M. J. de Oliveira, Phys. Rev. A 41, 4251 (1990).
- [6] W. S. Lo and R. A. Pelcovits, Phys. Rev. A 42, 7471 (1990).
- [7] M. Acharya and B. K. Chakrabarti, Phys. Rev. B 52, 6550 (1995).
- [8] S. W. Sides, R. A. Ramos, P. A. Rikvold and M. A. Novotny, J. Appl. Phys. (1996) (To be published); see also J. Appl. Phys. 79, 6482 (1996).
- [9] Y. L. He and G. C. Wang, Phys. Rev. Lett. 15, 2336 (1993).
- [10] G. C. Wang et al, Preprint, (1996).
- [11] M. Acharya, Phys. Rev. E (1997), (To be published); Physica A 235, 469 (1997).
- [12] K. Binder and D. W. Heermann, Monte Carlo Simulation in Statistical Physics, Springer-Verlag, Berlin, (1988)

Figure Captions

Fig.1 Temperature variations of $\langle Q \rangle$, C and Q^2 for two different values of field amplitudes h_0 : (a) For $h_0 = 0.2$; $\langle Q \rangle$ (solid line), C (circles) and Q^2 (triangles). (b) For $h_0 = 0.1$; $\langle Q \rangle$ (solid line), C (circles) and Q^2 (triangles). All the data points are plotted in arbitrary units. Here, $\Delta = 2 \pm 0.01$. Corresponding insets show the plots of $\log_e(C)$ (circles) and $\log_e(Q^2)$ (triangles) against $\log_e(T_d - T)$. Solid lines represent the linear best-fit in the region very closed to T_d .

Fig.2 Temperature variations of $\langle Q \rangle$, C and E^2 for two different values of field amplitudes h_0 : (a) For $h_0 = 0.2$; $\langle Q \rangle$ (solid line), C (circles) and E^2 (triangles). (b) For $h_0 = 0.1$; $\langle Q \rangle$ (solid line), C (circles) and E^2

Fig.1, EAJ634, Acharyya




Fig. 1, EAJ634, Acharyya




Fig. 2, EAJ634, Acharyya




Fig. 2, EAJ634, Acharyya

