

Mean field and corrections for the Euclidean Minimum Matching problem

Jacques H. Boutet de Monvel and Olivier C. Martin^y

Division de Physique Theorique, Institut de Physique Nucleaire, Universite Paris-Sud, F-91406 Orsay Cedex, France
(Submitted to Physical Review Letters)

Consider the length L_{MM}^E of the minimum matching of N points in d -dimensional Euclidean space. Using numerical simulations and the finite size scaling law $L_{MM}^E = L_{MM}^E(d) N^{1/(d+1)}$, we obtain precise estimates of $L_{MM}^E(d)$ for $2 \leq d \leq 10$. We then consider the approximation where distance correlations are neglected. This model is solvable [6] and gives at $d = 2$ an excellent "random link" approximation to $L_{MM}^E(d)$. Incorporation of three-link correlations further improves the accuracy, leading to a relative error of 0.4% at $d = 2$ and 3. Finally, the large d behavior of this expansion in link correlations is discussed.

PACS numbers: 75.50.Lk, 64.60.Cn

There has been a tremendous amount of work on mean field calculations for disordered systems in the past 20 years, in part driven by the exact solution provided by Parisi's replica symmetry breaking Ansatz. Although this solution was developed in the context of spin glasses, the formalism has been extremely useful for understanding other disordered systems. Generally, one expects mean field to give exact results as the dimensionality goes to infinity. One can then ask whether mean field leads to "acceptable" errors for systems of interest, e.g., in three dimensions, and whether it is possible to compute Euclidean corrections to the mean field formulae. Such computations might correspond to a $1/d$ expansion for the thermodynamic functions of interest. In frustrated disordered systems, however, this has turned out to be intractable. To date, such Euclidean corrections have been pushed furthest [8] for the Minimum Matching Problem (MMP) [10]. In what follows, we determine the accuracy of the mean field approximation and the effectiveness of the corrections thereto in the MMP by comparing with the actual properties of the d -dimensional Euclidean model. First, we find that the relative error introduced by the mean field approximation for the zero temperature energy density is less than 4% at $d = 2$ and 3% at $d = 3$. Second, the inclusion of the "leading" Euclidean corrections to the mean field approximation reduces the error by a factor of about 10 at $d = 2$ and $d = 3$. Third, we argue that the large d behavior of systems such as the MMP depends on arbitrarily high order correlations and is thus beyond all orders of the expansion proposed by Mezard and Parisi [8].

Consider N points (N even) and a specified set of link lengths $l_{ij} = l_{ji}$ separating the points, for $1 \leq i, j \leq N$. One defines a matching (a dimerization) of these points by combining them pairwise so that each point belongs to one and only one pair. Define also the energy or length of a matching as the sum of the lengths of the links associated with each matched pair. The Minimum Matching Problem is the problem of finding the matching of minimum energy. One can also consider the thermodynamics of this system, as proposed by Orland [9] and Mezard and Parisi [5], by taking all matchings but weighting

them with the Boltzmann factor associated with their energy. Here we concentrate on the $T = 0$ properties because there is no effective numerical method for extracting thermodynamic functions in this system, but exact energy minima can be obtained quite easily for any given instance. Indeed, the MMP belongs to the algorithmic class P of polynomial problems, and there are standard algorithms which solve any instance of size N using on the order of N^3 steps [1].

Physically, one is not interested in the properties of any particular instance of the MMP; more relevant are typical and ensemble properties such as the average energy when the lengths l_{ij} are random variables with a given distribution. One then speaks of the stochastic MMP. There are two frequently used ensembles for the l_{ij} , corresponding to the Euclidean MMP and the random link MMP. In the first, the N random points lie in a d -dimensional Euclidean volume and the l_{ij} are the usual Euclidean distances between pairs of points. The points are independent and identically distributed, so one speaks of a random point problem. In the second ensemble, it is the link lengths l_{ij} which are independent and identically distributed random variables. A connection between these two systems was first given by Mezard and Parisi [6]: they pointed out that the one and two-link distributions could be made identical in the two problems. A consequence is that the "Cayley tree" approximation for the random point and random link problems are the same. Mezard and Parisi were able to solve the random link MMP using an approach based on replicas [5,6]. One may then consider the random link MMP to be a "mean field model" for the Euclidean MMP. The mean field approximation consists of using the thermodynamic functions of the random link model as estimators for those of the Euclidean model. This approximation is applicable to all link based combinatorial optimization problems, such as the assignment problem and the traveling salesman problem; hereafter we shall refer to it as the random link approximation [4]. Finally, Mezard and Parisi have shown how to derive corrections systematically to the random link approximation using a connected-correlation link expansion. In [8], they have computed the leading corrections, these be-

ing associated with the triangle inequality (3-link correlations) in the Euclidean model. How accurate are these approximations? To answer this, we first give our results for the Euclidean problem, and then compare with the predictions of the random link approximation and of the link expansion method.

In the Euclidean M M P, let $L_{M M}^E$ be the energy or length of the minimum matching. Taking the points to be independent and uniformly distributed in a unit volume, Steele [12] has shown that as $N \rightarrow \infty$, $L_{M M}^E = N^{1/d}$ converges with probability one to a non-random, N -independent constant $\bar{L}_{M M}^E(d)$. In physics language, this result shows that $L_{M M}^E$ is self-averaging and that the zero-temperature energy density has an infinite volume limit when the density of points is kept fixed. To date, little has been done to compute $\bar{L}_{M M}^E(d)$. The best estimates seem to be due to Smith [11]: $\bar{L}_{M M}^E(2) = 0.312$ and $\bar{L}_{M M}^E(3) = 0.318$. Let us use a systematic procedure [4] to obtain $\bar{L}_{M M}^E(d)$ with quantifiable errors. First, in order to have a well defined dependence on N , we have used the ensemble average, $\overline{L_{M M}^E} = N^{1/d}$. Second, in order to reduce corrections to scaling in the extrapolation to the large N limit, we have placed the points randomly in the d -dimensional unit hypercube with periodic boundary conditions. This removes surface effects and empirically leads to the finite size scaling law

$$\frac{\overline{L_{M M}^E} N^{1/d}}{N^{1/d}} = \bar{L}_{M M}^E(d) \left(1 + \frac{A(d)}{N} + \frac{B(d)}{N^2} + \dots\right); \quad (1)$$

Finally, in order to reduce statistical fluctuations, we have used a variance reduction trick [4]. The improved estimator has the effect of reducing the variance of our estimates by more than a factor 4, and thus saves us a considerable amount of computer time. Using these methods, we have extracted from our numerical data $\bar{L}_{M M}^E(d)$ and its associated statistical error. The fits to Eq.(1) are good, with χ^2 values confirming the form of the finite size scaling law. The error bars on the extrapolated value $\bar{L}_{M M}^E(d)$ are obtained in the standard way by requiring that χ^2 increase by one from its minimum. We find in particular $\bar{L}_{M M}^E(2) = 0.3104 \pm 0.0002$, and $\bar{L}_{M M}^E(3) = 0.3172 \pm 0.00015$; values at higher dimensions are given in Table I. We have checked that these results are not significantly modified when using another random number generator to produce the instances, and that the fits are stable to truncation of the data.

Now we discuss how to use the random link model to approximate $\bar{L}_{M M}^E(d)$. For any two points $(i; j)$ placed at random in the unit d -dimensional hypercube, the density distribution of l_{ij} is given at short distances by $P_d(l_{ij} = r) = dB_d r^{d-1}$, where $B_d = \frac{1}{d!} = (d-1)!$ is the volume of the d -dimensional ball with unit radius. If we take the random link model where link lengths are independent and have the individual distribution $P_d(l)$, then the Euclidean and random link M M P have the same one and two-link distributions [6] because two Euclidean distances are independent. If correlations among three or

more link lengths are weak, then the properties of the two systems should be quantitatively close. Thus an analytic approximation to $\bar{L}_{M M}^E(d)$ is obtained by computing its analogue $\bar{L}_{M M}^{RL}(d)$ in the random link M M P. In references [5,6], Mezard and Parisi solved these random link models under the replica symmetry hypothesis. They showed further [7] that the replica symmetric solution is stable (at least for $d \geq 1$), and thus is most likely exact unless a first order phase transition occurs in this system. Their solution gives $\bar{L}_{M M}^{RL}(d)$ in terms of a function G_d related to the probability distribution of link lengths for matched pairs. In our Euclidean units their result can be written

$$\bar{L}_{M M}^{RL}(d) = \frac{D_1(d)}{2} \frac{d}{(1+d)!} \int_0^{Z+1} G_d(x) e^{-G_d(x)} dx \quad (2)$$

where G_d satisfies the integralequation

$$G_d(x) = d \int_x^{Z+1} (x+y)^{d-1} e^{-G_d(y)} dy \quad (3)$$

and where

$$D_1(d) = \lim_{N \rightarrow \infty} \overline{L_{11}^E} N^{1/d} = (1+d) B_d^{1/d} \quad (4)$$

is the average (rescaled) link length of the nearest neighbor graph in the limit $N \rightarrow \infty$.

Brunetti et al. [3] have used direct numerical simulations of these random link models to confirm the predictions to the level of 0.2% at $d = 1$ and 2, and we have done the same to the level of 0.1% at $1 \leq d \leq 10$, giving further evidence that the replica symmetric solution is exact. From the analytical side, solving the integralequation for G_d leads to $\bar{L}_{M M}^{RL}(1) = \frac{1}{2} = 0.5$, $\bar{L}_{M M}^{RL}(2) = 0.322580\dots$, and $\bar{L}_{M M}^{RL}(3) = 0.326839\dots$; values at higher dimensions are given in Table I. If we consider $\bar{L}_{M M}^{RL}(d)$ as a mean field prediction for $\bar{L}_{M M}^E(d)$, the accuracy is surprisingly good. Including the trivial value $\bar{L}_{M M}^E(1) = 0.5$, we see that the random link approximation leads to a relative error of 17.8% at $d = 1$, of 3.9% at $d = 2$, and of 3.0% at $d = 3$. Also, the error decreases with increasing dimension. It can be argued, for the M M P as well as for other link-based combinatorial optimization problems [4], that the random link approximation not only has a relative error tending towards 0 as $d \rightarrow \infty$, but that in fact this error is at most of order $1/d^2$. Given our high quality estimates, we are able to confirm this property numerically. In Figure 1 we plot the quantity $d(\bar{L}_{M M}^{RL} - \bar{L}_{M M}^E) = \bar{L}_{M M}^E$ along with a quadratic fit given to guide the eye. As expected, the data scales as $1/d$. Thus the random link approximation gives both the leading and $1/d$ subleading dependence of $\bar{L}_{M M}^E(d)$. In order to obtain analytic expressions for these coefficients, we have derived the $1/d$ expansion for $\bar{L}_{M M}^{RL}$ from Eqs.(2,3). We used two methods to do this. The first, straightforward but computationally lengthy, consists of setting $G_d(x) = G_d(x = x/d + 1/2)$ and then writing $G_d(x)$ as a power series in $1/d$. From this we find

$$\frac{R^L}{M^M}(d) = \frac{D_1(d)}{2} \left(1 + \frac{1}{d} + O(1/d^2)\right) \quad (5)$$

where $\gamma = 0.577 \dots$ is Euler's constant. If, as claimed, the random link approximation gives an error of order $1/d^2$, Eq.(5) gives an analytic expression for the leading and first subleading terms in the $1/d$ expansion of $\frac{E}{M^M}(d)$. This claim is strongly supported by the numerical results: performing a fit of four $\frac{E}{M^M}(d)$ values to a truncated $1/d$ series leads to 0.424 ± 0.008 for the coefficient of the $1/d$ term; this is to be compared to the theoretical prediction of $1 - \gamma = 0.422784 \dots$

We have been able to obtain the next coefficient of the series in $1/d$ for $\frac{R^L}{M^M}$ by using a second method. We introduce a modified random link model where the links are shifted and rescaled in such a way that the leading term of the $1/d$ expansion for this new model is exactly the $1/d$ coefficient for the initial one [2]. In fact it is possible to introduce a sequence of such "rescaled" models, where the k^{th} model is designed to produce the $1/d^k$ term of the expansion. We have computed the leading terms predicted by a replica symmetric analysis of these models for $k = 1$ and 2 , from which we find that the order $1/d^2$ coefficient in Eq.(5) is $\gamma^2 = 12 + \gamma^2 = 2$.

We now come to the main point of the paper: how well can one predict $\frac{E}{M^M}(d)$ by incorporating Euclidean corrections to the random link approximation? It is necessary here to review the work of Mezard and Parisi; for greater detail, we refer the reader to their article [8]. They begin with the partition function Z for an arbitrary stochastic MPM and write the quenched average for n replicas. In the Euclidean model, the l_{ij} have three and higher-link correlations. Mezard and Parisi keep the three-link correlations (arising only when the three links make a triangle) and neglect higher connected correlations. Note that it is not clear a priori whether these "higher order" terms are negligible compared to the three-link term. The resulting expression for the quenched average becomes

$$\overline{Z^n} = \frac{1}{P} \int \prod_{i,j} d\mathbf{u}_{ij} \frac{Z^n}{P^n} \frac{d}{2} e^{i \int \mathbf{u}_{ij} \cdot \mathbf{u}_{kl} \mathbf{u}_{mn} \dots} \quad (6)$$

where u_{ij} is a complicated nonlinear function of the link length l_{ij} . They then compute the limit $N \rightarrow \infty$, $n \rightarrow 0$ using the saddle point method while assuming that replica symmetry is not broken. In the zero temperature limit, just as in the standard random link model, the saddle point equations can be expressed in terms of G_d , but now G_d satisfies a more complicated integroequation (Eq.(34) in their paper). From this, one can calculate new estimates for $\frac{E}{M^M}(d)$, which we shall denote $\frac{E^C}{M^M}$, where EC stands for Euclidean corrections.

We have solved the equations numerically for this modified G_d , and have computed $\frac{E^C}{M^M}(d)$ for $2 \leq d \leq 10$. We find $\frac{E^C}{M^M}(2) = 0.30915$ and $\frac{E^C}{M^M}(3) = 0.31826$. The results for $d \leq 4$ are given in Table I. Comparing with

$\frac{E}{M^M}(d)$ and $\frac{R^L}{M^M}(d)$, we see that the new estimates are considerably more accurate. At $d = 2$, the random link approximation leads to an error of 3.9%; this error is decreased by nearly a factor 10 by incorporating these leading Euclidean corrections. Similarly at $d = 3$, the error is reduced from 3.0% to less than 0.4%. At larger d , the error continues to decrease, but the effect is less dramatic.

To interpret this last result, consider how the difference $\frac{E^C}{M^M} - \frac{R^L}{M^M}$ scales with d . Using Eq.(6), we see that it is sufficient to estimate the size of the 3-link correction term. Its dimensional dependence follows that of the probability of finding nearly equilateral triangles as d^{-1} . Since this probability goes to zero exponentially with d , the 3-link correlations give tiny corrections at large d (as confirmed by the numerics), and also the power series expansion in $1/d$ of $\frac{E^C}{M^M}$ is identical to that of $\frac{R^L}{M^M}$. This property continues to hold if one includes 4, 5, or any finite number of multi-link correlations in Eq.(6). This is due to the fact that the Euclidean and random link graphs have local properties that are identical up to exponentially small terms in d . In particular, the statistics of fixed sized (N -independent) loops connecting near neighbors are nearly identical.

Although this reasoning was given for the MPM, it applies equally well to other link-based problems. In such statistical mechanics systems, if the thermodynamic functions depend only on the local properties of the (short) link graph, then the random link approximation applied to the Euclidean system will have an error which is exponentially small in d . However, for combinatorial optimization problems such as the MPM, the assignment problem, and the traveling salesman problem, the $N \rightarrow \infty$ limit and the k -link expansion do not commute: k -link correlations with k growing with N remain important as $N \rightarrow \infty$. In particular, arbitrarily large loops matter and contribute to the thermodynamics at order $1/d^2$. In a polymer picture, we can say that the random link approximation is exponentially good in the dilute phase, while it leads to $1/d^2$ errors in the collapsed phase. The power corrections in this phase are beyond all orders in a k -link correlation expansion such as Eq.(6).

In summary, we have estimated by numerical simulation $\frac{E}{M^M}(d)$, the zero energy density in the Euclidean Minimum Matching problem at dimensions $2 \leq d \leq 10$. We have then computed two analytical estimates for these energy densities, namely $\frac{R^L}{M^M}(d)$ and $\frac{E^C}{M^M}(d)$. The first method uses the random link approximation where all link correlations are neglected. Using the "exact" mean field solution of Mezard and Parisi, we find that even at low dimensions, the error introduced by this approximation is small: 3.9% at $d = 2$, 3.0% at $d = 3$, and 2.0% at $d = 4$. In the second method, the connected three-link correlations are taken into account while higher ones are neglected. Using Mezard and Parisi's expressions, we find that this modification to the random link model gives excellent predictions at

$d = 2$ and 3 , with the error there being divided by almost 10 compared to the random link approximation. This provides a stringent quantitative test of a systematic expansion which goes beyond uncorrelated disorder variables, and suggests that even the leading such correction is enough to get predictions for thermodynamic functions precise to better than one percent. Finally, at high dimensions, we have seen that the k -link correlation expansion leads to corrections which vanish exponentially with d ; this expansion thus misses important $1=d$ power law corrections for problems such as the M M P. This leaves open the determination of the $1=d^2$ term in the expansion of the constants $E_{MM}^E(d)$. We have performed a fit on our data, imposing the leading and the $1=d$ term to be those of the random link model. We find the $1=d^2$ coefficient to be very small (smaller than 0.01 in absolute value). Clearly, it would be of major interest to obtain an analytical value for this term.

We are grateful to C. De Dominicis, J. Houdayer, W. Krauth, M. Mezard, and H. Orland for their interest and suggestions. JBdM acknowledges a fellowship from the M E N E S R, and O C M acknowledges support from the Institut Universitaire de France. The Division de Physique Theorique is an Unité de Recherche des Universités Paris XI et Paris VI associée au CNRS.

* Electronic address: boutet@ipno.in2p3.fr
 y Electronic address: martin@ipno.in2p3.fr




FIG. 1. Linear scaling with $1=d$ of the quantity $d(RL E) = E$.

- [1] M. O. Ball and U. Derigs. An analysis of alternate strategies for implementing matching algorithms. *Networks*, 13:517{549, 1983.
- [2] J. Boutet de Monvel. Physique statistique et modesles a liens aleatoires. PhD thesis, Institut de Physique Nucleaire, Universite Paris-Sud, Orsay, 1996.
- [3] R. Brunetti, W. Krauth, M. Mezard, and G. Parisi. Extensive numerical solutions of weighted matchings: Total length and distribution of links in the optimal solution. *"Europhys. Lett."*, 14(4):295{301, 1991.
- [4] N. J. Cerf, J. Boutet de Monvel, O. Bohigas, O. C. Martin, and A. G. Percus. The random link approximation for the Euclidean traveling salesman problem. *J. Phys. I France*, 7(1):117{136, 1997.
- [5] M. Mezard and G. Parisi. Replicas and optimization. *"J. Phys. Lett. France"*, 46:L771{L778, 1985.
- [6] M. Mezard and G. Parisi. Mean-field equations for the matching and the travelling salesman problems. *"Europhys. Lett."*, 2:913{918, 1986.
- [7] M. Mezard and G. Parisi. On the solution of the random link matching problems. *"J. Phys. France"*, 48:1451{1459, 1987.
- [8] M. Mezard and G. Parisi. The Euclidean matching problem. *"J. Phys. France"*, 49:2019{2025, 1988.
- [9] H. Orland. Mean-field theory for optimization problems. *"J. Phys. Lett. France"*, 46:L763{L770, 1985.

- [10] C. H. Papadimitriou and K. Steiglitz. *Combinatorial Optimization: Algorithms and Complexity*. Prentice Hall, Englewood Cliffs, NJ, 1982.
- [11] W. D. Smith. *Studies in Computational Geometry Motivated by Mesh Generation*. PhD thesis, Princeton University, Princeton, NJ, 1989.
- [12] M. Steele. Subadditive Euclidean functionals and nonlinear growth in geometric probability. *The Annals of Probability*, 9(3):365{376, 1981.

TABLE I. Comparison of the M M P constants for the three models: Euclidean, random link, and random link including 3-link Euclidean corrections ($d = 4$ to 10).

d	$E(d)$	$R L(d)$	$d \frac{R L E}{E}(d)$	$E^C(d)$	$d \frac{E^C E}{E}(d)$
4	0.3365	0.0003	0.343227	+ 0.080	0.33756 + 0.30%
5	0.3572	0.00015	0.362175	+ 0.070	0.35818 + 0.27%
6	0.3777	0.0001	0.381417	+ 0.059	0.37849 + 0.21%
7	0.3972	0.0001	0.400277	+ 0.054	0.39807 + 0.22%
8	0.4162	0.0001	0.418548	+ 0.045	0.41685 + 0.17%
9	0.4341	0.0001	0.436185	+ 0.042	0.43485 + 0.17%
10	0.4515	0.0001	0.453200	+ 0.037	0.45214 + 0.14%