

Expressing Products of Fermi Fields in terms of Fermi Sea Displacements

Girish S. Setlur

January 15, 2022

Abstract

An attempt is made to generalise the ideas introduced by Haldane and others regarding Bosonizing the Fermi surface. The present attempt involves introduction of Bose fields that correspond to displacements of the Fermi sea rather than just the Fermi surface. This enables the study of short wavelength fluctuations of the Fermi surface and hence the dispersion of single particle excitations with high energy. The number conserving product of two Fermi fields is represented as a simple combination of these Bose fields. It is shown that most (!) commutation rules involving these number conserving products are reproduced exactly, as are the dynamical correlation functions of the free theory. Also the work of Sharp, Meniko and Goldin has shown that the field operator may be viewed as a Unitary representation of the local current group. An explicit realisation of this unitary representation is given in terms of canonical conjugate of the density operator.

1 Introduction

Recent years have seen remarkable developments in many-body theory in the form an assortment of techniques that may be loosely termed bosonization. The beginnings of these types of techniques may be traced back to the work of Tomonaga [1] and later on by Luttinger [2] and by Lieb and Mattis [3]. In the 70's an attempt was made by Luther [4] at generalising these ideas to higher dimensions. Closely related to this is work by Sharp et. al. [11] in current algebra. Mention must also be made about the work of Feenberg and his collaborators [12] who have a theory of correlated electrons written down in the standard wavefunction approach familiar in elementary quantum mechanics. It seems that the approach adopted here has some similarities with this work although the details are not identical. More progress was made by Haldane [5] which culminated in the explicit computation of the single particle propagator by Castro-Neto and Fradkin [6] and by Houghton, Marston et. al. [7] and also by Kopietz et. al. [8]. Rigorous work by Frohlich and Marchetti [9] is also along similar lines.

The attempt made here is to generalise the concepts of Haldane [5] to accommodate short wavelengths fluctuations where the concept of a linearised bare fermion energy dispersion is no longer valid. To motivate progress in this direction one must first introduce the concept of the canonical conjugate of the Fermi density distribution. This concept is likely to be important in the construction of the field operator in terms of the Fermi sea displacements, although the latter is not completed in this article. The concept of the velocity operator being the canonical conjugate of the density has been around for a long time, and this has been exploited in the study of HeII by Sunakawa et. al. [10]. However, the author is not aware of a rigorous study of the meaning of this object, in particular, an explicit formula for the canonical conjugate of the density operator has to the best of my knowledge never been written down in terms of the field operators. The work by Sharp et. al. [11] comes close to what I am attempting here.

2 Unitary Representation of the Local Current Group

The work of Sharp et. al. [11] established that the current algebra itself does not convey the underlying particle statistics rather the statistics is hidden in a choice of a unitary representation of the local current group. Here, I try to write down some formulas that provide (possibly for the first time) an explicit and exact representation of the Fermi field operator in terms of the canonical conjugate of the density operator thereby providing an explicit realisation of the claims of Sharp et. al. [11]

2.1 Some Mathematical Identities

$$(\hat{x})^y = \hat{y}(\hat{x}) (\hat{x}) \quad (1)$$

is a bose-like object; namely it satisfies

$$[(\hat{x}); (\hat{x}^0)] = 0 \quad (2)$$

Notation:

$$N = \int dx (\hat{x})$$

$$n = \frac{N}{V} \\ = \ln i$$

is a c-number, whereas N and n are operators.

So it is natural to introduce the canonical conjugate (\hat{x}) .

$$[(\hat{x}); (\hat{x}^0)] = i^d (\hat{x} - \hat{x}^0); \quad (3)$$

$$[(\hat{x}); (\hat{x}^0)] = 0 \quad (4)$$

Propose the ansatz

$$(\hat{x}) = \exp(-i(\hat{x})) \exp(i([]; \hat{x})) (\hat{x})^{\frac{1}{2}} \quad (5)$$

The above Eq. (5) shall be called the DPVA ansatz. Here again,

$$(\hat{x}) = i \ln \left[\left(\frac{1}{2} + (\hat{x}) \right) \left(\frac{1}{2} + (\hat{x}) \right)^{\frac{1}{2}} \exp(-i([]; \hat{x})) \right] \quad (6)$$

$$(\hat{x}) = (\hat{x})^{\frac{1}{2}} \quad (7)$$

$$(\hat{x}) = (\hat{x}) \quad (8)$$

is a c-number given by $\ln i$. Where $([]; \hat{x})$ is some hermitian functional to be computed later. This ansatz automatically satisfies

$$^y(\hat{x}) (\hat{x}) = (\hat{x}) \quad (9)$$

and

$$[(\hat{x}); (\hat{x}^0)] = i^d (\hat{x} - \hat{x}^0); \quad (10)$$

Write

$$(\hat{x}) = n + \frac{1}{V} \sum_{q \neq 0}^X \exp(iq\mathbf{x}) \quad (11)$$

where $n = \frac{N}{V}$, and

$$(\hat{x}) = X_0 + \sum_{q \neq 0}^X \exp(iq\mathbf{x}) X_q \quad (12)$$

$$[X_0; N] = i; \quad (13)$$

$$[X_q; q^0] = i_{q^0 q^0}; \quad (14)$$

Therefore

$$(\hat{x}) = \exp(-i \sum_q^X \exp(iq\mathbf{x}) X_q) F([f_k^0 g]; \mathbf{x}) \quad (15)$$

where

$$F([f_k^0 g]; \mathbf{x}) = \exp(i([f_k^0 g]; \mathbf{x})) \left(n + \frac{1}{V} \sum_{q \neq 0}^X \exp(iq\mathbf{x}) \right)^{\frac{1}{2}} \quad (16)$$

It is possible to write these formulas in momentum space. This will be useful later on. First we write

$$([]; \mathbf{x}) = \sum_q^X ([]; q) \exp(iq\mathbf{x}) \quad (17)$$

Therefore

$$(\mathbf{x}) = \exp\left(i \sum_q \exp(iq\mathbf{x}) X_q\right) \exp\left(i \sum_q \exp(-iq\mathbf{x}) ([]; q)\right) (n + \frac{1}{V} \sum_{q \neq 0} \exp(-iq\mathbf{x}))^{\frac{1}{2}} \quad (18)$$

It is possible to write the corresponding formula in momentum space by making the identifications

$$\exp(iq\mathbf{x}) \rightarrow T_q(\mathbf{k}) \quad (19)$$

$$\exp(-iq\mathbf{x}) \rightarrow T_q(\mathbf{k}) \quad (20)$$

where

$$T_q(\mathbf{k}) = \exp(q\mathbf{x}_k) \quad (21)$$

$$(\mathbf{k}) = \exp\left(i \sum_q T_q(\mathbf{k}) X_q\right) \exp\left(i \sum_q T_q(\mathbf{k}) ([]; q)\right) (N + \sum_{q \neq 0} T_q(\mathbf{k})^{\frac{1}{2}})_{\mathbf{k};0} \quad (22)$$

The translation operators translate the Kronecker delta that appears in the extreme right. It may be verified that

$$[(\mathbf{k}); q; 0] = (\mathbf{k} - \mathbf{q})_{; 0} \quad (23)$$

$$[(\mathbf{k}); N; 0] = (\mathbf{k})_{; 0} \quad (24)$$

Here

$$(\mathbf{x}) = \frac{1}{V^{\frac{1}{2}}} \sum_k \exp(ik\mathbf{x}) (\mathbf{k}) \quad (25)$$

Also the following identities are going to be important.

$$(\mathbf{k}) = \exp\left(i \sum_q T_q(\mathbf{k}) X_q\right) f_k([]) \quad (26)$$

$$f_k([]) = \exp\left(i \sum_q T_q(\mathbf{k}) ([]; q)\right) (N + \sum_{q \neq 0} T_q(\mathbf{k})^{\frac{1}{2}})_{\mathbf{k};0} \quad (27)$$

Check by expansion that

$$\sum_k f_{k+q=2}^y([]) f_{k-q=2}([]) = \sum_q \quad (28)$$

2.2 Proof (??) of the Fermion Commutation Rules

We use the theory of distributions to (try) prove rigorously (good enough for a physicist!) the fermion anticommutation rules. Pardon the pretense of mathematical rigor. The approach is as follows. Let \mathcal{S} be the space of all smooth functions from $S = \mathbb{R}^3$ to \mathbb{C} . Further, on this space define the inner product (Schwartz space). In $L_C^2(S) \times L_C^2(S) \rightarrow \mathbb{C}$

$$(f, g) = \int dx \sum_i f(\mathbf{x}_i) g(\mathbf{x}_i)$$

Since the Fermi fields are operator-valued distributions we can construct operators

$$c_f = \int dx \sum_i (\mathbf{x}_i) f(\mathbf{x}_i)$$

and we then have to prove

$$f c_f; c_g g = 0 \quad (29)$$

$$c_f^2 = 0 \quad (30)$$

The above two relations shall be called the fermion anticommutation rules.

$$f c_f; c_g^y g = (f, g) \quad (31)$$

The above relation shall be called the $c \leftrightarrow c^\dagger$ anticommutation rule. for all f, g that belong to $L^2_C(S)$. The claim is, that these relations are satisfied provided ψ obeys the recursion below.

$$\begin{aligned} & ([f(y_1) - (y_1 - x)^0]_1; g; x) \\ & + ([f; x^0] - ([f; x]) \\ & ([f(y_1) - (y_1 - x)^0]_1; g; x^0) = m \end{aligned} \quad (32)$$

where m is an odd integer. This recursion is to be satisfied for all $(x) \in (x^0)$. At precisely $(x) = (x^0)$ the recursion obviously breaks down. But this is not a serious drawback as we shall soon find out. The proof involves working with a basis. Here the integer m has to be even for bosons and odd for fermions. The imposition of the commutation rules by themselves do not provide us with a formula for the hermitian functional ψ . Any redefinition of (x) consistent with $[f;] = i(\cdot\cdot\cdot)$ may be absorbed by a suitable redefinition of ψ (akin to gauge transformations pointed out by AHC Neto, private communication). Solution to $([f; q])$ by making contact with the free theory. A random choice of $([f; x])$ that satisfies the recursion is (A. J. Leggett : private communication) the Leggett ansatz (for fermions) (also F D M. Haldane [5]).

$$([f; x]) = \int_0^Z dx^0 X(t(x) - t(x^0)) (x^0)$$

redefinitions of ψ that leave the statistics invariant are the 'generalised gauge transformations'. Disclaim er: Please pardon my pretensions at rigor in the following paragraph. It is easy to get carried away. The relation that relates two fermi fields by a generalised gauge transformation is an equivalence relation. Therefore, analogous to the claim that a state of a system is a ray in Hilbert space, one is tempted to call the equivalence class of fermi fields under these transformations the fermi distribution. Therefore a fermi distribution is not just one fermi field but a whole bunch of equivalent ones. Therefore a state containing one fermion at a space point is obtained by acting the creation fermi distribution on the vacuum. Thus one may construct the Fock space by repeatedly acting these fermi distributions on the vacuum. Therefore there are two ways by which one can choose ψ . One is a random choice which satisfies the recursion. In which case ψ is determined by Eq.(6). The other choice is $\psi = i\psi$. In which case, ψ can no longer be chosen arbitrarily. It has to be determined by making contact with the free theory.

Here $\theta(x)$ is the Heaviside step function. The unpleasantness caused by the fact that the recursion for fermions does not hold when $(x) = (x^0)$ is probably remediable by multiplying the fermi fields which are operator-valued distributions by arbitrary smooth functions and proving properties about these latter objects. More importantly, ψ has to be invertible in order for the ansatz to satisfy the recursion. That such a bijective mapping exists is guaranteed by the theory of cardinals. However, this mapping is not continuous let alone differentiable and therefore of little practical value. (A continuous mapping would imply a homeomorphism between \mathbb{R} and \mathbb{R}^3 eg.)

The claim is that the state of the fermi system is prescribed by prescribing an amplitude for finding the system in a given configuration of densities. Let

$$W_{FS}([f]) = \psi_H([f]) \exp(U_{FS}[f]) \exp(i_{FS}([f]))$$

be the wavefunctional of the noninteracting fermi sea. The $U_{FS}[f]$ is uniquely determined by prescribing all moments of the density operator. The functional $i_{FS}[f]$ cannot be so determined. Moreover, since the amplitude for finding the system with negative densities is zero, we must also have a prefactor,

$$\psi_H([f]) = \prod_{y_1} \theta(y_1 - \psi_H((y_1 - 1)))$$

Where $\theta_H(x)$ is the Heaviside unit step function.

2.3 Making Contact With the Free Theory

Consider the operator

$$n(y) = \int_0^Z dx^y (x + y = 2) (x - y = 2) = \sum_k^X x^y(k) (k) \exp(-ik \cdot y) \quad (33)$$

$$n(y) W_{FS}([j]) = n_0(y) W_{FS}([j]) \quad (34)$$

$$n_0(y) = \sum_k (k_F - k) \exp(-ik \cdot y) \quad (35)$$

First assume $y \notin 0$. In eq. (34) there are two undetermined functionals. One is $([j]; x)$ and the other is $W_{FS}([j])$. We have remarked earlier that redefinitions of $([j]; x)$ are similar to 'gauge transformations'. These functional gauge transformations leave the local density of particles invariant but alter the statistics. There is a subgroup of these functional gauge transformations that also leaves the statistics invariant. A realisation of this subgroup is achieved by for example:

$$([j]; x) \rightarrow ([j]; x) + ([j])$$

This may be exploited to our advantage while solving eq.(34). In other words $W_{FS}([j])$ can be absorbed into $([j]; x)$ without altering the statistics. This leaves us with an equation for $([j]; x)$ in terms of the properties of the free theory exclusively. Exploiting the recursion relation for $([j]; x)$ we arrive at the result (Here $y \notin 0$.)

$$\begin{aligned} & \int_{-\infty}^{\infty} dx \left((x+y=2) \right)^{\frac{1}{2}} \left((x-y=2) \right)^{\frac{1}{2}} \exp[i([j]; (x-y=2)) - i([j]; (x+y=2))] \\ & \exp(U_{FS}([f(y^0) - (y^0(x-y=2))], g)) - U_{FS}([f(y^0) - (y^0(x+y=2))], g)) \\ & \frac{([f(y^0) - (y^0(x+y=2))], g))}{([f(y^0) - (y^0(x-y=2))], g))} = n_0(y) \end{aligned} \quad (36)$$

The terms involving the Heaviside are indeterminate unless we expand around

$$(y) = \delta(0) + (y) \quad (37)$$

and,

$$h(y) i = \delta(0) = \quad (38)$$

2.4 Point Splitting or no Point Splitting ?

The attempts made here are partly based on the work of Ligouri and Mintchev on Generalised statistics[13] and work of Goldin et. al.[11] and the series by Reed and Simon[15] Here I shall attempt to provide a framework within which questions such as the existence of the canonical conjugate of the fermi-density distribution may be addressed. For reasons of clarity we shall not insist on utmost generality. The philosophy being that the quest for utmost generality should not cloud the underlying basic principles. We start off with some preliminaries. Let H be an infinite dimensional separable Hilbert Space. We know from textbooks that such a space possesses a countable orthonormal basis $B = \{w_i; i \in \mathbb{Z}\}$. Here, \mathbb{Z} is the set of all integers. Thus $H =$ Set of all linear combinations of vectors chosen from B . We construct the tensor product of two such spaces

$$H^{\otimes 2} = H \otimes H$$

This is defined to be the dual space of the space of all bilinear forms on the direct sum. In plain English this means something like this. Let $f \in H$ and $g \in H$ define the object $f \otimes g$ to be that object which acts as shown below. Let $\langle v; w \rangle$ be an element of the Cartesian product $H \times H$.

$$(f \otimes g) \langle v; w \rangle = (f; v) (g; w)$$

Here, $(f; v)$ stands for the inner product of f and v . Define also the inner product of two $f \otimes g$ and $f' \otimes g'$

$$(f \otimes g; f' \otimes g') = (f; f') (g; g')$$

Construct the space of all finite linear combinations of objects such as $f \otimes g$ with different choices for f and g . Lump them all into a set. You get a vector space $H^{\otimes 2}$. It is still not the vector space $H^{\otimes 2}$. Because the space of all finite linear combinations of objects such as $f \otimes g$ is not complete. Not every Cauchy sequence converges.

Complete the space by appending the limit points of all Cauchy sequences from the space of all finite linear combinations of vectors of the type $f_1 \dots f_N g$. This complete space is the Hilbert space H^2 . Similarly one can construct H^n for $n = 0, 1, 2, 3, \dots$. Where we have set $H^0 = \mathbb{C}$ the set of complex numbers. Define the Fock Space over H as

$$F(H) = \bigoplus_{n=0}^{\infty} H^{\otimes n}$$

Physically, each element of it is an ordered collection of wavefunctions with different number of particles

$$(\psi_0; \psi_1(x_1); \psi_2(x_1, x_2); \dots; \psi_n; \dots)$$

is a typical element of $F(H)$. This is the Hilbert Space which we shall be working with. Let D^n be the space of all decomposable vectors.

$$D^n = \bigoplus_{f_1, \dots, f_n \in H} f_1 \otimes \dots \otimes f_n$$

For each $f \in H$ define

$$b(f) : D^n \rightarrow D^{n+1}; n \geq 1$$

$$b(f) : D^n \rightarrow D^{n+1}; n = 0$$

defined by

$$b(f) f_1 \otimes \dots \otimes f_n = \sqrt{n} (f \otimes f_1) \otimes f_2 \otimes \dots \otimes f_n$$

$$b(f) f_1 \otimes \dots \otimes f_n = \sqrt{n+1} f \otimes f_1 \otimes f_2 \otimes \dots \otimes f_n$$

We also define $b(f)H^0 = 0$. By linearity we can extend the definitions to the space of all finite linear combinations of elements of D^n namely $L(D^n)$. For any $\psi \in L(D^n)$ and $\phi \in L(D^{n+1})$

$$\langle \psi, b(f)\phi \rangle = \sqrt{n} \langle \psi, f \otimes \phi \rangle$$

$$\langle \psi, b(f)\phi \rangle = \sqrt{n+1} \langle \psi, f \otimes \phi \rangle$$

$$(\psi, b(f)\phi) = (b(f)\psi, \phi)$$

So long as $\|\phi\| < 1$, $b(f)$ and $b(f)$ are bounded operators. An operator O is said to be bounded if

$$\sup_{\|\phi\|=1} \|O\phi\| < \infty$$

O is unbounded otherwise. The norm of a bounded operator is defined as

$$\|O\| = \sup_{\|\phi\|=1} \|O\phi\|$$

In order to describe fermions, it is necessary to construct orthogonal projectors on $F(H)$. In what follows $c(f)$ will denote a fermi annihilation operator. $c^*(f)$ will denote a fermi creation operator. Physically, and naively speaking, these are the fermi operators in "momentum space" a_k and a_k^* . First define P to be the projection operator that projects out only the antisymmetric parts of many body wavefunctions. For example,

$$P f_1 \otimes f_2 = \frac{1}{2} (f_1 \otimes f_2 - f_2 \otimes f_1)$$

We now have

$$c(f) = P b(f) P$$

$$c^*(f) = P b^*(f) P$$

Let us take a more complicated example. Let us find out how $c(f)c(g)$ acts on a vector $v = f_1 \otimes \dots \otimes f_N$.

$$c(f)c(g)v = P b(f)P P b(g)P v$$

$$c(f)c(g)v = P b(f)P b(g)P v$$

$$c(f)c(g)v = P b(f)P b(g)P v$$

$$P^{-1} v = \frac{1}{2!} (f_1^0 f_2^0 f_2^0 f_1^0)$$

$$b(g)P^{-1} v = \frac{1}{2!} P^{-2} ((g; f_1) f_2^0 (g; f_2) f_1^0)$$

$$P^{-1} b(g)P^{-1} v = \frac{1}{2!} P^{-2} ((g; f_1) f_2^0 (g; f_2) f_1^0)$$

$$b(f)P^{-1} b(g)P^{-1} v = \frac{1}{2!} P^{-2} ((g; f_1) f_1^0 f_2^0 (g; f_2) f_1^0 f_1^0)$$

$$c(f)c(g)v = \left(\frac{1}{2!}\right)^2 P^{-2} ((g; f_1) [f_1^0 f_2^0 f_2^0 f_1^0] (g; f_2) [f_1^0 f_1^0 f_1^0 f_1^0])$$

Having had a feel for how the fermi operators behave, we are now equipped to pose some more pertinent questions. Choose a basis

$$B = \{w_i; i \in \mathbb{Z}^3\}$$

2.5 Definition of the Fermi Density Distribution

Here we would like to capture the notion of the fermi density operator. Physicists call it $\rho(x) = \psi^\dagger(x) \psi(x)$. Multiplication of two fermi fields at the same point is a tricky business and we would like to make more sense out of it. For this we have to set our single particle Hilbert Space:

$$H = L_p(\mathbb{R}^3) \otimes W$$

Here, $L_p(\mathbb{R}^3)$ is the space of all periodic functions with period L in each space direction. That is if $u \in L_p(\mathbb{R}^3)$ then

$$u(x_1 + L; x_2; x_3) = u(x_1; x_2; x_3)$$

$$u(x_1; x_2 + L; x_3) = u(x_1; x_2; x_3)$$

$$u(x_1; x_2; x_3 + L) = u(x_1; x_2; x_3)$$

W is the spin space spanned by two vectors. An orthonormal basis for W

$$f^\pm; g$$

A typical element of H is given by $f(x) \otimes g$. A basis for H is given by

$$B = \left\{ \frac{1}{L^3} \exp(iq_n \cdot x) \otimes f^\pm; n = (n_1; n_2; n_3) \in \mathbb{Z}^3; s \in \{f^\pm; g\} \right\}$$

$$q_n = \left(\frac{2\pi n_1}{L}; \frac{2\pi n_2}{L}; \frac{2\pi n_3}{L} \right) g$$

We move on to the definition of the fermi density distribution. The Hilbert Space H^N is the space of all N -particle wavefunctions with no symmetry restrictions. From this we may construct orthogonal subspaces

$$H_+^N = P_+ H^N$$

$$H_-^N = P_- H^N$$

Tensors from H_+^N are orthogonal to tensors from H_-^N . The only exceptions are when $n = 0$ or $n = 1$.

$$H_+^0 = H_-^0 = C$$

$$H_+^1 = H_-^1 = H$$

The space H_+^N is the space of bosonic wavefunctions and the space H_-^N is the space of fermionic wavefunctions. The definition of the fermi density distribution proceeds as follows. Let v be written as

$$v = \sum_{2f; \#g} a(\)$$

The Fermi density distribution is an operator on the Fock Space, given a vector $f \in H$ in the single particle Hilbert Space, and a tensor $'$ in the n -particle subspace of $F(H)$, there exists a corresponding operator $(f \cdot v)$ that acts as follows:

$$[(f \cdot v)']_h(x_1, x_2, \dots, x_n) = 0$$

if $' \in H_+^N$ and

$$[(f \cdot v)']_h(x_1, x_2, \dots, x_n) = \sum_{i=1}^n f(x_i) a(i)_n(x_1, x_2, \dots, x_n)$$

when $' \in H_-^N$. The physical meaning of this abstract operator will become clear in the next subsection.

2.6 Definition of the Canonical Conjugate of the Fermi Density

We introduce some notation. Let $g = \frac{1}{L^3} \exp(ik_m \cdot x) \in \mathcal{F}_r$

$$(k_m | r) = c(g)$$

$$(k_m | r) = (g)$$

This $(k_m | r)$ is nothing but the density operator in momentum space, familiar to Physicists

$$(k_m | r) = \sum_{q_n} c_{q_n + k_m}^\dagger c_{q_n} r$$

We want to define the canonical conjugate of the density operator as an operator that maps the Fock space (or a subset thereof) on to itself. If $' \in H^0 = \mathcal{C}$ then

$$X_{q_m} ' = 0$$

Let $' \in H_+^n, n = 2, 3, \dots$ then

$$X_{q_m} ' = 0$$

The important cases are when $' \in H^N, n = 2, 3, \dots$ or if $' \in H_-$. In such a case, we set $n = N_s^0 \neq 0$ for $s \in \mathcal{F}; \#g$. Let us introduce some more notation. $N_s = \sum_{q=0}^s$ is the number operator to be distinguished from the c-number N_s^0 . The eigenvalue of N_s is N_s^0 when it acts on a state such as $' \in H^N$. Some more notation.

$$(k_m | s) = (k_m | s) \prod_{N_s^0 = k_m; 0}$$

and

$$(k_m | s) = (k_m | s) N_s^0 = k_m; 0$$

The Canonical Conjugate of the density distribution in real space denoted by $\rho_s(x)$ is defined as follows.

$$\rho_s(x_1 + L; x_2; x_3) = \rho_s(x_1; x_2; x_3)$$

$$\rho_s(x_1; x_2 + L; x_3) = \rho_s(x_1; x_2; x_3)$$

$$\rho_s(x_1; x_2; x_3 + L) = \rho_s(x_1; x_2; x_3)$$

The definition is as follows.

$$\rho_s(x) = \sum_{q_m} \exp(iq_m \cdot x) X_{q_m} s$$

$$\begin{aligned}
X_{q_m s} &= i \ln \left[\left(1 + \frac{1}{N_s^0} \sum_{k_n} (k_n s) T_{k_n}(q_m) \right) \left(1 + \sum_{k_n} \frac{1}{N_s^0} (k_n s) T_{k_n}(q_m) \right) \right]_{q_m, 0} \\
\frac{1}{N_s^0} \sum_{k_n} (k_n s) T_{k_n}(q_m) &= \frac{1}{2} \exp \left(i \sum_{k_n} ([]; k_n s) T_{k_n}(q_m) \right) \\
([]; x s) &= \sum_{k_n} ([]; k_n s) \exp(i k_n x) \\
T_{k_n}(q_m) &= \exp(k_n x_{q_m})
\end{aligned} \tag{39}$$

The translation operator translates the q_m in the Kronecker delta that appears in the extreme right by k_n and $([]; x s)$ satisfies a recursion explained in detail in the previous manuscript. The logarithm is to be interpreted as an expansion around the leading term which is either N_s^0 or $\frac{1}{N_s^0}$. The question of existence of $X_{q_m s}$ now reduces to demonstrating that this operator (possibly unbounded) maps its domain of definition (densely defined in Fock space) on to the Fock space. Defining the limit of the series expansion is likely to be the major bottleneck in demonstrating the existence of $X_{q_m s}$. That this is the canonical conjugate of the density operator is not at all obvious from the above definition. A rigorous proof of that is also likely to be difficult. Considering that we arrived at this formula by first postulating the existence of $s(x)$, it is probably safe to just say "it is clear that" $s(x)$, is in fact the canonical conjugate of ρ . The way in which the above formula can be deduced may be motivated as follows:

$$\begin{aligned}
(x) &= \frac{1}{V^{\frac{1}{2}}} \sum_k \exp(i k x) (k) \\
&= \exp \left(i \sum_q \exp(i q x) X_q \right) \exp \left(i \sum_q \exp(-i q x) ([]; q) \right) \\
(n + \frac{1}{V} \sum_{q \neq 0} X_q \exp(-i q x))^{\frac{1}{2}} &
\end{aligned} \tag{40}$$

In the above Eq. (40) ONLY on the right side make the replacements,

$$\exp(i q x) \rightarrow T_q(k) \tag{41}$$

$$\exp(-i q x) \rightarrow T_q(k) \tag{42}$$

where

$$T_q(k) = \exp(q x_k) \tag{43}$$

and append a $k, 0$ on the extreme right. Also on the LEFT side of Eqn. (40) make the replacement (x) by (k) . And you get a formula for (k) . In order to get a formula for $X_{q_m s}$ we have to invert the relation and obtain,

$$\begin{aligned}
(x s) &= i \ln \left[\left(\frac{1}{N_s^0} + \sum_{k_n} \exp(i k_n x) (k_n s) \right) \right. \\
&\quad \left. \exp \left(i \sum_{k_n} \exp(-i k_n x) ([]; k_n s) \right) \left(\frac{1}{N_s^0} + \sum_{k_n} \exp(-i k_n x) (k_n s) \right)^{\frac{1}{2}} \right]
\end{aligned} \tag{44}$$

Make the replacements on the right side of Eqn. (44)

$$\exp(i k_n x) \rightarrow T_{k_n}(q_m) \tag{45}$$

$$\exp(-i k_n x) \rightarrow T_{k_n}(q_m) \tag{46}$$

where and append a $q_m, 0$ on the extreme right. Also on the LEFT side of Eqn. (44) replace $(x s)$ by $X_{q_m s}$. This results in formula given in Eqn. (39).

3 Expression in terms of Fermi Sea Displacements

The claim is that the following exact relation holds,

$$\begin{aligned} c_{k+q=2}^y c_{k-q=2} &= n_F(k) \frac{N}{hN i} c_{q=0} + \left(\frac{N}{hN i} \right) [\epsilon_k(q) - \epsilon_{k+q}(q) a_k(q) + \epsilon_k(q) - \epsilon_{k+q}(q) a_k^y(q)] \\ &+ \sum_{k_1, k_2} \sum_{q_1, q_2} \epsilon_{k_1, k_2}^{q_1, q_2}(k; q) a_{k_1}^y(q_1) a_{k_2}(q_2) \end{aligned} \quad (47)$$

where,

$$f c_k; c_k^0 g = 0 \quad (48)$$

and

$$f c_k; c_k^y g = \epsilon_{k, k^0} \quad (49)$$

and,

$$[a_k(q); a_{k^0}(q^0)] = 0 \quad (50)$$

$$[a_k(q); a_{k^0}^y(q^0)] = \epsilon_{k, k^0} c_{q, q^0} \quad (51)$$

and,

$$\epsilon_k(q) = \frac{P}{n_F(k+q=2) (1 - n_F(k-q=2))} \quad (52)$$

and

$$N = \sum_k c_k^y c_k \quad (53)$$

$$hN i = \sum_k n_F(k) \quad (54)$$

and $n_F(k) = \langle \epsilon_k - \epsilon_{k+q} \rangle$. The $\epsilon_k(q)$ is a remainder that $\epsilon_k(q)$ is nonzero only when $k+q=0$. Also,

$$[N; a_k(q)] = 0 \quad (55)$$

$$[c_p; a_k(q)] = \text{something complicated} \quad (56)$$

and the kinetic energy operator is,

$$K = \sum_k \epsilon_k c_k^y c_k$$

in terms of the Bose fields, it is postulated to be,

$$K = E_0 + \sum_{k, q} \epsilon_k(q) a_k^y(q) a_k(q) \quad (57)$$

$$\epsilon_k(q) = \epsilon_k(q) \frac{k+q}{m}$$

and $E_0 = \sum_k \epsilon_k n_F(k)$. Also, the filled Fermi sea is identified with the Bose vacuum. $\langle S \rangle = \langle J \rangle$. The fact that this is true will become clearer later. At this point it is sufficient to note that $\epsilon_k(q) = 0$ for all k and q . We have to make sure that for the right choice of $\epsilon_k(q)$,

$$h_{q_1}(t_1) c_{q_2}(t_2) c_{q_1+q_2}(t_3) i \neq 0 \quad (58)$$

The terms linear in the Fermi sea displacements in Eq. (47) are chosen in order to ensure that the correct dynamical four-point functions are recovered. Now for the quadratic terms. The best way to derive formulae for them is to compute the exact dynamical six point function and set it equal to the expression got from the free theory.

$$h c_{k+q=2}^y(t) c_{k-q=2}(t) c_{k^0+q^0=2}^y(t^0) c_{k^0-q^0=2}(t^0) c_{k^0+q^0=2}^y(t^0) c_{k^0-q^0=2}(t^0) i = I(k; q; t; k^0; q^0; t^0; k^0; q^0; t^0)$$

From the free theory we have

$$I(k; q; t; k^0; q^0; t^0; k^{\infty}; q^{\infty}; t^{\infty}) = \exp(i \frac{k \cdot q}{m} t) \exp(i \frac{k^0 \cdot q^0}{m} t^0) \exp(i \frac{k^{\infty} \cdot q^{\infty}}{m} t^{\infty})$$

$$[(1 - n_F(k - q = 2)) (1 - n_F(k^0 - q^0 = 2)) n_F(k + q = 2)]_{k+q=2; k^0+q^0=2} \exp(i \frac{k \cdot q}{m} t) \exp(i \frac{k^0 \cdot q^0}{m} t^0) \exp(i \frac{k^{\infty} \cdot q^{\infty}}{m} t^{\infty})$$

$$[(1 - n_F(k - q = 2)) n_F(k^0 + q^0 = 2) n_F(k + q = 2)]_{k+q=2; k^0+q^0=2} \exp(i \frac{k \cdot q}{m} t) \exp(i \frac{k^0 \cdot q^0}{m} t^0) \exp(i \frac{k^{\infty} \cdot q^{\infty}}{m} t^{\infty}) \quad (59)$$

Also the six-point function may be evaluated in terms of the Bose fields (summation over superfluous indices is implied),

$$I = h_k(q) a_k(q) \exp(i!_k(q) t) \exp(i!_{k_1}(q_1) t) \exp(i!_{k_2}(q_2) t) \exp(i!_{k^{\infty}}(q^{\infty}) t) \exp(i!_{k^0}(q^0) t^0) \exp(i!_{k^{\infty}}(q^{\infty}) t^{\infty}) \exp(i!_{k^0}(q^0) t^0) \exp(i!_{k^{\infty}}(q^{\infty}) t^{\infty}) \quad (60)$$

The reduced form of the six-point function may be written as,

$$I = h_k(q) \exp(i!_k(q) t) \exp(i!_{k_1}(q_1) t) \exp(i!_{k_2}(q_2) t) \exp(i!_{k^{\infty}}(q^{\infty}) t) \exp(i!_{k^0}(q^0) t^0) \exp(i!_{k^{\infty}}(q^{\infty}) t^{\infty})$$

$$h_k(q) a_{k_1}^y(q_1) a_{k_2}(q_2) a_{k^{\infty}}^y(q^{\infty}) i$$

This may be simplified to

$$I = \exp(i!_k(q) t) \exp(i!_{k^0}(q^0) t^0) \exp(i!_{k^{\infty}}(q^{\infty}) t) \exp(i!_{k^{\infty}}(q^{\infty}) t^{\infty}) h_k(q)$$

$$h_k(q) \exp(i!_{k^0}(q^0) t^0) \exp(i!_{k^{\infty}}(q^{\infty}) t^{\infty}) \quad (61)$$

Further, since the commutation rules,

$$[c_{k+q=2}^y, c_{k^0+q^0=2}^y] = c_{k+q=2}^y c_{k^0+q^0=2}^y - c_{k^0+q^0=2}^y c_{k+q=2}^y = 0 \quad (62)$$

This means, that the coefficients also have to satisfy,

$$h_k(q) = h_{k^0}(q^0) + h_{k^{\infty}}(q^{\infty})$$

$$= h_{k^0+q^0=2}(q^0) + h_{k^{\infty}+q^{\infty}=2}(q^{\infty}) \quad (63)$$

First notice that $h_k(q)$ is either zero or one. Notice that Eq.(59), Eq.(60) and Eq.(63) are such that they suggest to us,

$$h_{k_1}(q_1) = 0 \text{ unless } k_1(q_1) = 1 \text{ and } k_2(q_2) = 1 \quad (64)$$

$$h_{k_1}(q_1) = h_{k_1+q_1=2}(q_1) \text{ for } k_1(q_1) = 1$$

$$h_{k_2}(q_2) = h_{k_2+q_2=2}(q_2) \text{ for } k_2(q_2) = 1 \quad (65)$$

and

$$h_{k_1}(q_1) = 0 \text{ for } k_1(q_1) = 0 \quad (66)$$

From this one may verify that the formula for the kinetic energy operator written down previously, Eq.(57) is correct. Lastly, we must make sure that the following additional commutation rule holds.

$$h_{k_1}(q_1) h_{k_2}(q_2) = h_{k_1+q_1=2}(q_1) h_{k_2+q_2=2}(q_2)$$

$$= h_{k_1+q_1=2}(q_1) h_{k_2+q_2=2}(q_2) \quad (67)$$

Verification of the above rule completes the confirmation that the formula for $c_{k+q=2}^y$ in terms of the boson fields is exact. The algebra here goes through without a hitch except in this respect, namely we have to assume that the $h_k(q) = 1$.

To summarise let us write down the following:

THEOREM :

$$c_{k+q=2}^y c_{k-q=2} = n_F(k) \frac{N}{hN-1} c_{q=0} + \left(\frac{N}{hN-1} \right) [k(q) a_k(q) + k(q) a_k^y(q)] \\ + \sum_{k_1, k_2} \sum_{q_1, q_2}^{q_1+q_2} (k; q) a_{k_1}^y(q_1) a_{k_2}(q_2) \quad (68)$$

together with

$$a_{k_1, k_2}^{q_1, q_2}(k; q) = k_1(q_1) k_2(q_2) [k_1 q_1=2; k_2 q_2=2 \quad k_1+q_1=2; k+q=2 \quad k-q=2; k_2+q_2=2 \\ k_1+q_1=2; k_2+q_2=2 \quad k_1 q_1=2; k-q=2 \quad k+q=2; k_2 q_2=2] \quad (69)$$

is an exact transformation of products of Fermi fields into Bose fields. Namely, all dynamical moments of $c_{k+q=2}^y c_{k-q=2}$ are recovered exactly, provided we identify the filled Fermi sea with the Bose vacuum.

$$a_k(q) \Psi_S = 0 \quad (70)$$

and the above is also an operator identity, namely they satisfy proper mutual commutation rules.

This may also be written in a simplified form as,

$$c_{k+q=2}^y c_{k-q=2} = n_F(k) \frac{N}{hN-1} c_{q=0} + \left(\frac{N}{hN-1} \right) [k(q) a_k(q) + k(q) a_k^y(q)] \\ + \sum_{k+q=2} \sum_{q_1=2} (q_1) k_{q_1=2}(q-q_1) a_{k+q=2, q_1=2}^y(q_1) a_{k-q_1=2}(q+q_1) \\ + \sum_{k-q=2+q_1=2} (q_1) k_{k+q_1=2}(q-q_1) a_{k-q=2+q_1=2}^y(q_1) a_{k+q_1=2}(q+q_1) \quad (71)$$

The $\frac{N}{hN-1}$ is needed to ensure that, $\sum_k c_k^y c_k = N$ (and not $\sum_k c_k^y c_k = hN-1$) The $\left(\frac{N}{hN-1} \right)$ is needed to ensure that the commutation amongst the $c_{k+q=2}^y c_{k-q=2}$ come out right. However, there are other finer points that need to be discussed. For example, do the thermodynamic expectation values also come out right? That is, assuming that the temperature is finite, can we compute the dynamical density correlation function and show that it agrees with the free theory? The finite temp. case is most unusual. It seems that for dealing with fermions at finite temperature, we have to assume that the Bose like excitations are always at zero temperature with zero chemical potential, only the coefficients in the expansion acquire finite temp. values. All this is very strange and on shaky ground. Thus, I shall relegate the finite temp. case to the last section where the skeptical reader may have a field demolishing my ideas.

3.1 The Fermi Field Operator

Also how about expressing the Fermi fields themselves in terms of the Fermi sea displacements? For this one has to first construct a canonical conjugate of the density and then use it in the DPVA ansatz Eq.(5) [16] to compute the phase functional. I have tried to construct a canonical conjugate of (any canonical conjugate succeeds) but it has not been successful. Just for future reference, it is useful to write down a formula for the total current operator.

$$J_{tot} = \sum_k \left(\frac{k}{m} \right) c_k^y c_k = \sum_{k_1, k_2} \sum_{q_1, q_2} \frac{q_1, q_2}{k_1, k_2} a_{k_1}^y(q_1) a_{k_2}(q_2) \quad (72)$$

$$\frac{q_1, q_2}{k_1, k_2} = k_1(q_1) k_2(q_2) \left(\frac{q_1}{m} \right)_{k_1, k_2, q_1, q_2} \quad (73)$$

Therefore,

$$J_{tot} = \sum_k \left(\frac{k}{m} \right) c_k^y c_k = \sum_{k_1, q_1} k_1(q_1) \left(\frac{q_1}{m} \right) a_{k_1}^y(q_1) a_{k_1}(q_1) \quad (74)$$

Thus an alternative but less systematic approach seems to be in order. For this choose,

$$(\mathbf{x}) = \exp(-iU_0(\mathbf{x}))\exp(-iU_1(\mathbf{x}))\exp(-iU_2(\mathbf{x}))(\mathbf{x})^{\frac{1}{2}} \quad (75)$$

here $U_0(\mathbf{x})$ is a c-number and,

$$U_1(\mathbf{x}) = \sum_{\mathbf{k}, \mathbf{q}} [\alpha_{\mathbf{k}}(\mathbf{q})f(\mathbf{k}; \mathbf{q}; \mathbf{x}) + \alpha_{\mathbf{k}}^{\dagger}(\mathbf{q})f(\mathbf{k}; \mathbf{q}; \mathbf{x})] \quad (76)$$

$$U_2(\mathbf{x}) = \sum_{\mathbf{k}_1, \mathbf{q}_1} \sum_{\mathbf{k}_2, \mathbf{q}_2} U_2(\mathbf{k}_1; \mathbf{q}_1; \mathbf{k}_2; \mathbf{q}_2; \mathbf{x}) \alpha_{\mathbf{k}_1}^{\dagger}(\mathbf{q}_1) \alpha_{\mathbf{k}_2}(\mathbf{q}_2) \quad (77)$$

Now let us evolve this using the kinetic energy operator,

$$(\mathbf{x}; t) = e^{i t K} (\mathbf{x}) e^{-i t K} = \exp(-iU_0(\mathbf{x}))\exp(-iU_1(\mathbf{x}; t))\exp(-iU_2(\mathbf{x}; t))(\mathbf{x}; t)^{\frac{1}{2}} \quad (78)$$

here,

$$U_1(\mathbf{x}; t) = \sum_{\mathbf{k}, \mathbf{q}} [\alpha_{\mathbf{k}}(\mathbf{q})e^{i t \epsilon_{\mathbf{k}}(\mathbf{q})}f(\mathbf{k}; \mathbf{q}; \mathbf{x}) + \alpha_{\mathbf{k}}^{\dagger}(\mathbf{q})e^{i t \epsilon_{\mathbf{k}}(\mathbf{q})}f(\mathbf{k}; \mathbf{q}; \mathbf{x})] \quad (79)$$

$$U_2(\mathbf{x}; t) = \sum_{\mathbf{k}_1, \mathbf{q}_1} \sum_{\mathbf{k}_2, \mathbf{q}_2} U_2(\mathbf{k}_1; \mathbf{q}_1; \mathbf{k}_2; \mathbf{q}_2; \mathbf{x}) \alpha_{\mathbf{k}_1}^{\dagger}(\mathbf{q}_1) \alpha_{\mathbf{k}_2}(\mathbf{q}_2) e^{i t (\epsilon_{\mathbf{k}_1}(\mathbf{q}_1) - \epsilon_{\mathbf{k}_2}(\mathbf{q}_2))} \quad (80)$$

The basic philosophy it seems, involves observing that the commutation rules satisfied by number conserving objects such as $c_{\mathbf{k}+\mathbf{q}=2}^{\dagger} c_{\mathbf{k}-\mathbf{q}=2}$ amongst themselves is independent of the underlying statistics. Thus one is lead to postulate convenient forms for such objects in terms of other Bose fields. But not all approaches are likely to be correct. Take for example, the straightforward substitution,

$$c_{\mathbf{k}+\mathbf{q}=2}^{\dagger} c_{\mathbf{k}-\mathbf{q}=2} = b_{\mathbf{k}+\mathbf{q}=2}^{\dagger} b_{\mathbf{k}-\mathbf{q}=2} \quad (81)$$

where $c_{\mathbf{k}}$ are fermions and $b_{\mathbf{k}}$ are bosons. This no doubt makes the commutation rules come out right but is quite obviously wrong because all the correlation functions are wrongly represented.

4 Luttinger and Fermi Liquids

Consider an interaction of the type,

$$H_I = \sum_{\mathbf{q} \neq 0} \frac{V_{\mathbf{q}}}{2V} \sum_{\mathbf{k}, \mathbf{k}'} [\alpha_{\mathbf{k}}(\mathbf{q})\alpha_{\mathbf{k}'}(\mathbf{q}) + \alpha_{\mathbf{k}}^{\dagger}(\mathbf{q})\alpha_{\mathbf{k}'}^{\dagger}(\mathbf{q})][\alpha_{\mathbf{k}''}(\mathbf{q})\alpha_{\mathbf{k}'''}(\mathbf{q}) + \alpha_{\mathbf{k}''}^{\dagger}(\mathbf{q})\alpha_{\mathbf{k}'''}^{\dagger}(\mathbf{q})] \quad (82)$$

$$i \frac{\partial}{\partial t} \alpha_{\mathbf{k}}^{\dagger}(\mathbf{q}) = \epsilon_{\mathbf{k}}(\mathbf{q})\alpha_{\mathbf{k}}^{\dagger}(\mathbf{q}) + \left(\frac{V_{\mathbf{q}}}{V}\right) \sum_{\mathbf{k}''} [\alpha_{\mathbf{k}''}(\mathbf{q})\alpha_{\mathbf{k}}^{\dagger}(\mathbf{q}) + \alpha_{\mathbf{k}''}^{\dagger}(\mathbf{q})\alpha_{\mathbf{k}}^{\dagger}(\mathbf{q})] \quad (83)$$

The equations of motion for the Bose propagators read as,

$$\begin{aligned} \left(i \frac{\partial}{\partial t} - \epsilon_{\mathbf{k}}(\mathbf{q})\right) \frac{i \hbar \Gamma \alpha_{\mathbf{k}}^{\dagger}(\mathbf{q}) \alpha_{\mathbf{k}''}^{\dagger}(\mathbf{q}) i}{\hbar \Gamma 1 i} &= \sum_{\mathbf{k}', \mathbf{q}'} \alpha_{\mathbf{k}'}^{\dagger}(\mathbf{q}') \alpha_{\mathbf{k}''}^{\dagger}(\mathbf{q}') (t) \\ + \left(\frac{V_{\mathbf{q}}}{V}\right) \sum_{\mathbf{k}''} \alpha_{\mathbf{k}''}(\mathbf{q}) &\left[\alpha_{\mathbf{k}''}(\mathbf{q}) \frac{i \hbar \Gamma \alpha_{\mathbf{k}''}^{\dagger}(\mathbf{q}) \alpha_{\mathbf{k}''}^{\dagger}(\mathbf{q}) i}{\hbar \Gamma 1 i} + \alpha_{\mathbf{k}''}^{\dagger}(\mathbf{q}) \frac{i \hbar \Gamma \alpha_{\mathbf{k}''}^{\dagger}(\mathbf{q}) \alpha_{\mathbf{k}''}^{\dagger}(\mathbf{q}) i}{\hbar \Gamma 1 i} \right] \end{aligned} \quad (84)$$

$$\begin{aligned} &\left(i \frac{\partial}{\partial t} - \epsilon_{\mathbf{k}}(\mathbf{q})\right) \frac{i \hbar \Gamma \alpha_{\mathbf{k}}^{\dagger}(\mathbf{q}) \alpha_{\mathbf{k}''}^{\dagger}(\mathbf{q}) i}{\hbar \Gamma 1 i} \\ &= \left(\frac{V_{\mathbf{q}}}{V}\right) \sum_{\mathbf{k}''} \alpha_{\mathbf{k}''}(\mathbf{q}) \left[\alpha_{\mathbf{k}''}(\mathbf{q}) \frac{i \hbar \Gamma \alpha_{\mathbf{k}''}^{\dagger}(\mathbf{q}) \alpha_{\mathbf{k}''}^{\dagger}(\mathbf{q}) i}{\hbar \Gamma 1 i} + \alpha_{\mathbf{k}''}^{\dagger}(\mathbf{q}) \frac{i \hbar \Gamma \alpha_{\mathbf{k}''}^{\dagger}(\mathbf{q}) \alpha_{\mathbf{k}''}^{\dagger}(\mathbf{q}) i}{\hbar \Gamma 1 i} \right] \end{aligned} \quad (85)$$

The boundary conditions on these propagators may be written down as,

$$\frac{i\hbar\Gamma a_k^{yt}(q)a_{k^0}^y(q^0)i}{\hbar\Gamma 1i} = \frac{i\hbar\Gamma a_k^{y(t-i)}(q)a_{k^0}^y(q^0)i}{\hbar\Gamma 1i} \quad (86)$$

$$\frac{i\hbar\Gamma a_k^t(q)a_{k^0}^y(q^0)i}{\hbar\Gamma 1i} = \frac{i\hbar\Gamma a_k^{(t-i)}(q)a_{k^0}^y(q^0)i}{\hbar\Gamma 1i} \quad (87)$$

$$(t) = \left(\frac{1}{i}\right)^X \exp(!_n t) \quad (88)$$

$$(t) = \left(\frac{1}{i}\right)^X \frac{\exp(!_n t)}{!_n} \quad (89)$$

The boundary conditions imply that we may write,

$$\frac{i\hbar\Gamma a_k^t(q)a_{k^0}^y(q^0)i}{\hbar\Gamma 1i} = \sum_n \exp(!_n t) \frac{i\hbar\Gamma a_k^n(q)a_{k^0}^y(q^0)i}{\hbar\Gamma 1i} \quad (90)$$

$$\frac{i\hbar\Gamma a_k^{yt}(q)a_{k^0}^y(q^0)i}{\hbar\Gamma 1i} = \sum_n \exp(!_n t) \frac{i\hbar\Gamma a_k^{yn}(q)a_{k^0}^y(q^0)i}{\hbar\Gamma 1i} \quad (91)$$

and, $!_n = (2-n) =$. Thus,

$$(i!_n - !_k(q)) \frac{i\hbar\Gamma a_k^n(q)a_{k^0}^y(q^0)i}{\hbar\Gamma 1i} = \frac{k^0 k^0 q^0 q^0}{i} + \left(\frac{v_q}{V}\right)_k(q) \sum_{k^0} [k^0(q) \frac{i\hbar\Gamma a_{k^0}^n(q)a_{k^0}^y(q^0)i}{\hbar\Gamma 1i} + k^0(q) \frac{i\hbar\Gamma a_{k^0}^{yn}(q)a_{k^0}^y(q^0)i}{\hbar\Gamma 1i}] \quad (92)$$

$$(i!_n + !_k(q)) \frac{i\hbar\Gamma a_k^{yn}(q)a_{k^0}^y(q^0)i}{\hbar\Gamma 1i} = \left(\frac{v_q}{V}\right)_k(q) \sum_{k^0} [k^0(q) \frac{i\hbar\Gamma a_{k^0}^{yn}(q)a_{k^0}^y(q^0)i}{\hbar\Gamma 1i} + k^0(q) \frac{i\hbar\Gamma a_{k^0}^n(q)a_{k^0}^y(q^0)i}{\hbar\Gamma 1i}] \quad (93)$$

Define,

$$\sum_k k(q) \frac{i\hbar\Gamma a_k^n(q)a_{k^0}^y(q^0)i}{\hbar\Gamma 1i} = G_1(q; k^0; q^0; n) \quad (94)$$

$$\sum_k k(q) \frac{i\hbar\Gamma a_k^{yn}(q)a_{k^0}^y(q^0)i}{\hbar\Gamma 1i} = G_2(q; k^0; q^0; n) \quad (95)$$

Multiplying the above equations with $k^0(q)$ and summing over k one arrives at simple formulas for G_1 and G_2 .

$$G_1(q; k^0; q^0; n) = k^0(q) \frac{q^0 q^0}{i (i!_n - !_{k^0}(q))} + f_n(q) [G_1(q; k^0; q^0; n) + G_2(q; k^0; q^0; n)] \quad (96)$$

and,

$$G_2(q; k^0; q^0; n) = f_n(q) [G_1(q; k^0; q^0; n) + G_2(q; k^0; q^0; n)] \quad (97)$$

and,

$$f_n(q) = \left(\frac{v_q}{V}\right)_k \sum_k \frac{k(q)}{(i!_n - !_k(q))} \quad (98)$$

In 1D and 3D an analytical solution for $f_n(q)$ is possible. In 1D we have,

$$f_n(q) = v_q \left(\frac{m}{q} \right) \left(\frac{1}{2} \right) \ln \left(\frac{k_f \frac{\frac{m}{q} i!_n + \frac{q}{2}}{k_f \frac{\frac{m}{q} i!_n + \frac{q}{2}} + \frac{q}{2}} \right) \quad (b-a) v_q \left(\frac{1}{2} \right) \left(\frac{m}{q} \right) \ln \left(\frac{b \frac{\frac{m}{q} i!_n + \frac{q}{2}}{a \frac{\frac{m}{q} i!_n + \frac{q}{2}} + \frac{q}{2}} \right) \quad (99)$$

where, $b = \min(k_f; k_f - q)$ and $a = \max(k_f; k_f - q)$. The logarithm is to be interpreted as the principal value, that is, $\text{Im}(\ln(z)) = \text{Arg}(z) = \tan^{-1}(y/x)$, $x = \text{Re}(z)$; $y = \text{Im}(z)$, and $\text{Re}(\ln(z)) = \ln(r)$ and, $r = (x^2 + y^2)^{\frac{1}{2}}$. The nice thing about this definition (not shared by non-principal value defs.) is that $(\ln(z))^* = \ln(z^*)$. This is because $\text{Arg}(z) = \tan^{-1}(y/x)$. This may easily be solved and the final formulas for the Bose propagators may be written down as,

$$G_2(q; k^0; q^0; n) = \frac{f_n(q)}{(1 - f_n(q))} G_1(q; k^0; q^0; n)$$

$$G_1(q; k^0; q^0; n) + G_2(q; k^0; q^0; n) = G_1(q; k^0; q^0; n) = (1 - f_n(q))$$

$$G_1(q; k^0; q^0; n) = \left(\frac{1}{i} \right) \frac{(1 - f_n(q)) k^0(q) q; q^0}{(1 - f_n(q)) f_n(q) (i!_n i!_{k^0(q)})} \quad (100)$$

$$G_2(q; k^0; q^0; n) = \left(\frac{1}{i} \right) \frac{f_n(q) k^0(q) q; q^0}{(1 - f_n(q)) f_n(q) (i!_n i!_{k^0(q)})} \quad (101)$$

$$G_1(q; k^0; q^0; n) + G_2(q; k^0; q^0; n) = \left(\frac{1}{i} \right) \frac{k^0(q) q; q^0}{(1 - f_n(q)) f_n(q) (i!_n i!_{k^0(q)})}$$

$$\begin{aligned} \frac{i\hbar T a_k^n(q) a_{k^0}^y(q^0) i}{\hbar T i} &= \frac{k; k^0 q; q^0}{i (i!_n i!_k(q))} \\ &+ \left(\frac{1}{i} \right) \left(\frac{v_q}{V} \right) \frac{k(q)}{(i!_n i!_k(q))} \frac{k^0(q) q; q^0}{(1 - f_n(q)) f_n(q) (i!_n i!_{k^0(q)})} \end{aligned} \quad (102)$$

also,

$$\frac{i\hbar T a_k^{yn}(q) a_{k^0}^y(q^0) i}{\hbar T i} = \left(\frac{1}{i} \right) \left(\frac{v_q}{V} \right) \frac{k(q)}{(i!_n + i!_k(q))} \frac{k^0(q) q; q^0}{(1 - f_n(q)) f_n(q) (i!_n i!_{k^0(q)})} \quad (103)$$

The zero temperature correlation function of significance here is,

$$i\hbar a_{k^0}^y(q^0) a_k(q) i \quad (104)$$

This may be obtained from the above formulas as,

$$i\hbar a_{k^0}^y(q^0) a_k(q) i = \left(\frac{v_q}{V} \right) k(q) k^0(q) q; q^0 \oint_C \frac{d!}{2 i (i!_n i!_k(q)) (i!_n i!_{k^0(q)}) (1 - f_n(q)) f_n(q)} \quad (105)$$

where C is the positively oriented contour that encloses the upper half plane. Thus the problem now boils down to computing all the zeros of $(1 - f_n(q) f_n(q))$ that have positive imaginary parts. In 1D, we may proceed as follows,

$$\begin{aligned} 1 - f_n(q) f_n(q) &= 1 - v_q \left(\frac{m}{q} \right) \left(\frac{1}{2} \right) \ln \left[\frac{k_f \frac{\frac{m}{q} i! + \frac{q}{2}}{k_f \frac{\frac{m}{q} i! + \frac{q}{2}} + \frac{q}{2}} \right] \\ &+ (b-a) v_q \left(\frac{1}{2} \right) \left(\frac{m}{q} \right) \ln \left[\frac{b \frac{\frac{m}{q} i! + \frac{q}{2}}{a \frac{\frac{m}{q} i! + \frac{q}{2}} + \frac{q}{2}} \right] \\ &v_q \left(\frac{m}{q} \right) \left(\frac{1}{2} \right) \ln \left[\frac{k_f \frac{\frac{m}{q} i! + \frac{q}{2}}{k_f \frac{\frac{m}{q} i! + \frac{q}{2}} + \frac{q}{2}} \right] \\ &+ (b^0 - a^0) v_q \left(\frac{1}{2} \right) \left(\frac{m}{q} \right) \ln \left[\frac{b^0 \frac{\frac{m}{q} i! + \frac{q}{2}}{a^0 \frac{\frac{m}{q} i! + \frac{q}{2}} + \frac{q}{2}} \right] \end{aligned} \quad (106)$$

where $a = \max(k_f; k_f - q)$, $b = \min(k_f; k_f - q)$ and $a^0 = \max(k_f; k_f + q)$, $b^0 = \min(k_f; k_f + q)$. There are several regions of interest.

(A) $0 < q < k_f$ This means, $0 < q < k_f$ or $k_f - k_f - q < 0$, $k_f - k_f - q < 2k_f$, $k_f - k_f + q < 2k_f$, $k_f - k_f + q < 0$ Thus $a = k_f$, $b = k_f - q$, $a^0 = k_f + q$, $b^0 = k_f$. $(b - a) = 1$, $(b^0 - a^0) = 1$ Therefore,

$$1 - f_n(q) - f_n(q) = 1 + v_q \left(\frac{m}{q}\right) \left(\frac{1}{2}\right) \ln \left[\frac{k_f \frac{m-i!}{q} + \frac{q}{2}}{k_f \frac{m-i!}{q} + \frac{q}{2}} \right] + v_q \left(\frac{1}{2}\right) \left(\frac{m}{q}\right) \ln \left[\frac{k_f \frac{m-i!}{q} + \frac{q}{2}}{k_f \frac{m-i!}{q} + \frac{q}{2}} \right] \quad (107)$$

$$1 - f_n(q) - f_n(q) = 1 + v_q \left(\frac{1}{2}\right) \left(\frac{m}{q}\right) \ln \left[\frac{(k_f + q-2)^2 + \left(\frac{m-i!}{q}\right)^2}{(k_f - q-2)^2 + \left(\frac{m-i!}{q}\right)^2} \right] = 0 \quad (108)$$

$$\ln \left[\frac{(k_f + q-2)^2 + \left(\frac{m-i!}{q}\right)^2}{(k_f - q-2)^2 + \left(\frac{m-i!}{q}\right)^2} \right] = \left(\frac{2}{m}\right) \left(\frac{1}{v_q}\right) \quad (109)$$

$$\frac{(k_f + q-2)^2 + \left(\frac{m-i!}{q}\right)^2}{(k_f - q-2)^2 + \left(\frac{m-i!}{q}\right)^2} = \exp \left(\left(\frac{2}{m}\right) \left(\frac{1}{v_q}\right) \right) \quad (110)$$

$$\left(\frac{m-i!}{q}\right)^2 = [(k_f + q-2)^2 - (k_f - q-2)^2 \exp \left(\left(\frac{2}{m}\right) \left(\frac{1}{v_q}\right) \right)] = [1 - \exp \left(\left(\frac{2}{m}\right) \left(\frac{1}{v_q}\right) \right)] \quad (111)$$

We want to find a root of this that has a positive imaginary part.

$$! = i \left(\frac{q}{m}\right)^{\frac{1}{2}} \sqrt{\frac{(k_f + q-2)^2 - (k_f - q-2)^2 \exp \left(\left(\frac{2}{m}\right) \left(\frac{1}{v_q}\right) \right)}{1 - \exp \left(\left(\frac{2}{m}\right) \left(\frac{1}{v_q}\right) \right)}} \quad (112)$$

The next case is,

(B) $k_f - q < 0$. In this case, $0 < k_f + q < k_f$, $2k_f - k_f + q < k_f$, $k_f - k_f - q < 0$, $2k_f - k_f - q < k_f$, $0 < k_f - q < k_f$, $a = k_f - q$, $b = k_f$, $a^0 = k_f$, $b^0 = k_f + q$. Thus, $(b - a) = 1$, $(b^0 - a^0) = 1$ For case (B)

$$1 - f_n(q) - f_n(q) = 1 + v_q \left(\frac{m}{q}\right) \left(\frac{1}{2}\right) \ln \left[\frac{k_f \frac{m-i!}{q} + \frac{q}{2}}{k_f \frac{m-i!}{q} + \frac{q}{2}} \right] + v_q \left(\frac{1}{2}\right) \left(\frac{m}{q}\right) \ln \left[\frac{k_f \frac{m-i!}{q} + \frac{q}{2}}{k_f \frac{m-i!}{q} + \frac{q}{2}} \right] \quad (113)$$

Thus the pole for both cases (A) and (B) is given by,

$$! = i \left(\frac{q}{m}\right)^{\frac{1}{2}} \sqrt{\frac{(k_f + q-2)^2 - (k_f - q-2)^2 \exp \left(\left(\frac{2}{m}\right) \left(\frac{1}{v_q}\right) \right)}{1 - \exp \left(\left(\frac{2}{m}\right) \left(\frac{1}{v_q}\right) \right)}} \quad (114)$$

The quantity under the square root is positive in both cases. Third, we have the case,

(C) $k_f - q < 2k_f$, this means, $k_f - q < 2k_f$, also, $2k_f - k_f + q < 3k_f$, $0 < k_f + q < k_f$, $0 < k_f - q < k_f$, $2k_f - k_f - q < 3k_f$. $a = k_f$, $b = k_f - q$, $a^0 = k_f + q$, $b^0 = k_f$. $(b - a) = 1$ and $(b^0 - a^0) = 1$ This leads to the same root Eq.(114). In case

(D) $2k_f - q < k_f$, $k_f - k_f + q < 0$, $3k_f - k_f + q < 2k_f$, $2k_f - k_f - q < k_f$, $3k_f - k_f - q < 2k_f$, $k_f - k_f - q < 0$. $a = k_f - q$, $b = k_f$, $a^0 = k_f$, $b^0 = k_f + q$. $(b - a) = 1$ and $(b^0 - a^0) = 1$. In this case also we find the root given by Eq.(114). (E) $2k_f - q < 1$, $2k_f - q > 1$, $3k_f - k_f + q < 1$, $k_f - k_f + q < 1$, $k_f - k_f - q > 1$, $3k_f - k_f - q > 1$. $a = k_f$, $b = k_f - q$, $a^0 = k_f + q$, $b^0 = k_f$. $(b - a) = 0$, $(b^0 - a^0) = 0$. This case is somewhat different.

$$1 - f_n(q) - f_n(q) = 1 + v_q \left(\frac{m}{q}\right) \left(\frac{1}{2}\right) \ln \left[\frac{k_f \frac{m-i!}{q} + \frac{q}{2}}{k_f \frac{m-i!}{q} + \frac{q}{2}} \right]$$

$$v_q \left(\frac{m}{q} \right) \left(\frac{1}{2} \right) \ln \left[\frac{k_f \frac{m-i!}{q} \frac{q}{2}}{k_f \frac{m-i!}{q} \frac{q}{2}} \right] \quad (115)$$

This when solved gives the same result as Eq.(114). So too does the final case (F) which is
(F) $1 < q \leq 2k_f$, $1 < k_f + q \leq k_f$, $1 < k_f + q \leq 3k_f$, $1 > q \leq 2k_f$, $1 > k_f - q \leq 3k_f$,
 $1 > k_f - q \leq k_f$. This case also leads to the same result namely, Eq.(114). Therefore the final result may be written as,

$$h_{k^0}^y(q) a_k(q) i = \left(\frac{1}{V} \right) \frac{k(q) k^0(q) q; q^0}{(!_R(q) + !_k(q)) (!_R(q) + !_{k^0}(q)) \left(\frac{m}{q^2} \right) \left(\frac{1}{2k_f} \right) 2 \left(\frac{m}{q} \right)^2 !_R(q) (\cosh(q) - 1)} \quad (116)$$

Here we may write,

$$k(q) = \left(\frac{2q}{m} \right) \left(\frac{1}{v_q} \right) \quad (117)$$

$$!_R(q) = \left(\frac{m}{2} \right)^s \frac{(k_f + q/2)^2 - (k_f - q/2)^2 \exp(-q)}{1 - \exp(-q)} \quad (118)$$

In other words,

$$\begin{aligned} \langle h_{\mathbf{k}}^y \rangle &= n_F(\mathbf{k}) + \frac{1}{2} \frac{d q_1}{d \mathbf{k}} \frac{k_{q_1=2}(\mathbf{q}_1)}{2!_R(\mathbf{q}_1)(!_R(\mathbf{q}_1) + !_{k+q_1=2}(\mathbf{q}_1))^2 \left(\frac{m^3}{q_1^4}\right) (\cosh(\mathbf{q}_1) - 1)} \\ &= \frac{1}{2} \frac{d q_1}{d \mathbf{k}} \frac{k_{q_1=2}(\mathbf{q}_1)}{2!_R(\mathbf{q}_1)(!_R(\mathbf{q}_1) + !_{k+q_1=2}(\mathbf{q}_1))^2 \left(\frac{m^3}{q_1^4}\right) (\cosh(\mathbf{q}_1) - 1)} \end{aligned} \quad (119)$$

In order to see how good the present theory is, it is desirable to compare these results with the Calogero-Sutherland model or more specifically with the spin-spin correlation function of the Haldane-Shastry model. This is given by [14], $\beta = 1/2$, $\beta = 2$, $m = 1$, $k_F = \pi/2$.

$$\begin{aligned} \langle h_{\mathbf{j}}^y(\mathbf{x};0) h_{\mathbf{x}^0}^y(0) \rangle &= \frac{1}{2} \frac{d \mathbf{k}}{d \mathbf{x}} \exp(i \mathbf{k} \cdot (\mathbf{x} - \mathbf{x}^0)) \langle h_{\mathbf{k}}^y \rangle \\ &= \frac{1}{4} \frac{(3/2)^2}{(1/2)^2 (1)^2} \int_0^{\pi/2} dv_1 \int_0^{\pi/2} dv_2 (1 - v_1^2)^{-1/2} (1 - v_2^2)^{-1/2} j_1 - v_2 j_2 \\ &\quad \exp(i \frac{\pi}{2} (\mathbf{x} - \mathbf{x}^0) \cdot \mathbf{v}_1) \exp(i \frac{\pi}{2} (\mathbf{x} - \mathbf{x}^0) \cdot \mathbf{v}_2) \\ &= 4 \frac{(3/2)^2}{(1/2)^2 (1)^2} \sum_{n=0}^{\infty} \left(\frac{\pi}{2}\right)^n \frac{(2n)^2 (\mathbf{x} - \mathbf{x}^0)^{2n}}{(2n)!} \frac{1}{(2n+1)^2} \end{aligned} \quad (120)$$

For this we have to use the interaction given by $V(\mathbf{x}) = \frac{(1)}{x^2}$,

$$v_q = \frac{1}{q} \quad (1) \quad (121)$$

4.1 Qualitative Conclusions

It is clear that one of the features of the Luttinger liquid that is a result of the Mattis-Lieb solution is absent, namely the discontinuous dependence of the momentum distribution on the coupling strength as the latter goes to zero. This may therefore be an artifact of the Luttinger model and not a generic feature of all 1D systems. Also the highly nonperturbative manner in which the momentum distribution of the interacting system approaches the noninteracting one as the coupling goes to zero may be seen quite easily. The other point that may be seen is that $\langle h_{\mathbf{q}=0} \rangle = 0$ indicating that there is no Wigner crystallization at any density. Furthermore, the vanishing of the quasiparticle residue is seen only for sufficiently large values of the repulsion between the fermions. Thus the conventional view that the Fermi surface is destroyed for arbitrarily weak repulsion in case of 1d fermions does not seem to hold up in this case.

4.2 Momentum Distribution in 3D

For this one has to compute the expressions in 3D :

$$\begin{aligned} f_n(q) &= \frac{v_q}{(2\pi)^2} \int_0^{k_F} dk k^2 f\left(\frac{m}{k} \ln\left(\frac{i!_n}{i!_n} \frac{q + \frac{k j_j}{m}}{q + \frac{k j_j}{m}}\right)\right) \\ &\quad + \left(\frac{m}{k j_j}\right) (k_F - j_j - k) \ln\left(\frac{i!_n}{i!_n} \frac{q + \frac{k j_j}{m}}{q + \frac{k j_j}{m}}\right) \\ &\quad + \left(\frac{m}{k j_j}\right) (j_j + k - k_F) (k_F^2 - (k - j_j)^2) \ln\left(\frac{i!_n}{i!_n} \frac{k_F + \frac{k}{m}}{q + \frac{k j_j}{m}}\right) g \end{aligned} \quad (122)$$

This may be split into several cases

(A) $0 < \hbar j < k_f$

$$\begin{aligned}
 f_n(q) = & \frac{v_q}{(2)^2} \left(\frac{m}{\hbar j}\right) f \int_0^{Z_{k_f}} dk k \ln \left(\frac{i!_n q + \frac{k \hbar j}{m}}{i!_n q + \frac{k \hbar j}{m}} \right) \\
 & + \int_0^{Z_{k_f} \hbar j} dk k \ln \left(\frac{i!_n q + \frac{k \hbar j}{m}}{i!_n q + \frac{k \hbar j}{m}} \right) \\
 & + \int_{k_f \hbar j}^{Z_{k_f}} dk k \ln \left(\frac{i!_n k_f + k}{i!_n q + \frac{k \hbar j}{m}} \right) g
 \end{aligned} \tag{123}$$

(B) $k_f < \hbar j < 2 k_f$

$$\begin{aligned}
 f_n(q) = & \frac{v_q}{(2)^2} \left(\frac{m}{\hbar j}\right) f \int_0^{Z_{k_f}} dk k \ln \left(\frac{i!_n q + \frac{k \hbar j}{m}}{i!_n q + \frac{k \hbar j}{m}} \right) \\
 & + \int_{\hbar j k_f}^{Z_{k_f}} dk k \ln \left(\frac{i!_n k_f + k}{i!_n q + \frac{k \hbar j}{m}} \right) g
 \end{aligned} \tag{124}$$

(C) $\hbar j > 2 k_f$

$$f_n(q) = \frac{v_q}{(2)^2} \left(\frac{m}{\hbar j}\right) \int_0^{Z_{k_f}} dk k \ln \left(\frac{i!_n q + \frac{k \hbar j}{m}}{i!_n q + \frac{k \hbar j}{m}} \right) \tag{125}$$

(A)

$$\begin{aligned}
 f_n(q) = & \frac{v_q}{(2)^2} \left(\frac{m}{\hbar j}\right) \left[\frac{1}{2} k_f^2 \ln(i!_n q + \frac{k_f \hbar j}{m}) + \frac{1}{2} (k_f \hbar j)^2 \ln(i!_n + q + \frac{k_f \hbar j}{m}) \right. \\
 & + \left(\frac{m^2}{2 q^2}\right) f \frac{1}{2} (i!_n q + \frac{k_f \hbar j}{m})^2 - \frac{1}{2} (i!_n + q + \frac{k_f \hbar j}{m})^2 \\
 & + (i!_n q)^2 \ln \left(\frac{i!_n q + \frac{k_f \hbar j}{m}}{i!_n + q + \frac{k_f \hbar j}{m}} \right) \\
 & + \left(\frac{2 \hbar j^2}{m}\right) (i!_n q) g \\
 & \left. + m (i!_n) \ln(i!_n) - m i!_n - m \left(q + \frac{k_f \hbar j}{m} + i!_n \right) \ln \left(q + \frac{k_f \hbar j}{m} + i!_n \right) + m \left(q + \frac{k_f \hbar j}{m} + i!_n \right) \right]
 \end{aligned} \tag{126}$$

(B)

$$\begin{aligned}
 f_n(q) = & \frac{v_q}{(2)^2} \left(\frac{m}{\hbar j}\right) \left[\frac{1}{2} k_f^2 \ln(i!_n q + \frac{k_f \hbar j}{m}) + \frac{1}{2} (k_f \hbar j)^2 \ln(i!_n + q + \frac{k_f \hbar j}{m}) \right. \\
 & + \left(\frac{m^2}{2 q^2}\right) f \frac{1}{2} (i!_n q + \frac{k_f \hbar j}{m})^2 - \frac{1}{2} (i!_n + q + \frac{k_f \hbar j}{m})^2 \\
 & + (i!_n q)^2 \ln \left(\frac{i!_n q + \frac{k_f \hbar j}{m}}{i!_n + q + \frac{k_f \hbar j}{m}} \right) \\
 & + \left(\frac{2 \hbar j^2}{m}\right) (i!_n q) g \\
 & \left. + m (i!_n) \ln(i!_n) - m i!_n - m \left(q + \frac{k_f \hbar j}{m} + i!_n \right) \ln \left(q + \frac{k_f \hbar j}{m} + i!_n \right) + m \left(q + \frac{k_f \hbar j}{m} + i!_n \right) \right]
 \end{aligned} \tag{127}$$

(C)

$$f_n(q) = \frac{v_q}{(2)^2} \left(\frac{m}{\hbar^2} \right) \left[\frac{1}{2} k_f^2 \ln \left(\frac{i!_n}{i!_n} \frac{q + \frac{k_f \hbar^2}{m}}{\frac{k_f \hbar^2}{m}} \right) \right. \\ \left. + \frac{m^2}{2 \hbar^2} f \frac{1}{2} (i!_n \frac{q + \frac{k_f \hbar^2}{m}}{q})^2 - \frac{1}{2} (i!_n \frac{q + \frac{k_f \hbar^2}{m}}{q})^2 \right. \\ \left. + (i!_n \frac{q + \frac{k_f \hbar^2}{m}}{q})^2 \ln \left(\frac{i!_n}{i!_n} \frac{q + \frac{k_f \hbar^2}{m}}{\frac{k_f \hbar^2}{m}} \right) - \frac{4 k_f \hbar^2}{m} (i!_n \frac{q + \frac{k_f \hbar^2}{m}}{q}) g \right] \quad (128)$$

5 Fermions at Finite Temperature

It should be clear from the previous sections that fermions at finite temp. is a major headache. A straightforward generalisation is not working out for unknown reasons. In this section we shall not distinguish between N and N_i , things are complicated enough as it is! Take for example, the expectation value of the number density at finite temperature,

$$\langle n_k \rangle = n_F(k) + \sum_{q_1} \langle n_{k+q_1} \rangle \langle n_{k-q_1} \rangle \langle n_{k+q_1+q_2} \rangle \langle n_{k-q_1-q_2} \rangle \dots \quad (129)$$

Assume the $n_F(k)$ are all evaluated at zero temp. as in the previous sections. Now, taken at face value, one is obliged to compute the thermodynamic expectation values of the Bose occupation probabilities assuming the chemical potential for the Bosons is zero (this means that we are allowed to create and destroy any number of Bosons). Now such a calculation yields an infinite answer for $\langle n_k \rangle$ as the sum over all q_1 diverges (is proportional to the total number of fermions). This is the reason why a more nefarious approach may be necessary in dealing with fermions at finite temperature. For this the only guide is that all the finite temp. dynamical correlation functions involving the number conserving object $\langle n_{k+q_1} \rangle$ should be correctly reproduced. Hopefully the commutation rules involving these number conserving products are not damaged in the bargain. For this let us start with the simplest case namely, $\langle n_{k+q_1} \rangle$. We know what the answer should be, that is,

$$\langle n_{k+q_1} \rangle = n_F(k) \quad (130)$$

where,

$$n_F(k) = [\exp(\beta \epsilon_k) + 1]^{-1} \quad (131)$$

For this to happen we have to do the following,

- (1) Treat the bosons as before assuming that they are always at zero temperature and with zero chemical potential.
- (2) Assume that all finite temperature effects are lumped into the coefficients.
- (3) Fix the coefficients as before by demanding that the finite temp. dynamical moments of the number-conserving products come out right.

For the dynamical four-point function to come out right we must ensure that,

$$\langle n_k \rangle = n_F(k) \langle n_{k+q_1} \rangle \langle n_{k-q_1} \rangle \quad (132)$$

Unfortunately, the coefficient $\langle n_k \rangle$ is no longer either zero or one. This renders matters even more difficult. But the dynamical six-point function is still the same and has to be correctly given as in the zero-temp case.

$$I = \langle n_{k+q_1} \rangle \langle n_{k-q_1} \rangle \langle n_{k+q_1+q_2} \rangle \langle n_{k-q_1-q_2} \rangle \dots \\ = [(1 - n_F(k - q_1)) (1 - n_F(k + q_1)) n_F(k + q_1 + q_2) n_F(k - q_1 - q_2) \dots \\ (1 - n_F(k - q_1)) n_F(k + q_1 + q_2) n_F(k - q_1 - q_2) \dots] \quad (133)$$

In terms of the Bose fields we have,

$$I = \sum_{\mathbf{k}} \langle \mathbf{q} \rangle_{\mathbf{k}^0} \langle \mathbf{q}^0 \rangle_{\mathbf{k}; \mathbf{k}^0} \langle \mathbf{k}^0; \mathbf{q}^0 \rangle \quad (134)$$

Equating these two we arrive at a formula for the coefficient. It is somewhat different from the zero temp case. But we can remedy that by choosing,

$$\begin{aligned} \langle \mathbf{q}_1; \mathbf{q}_2 \rangle_{\mathbf{k}_1; \mathbf{k}_2} &= F(\mathbf{k}_1, \mathbf{q}_1) F(\mathbf{k}_2, \mathbf{q}_2) \left[\frac{1}{1 - n_F(\mathbf{k}_1 + \mathbf{q}_1 = 2)} \frac{1}{1 - n_F(\mathbf{k}_2 + \mathbf{q}_2 = 2)} \right. \\ &\quad \left. \frac{1}{n_F(\mathbf{k}_1 - \mathbf{q}_1 = 2) n_F(\mathbf{k}_2 - \mathbf{q}_2 = 2)} \right] \\ &\quad \left[\frac{1}{n_F(\mathbf{k}_1 - \mathbf{q}_1 = 2; \mathbf{k}_2 + \mathbf{q}_2 = 2)} \frac{1}{n_F(\mathbf{k}_1 - \mathbf{q}_1 = 2; \mathbf{k} + \mathbf{q} = 2)} \frac{1}{n_F(\mathbf{k} + \mathbf{q} = 2; \mathbf{k}_2 - \mathbf{q}_2 = 2)} \right] \end{aligned} \quad (135)$$

and,

$$F(\mathbf{x}) = 1 \text{ if } \mathbf{x} \neq 0 \text{ and } F(\mathbf{x}) = 0 \text{ for } \mathbf{x} = 0 \quad (136)$$

For any temperature above zero $F(\mathbf{x}) = 1$ always. But for exactly at zero temp, the original case is recovered. But more importantly, all the right sort of dynamical correlation functions are recovered if we choose in addition, the following formula for the kinetic energy operator,

$$K = \sum_{\mathbf{k}, \mathbf{q}} \langle \mathbf{q} \rangle_{\mathbf{k}} a_{\mathbf{k}}^y(\mathbf{q}) a_{\mathbf{k}}(\mathbf{q}) \quad (137)$$

and,

$$\langle \mathbf{k} \rangle_{\mathbf{q}} = F(\mathbf{k} - \mathbf{q}) \left(\frac{\mathbf{k} \cdot \mathbf{q}}{m} \right) \quad (138)$$

These choices ensure that all the finite temp. dynamical moments of $\sum_{\mathbf{k} + \mathbf{q} = 2} \langle \mathbf{q} \rangle_{\mathbf{k}} \langle \mathbf{q} = 2 \rangle$ are correctly recovered.

5.1 Demonstration of Emergence of Superconductivity

Consider two interactions, one the coulomb repulsion and the other electron-phonon interaction :

$$H_I = \sum_{\mathbf{q} \neq 0} \frac{v_{\mathbf{q}}}{2V} \sum_{\mathbf{k}, \mathbf{k}^0} [\langle \mathbf{k} \rangle_{\mathbf{q}} a_{\mathbf{k}}(\mathbf{q}) + \langle \mathbf{k} \rangle_{\mathbf{q}} a_{\mathbf{k}}^y(\mathbf{q})] [\langle \mathbf{k}^0 \rangle_{\mathbf{q}} a_{\mathbf{k}^0}(\mathbf{q}) + \langle \mathbf{k}^0 \rangle_{\mathbf{q}} a_{\mathbf{k}^0}^y(\mathbf{q})] \quad (139)$$

$$H_{e \text{ phon}} = \sum_{\mathbf{q} \neq 0} \frac{M_{\mathbf{q}}}{V^{\frac{1}{2}}} \sum_{\mathbf{k}} [\langle \mathbf{k} \rangle_{\mathbf{q}} a_{\mathbf{k}}(\mathbf{q}) + \langle \mathbf{k} \rangle_{\mathbf{q}} a_{\mathbf{k}}^y(\mathbf{q})] [b_{\mathbf{q}} + b_{\mathbf{q}}^y] \quad (140)$$

The displacement operators are,

$$P_{\mathbf{q}} = b_{\mathbf{q}} + b_{\mathbf{q}}^y \quad (141)$$

and,

$$X_{\mathbf{q}} = \left(\frac{i}{2} \right) (b_{\mathbf{q}} - b_{\mathbf{q}}^y) \quad (142)$$

The equations of motion for the Bose fields are,

$$\begin{aligned} i \frac{\partial}{\partial t} a_{\mathbf{k}}^t(\mathbf{q}) &= \langle \mathbf{k} \rangle_{\mathbf{q}} a_{\mathbf{k}}^t(\mathbf{q}) + \left(\frac{v_{\mathbf{q}}}{V} \right) \sum_{\mathbf{k}^0} [\langle \mathbf{k}^0 \rangle_{\mathbf{q}} a_{\mathbf{k}^0}^t(\mathbf{q}) + \langle \mathbf{k}^0 \rangle_{\mathbf{q}} a_{\mathbf{k}^0}^{yt}(\mathbf{q})] \\ &\quad + \frac{M_{\mathbf{q}}}{V^{\frac{1}{2}}} \sum_{\mathbf{k}} \langle \mathbf{k} \rangle_{\mathbf{q}} [b_{\mathbf{q}} + b_{\mathbf{q}}^y] \end{aligned} \quad (143)$$

$$i \frac{\partial}{\partial t} X_{\mathbf{q}} = \frac{i M_{\mathbf{q}}}{V^{\frac{1}{2}}} \sum_{\mathbf{k}^0} [\langle \mathbf{k}^0 \rangle_{\mathbf{q}} a_{\mathbf{k}^0}(\mathbf{q}) + \langle \mathbf{k}^0 \rangle_{\mathbf{q}} a_{\mathbf{k}^0}^y(\mathbf{q})] + \frac{i}{2} P_{\mathbf{q}} \quad (144)$$

$$i \frac{\partial}{\partial t} P_{\mathbf{q}} = \frac{2}{i} X_{\mathbf{q}} \quad (145)$$

$$i \frac{\partial}{\partial t} a_k^{yt}(\mathbf{q}) = \epsilon_k(\mathbf{q}) a_k^{yt}(\mathbf{q}) - \left(\frac{V_q}{V}\right) \sum_{k^0} [\epsilon_{k^0}(\mathbf{q}) a_{k^0}^t(\mathbf{q}) + \epsilon_{k^0}(\mathbf{q}) a_{k^0}^{yt}(\mathbf{q})]$$

$$\frac{M_q}{V^{\frac{1}{2}}} \epsilon_k(\mathbf{q}) [b_q + b_q^y] \quad (146)$$

$$\left(i \frac{\partial}{\partial t} - \epsilon_k(\mathbf{q})\right) \frac{i \hbar \Gamma a_k^t(\mathbf{q}) a_{k^0}^y(\mathbf{q}^0) i}{\hbar \Gamma 1 i} = \epsilon_{k; k^0, \mathbf{q}; \mathbf{q}^0}(\mathbf{t})$$

$$+ \left(\frac{V_q}{V}\right) \sum_{k^0} \epsilon_{k^0}(\mathbf{q}) \left[\epsilon_{k^0}(\mathbf{q}) \frac{i \hbar \Gamma a_{k^0}^t(\mathbf{q}) a_{k^0}^y(\mathbf{q}^0) i}{\hbar \Gamma 1 i} + \epsilon_{k^0}(\mathbf{q}) \frac{i \hbar \Gamma a_{k^0}^{yt}(\mathbf{q}) a_{k^0}^y(\mathbf{q}^0) i}{\hbar \Gamma 1 i} \right]$$

$$+ \frac{M_q}{V^{\frac{1}{2}}} \epsilon_k(\mathbf{q}) \frac{i \hbar \Gamma P_q^t a_{k^0}^y(\mathbf{q}^0) i}{\hbar \Gamma 1 i} \quad (147)$$

$$\left(i \frac{\partial}{\partial t} + \epsilon_k(\mathbf{q})\right) \frac{i \hbar \Gamma a_k^{yt}(\mathbf{q}) a_{k^0}^y(\mathbf{q}^0) i}{\hbar \Gamma 1 i} =$$

$$\left(\frac{V_q}{V}\right) \sum_{k^0} \epsilon_{k^0}(\mathbf{q}) \left[\epsilon_{k^0}(\mathbf{q}) \frac{i \hbar \Gamma a_{k^0}^t(\mathbf{q}) a_{k^0}^y(\mathbf{q}^0) i}{\hbar \Gamma 1 i} + \epsilon_{k^0}(\mathbf{q}) \frac{i \hbar \Gamma a_{k^0}^{yt}(\mathbf{q}) a_{k^0}^y(\mathbf{q}^0) i}{\hbar \Gamma 1 i} \right]$$

$$\frac{M_q}{V^{\frac{1}{2}}} \epsilon_k(\mathbf{q}) \frac{i \hbar \Gamma P_q^t a_{k^0}^y(\mathbf{q}^0) i}{\hbar \Gamma 1 i} \quad (148)$$

The corresponding propagators with phonons are,

$$i \frac{\partial}{\partial t} \frac{i \hbar \Gamma P_q^t a_{k^0}^y(\mathbf{q}^0) i}{\hbar \Gamma 1 i} = \left(\frac{2}{i} \frac{q}{i}\right) \frac{i \hbar \Gamma X_q^t a_{k^0}^y(\mathbf{q}^0) i}{\hbar \Gamma 1 i} \quad (149)$$

$$i \frac{\partial}{\partial t} \frac{i \hbar \Gamma X_q^t a_{k^0}^y(\mathbf{q}^0) i}{\hbar \Gamma 1 i} = \left(\frac{i M_q}{V^{\frac{1}{2}}}\right) \sum_{k^0} \left[\epsilon_{k^0}(\mathbf{q}) \frac{i \hbar \Gamma a_{k^0}^t(\mathbf{q}) a_{k^0}^y(\mathbf{q}^0) i}{\hbar \Gamma 1 i} + \epsilon_{k^0}(\mathbf{q}) \frac{i \hbar \Gamma a_{k^0}^{yt}(\mathbf{q}) a_{k^0}^y(\mathbf{q}^0) i}{\hbar \Gamma 1 i} \right]$$

$$+ \left(\frac{i}{2} \frac{q}{i}\right) \frac{i \hbar \Gamma P_q^t a_{k^0}^y(\mathbf{q}^0) i}{\hbar \Gamma 1 i} \quad (150)$$

Transform to frequency domain and we have the corresponding equations :

$$G_1(\mathbf{q}; k^0; \mathbf{q}^0; n) = \epsilon_{k^0}(\mathbf{q}) \frac{q; \mathbf{q}^0}{i (i!_n - \epsilon_{k^0}(\mathbf{q}))}$$

$$+ f_n(\mathbf{q}) [G_1(\mathbf{q}; k^0; \mathbf{q}^0; n) + G_2(\mathbf{q}; k^0; \mathbf{q}^0; n)] + g_n(\mathbf{q}) G_3(\mathbf{q}; k^0; \mathbf{q}^0; n) \quad (151)$$

$$G_2(\mathbf{q}; k^0; \mathbf{q}^0; n) = f_n(\mathbf{q}) [G_1(\mathbf{q}; k^0; \mathbf{q}^0; n) + G_2(\mathbf{q}; k^0; \mathbf{q}^0; n)] + g_n(\mathbf{q}) G_3(\mathbf{q}; k^0; \mathbf{q}^0; n) \quad (152)$$

$$i!_n G_3(\mathbf{q}; k^0; \mathbf{q}^0; n) = \left(\frac{2}{i} \frac{q}{i}\right) G_4(\mathbf{q}; k^0; \mathbf{q}^0; n) \quad (153)$$

$$i!_n G_4(\mathbf{q}; k^0; \mathbf{q}^0; n) = i [G_1(\mathbf{q}; k^0; \mathbf{q}^0; n) + G_2(\mathbf{q}; k^0; \mathbf{q}^0; n)] + \left(\frac{i}{2} \frac{q}{i}\right) G_3(\mathbf{q}; k^0; \mathbf{q}^0; n) \quad (154)$$

The solutions may be written down as,

$$G_3(\mathbf{q}; k^0; \mathbf{q}^0; n) = \left(-\frac{2}{\frac{2}{i} + i!_n^2}\right) [G_1(\mathbf{q}; k^0; \mathbf{q}^0; n) + G_2(\mathbf{q}; k^0; \mathbf{q}^0; n)] \quad (155)$$

$$G_2(q; k^0; q^0; n) = \left(\frac{n(q)}{1 - n(q)} \right) G_1(q; k^0; q^0; n) \quad (156)$$

$$n(q) = f_n(q) - \frac{2}{q} \frac{q}{2 + n^2} g_n(q) \quad (157)$$

and finally,

$$G_1(q; k^0; q^0; n) = \frac{F_n(q) k^0(q)}{q q^0 \frac{1}{i} (i!_n!_k(q))} \quad (158)$$

and,

$$F_n(q) = \left[\frac{f_n(q)}{1 - n(q)} + \left(\frac{2}{q} \frac{q}{2 + n^2} \right) \left(\frac{g_n(q)}{1 - n(q)} \right) \right]^{-1} \quad (159)$$

here,

$$f_n(q) = \left(\frac{v_q}{V} \right)^X \frac{(k(q))^2}{(i!_n!_k(q))} \quad (160)$$

$$g_n(q) = \left(\frac{M_{qj}}{V} \right)^X \frac{(k(q))^2}{(i!_n!_k(q))} \quad (161)$$

For the four-point function we have to proceed as follows, the quantity of interest is (with phonons and coulomb),

$$\frac{1}{i^2} h\Gamma(a_{k_1}^{t_1}(q_1) a_{k_2}^{t_2}(q_2) a_{k_2^0}^{yt_2^0}(q_2^0) a_{k_1^0}^{yt_1^0}(q_1^0)) i$$

for this we as usual decompose as follows,

$$\begin{aligned} & \frac{1}{i^2} h\Gamma(a_{k_1}^{t_1}(q_1) a_{k_2}^{t_2}(q_2) a_{k_2^0}^{yt_2^0}(q_2^0) a_{k_1^0}^{yt_1^0}(q_1^0)) i \\ &= \sum_n \exp(i n t_1) \frac{1}{i^2} h\Gamma(a_{k_1}^n(q_1) a_{k_2}^{t_2}(q_2) a_{k_2^0}^{yt_2^0}(q_2^0) a_{k_1^0}^{yt_1^0}(q_1^0)) i \end{aligned} \quad (162)$$

$$(i!_n!_{k_1}(q_1)) \frac{1}{i^2} h\Gamma(a_{k_1}^n(q_1) a_{k_2}^{t_2}(q_2) a_{k_2^0}^{yt_2^0}(q_2^0) a_{k_1^0}^{yt_1^0}(q_1^0)) i$$

$$= F_1(k_1 q_1 n; k_2 q_2 t_2; k_2^0 q_2^0 t_2^0; k_1^0 q_1^0 t_1^0)$$

$$+ \left(\frac{k_1(q_1)}{V} \right) [v_{q_1} - \left(\frac{2}{n^2 + q_1^2} \right) M_{q_1 j}]$$

$$F_1(q_1 n; k_2 q_2 t_2; k_2^0 q_2^0 t_2^0; k_1^0 q_1^0 t_1^0) + F_2(q_1 n; k_2 q_2 t_2; k_2^0 q_2^0 t_2^0; k_1^0 q_1^0 t_1^0) = (1 - n(q_1) - n(q_1)) \quad (163)$$

$$\begin{aligned} & F_1(q_1 n; k_2 q_2 t_2; k_2^0 q_2^0 t_2^0; k_1^0 q_1^0 t_1^0) \\ &= \sum_{k_1} \frac{k_1(q_1)}{(i!_n!_{k_1}(q_1))} F_1(k_1 q_1 n; k_2 q_2 t_2; k_2^0 q_2^0 t_2^0; k_1^0 q_1^0 t_1^0) \end{aligned} \quad (164)$$

$$\begin{aligned} & F_2(q_1 n; k_2 q_2 t_2; k_2^0 q_2^0 t_2^0; k_1^0 q_1^0 t_1^0) \\ &= \sum_{k_1} \frac{k_1(q_1)}{(i!_n + !_{k_1}(q_1))} F_2(k_1 q_1 n; k_2 q_2 t_2; k_2^0 q_2^0 t_2^0; k_1^0 q_1^0 t_1^0) \end{aligned} \quad (165)$$

5.2 Bosons at zero temperature

In this subsection we do what we did earlier except that here we are bosonizing the bosons. Let b_k and b_k^y be the Bose fields in question. Analogous to Fermi sea displacements we introduce Bose condensate displacements.

$$\begin{aligned}
 b_{k+q=2}^y b_{k-q=2} &= N \delta_{k=0; q=0} + \left(\frac{p}{N} \right) [\delta_{k+q=2; 0} d_k(q) + \delta_{k-q=2; 0} d_k^y(q)] \\
 &+ \sum_{q_1} T_1(k; q; q_1) d_{k+q=2-q_1=2}^y(q_1) d_{k-q_1=2}(q+q_1) \\
 &+ \sum_{q_1} T_2(k; q; q_1) d_{k-q=2+q_1=2}^y(q_1) d_{k+q_1=2}(q+q_1)
 \end{aligned} \quad (166)$$

To make all the dynamical moments of $b_{k+q=2}^y b_{k-q=2}$ come out right and the commutation rules amongst them also come out right provided we choose T_1 and T_2 such that,

$$\begin{aligned}
 &\left(\frac{p}{N} \right)^2 \delta_{k+q=2; 0} \delta_{k^0=0; q^0=2; 0} \sum_{q_1} T_1(k^0; q^0; q_1) d_k(q) d_{k^0+q^0=2-q_1=2}^y(q_1) d_{k^0-q_1=2}(q^0+q_1) d_{k^0}^y(q^0) i \\
 &+ \left(\frac{p}{N} \right)^2 \delta_{k+q=2; 0} \delta_{k^0=0; q^0=2; 0} \sum_{q_1} T_2(k^0; q^0; q_1) d_k(q) d_{k^0-q^0=2+q_1=2}^y(q_1) d_{k^0+q_1=2}(q^0+q_1) d_{k^0}^y(q^0) i \\
 &= N \delta_{k+q=2; 0} \delta_{k+q=2; k^0=0; q^0=2} \delta_{k-q=2; k^0+q^0=2} \delta_{k^0-q^0=2; k^0=0; q^0=2}
 \end{aligned} \quad (167)$$

This means,

$$T_1(q^0=2; q^0; q) = 1; q \notin 0; q^0 \notin 0 \quad (168)$$

$$T_2(q^0=2; q^0; q) = 0; q \notin 0; q^0 \notin 0 \quad (169)$$

in order for the kinetic energy operator to have the form,

$$K = \sum_k d_{(1=2)k}^y(k) d_{(1=2)k}(k) \quad (170)$$

we must have,

$$\delta_{k+q_1=2} T_1(k+q_1=2; 0; q_1) + \delta_{k-q_1=2} T_2(k-q_1=2; 0; q_1) = \delta_{k; q_1=2} \quad (171)$$

In order for,

$$\sum_k b_k^y b_k = N$$

we must have,

$$T_1(k+q_1=2; 0; q_1) + T_2(k-q_1=2; 0; q_1) = 0 \quad (172)$$

In order for the commutation rules amongst the $b_{k+q=2}^y b_{k-q=2}$ to come out right, we must have in addition to all the above relations,

$$T_2(q=2; q; q^0) = 0 \quad (173)$$

$$T_2(q=2; q; q+q^0) = 0 \quad (174)$$

Any choice of T_1 and T_2 consistent with the above relations should suffice. From the above relations (Eq. (171) and Eq. (172)), we have quite unambiguously,

$$T_1(k+q_1=2; 0; q_1) = \delta_{k; q_1=2} \quad (175)$$

$$T_2(k-q_1=2; 0; q_1) = \delta_{k; q_1=2} \quad (176)$$

$$T_1(q^0=2; q^0; q) = 1 \quad (177)$$

and,

$$T_1(k; q; q_1) = 0; \text{ otherwise} \quad (178)$$

$$T_2(k; q; q_1) = 0; \text{ otherwise} \quad (179)$$

This means that we may rewrite the formula for $b_{k+q=2}^y b_{k-q=2}$ as follows,

$$b_{k+q=2}^y b_{k-q=2} = N_{k;0-q;0} + \left(\frac{p}{N} \right) [k+q=2;0 d_k(q) + k-q=2;0 d_k^y(q)] \\ + d_{(1=2)k+q=2}^y (k+q=2) d_{(1=2)k-q=2} (k-q=2) \\ \sum_{q_1} d_{q_1=2}^y(q_1) d_{q_1=2}(q_1) \quad (180)$$

Consider an interaction of the type,

$$H_I = \left(\frac{0}{2} \right) \sum_{q \neq 0} \sum_{k;k^0} v_q [k+q=2;0 d_k(q) + k-q=2;0 d_k^y(q)] [k^0-q=2;0 d_{k^0}(q) + k^0+q=2;0 d_{k^0}^y(q)] \quad (181)$$

or,

$$H_I = \left(\frac{0}{2} \right) \sum_{q \neq 0} v_q [d_{q=2}(q) + d_{q=2}^y(q)] [d_{q=2}(q) + d_{q=2}^y(q)] \quad (182)$$

and the free case is given by,

$$H_0 = \sum_q q d_{(1=2)q}^y(q) d_{(1=2)q}(q) \quad (183)$$

The full hamiltonian may be diagonalised as follows,

$$H = \sum_q !_q f_q^y f_q \quad (184)$$

and,

$$f_q = \left(\frac{!_q + q + 0v_q}{2 !_q} \right)^{\frac{1}{2}} d_{q=2}(q) + \left(\frac{!_q + q + 0v_q}{2 !_q} \right)^{\frac{1}{2}} d_{q=2}^y(q) \quad (185)$$

$$f_q^y = \left(\frac{!_q + q + 0v_q}{2 !_q} \right)^{\frac{1}{2}} d_{q=2}(q) + \left(\frac{!_q + q + 0v_q}{2 !_q} \right)^{\frac{1}{2}} d_{q=2}^y(q) \quad (186)$$

$$d_{q=2}(q) = \left(\frac{!_q + q + 0v_q}{2 !_q} \right)^{\frac{1}{2}} f_q - \left(\frac{!_q + q + 0v_q}{2 !_q} \right)^{\frac{1}{2}} f_q^y \quad (187)$$

$$d_{q=2}^y(q) = \left(\frac{!_q + q + 0v_q}{2 !_q} \right)^{\frac{1}{2}} f_q^y - \left(\frac{!_q + q + 0v_q}{2 !_q} \right)^{\frac{1}{2}} f_q \quad (188)$$

$$!_q = \frac{q}{\frac{2}{q} + 2 + 0v_q} \quad (189)$$

From this we may deduce,

$$hd_{(1=2)q}^y(q) d_{(1=2)q}(q) i = \frac{!_q + q + 0v_q}{2 !_q} \quad (190)$$

From this it is possible to write down the filling fraction.

FILLING FRACTION

$$f_0 = N_0/N = 1 - \sum_{q=1}^N hd_{(1=2)q}^y(q) d_{(1=2)q}(q) i \quad (191)$$

or,

$$f_0 = N_0/N = 1 - \sum_{q_{min}}^{Z-1} d_q q^2 \left(\frac{!_q + q + 0v_q}{2 !_q} \right) \quad (192)$$

Here the lower limit is necessary to ensure that the state

$$d_{(1=2)q}(q) \neq i$$

has positive norm. and the value of q_{min} is given by,

$$!_{q_{min}} + q_{min} + 0v_{q_{min}} = 0 \quad (193)$$

First define,

$$A_q = \left(\frac{q! + q + 0V_q}{2q!} \right)^{\frac{1}{2}} \quad (194)$$

$$B_q = \left(\frac{q! + q + 0V_q}{2q!} \right)^{\frac{1}{2}} \quad (195)$$

The density operator is,

$$\rho_q(t) = \frac{1}{N} [\rho_{(1=2)q}(q)(t) + d_{(1=2)q}^y(q)(t)] + \sum_k d_{(1=2)k+q=2}^y(k+q=2)(t) d_{(1=2)k-q=2}(k-q=2)(t) \quad (196)$$

$$\rho_{(1=2)q}(q)(t) = A_q f_q e^{i!_q t} B_q f_q^y e^{i!_q t} \quad (197)$$

$$d_{(1=2)q}^y(q)(t) = A_q f_q^y e^{i!_q t} B_q f_q e^{i!_q t} \quad (198)$$

$$d_{(1=2)k-q=2}(k-q=2)(t) = A_{k-q=2} f_{k-q=2} e^{i!_{k-q=2} t} B_{k-q=2} f_{k+q=2}^y e^{i!_{k-q=2} t} \quad (199)$$

$$d_{(1=2)k+q=2}^y(k+q=2)(t) = A_{k+q=2} f_{k+q=2}^y e^{i!_{k+q=2} t} B_{k+q=2} f_{k-q=2} e^{i!_{k+q=2} t} \quad (200)$$

Now define,

$$\begin{aligned} S^>(q;t) &= \langle h_q(t) \rho_q(0) \rangle = \frac{1}{N} \langle h [\rho_{(1=2)q}(q)(t) + d_{(1=2)q}^y(q)(t)] \rho_{(1=2)q}(q)(0) + d_{(1=2)q}^y(q)(0) \rangle \\ &= \sum_{k,k^0} \langle h d_{(1=2)k+q=2}^y(k+q=2)(t) d_{(1=2)k-q=2}(k-q=2)(t) d_{(1=2)k^0+q=2}^y(k^0+q=2)(0) d_{(1=2)k^0-q=2}(k^0-q=2)(0) \rangle \\ &= N \left(\frac{q}{q!} \right) \exp(-i!_q t) \end{aligned}$$

$$\begin{aligned} &+ \sum_{k,k^0} \langle h B_{k+q=2} f_{k-q=2} e^{i!_{k+q=2} t} A_{k-q=2} f_{k-q=2} e^{i!_{k-q=2} t} A_{k^0+q=2} f_{k^0-q=2}^y B_{k^0+q=2} f_{k^0-q=2}^y \rangle \\ &+ \sum_{k,k^0} \langle h B_{k+q=2} f_{k-q=2} e^{i!_{k+q=2} t} B_{k-q=2} f_{k+q=2}^y e^{i!_{k-q=2} t} B_{k^0+q=2} f_{k^0+q=2} B_{k^0+q=2} f_{k^0-q=2}^y \rangle \end{aligned}$$

$$S^>(q;t) = N \left(\frac{q}{q!} \right) \exp(-i!_q t)$$

$$+ \sum_k \exp(-i!_{k+q=2} - i!_{k-q=2} t) [B_{k+q=2}^2 A_{k-q=2}^2 + B_{k+q=2} B_{k-q=2} A_{k-q=2} A_{k+q=2}]$$

$$S^<(q;t) = N \left(\frac{q}{q!} \right) \exp(i!_q t) + \sum_k \exp(i!_{k+q=2} + i!_{k-q=2} t) [B_{k-q=2}^2 A_{k+q=2}^2 + B_{k-q=2} B_{k+q=2} A_{k+q=2} A_{k-q=2}]$$

and,

$$S^>(q;t) = \langle h_q(t) \rho_q(0) \rangle$$

$$S^<(q;t) = \langle h_q(0) \rho_q(t) \rangle$$

From the above equations, it is easy to see that there is a coherent part corresponding to the Bogoliubov spectrum and an incoherent part which is due to correlations and is responsible (hopefully) for the roton minimum.

6 Acknowledgements

The author is deeply indebted to Prof. A. J. Leggett and Prof. A. H. Castro-Neto for very illuminating conversations and indeed implicitly hinting at the next step in the developments. They should have been the coauthors but are not due to their own choosing.

References

- [1] S. Tomonaga, *Prog. Theor. Phys.* 5, 544 (1950)
- [2] J. M. Luttinger, *J. Math. Phys.* 4, 1154 (1963)
- [3] D. C. Mattis and E. H. Lieb, *J. Math. Phys.* 6, 304 (1965)
- [4] A. Luther, *Phys. Rev. B* 19, 320 (1979)
- [5] F. D. M. Haldane, *J. Phys. C* 14, 2585 (1981); *Helv. Phys. Acta* 65, 152 (1992); *Perspectives in Many-Particle Physics*, Proc. of the Intl. School of Physics "Enrico Fermi" Course CXXI, Varenna, 1992, eds. R. Schrieffer and R. A. Broglia (North-Holland, New York, 1994).
- [6] A. H. Castro-Neto and E. Fradkin, *Phys. Rev. Lett.* 72 1393 (1994); *Phys. Rev. B* 49, 10 877 (1994); *Phys. Rev. B* 51, 4084 (1995)
- [7] A. Houghton and J. B. Marston, *Phys. Rev. B* 48, 7790 (1993); A. Houghton, H.-J. Kwon and J. B. Marston, *Phys. Rev. B* 50, 1351 (1994); A. Houghton, H.-J. Kwon, J. B. Marston and R. Shankar *J. Phys. C* 6, 4909 (1994)
- [8] P. Kopietz, J. Hermisson and K. Schonhammer, *Phys. Rev. B* 52 10 877 (1995)
- [9] J. Frohlich, R. Gotschmann and P. A. Marchetti, *J. Phys. A* 28, 1169 (1995); J. Frohlich and R. Gotschmann *Phys. Rev. B* 55 no. 11 (1997)
- [10] S. Sunakawa, S. Yamasaki and T. Kebukawa *Prog. Theor. Phys.* 38, 804 (1967); 41, 919 (1969); 44 565 (1970); 49 1802 (1973)
- [11] G. A. Goldin, R. Meniko and D. H. Sharp, *J. Math. Phys.* 21 (1980); R. Meniko and D. H. Sharp, *J. Math. Phys.* 18, 471 (1977); R. F. Dashen and D. H. Sharp, *Phys. Rev.* 165, 1857 (1968)
- [12] E. Feenberg, "Theory of Quantum Fluids"
- [13] A. Liguori and M. Mintchev, *Comm. Math. Phys.* 169, 635-52 (1995)
- [14] F. Lesage, V. Pasquier and D. Serban, *Nucl. Phys. B* 435 [FS] (1995) 585-603
- [15] M. Reed and B. Simon, "Methods of Modern Mathematical Physics" vols. I and II, Academic Press Inc. c1980
- [16] G. S. Setlur, "Density Phase Variable Approach for Nonrelativistic Fermions" UIUC preprint (1996)