

Finite-Size Studies on the $SO(5)$ Symmetry of the Hubbard Model

Stefan Meixner, Werner Hanke

Institut für Theoretische Physik, Am Hubland, D-97074 Würzburg, Federal Republic of Germany

Eugene Demler and Shou-Cheng Zhang

Department of Physics, Stanford University, Stanford, CA 94305

(November 13, 2021)

We present numerical evidence for the approximate $SO(5)$ symmetry of the Hubbard model on a 10 site cluster. Various dynamic correlation functions involving the π operators, the generators of the $SO(5)$ algebra, are studied using exact diagonalisation, and are found to possess sharp collective peaks. Our numerical results also lend support on the interpretation of the recent resonant neutron scattering peaks in the $YBCO$ superconductors in terms of the Goldstone modes of the spontaneously broken $SO(5)$ symmetry.

PACS numbers: 7.10.Bg, 71.55.Jv

Although the single band Hubbard model has been extensively studied in recent years in connection with high- T_c superconductivity, its low-energy content has so far eluded both analytical and numerical investigations. While being deceptively simple, this model may have many possible competing ground states, and it has proved to be very difficult to organize the low-energy degrees of freedom. Recently, a new analytic approach based on a symmetry principle has been suggested. It was noticed that the Hubbard model enjoys an approximate $SO(5)$ symmetry, unifying antiferromagnetism (AF) with d -wave superconductivity (SC) [1]. This symmetry principle gives a simple description of the transition from an AF ground state to a SC ground state as the chemical potential is varied, and it gives a unified treatment of the low-energy collective degrees of freedom of the Hubbard model.

Motivated by its potential importance to the high- T_c problem, we undertake a numerical finite size study to test this approximate symmetry in the single band Hubbard model. The central object of our study is the so-called π operator defined as follows,

$$\pi_{s/d}^\dagger = \sum_{\mathbf{k}} (\cos k_x \pm \cos k_y) c_{\mathbf{k}+\mathbf{Q},\uparrow}^\dagger c_{-\mathbf{k},\uparrow}^\dagger \quad \mathbf{Q} = (\pi, \pi) \quad (1)$$

This operator carries spin 1, charge 2 and total momentum (π, π) . Charge conjugation and spin lowering operation gives five other similar operators. Here the subscript $\alpha = s, d$ refers to the internal s -wave or d -wave symmetry of this composite operator. The π operators were first introduced by Demler and Zhang [2] to explain the resonant neutron scattering peaks in the $YBCO$ superconductors [3], and are constructed by following the analogy with the η operators considered by Yang [4]. More recently, it was found [1] that together with the total spin and charge operators, the six π operators form the generators of the $SO(5)$ algebra, and furthermore, they rotate AF and SC order parameters into each other. In reference [2] and [1], it was argued that the π_d operators are approximate eigen-operators of the Hubbard Hamiltonian. This implies that the $SO(5)$ symmetry is an approximate symmetry of the Hubbard model [1], and one can use it to constrain the form of the low-energy effective Hamiltonian.

In this work, we present exact numerical diagonalisation studies of the dynamic correlation functions involving the π operators and the AF and SC order parameters. We verify that the π operators are approximate eigen-operators of the Hubbard model, and show that the properties of the various mixed correlation functions are consistent with the anticipated pattern of the $SO(5)$ symmetry breaking. Our exact numerical calculation is performed on a $\sqrt{10} \times \sqrt{10}$ site Hubbard cluster, using standard exact diagonalisation (ED) methods based on the Lanczos algorithm [5]. Mixed dynamical correlation functions are calculated utilizing a technique previously used to obtain anomalous Green's functions [6].

As a check to our numerical algorithm and the general framework, we first performed our calculations on the correlation functions involving the η operator [4], its mixed correlation functions with the charge-density-wave (CDW) order parameter [7] and the dynamic auto-correlation function of the CDW order parameter for the negative U Hubbard model. The relationship between the η and the π operators is explained in reference [1]. Since the η operator is a exact eigen-operator and the negative U Hubbard model is known to have a s -wave SC ground state away from half-filling, various exact results can be derived for these correlation functions [7]. Our numerical results are consistent with these exact results. However, due to the space limitations, we will not present these calculations here.

Let us first consider the auto-correlation function of the π operators, defined as follows,

$$\pi_\alpha^+(\omega) = -\frac{1}{\pi} \Im \langle \Psi_0^N | \pi_\alpha \frac{1}{\omega - H + E_0^{N+2} + i\eta} \pi_\alpha^\dagger | \Psi_0^N \rangle. \quad (2)$$

Here H is the standard Hubbard Hamiltonian, $|\Psi_0^N\rangle$ its ground state with N electrons and E_0^N the corresponding ground state energy. \Im takes the imaginary part of a correlation function. Throughout this paper, we measure the energy of a spectral function from the ground state energy of the intermediate state. Since $\pi_{s/d}^\dagger$ changes particle number by two, the natural energy scale of the intermediate states is the ground state energy of a $N + 2$ electron system.

Figures 1(a), 1(b), 2(a) and 2(b) plot the $\pi_d^+(\omega)$ correla-

tion function for $U = 4t$ and $U = 8t$, with electron densities $\langle n \rangle = 0.6$ and 0.8 respectively. We see that these dynamic correlation functions are dominated by a sharp peak, well separated from a higher energy continuum. If π_d^\dagger was an exact eigen-operator of the Hubbard Hamiltonian, its dynamical auto-correlation function should consist of a single peak. Here we see that it is only an approximate eigen-operator in the sense that there is also a high-energy continuum in addition to the resonance peak. The ratio of the spectral weight under the peak to the total spectral weight is found to be 0.63, 0.68, 0.46 and 0.31 in figs 1(a) to 2(b). The separation between the peak and the continuum, and the large relative spectral weight of the peak demonstrate that the π_d^\dagger operator is an eigen-operator of the Hubbard model to a good degree of approximation. The energies of these peaks are $1.06t$, $0.66t$, $0.55t$ and $0.24t$, respectively, and are low compared to U . This can easily be understood from the fact that the π operators are spin triplet operators, therefore, the bare U interaction should not enter its energy scale.

In reference [2], the dynamic auto-correlation function of the π_d operators was calculated within the t -matrix approximation in the particle-particle channel. Within this approximation, the dynamic auto-correlation function consists of a single peak, with a spectral weight proportional to $1 - n$, *i.e.* the hole density, and an energy of $J(1 - n)/2 - 2\mu$, where $J \approx 4t^2/U$ is the AF exchange constant and μ is the chemical potential measured from half-filling. Strictly speaking, the t -matrix approximation should be valid only in the low electron density limit. Here we see that the qualitative features of the t -matrix calculation are still valid near half-filling. Our exact numerical calculations show that the $\pi_d^+(\omega)$ correlation function is indeed dominated by a single peak whose spectral weight is proportional to the hole density $1 - n$. Furthermore, the energy of the π resonance peak is also proportional to the hole density, with a scale comparable to J .

The π resonance is a composite particle made out of two electrons. Near half-filling, it is very difficult for a single electron or hole to propagate coherently. In fact it is not clear if there is a quasi-particle peak in the one particle Green's function at all. In view of this fact, it is rather surprising that a two electron Green's function $\pi_d^+(\omega)$ has a coherent peak while the individual electrons propagate incoherently. To demonstrate that the π peak is a genuine collective behavior, we plot in figures 1(c) and 2(c) the bubble approximation to the $\pi_d^+(\omega)$ correlation function. This approximation to $\pi_d^+(\omega)$ consists of a particle-particle bubble with fully dressed one particle Green's function of the $\sqrt{10} \times \sqrt{10}$ site cluster inside the bubble. It fully takes into account the single-particle dressing effects, only the vertex correction is neglected. From these figures, we see that the bubble approximation yields a broad spectral distribution, without any identifiable resonance peak. The height of the broad spectral distribution is one order of magnitude less than the height of the π peak found in the full calculation. *This calculation demonstrates unambiguously that the collective behavior, or in other words, the vertex correction, is responsible for the existence of the π resonance.*




FIG. 1. Dynamic correlation functions of the $\sqrt{10} \times \sqrt{10}$ Hubbard model with $U = 4t$: (a) $\pi_d^+(\omega)$ spectrum at $\langle n \rangle = 0.6$, (b) $\pi_d^+(\omega)$ spectrum at $\langle n \rangle = 0.8$, (c) bubble approximation to $\pi_d^+(\omega)$ as shown in (b), (d) $\pi_s^+(\omega)$ at $\langle n \rangle = 0.8$, (e) off-diagonal correlation function $S_d(\omega)$ at $\langle n \rangle = (0.8/1.0)$, as defined in the text, (f) spin correlation function $\chi_Q(\omega)$ and (g) bubble approximation $\bar{\chi}_Q(\omega)$.

Figures 1(d) and 2(d) give results for $\pi_s^+(\omega)$ at $U = 4t$ and $U = 8t$, respectively, at density $n = 0.8$. For $U = 8t$, the spectrum is broadly distributed and no resonance peak can be identified. For $U = 4t$, although there is an isolated spectral feature, it is at a much higher energy ($\approx 2t$) compared to the $\pi_d^+(\omega)$ counterpart, and the height of the peak is almost one order of magnitude less. We interpret these results as evidence that in contrast to π_d^\dagger , π_s^\dagger is not an approximate eigen-operator of the Hubbard model near half-filling. This result is consistent with the conclusion of the t -matrix calculation [2]. The crucial difference between the $\pi_d^+(\omega)$ and $\pi_s^+(\omega)$ correlation function has recently been used by Zhang [1] to argue that *there is an approximate symmetry between AF and d-wave SC, but no symmetry between AF and extended s-wave SC near half-filling.*

Originally, the π operators were introduced [2] in connection with the resonant neutron scattering peaks recently found

in the *YBCO* superconductors [3]. Below the superconducting transition temperature T_c , a resonance peak appears in the polarized neutron scattering amplitude with momentum transfer (π, π) , and a resonance energy of 25meV , 33meV and 41meV , depending on doping. Demler and Zhang [2] identified this experimental feature with the π resonance, and showed that the particle-particle π resonance can be mixed into the particle-hole channel below T_c , and therefore appear in the neutron scattering cross-section. A number of other theoretical papers [8] explain the experimental feature in terms of a particle-hole threshold behavior near the superconducting gap or possible excitonic states inside the gap. More recently, the π resonance has been interpreted more broadly by Zhang [1] as the Goldstone boson associated with the spontaneous breaking of the $SO(5)$ symmetry below T_c .




FIG. 2. Same as in Fig. 1, but here for $U = 8t$.

In this work, we test the ideas of references [2] and [1] by studying the mixed correlation functions involving the spin-density-wave order parameter

$$S_{\mathbf{Q}}^+ = \sum_{\mathbf{k}} c_{\mathbf{k}+\mathbf{Q},\uparrow}^\dagger c_{\mathbf{k},\downarrow} \quad S_{\mathbf{Q}}^- = S_{\mathbf{Q}}^{+\dagger} \quad (3)$$

and the π_d operator, defined by

$$S_d(\omega) = -\frac{1}{\pi} \Im \langle \Psi_0^{N-2} | (\pi_d \frac{1}{\omega - (H - E_0^N) + i\eta} S_{\mathbf{Q}}^+ - S_{\mathbf{Q}}^+ \frac{1}{\omega - (H - E_0^{N-2}) + i\eta} \pi_d) | \Psi_0^N \rangle \quad (4)$$

and the dynamical spin correlation function itself, defined by

$$\chi_{\mathbf{Q}}^+(\omega) = -\frac{1}{\pi} \Im \langle \Psi_0^N | S_{\mathbf{Q}}^- \frac{1}{\omega - (H - E_0^N) + i\eta} S_{\mathbf{Q}}^+ | \Psi_0^N \rangle \quad (5)$$

The correlation function $S_d(\omega)$ satisfies an important exact sum rule

$$\int_{-\infty}^{\infty} d\omega S_d(\omega) = 2 \langle \Psi_0^{N-2} | \Delta_d | \Psi_0^N \rangle \quad (6)$$

where $\Delta_d = \sum_{\mathbf{k}} (\cos k_x - \cos k_y) c_{\mathbf{k},\uparrow} c_{-\mathbf{k},\downarrow}$ is the d -wave superconducting order parameter. This sum rule follows from the fact that the π_d operator is a $SO(5)$ symmetry generator which rotates the AF order parameter into the d -wave SC order parameter [1]. Mixed correlation functions like $S_d(\omega)$, which involve a symmetry generator and an order parameter, are commonly used to prove the Goldstone theorem on the Goldstone bosons associated with spontaneous symmetry breaking [9].

In figures 1(e) and 2(e), $S_d(\omega)$ is plotted for $U = 4t$ and $U = 8t$, for electron density between $n = 0.8$ and $n = 1$. We see that the spectrum is dominated by a single peak with relative weight 0.77 and 0.72 respectively, located at the same energy as the π resonance. We also verified that the sum rule (6) is satisfied by explicitly computing the d -wave order parameter defined on the right hand side. Therefore, our result shows that the 10 site Hubbard cluster at densities between $n = 0.8$ and $n = 1$ possesses considerable d -wave SC fluctuations, and furthermore, there is indeed a Goldstone pole associated with this spontaneous symmetry breaking, consistent with the $SO(5)$ theory [1]. The finite energy of the Goldstone pole results from the fact that the $SO(5)$ is not only broken spontaneously by the SC order parameter, but also broken explicitly by the chemical potential [1]. (Sometimes, a Goldstone boson associated with a symmetry which is broken both explicitly and spontaneously is called a pseudo Goldstone boson. The pion in particle physics is such an example.) We also calculated $S_d(\omega)$ between the densities $n = 0.6$ and $n = 0.8$ and found it to be nearly zero. Similarly, the mixed correlation function $S_s(\omega)$ involving the π_s and $S_{\mathbf{Q}}^+$ operators vanishes for all densities. This result is consistent with other numerical evidence for d -wave pairing fluctuations in the Hubbard model [10].

Finally we show our calculation for the spin correlation function $\chi_{\mathbf{Q}}^+(\omega)$ in figures 1(f) and 2(f), for $U = 4t$ and $U = 8t$ respectively at density $n = 1$. We see that there is a sharp resonance feature at the same energy as the π resonance. The fact that all three correlation functions $\pi_d^+(\omega)$, $S_d(\omega)$ and $\chi_{\mathbf{Q}}^+(\omega)$ have resonance peaks at the same energy

clearly demonstrates that the peak in the spin correlation function $\chi_{\mathbf{Q}}^+(\omega)$ is due to a particle-particle intermediate state $\pi_d^\dagger|\Psi_0^{N-2}\rangle$. Because of the finiteness of the mixed correlation function $S_d(\omega)$ and the right-hand-side of equation (6), a particle-particle excitation at density $n = 0.8$ makes a finite contribution to the spin correlation function at $n = 1$. This feature confirms the argument of reference [2], which explains the experimental resonant neutron scattering cross section in terms of the collective particle-particle π resonance.

To exhibit the collective nature of the resonance peak in $\chi_{\mathbf{Q}}^+(\omega)$, and to contrast it with single-particle behavior, we plot in figures 1(g) and 2(g) the bubble approximation to $\chi_{\mathbf{Q}}^+(\omega)$ on the $\sqrt{10} \times \sqrt{10}$ site cluster. In contrast to the full spin correlation function, the bubble approximation does not possess any low-energy collective poles. These results show that it is necessary to include the vertex corrections and the mixing into the particle-particle channels to calculate $\chi_{\mathbf{Q}}^+(\omega)$, in order to exhibit its resonance peak. These calculations have been performed recently and will be published elsewhere [11].

In conclusion, we have numerically verified one of the most fundamental assumptions of the $SO(5)$ theory of the Hubbard model [1], which asserts that the $SO(5)$ symmetry generators, the π_d operators, are approximate eigen-operators of the Hubbard model both near and away from half-filling. Their dynamic auto-correlation function shows well separated resonance peaks with large relative spectral weight and low-energy. This behavior is intrinsically collective, and can not be reproduced by any calculations which neglects vertex corrections. In contrast to the π_d operators, the π_s operators do not show well separated peaks near half-filling. This result shows that there is only an approximate $SO(5)$ symmetry between the AF and the d -wave SC order parameters, no symmetry between the AF and the s -wave SC order parameters. Close to half-filling, between the densities $n = 0.8$ and $n = 1$, the 10 site Hubbard cluster has finite d -wave superconducting fluctuations, which lead to a non-vanishing mixed correlation function involving the $SO(5)$ symmetry generator and the AF order parameter. Because of the finiteness of this mixed correlation function, there is a contribution to the dynamic spin correlation function from the particle-particle π resonance. These observations are consistent with the proposed explanation of the resonant neutron scattering peaks in the $YBCO$ superconductors [2].

We would like to acknowledge useful discussions with Prof. D. Scalapino and P. Hedegard who urged us to contrast the collective behavior of the π resonance with the single-particle bubble approximation. This work is supported by FOR-SUPRA II, BMBF (05 605 WWA 6), ERB CHRXCT940438 and in part by the NSF under grant numbers DMR-9400372 and DMR-9522915.

- [1] S.C. Zhang, *Science*, to be published.
- [2] E. Demler and S.C. Zhang, *Phys. Rev. Lett.*, 75:4126, 1995.
- [3] J. Rossat-Mignod et al. *Physica C*, 185:86, 1991; H. Mook et al. *Phys. Rev. Lett.*, 70:3490, 1994; H.F. Fong et al. *Phys. Rev. Lett.*, 75:316, 1995; H.F. Fong et al. *Phys. Rev. Lett.*, 78:713, 1997; P. Dai et al. *Phys. Rev. Lett.*, 77:5425, 1996.
- [4] C.N. Yang, *Phys. Rev. Lett.*, 63:2144, 1989; C.N. Yang and S.C. Zhang, *Mod. Phys. Lett. B*, 4:759, 1990.
- [5] "Exact diagonalisation methods for models of strongly correlated fermions", D. Poilblanc in "Numerical methods for strongly correlated systems", Ed. D. Scalapino, Frontiers in Physics (1997).
- [6] Y. Otha, T. Shimoizato, R. Eder and S. Maekawa, *Phys. Rev. Lett.*, 73:324, 1994;
- [7] S. C. Zhang, *Phys. Rev. Lett.*, 65:120, 1990; S. C. Zhang, *Int. J. Mod. Phys. B*, 5:153, 1991; E. Demler, S.C. Zhang, N. Bulut, and D. Scalapino, *Int. J. Mod. Phys. B*, 10:2137, 1996.
- [8] N. Bulut and D. Scalapino, *Phys. Rev. B*, 53:5149, 1996; H.F. Fong et al. *Phys. Rev. Lett.*, 75:316, 1995; I.I. Mazin and V.M. Yakovenko, *Phys. Rev. Lett.*, 75:4134, 1995; D.Z. Liu et al. *Phys. Rev. Lett.*, 75:4130, 1995; A. Millis and H. Monien, On the bilayer coupling in the yttrium-barium family of high temperature superconductors. preprint; L. Yin, S. Chakravarty, P. W. Anderson, The Neutron Peak in the Interlayer Tunneling Model of High Temperature Superconductors. preprint, cond-mat/9606139.
- [9] J. Goldstone, A. Salam, and S. Weinberg, *Phys. Rev.*, 127:965, 1962.
- [10] D. Scalapino *Phys. Reports*, 250:329, 1995 and references found therein; E. Dagotto *Rev. Mod. Phys.*, 66:763, 1994 and references found therein.
- [11] H. Kohno, B. Normand, and H. Fukuyama, *to be published*; E. Demler and S.C. Zhang, *to be published*; H. Kohno, B. Normand, and H. Fukuyama, *Proceedings of 10th Anniversary HTS Workshop*. World Scientific Publishing Company.