

Comment on “Duality relations for Potts correlation functions”

Jesper Lykke Jacobsen^{1,2}

¹*Institute of Physics and Astronomy, University of Aarhus, DK-8000 Aarhus C, Denmark*

²*Somerville College and Department of Theoretical Physics, University of Oxford, 1 Keble Road, Oxford OX1 3NP, United Kingdom*

(March 23, 2022)

Abstract

In a recent paper by Wu [1] the three-point correlation of the q -state Potts model on a planar graph was related to ratios of dual partition functions under fixed boundary conditions. It was claimed that the method employed could straightforwardly be applied to higher correlations as well; this is however not true. By explicitly considering the four-point correlation we demonstrate how the appearance of non-well-nested connectivities invalidates the method.

Consider the q -state Potts model [2] on a two-dimensional planar graph L having a free boundary. In accordance with Fig. 2 in Ref. [1] we let i, j, k and l be four sites on the boundary of L , following one another in a clockwise fashion, and we define exterior dual spins s_1, s_2, s_3 and s_4 so that all boundary spins between sites l and i of L interact with a spin state s_1 , boundary spins between i and j interact with s_2 , spins between j and k with s_3 , and finally spins between k and l with s_4 . The partition function with the four Potts spins $\sigma_i, \sigma_j, \sigma_k$ and σ_l fixed in definite states is called $Z_{\sigma_i\sigma_j\sigma_k\sigma_l}$, and similarly the dual partition function for fixed exterior dual spins is denoted by $Z_{s_1s_2s_3s_4}^*$.

Up to the q -fold permutation symmetry of the Potts spin labels there exist 15 different boundary conditions for $Z_{\sigma_i\sigma_j\sigma_k\sigma_l}$ out of which we can form five combinations

$$\begin{aligned}
Z_4 &= Z_{1111}, \\
Z_3 &= Z_{2111} + Z_{1211} + Z_{1121} + Z_{1112}, \\
Z_{2p} &= Z_{2211} + Z_{2121} + Z_{2112}, \\
Z_p &= Z_{1123} + Z_{1213} + Z_{1231} + Z_{2113} + Z_{2131} + Z_{2311}, \\
Z_0 &= Z_{1234},
\end{aligned} \tag{1}$$

which are symmetric under permutations of the four sites i, j, k and l . We introduce them here in order to simplify the notation in subsequent equations.

Following Wu's strategy [1] we should extract equations relating the $Z_{\sigma_i \sigma_j \sigma_k \sigma_l}$ and the $Z_{s_1 s_2 s_3 s_4}^*$ by running through all possible ways of connecting the sites i, j, k and l by auxiliary bonds and using the fundamental duality relation, Wu's Eq. (3). Each time a bond is added one of the exterior dual spins s is separated from its neighbours and thus allowed to take a different value.

First, using the 'empty' connection (i.e., introducing no auxiliary bonds) we find the relation

$$Z_4 + (q-1)[Z_3 + Z_{2p}] + (q-1)(q-2)Z_p + (q-1)(q-2)(q-3)Z_0 = qCZ_4^*, \tag{2}$$

corresponding to Wu's Eq. (19).

Next, consider adding a bond between sites k and l . Summing over the two 'free' sites, i and j , we obtain a reduction to the two-point case:

$$\begin{aligned}
Z_{11} &= \sum_{i=1}^q \sum_{j=1}^q Z_{ij11} \\
&= Z_{1111} + (q-1)[Z_{2111} + Z_{1211} + Z_{2211}] + (q-1)(q-2)Z_{2311}.
\end{aligned} \tag{3}$$

Among the exterior dual spins, s_4 has been separated from $s_1 = s_2 = s_3$, so that $Z_{11}^* = Z_{1111}^*$ and $Z_{12}^* = Z_{1112}^*$. Using the known result for Z_{11} , Wu's Eq. (12), along with the duality we find that

$$Z_{1111} + (q-1)[Z_{2111} + Z_{1211} + Z_{2211}] + (q-1)(q-2)Z_{2311} = C[Z_{1111}^* + (q-1)Z_{1112}^*]. \tag{4}$$

The equation is one out of a set of six obtained by connecting two of the sites i, j, k and l with a bond. Further equations can be found by considering Z_{12} instead of Z_{11} , but they can be shown to be linear combinations of Eq. (2) and the six equations just obtained.

Another set of four equations can be found by letting one of the sites i, j, k and l be ‘free’ and connecting the remaining three. This corresponds to a reduction to the three-point case. For example, adding a bond between sites j and k and another between k and l , we find that $Z_{111} = \sum_{i=1}^q Z_{i111} = Z_{1111} + (q-1)Z_{2111}$. The exterior dual spins now satisfy $s_2 = s_1$, and from Wu’s Eq. (25) for Z_{111} and the duality we obtain

$$Z_{1111} + (q-1)Z_{2111} = \frac{C}{q} \{Z_{1111}^* + (q-1)[Z_{2211}^* + Z_{1121}^* + Z_{1112}^*] + (q-1)(q-2)Z_{1123}^*\}. \quad (5)$$

Again, the equations found by reducing to Wu’s expressions for Z_{123} , Z_{211} , Z_{121} or Z_{112} do not contain any new information.

The remaining equations can be found by connecting all of the sites i, j, k and l with auxiliary bonds of strength K . One way of doing this corresponds to Wu’s Eq. (24) and implies connecting all the four sites to a common new point n outside the boundary of L . All of the exterior dual spins are now separated from one another. Summing over n we can express the partition function \tilde{Z} of this system through the $Z_{\sigma_i\sigma_j\sigma_k\sigma_l}$ and relate it by duality to its corresponding dual partition function \tilde{Z}^* . The result is

$$\begin{aligned} & Z_4(e^{4K} + q - 1) + (q - 1)Z_3(e^{3K} + e^K + q - 2) + (q - 1)Z_{2p}(2e^{2K} + q - 2) \\ & + (q - 1)(q - 2)Z_p(e^{2K} + 2e^K + q - 3) + (q - 1)(q - 2)(q - 3)Z_0(4e^K + q - 4) \\ & = \frac{C}{q^2}(e^K - 1)^4 \{ Z_4^*e^{4K^*} + (q - 1)[Z_3^* + Z_{2p}^* - Z_{2121}^*]e^{2K^*} + (q - 1)Z_{2121}^* \\ & + (q - 1)(q - 2)[Z_p^* - Z_{1213}^* - Z_{2131}^*]e^{K^*} + (q - 1)(q - 2)[Z_{1213}^* + Z_{2131}^*] \\ & + (q - 1)(q - 2)(q - 3)Z_0^* \}. \end{aligned} \quad (6)$$

The other way of connecting all four sites is to connect two of them to a new point m and the remaining two to another new point n . For instance, connecting each of i and j to m with a bond of strength K , and similarly k and l to n , we find that s_2 and s_4 are separated whilst $s_3 = s_1$. Summing over m and n as before we arrive at

$$\begin{aligned}
& \Sigma_{11}^2 \{Z_4 + (q-1)Z_{2211}\} + \Sigma_{11}\Sigma_{21} \{(q-1)Z_3 + (q-1)(q-2)[Z_{1123} + Z_{2311}]\} \\
& + \Sigma_{21}^2 \{(q-1)[Z_{2112} + Z_{2121}] + (q-1)(q-2)[Z_p - Z_{1123} - Z_{2311}] + (q-1)(q-2)(q-3)Z_0\} \\
& = \frac{C}{q}(e^K - 1)^4 \left\{ Z_4^* e^{4K^*} + (q-1)[Z_{1211}^* + Z_{1112}^*]e^{2K^*} + (q-1)Z_{2121}^* + (q-1)(q-2)Z_{1213}^* \right\},
\end{aligned} \tag{7}$$

where we have defined $\Sigma_{11} = e^{2K} + (q-1)$ and $\Sigma_{21} = 2e^K + (q-2)$. A similar equation can be obtained by connecting i and l to m , and j and k to n .

Exactly at this point we get into trouble. For the 15 unknown quantities appearing on the right-hand side of Eq. (1) we have obtained 14 independent equations, namely Eq. (2), six equations of the type (4), four equations of the type (5), Eq. (6), and two equations of the type (7). Clearly, what is missing is a third equation of the type (7) in which i and k are connected to m , and j and l are connected to n . But the auxiliary bonds making such a connection would necessarily intersect, thus violating the planarity of the graph L . In other words, it would be impossible to define the four exterior dual spins.

This problem has to do with the connectedness [3] of the four points. Generally, for an n -point correlation the number of partition functions $Z_{\sigma_1\sigma_2\dots\sigma_n}$ with fixed values of the n boundary spins equals the number of ways \tilde{c}_n in which n points can be interconnected, as is easily seen by interpreting $\sigma_i = \sigma_j$ as a connection between sites i and j . But the number of equations obtainable using the method of Wu is only equal to the number c_n of well-nested n -point connectivities [3], which is in general less than \tilde{c}_n . For $n = 1, 2, 3, 4, 5, \dots$ we have $\tilde{c}_n = 1, 2, 5, 15, 51, \dots$, whilst $c_n = 1, 2, 5, 14, 42, \dots$

In an attempt to obtain more equations one could imagine introducing connections between spins in the infinite face, thus converting any spin of valence four or higher into interconnected spins of valence three [4]. In the case of the four-point function this would mean discarding Eq. (6) and in addition to the two equations of type (7) considering two new equations obtained from these by connecting spins m and n with a fifth auxiliary bond. The first of these, corresponding to Eq. (7), looks like

$$\begin{aligned}
& Z_4(\Sigma_{11}^2 + \Sigma_4) + (q-1)Z_3(\Sigma_{11}\Sigma_{21} + \Sigma_3) + (q-1)Z_{2211}(\Sigma_{11}^2 + \Sigma_{2p}) \\
& + (q-1)(q-2)[Z_{1123} + Z_{2311}](\Sigma_{11}\Sigma_{21} + \Sigma_p) + (q-1)[Z_{2121} + Z_{2112}](\Sigma_{21}^2 + \Sigma_{2p})
\end{aligned}$$

$$\begin{aligned}
& + (q-1)(q-2)[Z_p - Z_{1123} - Z_{2311}](\Sigma_{21}^2 + \Sigma_p) + (q-1)(q-2)(q-3)Z_0(\Sigma_{21}^2 + \Sigma_0) \\
& = \frac{C}{q^3}(e^K - 1)^5 \left\{ Z_4^* e^{5K^*} + (q-1)[Z_{1211}^* + Z_{1112}^*] e^{3K^*} \right. \\
& + (q-1)[Z_{2111}^* + Z_{1121}^* + Z_{2211}^* + Z_{2112}^*] e^{2K^*} + (q-1)Z_{2121}^* e^{K^*} \\
& \left. + (q-1)(q-2)[Z_p^* - Z_{2131}^*] e^{K^*} + (q-1)(q-2)Z_{2131}^* + (q-1)(q-2)(q-3)Z_0^* \right\}, \quad (8)
\end{aligned}$$

where we have defined $\Sigma_4 = (e^K - 1)(e^{4K} + q - 1)$, $\Sigma_3 = (e^K - 1)(e^{3K} + e^K + q - 2)$, $\Sigma_{2p} = (e^K - 1)(2e^{2K} + q - 2)$, $\Sigma_p = (e^K - 1)(e^{2K} + 2e^K + q - 3)$, and $\Sigma_0 = (e^K - 1)(4e^K + q - 4)$. Now, subtracting the sum of Eqs. (8) from the sum of Eqs. (7) we regain, after some tedious algebra, $2(e^K - 1)$ times Eq. (6). But, on the other hand, subtracting the difference of Eqs. (8) from the difference of Eqs. (7) we find the result *zero*, i.e., out of the four equations where all the four spins are connected to auxiliary bonds there are still only three that are linearly independent.

Although the method thus fails to isolate the $Z_{\sigma_i \sigma_j \sigma_k \sigma_l}$, which would enable us to construct the generalised correlation function $P_4(\sigma, \sigma', \sigma'', \sigma''')$ as defined in Wu's Eq. (1), we could still hope to obtain the ordinary correlation function $\Gamma_4(\sigma_i, \sigma_j, \sigma_k, \sigma_l)$ (see Wu's Eq. (2)). Evidently, to do so it suffices to isolate $Z_4 \equiv Z_{1111}$, which should be possible by finding linear combinations of the above set of equations for which the left-hand sides are symmetric in the sense of Eq. (1).

Clearly, Eqs. (2) and (6) are already in this form. The sum of the six equations of type (4) yields

$$\begin{aligned}
6Z_4 + 3(q-1)Z_3 + 2(q-1)Z_{2p} + (q-1)(q-2)Z_p = \\
C \left\{ 6Z_4^* + (q-1)[Z_3^* + Z_{2p}^* - Z_{2121}^*] \right\}, \quad (9)
\end{aligned}$$

while the four equations of type (5) sum up to

$$\begin{aligned}
4Z_4 + (q-1)Z_3 = \\
\frac{C}{q} \left\{ 4Z_4^* + (q-1)(q-2)[Z_p^* - Z_{1213}^* - Z_{2131}^*] + (q-1)[2Z_3^* + 2(Z_{2p}^* - Z_{2121}^*)] \right\}. \quad (10)
\end{aligned}$$

As before, employing the two-point reduction Z_{12} or the three-point reductions Z_{123} , Z_{211} , Z_{121} and Z_{112} merely provides us with linear combinations of what we already know.

But again, the two equations of the type (7) lead us into trouble. When taking their sum it can easily be seen that the left-hand side cannot be written in terms of the symmetrical combinations (1). To symmetrise we would again need the non-well-nested equation, which however leaves the exterior dual spins undefined.

In conclusion we have shown that the method of Wu breaks down for the four-point correlation, and in fact for all higher correlations as well. This is reminiscent of the fact that in conformal field theory it is necessary to consider non-planar, as well as planar, monodromies in order that the four- and higher-point correlation functions be well determined [5]. The failure to relate the four-point function to generalised surface tensions is especially regrettable in so far as such a relation might prove useful in calculating the energy-energy correlation of the Potts model.

Useful discussions with J. L. Cardy and F. Y. Wu are gratefully acknowledged. This research was supported in part by the Engineering and Physical Sciences Research Council under Grant GR/J78327, and by the National Science Foundation under Grant PHY94-07194.

REFERENCES

- [1] F. Y. Wu, Phys. Lett. A **228**, 43-47 (1997).
- [2] R. B. Potts, Proc. Cambr. Philos. Soc. **48**, 106 (1952).
- [3] H. W. J. Blöte and M. P. Nightingale, Physica A **112**, 405-465 (1982).
- [4] F. Y. Wu, private communication.
- [5] Vl. S. Dotsenko and V. A. Fateev, Nucl. Phys. B **251**, 691 (1985).