

Tunneling Conductance and Coulomb Blockade Peak Splitting of Two Quantum Dots Connected by a Quantum Point Contact

Yu-Liang Liu

Max-Planck-Institut für Physik Komplexer Systeme, Bayreuther Str. 40, D-01187 Dresden,
Germany

Abstract

By using bosonization method and unitary transformation, we give a general relation between the dimensionless tunneling conductance and the fractional Coulomb blockade conductance peak splitting which is valid both for weak and strong transmission between two quantum dots, and show that the tunneling conductance has a linear temperature dependence in the low energy and low temperature limit.

73.20.Dx, 73.40.Gk

Electron tunneling in a mesoscopic structure may be drastically influenced by the charging effects. If the charge spreading is impeded by weak links, or by a special geometry of the structure, the charging suppresses the electron tunneling. Such a suppression of the electron tunneling is commonly called as the Coulomb blockade [1,2,3]. It has become possible to observe the Coulomb blockade effect in semiconductor heterostructures where the geometry of the system can be easily modified by adjusting the voltages on special gate electrodes. Recently, the electron tunneling through two quantum dots connected by single quantum point contact which is controlled by adjusting voltage on additional gate electrode has been extensively studied, both experimental [4,5,6,7,8] and theoretical [9,10,11,12,13]. The electron tunneling between two quantum dots leads to a decay of the Coulomb blockade of the individual dot. For a pair of electrostatically identical quantum dots, the progress of this decay can be described by tracking the splitting of the Coulomb blockade conductance peaks as they evolve from doubly degenerate single-dot conductance resonances to that of nondegenerate double-dot peaks with twice the original periodicity. The Coulomb blockade peak splitting significantly depends upon the transmission coefficient T of the quantum point contact, i.e., the tunneling conductance. For the case of $T = 1$, the Coulomb blockade between two quantum dots disappears, they become a double-quantum-dot. For the case of $T = 0$, the Coulomb blockade is maximum which completely suppresses the electron tunneling between two quantum dots, therefore, they are separated from each other.

In this paper, by using bosonization method and unitary transformation developed in Ref. [14], we can effectively treat the system of two electrostatically identical quantum dots connected by one quantum point contact which is controlled by adjusting voltage on additional gate electrode, and give a general relation between the dimensionless tunneling conductance g and the fractional Coulomb blockade peak splitting f , which is valid both for weak ($T \ll 0$) and strong ($T \ll 1$) transmission coefficients. We also study the temperature dependence of the tunneling conductance in the low energy and low temperature limit.

In generally, each individual quantum dot can be described by two-dimensional electron gas, and the Coulomb charging energy $E_c = e^2/(2C)$, where C is the capacitance of the

individual quantum dot, is very large compared to the single-particle level spacing but small compared to the tunneling channel bandwidths. For the case of only a few modes can tunnel from one quantum dot to another one, the quantum point contact can be described by a quantum wire [15,16]. We can also use a potential barrier to adjust the transmission coefficient. Based upon above considerations, we consider the following Hamiltonians [15,16,17]

$$H_0 = \hbar v_F \int dx [\psi_R^\dagger(x) \partial_x \psi_R(x) + \psi_L^\dagger(x) \partial_x \psi_L(x)] \quad (1)$$

$$H_c = E_c \left(\hat{n} - \frac{N}{2} \right)^2 \quad (2)$$

$$H_I = V_{2k_f} \int dx [\psi_R^\dagger(0) \psi_L(0) + \psi_L^\dagger(0) \psi_R(0)] \quad (3)$$

where $\psi_R(x)$ are right-moving electron operators, $\psi_L(x)$ are left-moving electron operators; $E_c = e^2/(2C)$, $eN = C_g(V_{gR} - V_{gL})$ is the gate voltage parameter of two quantum dots, and C_g is a gate-to-dot capacitance. $e\hat{n} = (e/2) \int_0^L dx [\psi_R^\dagger(x) \psi_R(x) - \psi_L^\dagger(x) \psi_L(x)]$ is the charge difference between two quantum dots; V_{2k_f} is a backward scattering potential which controls the transmission coefficient T . The system described by these Hamiltonians (1), (2) and (3) has been extensively studied by directly using the bosonization representation of the electron fields $\psi_{R(L)}(x)$ in the strategy of perturbation methods. However, the backward scattering term is relevant in the terminology of the renormalization group, the perturbation method may fail. To more effectively study this problem, we adopt another way developed in Ref. [14] where the backward scattering term can be exactly cancelled by an unitary transformation, and its effect is reflected on the correlation functions of the electrons and the expression of the charge operator $e\hat{n}$. We define the following new fermion fields

$$\psi_1(x) = \frac{1}{\sqrt{2}} [\psi_R(x) + \psi_L(x)]; \quad \psi_2(x) = \frac{1}{\sqrt{2}} [\psi_R(x) - \psi_L(x)] \quad (4)$$

while the bosonic representation of these fermion fields can be written in the usual way [18,19,20] as

$$\psi_{1(2)}(\mathbf{x}) = \left(\frac{D}{2\hbar v_F}\right)^{1/2} e^{i\phi_{1(2)}(\mathbf{x})} \quad (5)$$

where D is the band width of the conduction electrons. The boson fields $\psi_{1(2)}(\mathbf{x})$ have the relation with the density operators $\rho_{x-1(2)}(\mathbf{x}) = 2\psi_{1(2)}^\dagger(\mathbf{x})\psi_{1(2)}(\mathbf{x})$, and satisfy the commutation relations $[\rho_{x-1}(\mathbf{x}), \phi_{1(2)}(\mathbf{y})] = i2\delta(\mathbf{x}-\mathbf{y})$, $[\rho_{x-2}(\mathbf{x}), \phi_{2(2)}(\mathbf{y})] = i2\delta(\mathbf{x}-\mathbf{y})$, where the density operators are defined as $\rho_{1(2)}(\mathbf{x}) = \psi_{1(2)}^\dagger(\mathbf{x})\psi_{1(2)}(\mathbf{x})$. The Hamiltonians (1) and (3) can be written in the bosonic representation of the fermion fields $\psi_{1(2)}(\mathbf{x})$ as

$$H_0 = \frac{\hbar v_F}{4} \int dx [(\rho_{x-1})^2 + (\rho_{x-2})^2] \quad (6)$$

$$H_I = \frac{\hbar v_F}{2} \int dx [\rho_{x-1}(\mathbf{x}) - \rho_{x-2}(\mathbf{x})] j_{x=0} \quad (7)$$

where $\phi = \arctan(V_{2k_F}/(\hbar v_F))$ is phase shift induced by the backward scattering potential.

According to Eq.(4), the charge operator \hat{n} can be written as

$$\hat{n} = \frac{1}{2} \int_0^L dx [\psi_{1-}^\dagger(\mathbf{x})\psi_{2-}(\mathbf{x}) + \psi_{2-}^\dagger(\mathbf{x})\psi_{1-}(\mathbf{x}) - \psi_{1+}^\dagger(\mathbf{x})\psi_{2+}(\mathbf{x}) - \psi_{2+}^\dagger(\mathbf{x})\psi_{1+}(\mathbf{x})] \quad (8)$$

which only includes the cross terms of the fermion fields $\psi_{1-}(\mathbf{x})$ and $\psi_{2-}(\mathbf{x})$.

In order to cancel the ϕ -term in (7), we define the following unitary transformation

$$U = \exp i \frac{\phi}{2} \int dx [\rho_{x-1}(0) - \rho_{x-2}(0)] j_{x=0} \quad (9)$$

Under this unitary transformation, we can obtain the following relations

$$\begin{aligned} U^\dagger (H_0 + H_I) U &= H_0 \\ H_c &= U^\dagger H_c U = E_c \left[U^\dagger \hat{n} U - \frac{N}{2} \right]^2 \end{aligned} \quad (10)$$

It is worth notice that the backward scattering term completely disappears, its effect is only reflected on the charge operator \hat{n} for this problem. It has not any influence on the correlation functions of the fermion fields $\psi_{1(2)}(\mathbf{x})$ because we do not consider the electron-electron interactions in the quantum wire. The calculation of $U^\dagger \hat{n} U$ is very simple. By using the following formulae

$$[\psi_1(\mathbf{x}); \psi_1^\dagger(\mathbf{y})] = i \delta(\mathbf{x} - \mathbf{y}); [\psi_2(\mathbf{x}); \psi_2^\dagger(\mathbf{y})] = i \delta(\mathbf{x} - \mathbf{y})$$

$$U^\dagger \psi_1(\mathbf{x}) U = \psi_1(\mathbf{x}) + \frac{1}{2} \text{sgn}(\mathbf{x}); U^\dagger \psi_2(\mathbf{x}) U = \psi_2(\mathbf{x}) - \frac{1}{2} \text{sgn}(\mathbf{x}) \quad (11)$$

and taking the following gauge transformations

$$\psi_1(\mathbf{x}) = \psi_1(\mathbf{x}) e^{i\phi_1}; \quad \psi_2(\mathbf{x}) = \psi_2(\mathbf{x}) e^{i\phi_2}; \quad \psi_1^\dagger \psi_2 = \quad (12)$$

we can easily obtain the following expression of the charge operator $eU^\dagger \hat{N} U$

$$U^\dagger \hat{N} U = \hat{N} + \frac{\cos(2\phi)}{2} \int d\mathbf{x} [\psi_1^\dagger(\mathbf{x}) \psi_2(\mathbf{x}) + \psi_2^\dagger(\mathbf{x}) \psi_1(\mathbf{x})]$$

$$+ i \frac{\sin(2\phi)}{2} \int d\mathbf{x} [\psi_1^\dagger(\mathbf{x}) \psi_2(\mathbf{x}) - \psi_2^\dagger(\mathbf{x}) \psi_1(\mathbf{x})] \quad (13)$$

It is noted that although the backward scattering term is eliminated by the unitary transformation U (9), the charge operator $eU^\dagger \hat{N} U$ becomes complex. To simplify it, we redefine the right- and left-moving electronic fields

$$\psi_R(\mathbf{x}) = \frac{1}{\sqrt{2}} [\psi_1(\mathbf{x}) + \psi_2(\mathbf{x})]; \quad \psi_L(\mathbf{x}) = \frac{1}{\sqrt{2}} [\psi_1(\mathbf{x}) - \psi_2(\mathbf{x})] \quad (14)$$

while their bosonic representations reads $\psi_{R(L)}(\mathbf{x}) = \left(\frac{D}{2\hbar v_F}\right)^{1/2} \exp[i\phi_{R(L)}(\mathbf{x})]$. These new right- and left-moving electronic fields are different from original ones due to we have taken the unitary and gauge transformations of the fermion fields $\psi_{1(2)}(\mathbf{x})$. In terms of these new electronic fields, the charge operator $eU^\dagger \hat{N} U$ can be rewritten as

$$U^\dagger \hat{N} U = n + \frac{\cos(2\phi) + 1}{4} (\psi_L(0) - \psi_R(0))$$

$$+ i \frac{\sin(2\phi)}{2} \int_0^{\infty} d\mathbf{x} [\psi_L^\dagger(\mathbf{x}) \psi_R(\mathbf{x}) - \psi_R^\dagger(\mathbf{x}) \psi_L(\mathbf{x})] \quad (15)$$

where $n = (1/(4\pi)) \int_0^P [\cos(2\phi) \psi_R(1) + \psi_R(1) - \psi_L(1) - \cos(2\phi) \psi_L(1)]$. If we take the value of the phase shift as $\phi = \pi/2$, we can obtain the following relation

$$U^\dagger \hat{N} U j_c = n_c = \frac{1}{2} (N_L - N_R) \quad (16)$$

where $N_{R(L)} = \int_0^{\infty} d\mathbf{x} \psi_{R(L)}^\dagger(\mathbf{x}) \psi_{R(L)}(\mathbf{x})$ are the right- and left-moving electron numbers, respectively. Therefore, the value of the phase shift $\phi = \pi/2$ corresponds to strong

coupling critical point of the system induced by the backward scattering potential. At this strong coupling critical point, the transmission coefficient T of the electrons from the left quantum dot to the right one (or vice versa) is zero, i.e., the electrons are completely reflected on the impurity site $x = 0$. These two quantum dots are completely separated, therefore, we can observe a series of the double degenerate Coulomb blockade conductance peaks at this strong coupling critical point.

We now study the low temperature behavior of the tunneling conductance and the fractional Coulomb blockade conductance peak splitting away from this strong coupling critical point ($\nu = 2$). Because the directly hopping of the electrons from the right of the impurity to its left or vice versa is very weak, we can approximately replace the quantity

$$\int_0^Z dx [\psi_L^\dagger(x) \psi_R(x) + \psi_R^\dagger(x) \psi_L(x)]$$

by

$$a [\psi_L^\dagger(0) \psi_R(0) + \psi_R^\dagger(0) \psi_L(0)]$$

where a is a small constant. For simplicity, we define the following new boson fields

$$\phi_c(x) = \frac{1}{2} [\phi_L(x) + \phi_{L\#}(x) - (\phi_R(x) + \phi_{R\#}(x))]$$

$$\phi_s(x) = \frac{1}{2} [\phi_L(x) - \phi_{L\#}(x) - (\phi_R(x) - \phi_{R\#}(x))]$$

In terms of these new boson fields, the charge operator $eU^\dagger \hat{n} U$ is

$$U^\dagger \hat{n} U = n + \frac{\cos(2\theta) + 1}{2} \phi_c(0) + \frac{aD \sin(2\theta)}{h\nu_F} \sin(\phi_c(0)) \cos(\phi_s(0)) \quad (17)$$

and the Hamiltonian (6) reads

$$H_0 = \frac{h\nu_F}{4} \int_0^Z dx [(\partial_x \phi_c(x))^2 + (\partial_x \phi_s(x))^2 + (\partial_x \phi_{c\#}(x))^2 + (\partial_x \phi_{s\#}(x))^2] \quad (18)$$

However, due to the quantity $a \sin(2\theta)$ is very small, the last term in (17) can be perturbatively treated, the electrostatic Hamiltonian (10) can be written as by using Eq.(17)

$$H_c = E_c \left[\frac{\cos(2\theta) + 1}{2} \langle c(0) \rangle^2 + E_c \left(\frac{aD \sin(2\theta)}{2\hbar v_F} \right)^2 \sin^2(\theta_c(0)) \cos^2(\theta_s(0)) \right] \quad (19)$$

where $N = n + N = 2$. It is worth noting that for the case of weak backward scattering $\theta = 0$, i.e., the transmission coefficient $T = 1$, the low temperature behavior of the system is different from that for the strong backward scattering. We first consider the case of the weak backward scattering $\theta = 0$. Because the factor $(\cos(2\theta) + 1) = (2)$ is finite, the boson field $c(0)$ has a non-zero expectation value

$$\frac{\cos(2\theta) + 1}{2} \langle c(0) \rangle = N \quad (20)$$

which suppresses its low energy excitations. Therefore, we can safely integrate out the boson field $c(0)$ and obtain the following effective Hamiltonian

$$H_c^0 = E_c \left(\frac{aD \sin(2\theta)}{2\hbar v_F} \right)^2 \left[1 - \left(\frac{4eE_c}{D} \right)^2 \cos(2\theta_c(0)) \right] [1 + \cos(2\theta_s(0))] \quad (21)$$

where $E_c = E_c(1 + \cos(2\theta))^2 = (2)^2$, and $\gamma = 0.577$ is Euler constant. However, in the fermionization representation, $\cos(2\theta_s(0))$ can be written as: $\cos(2\theta_s(0)) = \frac{1}{2} (\psi^\dagger + \psi)(\psi^\dagger - \psi)$, where $\psi = \psi_0 + i\psi_1$ is an infinitesimal quantity, which produces a gap in the spectrum of fermion $\psi(0)$. Therefore, the $\cos(2\theta_s(0))$ -term in (21) is irrelevant in the terminology of the renormalization group.

For the case of the strong backward scattering $\theta = \pi/2$, the electrostatic energy significantly depends upon the quantity N . If N is zero, $N = 0$, after integrating out the boson field $c(0)$, we can obtain the following electrostatic Hamiltonian

$$H_c^0 = E_c \left(\frac{aD \sin(2\theta)}{2\hbar v_F} \right)^2 \left[1 - \left(\frac{4eE_c}{D} \right)^2 \right] [1 + \cos(2\theta_s(0))] \quad (22)$$

For the case of $N \neq 0$, because of $\max(\langle c(0) \rangle) = N$, if the phase shift satisfies the relation: $\theta < \theta_0$, where θ_0 is defined as: $1 + \cos(2\theta_0) = 2N$, after integrating out the boson field $c(0)$, we can obtain the same Hamiltonian as that in (21). If the phase shift satisfies the relation: $\theta > \theta_0$, we can obtain the following effective Hamiltonian

$$H_c^0 = E_c \left[\frac{\cos(2\theta) + 1}{2} \langle c(0) \rangle^2 + E_c \left(\frac{aD \sin(2\theta)}{2\hbar v_F} \right)^2 \left[1 - \left(\frac{4eE_c}{D} \right)^2 \right] [1 + \cos(2\theta_s(0))] \right] \quad (23)$$

where $\langle \phi_c(0) \rangle = \phi_c$ for $N > 0$, and $\langle \phi_c(0) \rangle = -\phi_c$ for $N < 0$. It is worth notice that the present method used to get the effective Hamiltonians in (21), (22) and (23) is completely different from that in Refs. [15] and [21]. After performing the unitary transformation (9), all information of the conduction electron scattering on the impurity is incorporated into the electrostatic Hamiltonian (19) which only contains higher order terms of the boson fields $\phi_{c(s)}(0)$. However, in Refs. [15] and [21], the authors directly take the approximation on the backward scattering potential which leads to a lower order terms of the boson fields $\phi_{c(s)}(0)$. This difference may make us obtain a different transport behavior from that in Ref. [21] in the low temperature limit. On the other hands, in terms of the terminology of the renormalization group, the former only gives some irrelevant terms, but the latter gives some relevant terms. Based upon above discussions, for the case of $0 < j < j_0$, we can obtain the following ground state energy of the system

$$E = E_0 - E_c \left(\frac{2ae E_c \sin(2\phi_c)}{h\nu_F} \right)^2 \cos(2\langle \phi_c(0) \rangle) \quad (24)$$

For the case of $j_0 < j < j_0 + 2$, the ground state energy of the system reads

$$E = E_0 + E_c [j - N] \frac{1}{2} (\cos(2\phi_c) + 1) + E_c \left(\frac{2ae E_c \sin(2\phi_c)}{h\nu_F} \right)^2 \quad (25)$$

Now we can obtain the fractional Coulomb blockade conductance peak splitting [13]. For the case of $0 < j < j_0$, it can be written as

$$\begin{aligned} f &= [E(N = \frac{1}{2}) - E(N = 0)] / [E(N = \frac{1}{2}) - E(N = -\frac{1}{2})] \\ &= 4 \left(\frac{2ae E_c \sin(2\phi_c)}{h\nu_F} \right)^2 \left(\frac{1 + \cos(2\phi_c)}{2} \right)^4 \left(1 - \cos\left(\frac{2\phi_c}{1 + \cos(2\phi_c)} \right) \right) \end{aligned} \quad (26)$$

In the case of $j_0 < j < j_0 + 2$, it can be written as

$$f = 4 \left[\frac{1}{2} - \frac{1}{2} (1 + \cos(2\phi_c)) \right] \quad (27)$$

It is noted that the fractional Coulomb blockade conductance peak splitting significantly depends upon the phase shift ϕ_c , i.e., the transmission coefficient T . However, in the case of

the weak backward scattering $0 < j < j_0$, it also depends upon the parameter a , the Fermi velocity v_F and the electrostatic Coulomb energy $E_c = e^2/(2C)$. In the case of the strong backward scattering $j_0 < j < j_0 = 2$, the fractional Coulomb blockade conductance peak splitting f only depends upon the phase shift ϕ , therefore, it is universal.

We now study the low temperature behavior of the tunneling conductance. To this end, we define the following current operator

$$I = e \langle U^\dagger \hat{n} U \rangle = e \frac{1 + \cos(2\phi)}{2} \langle \hat{n}_c(0) \rangle + \frac{aeD \sin(2\phi)}{h v_F} [\langle \hat{n}_c(0) \cos(\phi_c(0)) \cos(\phi_s(0)) \rangle - \langle \hat{n}_s(0) \sin(\phi_c(0)) \sin(\phi_s(0)) \rangle] \quad (28)$$

Using the Kubo formula of the conductance, we can easily obtain the following tunneling conductance in the low energy and low temperature limit

$$G(T) = G_0(\phi) [1 + AT (\frac{\sin(2\phi)}{2})^2 \cos(2\phi_c(0) > \phi)] \quad (29)$$

where $G_0(\phi) = e^2/(2h) (1 + \cos(2\phi))^2$, T is the temperature, and A is a constant. It is noted that the tunneling conductance has a linear temperature dependence, and at the strong coupling critical point $\phi_c = \phi_s = 2$, it is equal to zero. It is necessary to mention that as the frequency ω and the temperature T tend to zero, the system approaches to the strong coupling critical point $\phi_c = \phi_s = 2$ because the backward scattering term is relevant, the renormalized backward scattering potential V_{2k_F} tends to infinity in the low energy limit [22].

We define the following dimensionless tunneling conductance

$$g = \frac{G(T)}{G_0} = \frac{1}{2} (1 + \cos(2\phi))^2 \quad (30)$$

where $G_0 = e^2/(2h)$ is a unit quantum conductance of each tunneling channel. From Eqs.(26) and (27), we can obtain the following relation between the dimensionless tunneling conductance g and the fractional Coulomb blockade peak splitting f

$$f = \begin{cases} \frac{8}{\pi} \frac{P}{2B} g^{5/2} (1 - \frac{g}{2}) (1 - \cos(\frac{P}{2g})); & 0 < j < j_0 \\ 4 (\frac{1}{2} - \frac{g}{2})^2; & j_0 < j < j_0 = 2 \end{cases} \quad (31)$$

where $B = [ae E_c = (\hbar^2 v_F)^2]$. It qualitatively agrees with the experimental data in Ref. [4,5]. Because the factor a depends upon the structure of the quantum point contact, and the electrostatic energy E_c is determined by the structure of the quantum dot, the fractional Coulomb blockade peak splitting is not universal in the weak backward scattering $0 < j < j_0$. However, in the strong backward scattering $j_0 < j < j_0 + 2$, it is universal.

In summary, by using the bosonization method and the unitary transformation, we have shown that the tunneling conductance between two quantum dots has the linear temperature dependence in the low energy and low temperature limit, and given a general relation between the dimensionless tunneling conductance g and the fractional Coulomb blockade peak splitting f which is valid both for the weak and strong backward scattering of the electrons on the impurity. Our treatment of this kind of system is simple and effective, and can be used to study other similar problems because it can exactly treat the backward scattering potential which is a relevant term in the low energy region, the usual perturbation expansion methods of this term may fail because the high orders are divergent in the low energy limit.

We are very grateful to Prof. Peter Fulde for his encouragement.

REFERENCES

- [1] G. Schon and A. D. Zaikin, *Phys. Rep.* 198, 237 (1990); K. Flensberg and M. Jonson, *Phys. Rev. B* 43, 7586 (1991).
- [2] D. V. Averin and K. K. Likharev, in *Mesoscopic Phenomena in Solids*, eds. B. L. Altshuler, P. A. Lee and R. A. Webb (North-Holland, Amsterdam, 1991) p.173.
- [3] *Single Charge Tunneling*, eds. H. Grabert and M. H. Devoret (Plenum Press, New York, 1992); M. A. Kastner, *Rev. Mod. Phys.* 64, 849 (1992).
- [4] F. R. Waugh et al, *Phys. Rev. Lett.* 75, 705 (1995); *Phys. Rev. B* 53, 1413 (1996).
- [5] L. W. Molenkamp, K. Flensberg and M. Kemmerink, *Phys. Rev. Lett.* 75, 4282 (1995).
- [6] N. C. van der Vaart et al, *Phys. Rev. Lett.* 74, 4702 (1995).
- [7] F. Hofmann et al, *Phys. Rev. B* 51, 13872 (1995).
- [8] C. H. Crouch et al, *Surface Science* 361/362, 631 (1996).
- [9] I. M. Ruzin, V. Chandrasekhar, E. I. Levin and L. I. Glazman, *Phys. Rev. B* 45, 13469 (1992).
- [10] C. A. Stafford and S. Das Sarma, *Phys. Rev. Lett.* 72, 3590 (1994).
- [11] G. Klimck, G. Chen and S. Datta, *Phys. Rev. B* 50, 2316 (1994).
- [12] K. A. Matveev, L. I. Glazman and H. J. Baranger, *Phys. Rev. B* 53, 1034 (1996); *ibid*, B 54, 5637 (1996).
- [13] J. M. Golden, B. I. Halperin, *Phys. Rev. B* 53, 3893 (1996); cond-mat/9604063.
- [14] Y. L. Liu, Effective treatment of single impurity scattering in Tomonaga-Luttinger liquid, preprint.
- [15] K. Flensberg, *Phys. Rev. B* 48, 11156 (1993); *Physica B* 203, 432 (1994).
- [16] K. A. Matveev, *Physica B* 203, 404 (1994); *Phys. Rev. B* 51, 1743 (1995).

- [17] F Guinea and M Ueda, *Physica B* 203, 440 (1994).
- [18] A Luther, and I Peschel, *Phys. Rev. B* 9, 2911 (1974).
- [19] V J Emery, in *Highly Conducting One-Dimensional Solids*, eds. J.T Devreese et al, (Plenum Press, New York, 1979); J Solyom, *Adv. Phys.* 28, 201 (1979).
- [20] F D M Haldane, *J. Phys. C* 14, 2585 (1981).
- [21] A Furusaki, and K A Matveev, *Phys. Rev. Lett.* 75, 709 (1995).
- [22] C L Kane, and M P A Fisher, *Phys. Rev. Lett.* 68, 1220 (1992); *Phys. Rev. B* 46, 15233 (1992).