

Two interacting particles in a disordered chain I : Multifractality of the interaction matrix elements

Xavier Waintal and Jean-Louis Pichard

CEA, Service de Physique de l'Etat Condensé,
Centre d'Etudes de Saclay, F-91191 Gif-sur-Yvette, France

today

Abstract. For N interacting particles in a one dimensional random potential, we study the structure of the corresponding network in Hilbert space. The states without interaction play the role of the "sites". The hopping terms are induced by the interaction. When the one body states are localized, we numerically find that the set of directly connected "sites" is multifractal. For the case of two interacting particles, the fractal dimension associated to the second moment of the hopping term is shown to characterize the Golden rule decay of the non interacting states and the enhancement factor of the localization length.

PACS. 05.45.+b Theory and models of chaotic systems { 72.15.Rn Quantum localization { 71.30.+h Metal-insulator transitions and other electronic transitions

The wave functions of one particle in a random potential have been extensively studied. In two dimensions within the localization domains [1] the large fluctuations of their amplitudes have a multifractal character. In one dimension, the elastic mean free path l and the localization length L_1 coincide, preventing a single one particle wave function to be multifractal over a significant range of scales. The description of the correlations existing between the localized eigenstates is more difficult. This is quite unfortunate, since a local two-body interaction reorganizes the non interacting electron gas in a way which depends on the spatial overlap of (four) different one particle states. When one writes the N -body Hamiltonian in the basis built out from the one particle states (eigenbasis without interaction), this overlap determines the interaction matrix elements, i.e. the hopping terms of the corresponding network in Hilbert space. In this work, we numerically study the distribution of the hopping terms in one dimension, when the one body states are localized. It has been observed [2] that this distribution is broad and non Gaussian. We give here numerical evidence that this distribution is multifractal. Moreover, since the obtained Renyi dimensions do not depend on L_1 , simple power laws describe how the moments scale with the characteristic length L_1 of the one body problem. Since the main applications we consider (Golden rule decay of the non interacting states, enhancement factor of the localization length for two interacting particles) depend on the square of the hopping terms, we are mainly interested by the scaling of the second moment. For a size $L \gg L_1$, we show that, contrary to previous assumptions, the N -body eigenstates without interaction directly coupled by the square of the hopping terms have not a density of the order of

the two-body density $\rho_2(L_1) / L_1^2$, but a smaller density $\rho_2(L_1) / L_1^{f(q=2)}$. The dimension $f(q=2)$ (1.75 for hopping terms involving four different one body states) characterizes the fractal set of N -body eigenstates without interaction which are directly coupled by the square of the hopping terms.

We consider N electrons described by an Hamiltonian including the kinetic energy and a random potential, plus a two-body interaction:

$$H = \sum_i d_i^\dagger d_i + U \sum_{i,j} Q_{ij} d_i^\dagger d_j d_j^\dagger d_i \quad (1)$$

The operators d_i^\dagger (d_i) create (destroy) an electron in a one body eigenstate $|j\rangle$ of spin \uparrow (\downarrow). Noting $\psi_j(n)$ the amplitude on site n of the state $|j\rangle$ with energy ϵ_j , the interaction matrix elements are proportional to the Q given by:

$$Q_{ij} = \sum_n \psi_j(n) \psi_i(n) \psi_j(n) \psi_i(n) \quad (2)$$

This comes from the assumption that the interaction $U \sum_n c_n^\dagger c_n c_n^\dagger c_n$ is local. The c_n^\dagger (c_n) create (destroy) an electron on the site n and $d_i^\dagger = \sum_n \psi_j(n) c_n^\dagger$. When $U = 0$, the Hamiltonian is diagonal in the basis built out from the one particle states, and the N -body states ($|\psi\rangle = \sum_{i=1}^N d_i^\dagger |i\rangle$) can be thought as the "sites" with energy ϵ_i of a certain network which is not defined in the real space, but in the N -body Hilbert space. When $U \neq 0$, different "sites" can be directly connected by off-diagonal interaction matrix elements. Therefore, one can map [3] this complex N -body problem onto an Anderson

localization problem defined on a particular network in the N -body Hilbert space. Since the interaction is two body, only the "sites" differing by two quantum numbers can be directly coupled. This restriction will not matter [4] for $N = 2$ and (under certain approximations) may yield a Cayley tree topology [3] for the resulting network, if N is large. We study the additional restrictions coming from one body dynamics.

We summarize a few evaluations of the second moment ($q = 2$) of Q which have been previously used. Case (i): The one body Hamiltonian is described by random matrix theory (RMT). The statistical invariance under orthogonal transformations $O(M)$ implies that $\langle Q^2 \rangle_i = 1/M^3$ where M is the number of one body states. Case (ii): The system is a disordered conductor of conductance g . An estimate [3] based on perturbation theory gives $\langle (UQ)^2 \rangle_i / \langle g \rangle^2$. Since the one particle mean level spacing $\Delta / 1 = M$, this perturbative result coincides with the previous RMT results if one takes $M = g^2$. Moreover, it is valid only if all the one particle states appearing in Eq.(2) are taken from a sequence of g consecutive levels in energy. Otherwise, Q can be neglected. Case (iii): The system is a disordered insulator. Shepelyansky [4] in his first study of the two interacting particles (TIP), assumes a RMT behavior for the $M = L_1^d$ components of the wave function inside the localization domain, and neglects the exponentially small components outside this domain. When the dimension $d = 1$, one gets a term $\langle Q^2 \rangle_i = L_1^3$ for the terms coupling a TIP state $j >$ to L_1^2 TIP states $j >$. This estimate for $g < 1$ differs from the one valid when $g > 1$ under two important aspects: not only $M = L_1^d$ instead of g^2 , but the condition for a large hopping term is not given by four one particle states close in energy, but by four states close in real space, i.e. located inside the same localization domain. Ponomarev and Silvestrov have criticized [5] this estimate, using an approximate description of a localized state for weak disorder. They note that the density of TIP states coupled by the interaction is sensibly smaller.

For a more accurate study of Q in one dimension, we consider a spin independent one particle Anderson tight binding model with L sites and nearest neighbor hopping ($t = 1$). The on-site potentials V_n are taken at random in the interval $[-W, W]$ and the boundary conditions are periodic. L_1 is estimated from the weak disorder formula $L_1 = 25W^{-2}$. The $Q_{j_0 j}$ are calculated using Eq.(2) and numerical diagonalization of the one particle Hamiltonian. Q , for fixed d and L is a two-dimensional object which is not defined in the real 2d space, but in the space of two one particle quantum numbers j_0 and j . Those states $j >$ (and $j_0 >$) can be ordered in different ways: (a) spectral ordering by increasing eigenenergy, (b) spatial ordering by the location n of their maximum amplitude, from one side of the sample to the other, (c) momentum ordering if $W = 0$. Let us note that ordering (b) becomes meaningful only in the localized regime ($L > L_1$).

We first study the matrix element $Q_{j_0 j_0}$, characterizing two electrons with opposite spins in the same state $j_0 >$ hopping to an arbitrary state $j >$. Hopping is very unlikely over scales larger than L_1 . The L_1^2 large values of the hopping term are concentrated inside a square of size L_1^2 , as shown in Fig.1 for a given sample using ordering (b) and a rainbow color code. Fig.1 is not homogeneously colored, but exhibits a complex pattern which reminds us another bidimensional object: the one particle wave function in a two dimensional disordered lattice. This suggests us to analyze its fluctuations as for the 2d one body states, and to check if this pattern is not the signature of a multifractal structure.




Fig. 1. $Q_{j_0 j_0}$ with $j_0 >$ taken in the bulk ($n_0 = 50$) of a sample of size $L = L_1 = 100$. Spatial ordering (b). The color code goes from red (small value) to violet (large value) through yellow, green and blue.

In analogy with the 2d one body problem, we do not expect that this multifractality will be valid in the whole $(j; j_0)$ Hilbert space, but only in a limited but parametrically large domain.

We proceed as usual (see references [6,7]) for the multifractal analysis. For L_1 and L fixed, we divide the plane $(j; j_0)$ into $(L/D)^2$ boxes of size D and we calculate the ensemble averaged function for different values of q

$$I_q(D) = \left(\sum_{i=1}^{N_{\text{boxes}}} |Q_{j_0 j}|^q \right)^{1/q} \quad (3)$$

The existence of a multifractal measure defined in the $(j; j_0)$ -plane by the interaction matrix elements is established in the next figures. In Fig.2, a single sample has been used and power laws $I_q(D) / D^{-q/d}$ are obtained over many orders of magnitude for different values of q .

The limits of validity of these power laws are shown in Fig.3.




Fig. 2. Power laws showing that Fig. 1 corresponds to a multifractal measure in the (i, j) two dimensional plane. $I_q(D)$ are calculated for a single sample with $L_1 = L = 2500$. i, j in the bulk of the spectrum. The states j, i and i are ordered by increasing eigenenergy (ordering a). The dashed line corresponds to the RMT prediction (case (i)).

On the left side, spatial ordering (b) for different values of L_1 is used for the states (i, j) . One can see that $I_q(D) / D^{-q}$, for scales $1 < D < L_1$, as indicated by the arrows. The lower scale is given by the lattice spacing of the (n, n) network in Hilbert space. The upper scale L_1 is the largest scale compatible with a spatial overlap of the states i and j , for a fixed j_0 . This means that the multifractality of the interaction matrix elements $Q_{j_0 j}$ in the two dimensional Hilbert space (i, j) has the same parametrically large range of validity as the one body wave function $[\psi]$ in two dimensions (scale $1 < D < L_1$). Here, multifractality is valid for L_1^2 matrix elements as multifractality is valid in the 2d one body problem for L_1^2 sites.

On the right side of Fig. 3 spectral ordering (a) is used for the same samples, giving the same power laws as with ordering (b), inside the corresponding energy range $(L_1) < D < L$ indicated by the arrows.

$(\lambda) / x^{-1}$ is the level spacing of a segment of size x , and 1 is the band width. The exponents $d(q)$ are independent of the ordering when $L > L_1$ (i.e. when the ordering (b) becomes meaningful) and the small fluctuations from sample to sample are removed by ensemble averaging.

The corresponding Renyi dimensions

$$d(q) = (q-1)$$

are shown in Fig. 4 for different L and L_1 , using ordering (a) and ensemble averaging.

For an infinite L_1 (no disorder), the eigenstates are plane waves of momentum k and $Q \neq 0$ only if $k + k = k - k = 0$. This gives $d(0) = 2$ and $d(q > 0) = 1$ with ordering (c). The dimensions calculated with ordering (a)




Fig. 3. Left: Spatial ordering (b). $I_2(D)$ for $\mathcal{D}_{j_0 j}$ with j_0 in the bulk of the spectrum and for L_1 (indicated by the arrows) = 25; 200; 2500. $L = 2500$. The power law behaviors are obtained for a given sample if $1 < D < L_1$. Right: Same samples using spectral ordering (a). The arrows indicate the lower scales associated with (L_1) . The power laws are valid for $L = L_1 < D < L$.

are close to this limit. For a finite L_1 , $d(q)$ goes from the clean limit ($L < L_1$) to an L_1 -independent regime when $L > L_1$. In the crossover regime ($L \approx L_1$) the $d(q)$ depend on L_1 . In the limit $L \gg L_1$, the $d(q)$ (using orderings (a) or (b)) do not depend on L and L_1 . For $0 < q \leq 3$,

$$d(q) = 2 - q$$

with a slope -0.135 . The L_1 -independence of $d(q)$ is shown in the insert of Fig. 4 for $L_1 = L$ up to $L_1 = 600$.

A multifractal distribution has scaling behavior described by the $f(q)$ -spectrum, given by the relations:

$$d(q) = \frac{d}{dq} \text{ and } f(d(q)) = (q-1) d(q): \quad (4)$$

We obtain

$$f(d(q)) = 2 - q^2 \quad (5)$$

for $q \leq 3$, i.e. a parabolic shape $f(d) = 2 - d^2$ around the maximum $2 + 1$. We have mainly studied the first positive moments, since we are mainly interested by $f(d(q=2))$. Indeed, when one uses Fermi golden rule to calculate the interaction-induced decay of a non-interacting state, one needs to know the density of states directly coupled by the second moment ($q = 2$) of the hopping term. The fractal dimension of the support of this density is given by $f(d(q=2))$. For greater values of q , there are deviations around the parabolic approximation, indicating deviations around simple lognormal distributions. From a study of the large and small




Fig. 4. $d(q)$ with ϵ_0 in the bulk of the spectrum using ordering (a) and after ensemble averaging. Filled symbols: $L = 240$ and $L_1 = 1$ (diamonds), $25 \cdot 10^4$ (circles) and 2500 (squares). Open symbols: $L_1 = 70$ and $L = 960$ (diamonds), 480 (circles) and 240 (squares). Insert: the slope $\Delta(L_1)$ showing that $d(q)$ is disorder independent for $q \geq 3$ and $L_1 < L$.

values of $\mathcal{D}_{\epsilon_0, j}$ one can obtain $d(q \rightarrow 1)$. We find $d(+1) = 1.33$ and $d(-1) = 3.15$, giving the limits of the support of $f(\cdot)$.

We have also checked that our results for $\mathcal{Q}_{\epsilon_0, j}$ do not depend on the chosen ϵ_0 and studied the general case where $j > \epsilon_0$ and $j < \epsilon_0$ are not the same. In Fig. 5, one can see that the $\mathcal{Q}_{\epsilon_0, j}$ studied for different ϵ_0 give the same curves $d(q)$. Using energy ordering (a) and imposing an energy separation $j - j' > (L_1)$ in order to have a good overlap between the fixed states and ϵ_0 , we find also power law behaviors for $I_q(\mathcal{D})$. The corresponding dimensions $d(q)$ are given in Fig. 5, characterized by a slope

$$(\epsilon_0) = (\epsilon_0) = 2 \quad 0.065:$$

Therefore, the multifractal character of \mathcal{Q} is less pronounced when $j > \epsilon_0$, but remains relevant.

So far, we have discussed the hopping terms of the general N -body problem. We now discuss how our results modify previous assumptions for two interacting particles (TIP). As pointed out by Shepelyansky, the interaction induced hopping mixes nearby in energy TIP states $j > \epsilon_0$ and $j' > \epsilon_0$. The decay width [8,9,10] of a TIP state $j > \epsilon_0$, built out from two one particle states localized within L_1 , can be estimated using Fermi golden rule. If one assumes RMT wave functions inside L_1 for the one particle states, (case (iii)) the \mathcal{Q} $U \cdot L_1^{3/2}$ couple the TIP state $j > \epsilon_0$ to all the TIP states $j' > \epsilon_0$ inside L_1 . Around the band center, they have a density $\rho_2(L_1) / L_1^2$




Fig. 5. $d(q)$ calculated for $L_1 = 70$ and $L = 240$ using ordering (a). The filled symbols correspond to $\epsilon_0 = 0$, averaged over 10 consecutive ϵ_0 chosen in various parts of the spectrum: down triangle ($130 \leq \epsilon_0 \leq 140$), square ($120 \leq \epsilon_0 \leq 130$), diamond ($145 \leq \epsilon_0 \leq 155$) and up triangle ($170 \leq \epsilon_0 \leq 180$). Empty symbols correspond to the case $\epsilon_0 = 0$. $d(q)$ is averaged for a few ϵ_0 : circle ($\epsilon_0 = 100; 123; 133$), cross ($\epsilon_0 = 80; 123; 133$) and plus ($101; 112$ and $123; 133$).

and Fermi golden rule gives

$$(\mathcal{E} = 0) / \frac{U^2}{L_1^3} \rho_2(L_1) = \frac{U^2}{L_1} \quad (6)$$

We have shown that all the TIP states which can be coupled by the interaction within the localization domains are not equally coupled. Since the square of the hopping term appears in the Golden rule, our multifractal analysis gives a reduced effective TIP density $\rho_2^{\text{eff}} / L_1^{f(q=2)}$ which should replace the total TIP density $\rho_2(L_1)$. The resulting expression

$$/ \frac{U^2}{L_1^3} L_1^{f(q=2)} \quad (7)$$

can be compared to the direct numerical evaluation:

$$= U^2 \sum_{j, j'} \mathcal{D}_{\epsilon_0, j}^2 (\epsilon_0 + \epsilon_0') \quad (8)$$

of the Golden rule decay.

In Fig. 6, we show for three different sizes L how the decay rate numerically calculated using Eq.(8) depends on L_1 , for a TIP state $j > \epsilon_0$ where ϵ_0 is taken in the bulk of the spectrum. From Fig. 4, one gets $(2) = 1.52$ and $(2) = 1.69$, and Eq.(4) gives $f(q=2) = 1.35$. For this value, one can see in Fig. 6 that Eq.(7) and Eq.(8) give indeed the same L_1 -dependence. This observed $L_1^{1.65}$ law




Fig. 6. $\Gamma_{\alpha\alpha}(L_1)$ for three different sizes L . Dashed line: $y = 2.5L_1^{-1.65}$. Dotted line: $y = 0.5L_1^{-1}$.

clearly differs from the L_1^{-1} law implied by the RMT assumption (case (iii)). We can also see that $\Gamma_{\alpha\alpha}(L_1)$ does not depend on L when $L_1 < L$, since there are no significant hopping terms for range larger than L_1 .

Another interesting issue is the enhancement of the localization length L_2 , which is induced by the interaction and characterizes a restricted set of TIP states which have a sufficient overlap to be re-organized by a local interaction [11,12]. Using the Thouless block scaling analysis [13], one finds $\frac{L_2}{L_1} \propto (U = L_1^{3/2} \Gamma_{\alpha\alpha}(L_1))^2$. If the density $\Gamma_{\alpha\alpha}(L_1)$ of states coupled by the interaction is the total TIP density for a size L_1 , one finds the original estimate [4,13] L_2 / L_1^2 . The multifractality yields a reduced effective density $\rho_e / L_1^{f(q=2)}$ instead of the total TIP density. Since the contribution of TIP states $j > \epsilon$ with ϵ dominates, we use $f(q=2) = 1.75$ valid when ϵ and we find $L_2 / L_1^{1.5}$. This L_1 -dependence is in agreement with recent numerical results [2,14]. So there is an enhancement, though weaker than the original estimate [4] (L_2 / L_1^2), due to the multifractal distribution of the hopping terms.

In summary, we have studied how one particle dynamics (one dimensional localization) can affect the many body problem through non trivial properties of the distribution of the two-body interaction. In a clean system, one has $f(q=2) = 1$ and the density of states which are effectively coupled by the interaction is the one particle density ρ_1 / L . The disorder, as it is well known, enhances the effect of the interaction, since the effective density $\rho_e / L_1^{f(q=2)}$, with $1 < f(q=2) < 2$ for $L = L_1$. This enhancement of the density of states coupled by the interaction inside a system of size L_1 is nevertheless smaller than the one (ρ_2 / L_1^2) given by fully chaotic one body states inside their localization domains. In a second paper [15], a study of the TIP spectral fluctuations will be presented, showing that statistics is critical (as for the one body spectrum at a mobility edge) if U is large enough, accompanied by multifractal wavefunctions in the

TIP eigenbasis for $U = 0$. In a third paper [16], a study of the dynamics of a TIP wave packet will be presented, showing that the center of mass exhibits anomalous diffusion between L_1 and L_2 . These three studies provide consistent and complementary observations supporting our claim: multifractality and criticality are relevant concepts for a TIP system with on site interaction in one dimension. Our results go beyond the TIP problem and show that oversimplified two-body random interaction matrix models [17,18,19] which ignore multifractality in the hopping cannot properly describe the many body quantum motion in Anderson insulators.

We are indebted to S.N. Evangelou for very useful comments.

References

1. V.I. Fal'ko and K.B. Efetov, Euro. Phys. Lett. 32, 627 (1995); Phys. Rev. B 52, 17413 (1995).
2. K. Frahm, A. Müller-Groeling, J.-L. Pichard and D. Weinmann, Europhys. Lett. 31, 169 (1995).
3. B.L. Altshuler, Y. Gefen, A. Kamenev, and L.S. Levitov, Phys. Rev. Lett. 78, 2803 (1997).
4. D.L. Shepelyansky, Phys. Rev. Lett. 73, 2067 (1994).
5. I.V. Ponomarev and P.G. Silvestrov, Phys. Rev. B 56, 3742 (1997).
6. G. Paladin and A. Vulpiani, Phys. Rep. 156, 147 (1987).
7. T.C. Halsey, M.H. Jensen, L.P. Kadano, I. Procaccia and B.I. Shraiman, Phys. Rev. A 33, 1141 (1986).
8. Ph. Jacquod and D.L. Shepelyansky, Phys. Rev. Lett. 75, 3501 (1995).
9. D. Weinmann and J.-L. Pichard, Phys. Rev. Lett. 77, 1556 (1996).
10. D. Weinmann, J.-L. Pichard and Y. Imry, J. Phys. 1. France 7 1559 (1997).
11. F. von Oppen, T. Wettig and J. Müller, Phys. Rev. Lett. 76, 491 (1996).
12. D. Weinmann, A. Müller-Groeling, J.-L. Pichard and K. Frahm, Phys. Rev. Lett. 75, 1598 (1995).
13. Y. Imry, Europhys. Lett. 30, 405 (1995).
14. P.H. Song and Doochul Kim, cond-mat 9705081 (1997).
15. X. Waintal, D. Weinmann and J.-L. Pichard, cond-mat 9801134.
16. S. de Toro Arias, X. Waintal and J.-L. Pichard, in preparation.
17. O. Bohigas and J. Flores, Phys. Lett. 34B, 261 (1971); 35B, 383 (1971).
18. V.V. Flambaum, F.M. Izrailev and G. Casati, Phys. Rev. E 54, 2136 (1996).
19. Ph. Jacquod and D.L. Shepelyansky, Phys. Rev. Lett. 79, 1837 (1997).