

# A Universal Formula for Percolation Thresholds II. Extension to Anisotropic and Aperiodic Lattices

Serge Galam and Alain Mauger

Laboratoire des Milieux Désordonnés et Hétérogènes  
Tour 13 – Case 86, 4 place Jussieu,  
75252 Paris Cedex 05, France

.

Phys. Rev. E, 1997

PA Classification Numbers: 64.60 A, 64.60 C, 64.70 P

## Abstract

In a recent paper, we have reported a universal power law for both site and bond percolation thresholds in any Bravais lattice with  $q$  equivalent nearest neighbors in dimension  $d$ . We now extend it to three different classes of lattices which are, respectively, anisotropic lattices with not equivalent nearest neighbors, non-Bravais lattices with two atom unit cells, and quasicrystals. The investigation is focussed on  $d = 2$  and  $d = 3$ , due to the lack of experimental data at higher dimensions. The extension to these lattices requires the substitution of  $q$  by an effective (non integer) value  $q_{\text{eff}}$  in the universal law. For each out of 17 lattices which constitute our sample, we argue the existence of one  $q_{\text{eff}}$  which reproduces both the site and the percolation threshold, with a deviation with respect to numerical estimates which does not exceed 0.01.

---

Laboratoire de l'Université P. et M. Curie – Paris 6, associé au CNRS (URA n° 800)

# 1 Introduction

Very recently we have posulated a universal power law for both site and bond percolation thresholds [1]. The formula yields thresholds for any Bravais lattice, at any dimension with an impressive accuracy. It writes

$$p_c = p_0 [(d-1)(q-1)]^{-a} d^b \quad (1)$$

with  $d$  the space dimension and  $q$  the coordination number. While  $b = a$  for bond dilution, it is  $b = 0$  for site dilution. Three different classes were found with three distinct parameter sets  $fp_0; a; b$ . The first class includes two-dimensional triangle, square and honeycomb lattices. It is characterized by  $fp_0 = 0.8889; a = 0.3601; b = 0.6558$  for site dilution and by  $fp_0 = 0.6558; a = 0.6897; b = 0.6897$  for bond dilution. Two-dimensional Kagome and all other lattices of cubic symmetry (for  $3 \leq d \leq 6$ ) constitute the second class which is characterized by  $fp_0 = 1.2868; a = 0.6160; b = 0.7541$  for sites and  $fp_0 = 0.7541; a = 0.9346; b = 0.9346$  for bonds respectively. A third class has been found at high dimensions ( $d > 6$ ), which recovers the infinite Cayley tree limit, but is not relevant to present investigation which deals with lattices only at  $d = 2$  and  $d = 3$ .

Besides the dimension  $d$ , percolation thresholds within a class depend only on  $q$ . This is understood in lattices where the  $q$  nearest neighbors of any site are equivalent, which is indeed the case for all lattices above mentioned. This is however a drastic restriction since many percolation problems in physics deal with lattices which do not have this property. It is the purpose of the present work to investigate the extension of Eq. (1) to other lattices, via the substitution of  $q$  by an effective parameter  $q_{eff}$ .

We have checked on a sample of 17 lattices that, for each of them, there exists one value of  $q_{eff}$  which reproduces both the site and the bond percolation thresholds. The error is within 0.006 for all the lattices, except for two of them (dual of Penrose, and dodecagonal lattice with ferromagnetic links), where the error reaches 0.01 for reasons we also discuss in this paper. We consider this sample is large enough to be representative of a general trend, and then conclude in the existence of such a  $q_{eff}$  in any lattice. For a lattice which does not belong to our sample, this parameter could be used as an intermediate quantity to predict bond from site percolation thresholds, or vice versa, with the same accuracy.

The paper is organised as follows. Section 2 discusses a series of lattices with non-equivalent neighbors. The introduction of an effective number of nearest neighbors in our universal formula for percolation is argued in Section 3. The results are shown and discussed in Section 3. The last section contains some remarks for future work.

## 2 Non-equivalent neighbors

Some periodic Bravais lattices are anisotropic, for instance the hexagonal lattice at  $d = 3$ . In this case, any lattice site has 6 equivalent nearest neighbors in the  $a$ - $b$  plane (bonding angle is  $60^\circ$ ) and 2 non-equivalent sites along the  $c$  axis (bonding angle is  $90^\circ$ ). Actually, the anisotropic lattice percolation threshold should depend on the degree of anisotropy. For the hexagonal lattice, it means  $p_c$  should be different from that of the corresponding isotropic lattice with the same set  $d = 3; q = 8$ , i.e., the bcc lattice. This difference has been observed indeed recently [2] in the particular case of the stacked triangular lattice, which becomes the hexagonal lattice when  $a = b = c$ .

There also exist non-Bravais lattices which are periodic like fcc and bcc. Another case is the hexagonal close packed (hcp) lattice which, on a topologic view point, is a simple hexagonal lattice with two atoms per unit cell. Percolation thresholds for the hcp were obtained long ago [3].

Some lattices are not even periodic. This is the case of quasicrystals which are aperiodic lattices with long-range order. Besides their own interest such structures can serve as models for alloy-like materials making a growing interest in the determination of the quasicrystalline lattice percolation thresholds. First determinations of percolation thresholds on 2-d quasilattices have been made on the Penrose tiling and its dual [4, 5, 6]. Recently, percolation thresholds have also been computed in two of the most important quasilattices, the simple octagonal and dodecagonal tilings [7], which belong to the Penrose local isomorphism class.

By analogy with other lattices, one can require that lattice sites are connected only via the tile edges. The corresponding percolation problem is named chemical percolation. Adding connection through the diagonals of the tiles which are shorter than the tile edge leads to the so-called ferromagnetic percolation [8]. Both these percolations will be considered here. These lattices are reproduced in Fig. (1).

At last, some lattices do not have a single-valued coordination number  $q$ .

Lattices with mixed-valued coordination can be either periodic or non-periodic. An example of periodic lattice at  $d = 2$  is provided by the dice lattice which mixes  $q = 3$  and  $6$ . An example of non-periodic lattice is the Penrose tiling, which mixes  $q = 3; 4; 5; 6$  and  $7$ . Note the dual of dice is the Kagome lattice and vice versa, while the dual of the Penrose tiling is named dual lattice of Penrose. Both the Kagome and the dual of Penrose are lattices with the single-valued coordination  $q = 4$ . Octagonal and dodecagonal lattices have mixed-value coordinations. Numerical estimates of the percolation thresholds of dice can be found in [6].

### 3 An effective number of nearest neighbors

In the past, attempts have been made to generalize empirical relations such as the Scher and Zallen approximate [9] which depends on dimension. For instance at  $d = 2$ , these are  $q_{pc}(\text{bond}) \approx 2.0$  and  $q_{pc}(\text{site}) \approx 0.45$ , with  $f$  the lattice filling factor. Extension to quasilattices [6, 7] consists in replacing  $q$  by mean coordination number  $\bar{z}$  in the invariant appropriate to bonds (the problem for sites is not solved, because there is no ambiguous definition of the filling factor in quasicrystals [7]).

In the same spirit, we propose to extend Eq. (1), replacing  $q$  by an effective value  $q_{eff}$ . The dependence of the percolation thresholds on site connectivity does not imply that the relevant variable should be the arithmetic average  $\bar{z}$ . Therefore, we regard  $q_{eff}$  as a parameter which has to be determined, rather than arbitrarily forced to be equal to  $\bar{z}$ .

The need of substituting  $q$  by  $q_{eff}$  in the equations is easily understood in both the case of quasilattices and the case of lattices which mix different values of  $q$ . It is also needed in anisotropic lattices. Let us again consider the hexagonal lattice with spacing  $a$ ,  $b$  and  $c$  in the three lattice directions. In the limit  $c \rightarrow +\infty$ , one is left with decoupled  $ab$  planes for physical systems having a finite range of interaction. Therefore, the percolation threshold of the hexagonal lattice must depend somehow on the ratio  $c/a$ . It is expected to range between that of isotropic bcc with  $d = 3; q = 8$  and that of the  $d = 2$  triangular lattice, associated to the limit  $c/a \rightarrow +\infty$ . Note in this limit one recovers an isotropic lattice with  $q = 6$  instead of  $q = 8$ , which suggests an effective coordination  $6 \leq q_{eff} \leq 8$ . In view of such considerations, we propose to extend Eq. (1) substituting  $q$  by some  $q_{eff}$  for any lattice.

Note within the two classes defined by the set  $(a; p_0)$ ,  $q_{eff}$  is the unique

unknown parameter. For each lattice, we find a value of this fitting parameter which, when inserted into Eq.1, reproduces both the site and the bond percolation threshold within 0.01. Values of  $q_{\text{eff}}$  are reported for different lattices in Table (1), together with site and bond percolation thresholds  $p_c^s$ ,  $p_c^b$  obtained when  $q$  is replaced by  $q_{\text{eff}}$  in Eq. (1). Mean coordination  $\bar{z}$  and  $p_c^e$  for site and bond percolation are also reported for comparison. Exact or numerical estimates  $p_c^e$  are from Refs [6, 7, 10].

## 4 The results

For lattices at  $d = 2$ ,  $q_{\text{eff}}$  differs from  $\bar{z}$  by 1% in the case of the dual lattice of Penrose, and is smaller than 0.5% for all the other lattices. These data corroborate that two dimensional lattices are divided into the two distinct classes defined in [1]. This is illustrated in Figs. (2) and (3), where  $p_c^e$ 's are reported in a log-log plot such that the experimental points for lattices in the same class align on a straight line according to Eq.1.

The pertinent variable after Eq. (1) is  $(d-1)(q_{\text{eff}}-1)$  for sites and  $(d-1)(q_{\text{eff}}-1)=d$  for bonds. Those are the variables in abscissa in Fig. (3) which reports data for lattices in dimensions  $d = 2$  and  $d = 3$  (data in higher dimensions have been already displayed in a similar plot in [1] with  $q_{\text{eff}} = q$ ). Since the first class concerns only lattices which are all at  $d = 2$ , above variables for respectively sites and bonds may reduce to one single common variable  $q_{\text{eff}}$  as shown in Fig. (2).  $\delta$  reaches 0.01 only in the dodecagonal lattice with ferromagnetic links, and in the dual lattice of Penrose. In all the other cases, the error in the percolation threshold estimates is only on the third decimal.

Note the larger error in the dodecagonal lattice. We attribute it to an actual bond percolation threshold  $p_c^e(\text{bond}) = 0.495$  [6] larger than a priori expected. One would indeed have expected  $p_c(\text{bond}) = 0.475$  from the straight line in Figure 3, in which case a value  $q_{\text{eff}} = 4.289$ , close to  $\bar{z} = 4.27$ , would have reproduced both site and bond percolation thresholds within  $\delta \approx 0.001$ . The problem with the estimate  $p_c^e(\text{bond}) = 0.495$  is also evident from the Scher and Zallen invariant. It yields  $\bar{z}p_c^e(\text{bond}) = 2.11$ , the largest value of this parameter among all the lattices investigated [6]. At the opposite,  $p_c(\text{bond}) = 0.475$  would yield  $\bar{z}p_c(\text{bond}) = 2.028$ , close to the invariant value 2.0 at  $d = 2$ .

At  $d = 3$ , percolation values are reproduced with a very good accuracy, since  $\delta \approx 0.004$  for all the lattices. The difference between  $q_{\text{eff}}$  and  $\bar{z}$  is not

negligible as it reaches few per cent in some cases. This is easily understood in the case of the hexagonal (stack triangle) case, where we have argued earlier that a value  $q_{eff}$  smaller than 8 is expected. Actually, we find  $q_{eff} = 7.66$ , as a consequence of the anisotropy. The hexagonal close packed (hcp) lattice has percolation thresholds which are different from those of the fcc lattice, as expected since the hcp lattice is not a Bravais lattice.

However, differences are small which may be attributed to the fact that both lattices are indeed isotropic, each site being surrounded by 12 equivalent neighbors. In this context, the small value of  $q_{eff}$  close to 11 in hcp lattice is not only due to the non-Bravais nature of the lattice. It is also related to the fact that in the fcc lattice,  $q_{eff}$  is only 11.6, significantly smaller than  $q = 12$ , although the coordination number is single-valued.

On the other hand,  $q_{eff}$  differs from  $q$  by only 1% in the other lattices (sc, diamond, bcc). Yet one would expect  $q_{eff} = q$  in such isotropic lattices with single-valued coordination number. The difference between  $q_{eff}$  and  $q$  in this case illustrates that our formula for the percolation thresholds is not exact as we already stated in [1], and shown convincingly in ref. 2. Nevertheless, both site and bond percolation thresholds for all the lattices in any dimension are provided within 1% by our universal law involving only two parameters: the dimension  $d$  and a parameter  $q_{eff}$  which contains more information on the geometry than the mean coordination.

## 5 Conclusion

We have shown that our universal formula for percolation thresholds we reported earlier [1] for periodic Bravais lattices with equivalent nearest neighbors does extend to any kind of lattice, provided the coordination number is replaced by an effective value  $q_{eff}$ . We then conclude that a good estimate of both site and bond percolation thresholds can be obtained from the formula in Eq. 1 involving only the dimension  $d$  of the lattice, and one parameter  $q_{eff}$  which contains the geometric information of the lattice. This parameter however, does not reduce to the mean coordination number  $\bar{z}$ , although  $\bar{z}$  and  $q_{eff}$  differ by few per cent only. Indeed, we find the universality does include different numerical values for  $q_{eff}$  and thus different percolation thresholds for lattices which have the same set  $(d; \bar{z})$ . This is evident from Table (1) which reports results for as many as six lattices with  $d = 2; \bar{z} = 4$ , two lattices with  $(d = 3; \bar{z} = 8)$ , two other ones with  $(d = 3; \bar{z} = 12)$ .

In our previous work, we had only three lattices belonging to the first class. The present work extends this class from 3 to 8 lattices. We might have invoked chance for three points aligned on a same line, but not for 8 points like in Fig.1. Therefore, we confirm the existence of two different classes, one for some of the two-dimensional lattices, the other one for all the other two-dimensional lattices and all the lattices up to  $d=7$ .

We do not have a scheme to derive the relevant variable  $q_{eff}$  from the geometry of the lattice. However, an important result of this work is that this variable does exist. It means that there exists a single value for  $q_{eff}$  which accounts for both the site and the bond percolation thresholds for any given lattice within  $\pm 0.01$ . This result is sufficient to give our formula a prediction ability for percolation thresholds of other lattices which have not been computed yet. For example, the knowledge of one site (bond) percolation threshold for a given lattice is sufficient to determine a point on the relevant straight line in Fig. 2 or 3. Then  $q_{eff}$  can be found from the abscissa of this point, which in turn allows for the determination of the bond (site) percolation threshold from our universal formula, within  $\pm 0.01$ . Depending on whether  $p_c^s$  or  $p_c^b$  is known, the value of  $q_{eff}$  deduced from Fig. (2) or (3) will be different. However, this difference only corresponds to the deviation of the  $p_c^e$ 's with respect to the universal law, i.e. less than 1% with the exception of only few outliers.

The robustness of our formula suggests the extension to more complex problems such as directed percolation. Also, anisotropic percolations with different bond probabilities in different directions may be addressed in the near future.

Acknowledgments.

We would like to thank Dietrich Stauffer for stimulating comments.

## References

1. S. Galam and A. Mauger, Phys. Rev. E 53, 2177 (1996).
2. Van der Marck, Phys. Rev. E 55, 1228 (1997); S. Galam and A. Mauger, Phys. Rev. E 55, 1230 (1997).
3. V.K.S. Shante and S. Kirkpatrick, Adv. Phys. 20, 326 (1971).
4. J.P. Lu and J.L. Bim an, J. Stat. Phys. 40, 1057 (1987).
5. F. Babalievski and O. Pesshev, C. R. Acad. Sci. Bulg. 41, 85 (1988).
6. F. Yonesava, S. Sakamoto, K. Aoki, S. Nose and M. Hori, J. Non-Cryst. Solids 123, 73 (1988).
7. F. Babalievski, Physica 220, 245 (1995).
8. D. Ledue and J. Teillet, J. Non-Cryst. Solids 191, 216 (1995).
9. H. Scher and R. Zallen, J. Chem. Phys. 53, 3759 (1970).
10. D. Stauffer and A. Aharony, Introduction to Percolation Theory, 2nd ed. (Taylor and Francis, London, 1994).



## Figure captions

Fig. 1. Less common lattices studied in this paper : Kagome (a) and its dual, Dice lattice (b); Penrose quasicrystalline lattice (c) and its dual (d); octagonal quasicrystalline lattice with ferromagnetic links (e) and chemical links (f); dodecagonal quasicrystalline lattice with ferromagnetic links (g) and chemical links (h).

Fig. 2. Inverse of percolation thresholds as a function of the variables  $(d-1)(q_{\text{eff}}-1)$  and  $(d-1)(q_{\text{eff}}-1)=d$  which reduce here ( $d=2$ ) to the single variable  $q_{\text{eff}}$  in logarithmic scales for lattices belonging to the first class.

Fig. 3. Inverse of percolation thresholds as a function of the variables  $(d-1)(q_{\text{eff}}-1)$  and  $(d-1)(q_{\text{eff}}-1)=d$  appropriate to site and bonds, respectively, in logarithmic scales for lattices belonging to the second class.





rst class								
Lattice	$\bar{z}$	$q_{eff}$	site			bond		
			$p_c^e$	$p_c$		$p_c^e$	$p_c$	
Square	4	4.02	0.5928	0.5970	+ 0.0042	0.5	0.4935	-0.0064
Honeycomb	3	2.99	0.6962	0.6938	-0.0024	0.6527	0.6581	+ 0.0054
Triangular	6	5.98	0.5	0.4986	-0.0014	0.34729	0.34955	+ 0.0023
Dice	4	4.189	0.5851	0.5854	+ 0.0003	0.476	0.4754	-0.0006
Penrose	4	4.194	0.5837	0.5851	+ 0.0014	0.477	0.4748	-0.0022
Octahedron links	4	4.170	0.585	0.5867	+ 0.0017	0.48	0.4773	-0.0027
Octahedron links	5.17	5.013	0.543	0.5389	-0.0041	0.40	0.406	+ 0.0057
Dodecahedron links	3.63	3.638	0.628	0.6269	-0.0011	0.54	0.5419	+ 0.0019
second class								
Lattice	$\bar{z}$	$q_{eff}$	site			bond		
			$p_c^e$	$p_c$		$p_c^e$	$p_c$	
Kagome	4	3.980	0.6527	0.6567	+ 0.0040	0.5244	0.5195	-0.0049
dual Penrose	4	4.04	0.6381	0.6487	+ 0.0106	0.5233	0.5099	-0.01341
Dodecahedron links	4.27	4.218	0.617	0.6264	+ 0.0094	0.495	0.4835	-0.0115
hex. compact	12	11.146	0.204	0.2015	-0.0025	0.124	0.1263	+ 0.0023
stac. triangle	8	7.661	0.2623	0.2611	-0.0012	0.1859	0.1872	0.0013
Diamond	4	4.0087	0.43	0.4260	-0.0040	0.1859	0.3935	+ 0.0055
sc	6	5.9558	0.3116	0.3132	+ 0.0016	0.2488	0.2468	-0.0020
bcc	8	8.1355	0.246	0.2502	+ 0.0042	0.1803	0.1755	-0.0047
fcc	12	11.626	0.198	0.1958	-0.0022	0.119	0.1210	0.0020

Table 1: percolation thresholds from this work  $p_c$  compared to "exact estimates"  $p_c^e$  taken from [6, 7 and 10].  $p_c - p_c^e$  refers to multi-valued coordination number (mean coordination  $\bar{z}$ ). All the lattices are in dimension  $d = 2$ , except the 6 last ones in the second class which have  $d = 3$ . We have completed the table caption as follows:  $q_{eff}$  has been chosen as the arithmetic average of the parameters  $q$  which reproduces  $p_c^e$  for respectively site and bond when inserted in Eq. (1).