December 15, 2021

a constriction in a FQH annulus

Stefan Kettem ann

Max-Planck Institute f. Physik Komplexer Systeme, Dresden, Germany

The ux periodicity of them odynam ic properties of an annulus in the fractional quantum hall state with a constriction is considered. It is found that $_0$ -periodicity is obtained due to transfer of fractionally charged particles or composite ferm ions between the edges of the annulus, respectively. The result for the nite magnitude of the persistent current across a very strong constriction is presented, as obtained with an extension of W en's edge state theory.

Keywords: Fractional quantum Halle ect, Persistent Current, Tunneling

Corresponding Author:

Stefan K ettem ann, M ax-P lanck Institute for Physics of C om plex System s, N oethnitzerstr. 38, 01187 D resden, G erm any Fax: 49-351-871-1199 e-m ail: kettem an@m pipksm pi-dresden m pg.de

I. IN TRODUCTION

The Fractional Qunatum Halle ect, discovered by Stoermer, Tsui and Gossard, [1] was explained by Laughlin in a theory based on a trial wave function [2]. Since then, the question was debated if quasiparticle excitations with fractional charge and statistics can be observed from such a state.

Thouless and Gefen considered an annulus in a strong magnetic eld which condenses uid within the theory of composite ferm ions. the electrons to a fractional quantum hall W hen a composite ferm ion is moved around state. They predicted that therm odynamic a circle in the annulus, which is pierced by an properties of such a system have ux peri-additional ux , it aquires a phase odicity 0 due to the existence of fam ilies of states of the FOH - annulus. These were connected by a physical mechanism, the - where S is the area enclosed by the circle nite back scattering amplitude of fractionally [8]. x is the number of composite ferm ions charged quasiparticles [3].

persistent current in a FQH - annulus with annulus, we have x = (S = S)B = 0, giving a weak constriction [4]. There, an extension of W en's edge state theory [5] was used, which takes into account the dynamics of so that the composite ferm ion sees a reduced zero-modes [6].

There exists the alternative theory of the fractional quantum halle ect based on com posite ferm ions as proposed by Jain [7].

A composite Fermion of a FQH state of lling factor = 1 = (2m + 1) has charge (-e), and 2m vortices attached to it, corresponding to 2 m ux quanta $_0 = 2$ c=e aligned oppositely to the external magnetic eld B.

II. THE FQH ANNULUS IN THE COMPOSITE FERM ION PICTURE

Let us consider the annulus of a FQH liq-

$$= \frac{2}{0} \int_{0}^{I} dr A = \frac{SB}{0} + \frac{1}{0} \quad 4m \quad x; \quad (1)$$

within that circle. Since there are no com -Recently, this was veried by deriving the posite ferm ions in the inner area S_i of the

$$= 2 \frac{SB'}{0} + 2 - \frac{1}{0}; \qquad (2)$$

magnetic eld

$$B^{\sim} = (1 \ 2m) B$$
 (3)

and an enhanced ux, piercing the annulus,

1

$$= + 2m S_{i}B$$
: (4)

We obtain the guiding centers of the Eigenstates of the CF's by dem and ing quantization of the phase = 2 1, 1 an integer [9].

An adiabatic increase of the ux reduces the positions of the CF's. Thus, the FQH liquid moves up the inner edge. The area S_i where there are no CF's becomes smaller, so that there are x_i CF's added to the inner edge with,

$$x_i = -$$
: (5

An equal amount of CF's is removed from the outer edge. A fler a change of the ux by one ux quantum $_0$, a fraction of a CF is added to the inner edge [10]. This is in agreement with the fact that the Laughlin wave function yields the addition of a charge e to the inner edge, when one ux quantum is added adiabatically into the annulus. If only electrons could tunnel between the edges of the FQH annulus, its periodicity would be enhanced to 1= $_0$.

Is the 0-periodicity as required by the Byers- and Young theorem [11] restored by the transport of composite ferm ions between the edges?

The rem oval of a CF from the inner edge adds 2m ux quanta to the ux piercing the annulus, since the two vortices attached to the rem oved electron did cancel 2 ux quanta of the magnetic eld B. Thus, the total num – ber of com posite ferm ions added to the inner edge, when the ux piercing the annulus is changed by , and one com posite ferm ion tunnels to the outer edge, is:

$$x_{i} = \frac{+2m_{0}}{0}$$
 1: (6)

Thus, after a ux change of 1 ux quantum,
 the tunneling of the composite ferm ion to the
 outer edge brought the system to its state at
 y = 0, restoring the 0-periodicity.

Therefore, in order that the ux periodicity of the FQH – annulus can be understood from the composite ferm ion perspective, there has to be tunneling of composite ferm ions between the edges of the annulus.

It is now natural to ask if the tunneling am plitude of composite ferm ions is renorm alized, how this is compared with the renormalization of fractionally charged particles [12], and if it yields the same temperature and size dependence of the persistent current in the presence of a constriction as was obtained in R ef. [4], using the edge state theory

2

one has to formulate the edge state theory tunnel across the barrier. of composite ferm ions which is complicated by the dynamics of the vortices attached to search.

Here, rather we want to conclude by presenting results on the persistent current through a very strong constriction using the edge state theory as presented in Ref. [4].

> III. PERSISTENT CURRENT THROUGH A VERY STRONG CONSTRICTION

W e consider again an annulus in the FQ H state at odd inverse lling factor. There is a strong constriction at one point in the annulus, producing a weak link through which particles can tunnel, only. The potential barrier is taken to be much larger than the energy gap in the bulk of the fractional quantum hall liquid, $V_0 >> bulk$. Then, the many-body wave function of the FQH state in the annulus decays exponentially. Thus, the strong correlations in the FQH state can

of fractionally charged particles. To this end, not extend through it and only electrons can

We obtain the persistent current in secthe ferm ions. Then, one can use this theory ond order perturbation theory in the most to nd a bosonized form of composite edge relevant electron tunneling amplitude $w_{1=}$. ferm ions. This is the subject of ongoing re-We use the edge state theory of a single chiral edge connected at the constriction.

$$I(') = \frac{e}{angmom : l} X (0, W_{l=eff}(l; '))$$

$$cos[2 (l+ ')]; (7)$$

where $\prime = = 0$, being the ux penetrating the annulus in addition to the constant background magnetic eld B... For tem peratures below the level spacing, T v=L, the renorm alized tunneling am plitude of electrons w_{eff} is given by

$$w_{1=} (l;')_{eff} = w_{1=} (l;') (v=(2L))^{1=}$$

$$(2(1 \cos((x_{R} x_{L})=L)))^{1=(2)}: (8)$$

Here, L is the total length of the single chiral edge, and $x_{\rm R}\,$ and $x_{\rm L}\,$ are the two positions at which the tunneling takes place. Their distance is taken to be of the order of L=2. w (l;') are the ux dependent coe cients of an expansion of the tunneling am plitude of electrons across the barrier in terms

3

of the Eigen states of angularm on entum lof ite fermions. The ux periodicty $_0$ is rethe annulus of noninteracting electrons with- stored due to the existence of back scattering out constriction. They satisfy the relation, of composite ferm ions between the edges. $w_{1=} (l; ' + 1) = w_{1=} (l + 1; ').$ There is a nite persistent current of elec-

strong constriction a nite operiodic per- the barrier height exceeds the bulk energy sistent current of electrons across the weak gap, at tem peratures not exceeding the level link and its amplitude decreases with in-spacing. creasing circum ference L of the annulus only slow ly, like a power law.

Thus, there is in the presence of a very trons across a very strong constriction, when

The author would like to thank Yuval Gefen for many useful discussions and Kon-When the temperature exceeds the level stantin Efetov and Benham Farid for usefull spacing, T = v=L, the elective tunneling comments on the tunneling through a barrier.

am plitude is

w_{1= eff} (1;') = $w_{1=}$ (1;')(T=)¹⁼ exp[1= TL=v $(1 \exp(j_{\underline{x}} \underline{x} \underline{j}=L)) \cos(v=(2TL))]$ $(1 \exp(2 T = v j x x j))^{1=}$ (9)

and the persistent current has an exponentially sm all am plitude.

The derivation, as well as the result for a potential barrier sm aller than the bulk energy $gap_{I}V_{0} < bulk$ will be presented elsewhere.

The ux periodicity of a FQH annulus can be understood within the picture of com pos-

REFERENCES

- [1] H.L.Stormer, D.C.Tsui, A.C.Gossard, Surface Science 113, 32 (1982),
- [2] R.B. Laughlin, Phys. Rev. Lett. 50, 1395 (1983),
- [3] D. J. Thouless, Phys. Rev. B 40, 12034 (1989), D. J. Thouless, Y. Gefen, Phys. Rev. Lett. 66, 806 (1991), Y. Gefen, D. J. Thouless Phys. Rev. B 47, 10423 (1993),
- [4] S.Kettem ann, Phys. Rev. B 55, 2512-22 (1997),
- [5] X.G.Wen, Phys. Rev. B 43, 11025 (1991), Int. J. Mod. Phys. B 6, 1711 (1992),
- [6] F.D.M. Haldane, J. Phys. C 14, 2585 (1981),
- [7] J.K. Jain, Phys. Rev. Lett. 63, 199 (1989),
- [8] A.S.Goldhaber, J.K. Jain, unpublished (1995),
- [9] B. I. Halperin, Phys. Rev. B 25, 2185 (1982).
- [10] D.B.Chklovskii, B.I.Halperin, unpublished (1997),
- [11] N.Byers and C.N.Young, Phys. Rev. Lett. 7,46(1961),
- [12] C.de C.Cham on and X.G.W en, Phys. Rev. Lett. 70, 2605 (1993), K.M oon, H.Yi, C. L.Kane, S.M.Girvin, and Matthew P.A.Fisher, Phys. Rev. Lett. 71, 4381 (1993),