

## Sound Propagation and Transport Properties of Liquid $^3\text{He}$ in Aerogel

D. Rainer<sup>1</sup> and J. A. Sauls<sup>2</sup>

<sup>1</sup>*Physikalisches Institut, Universität Bayreuth, Bayreuth, Germany*

<sup>2</sup>*Department of Physics & Astronomy, Northwestern University, Evanston, Illinois*

(Received August 31, 1997)

### Abstract

*Superfluid  $^3\text{He}$  confined in aerogel offers a unique chance to study the effects of a short mean free path on the properties of a well defined superfluid Fermi liquid with anisotropic pairing. Transport coefficients and collective excitations, e.g. longitudinal sound, are expected to react sensitively to a short mean free path and to offer the possibility for testing recently developed models for quasiparticle scattering at aerogel strands. Sound experiments, together with a theoretical analysis based on Fermi liquid theory for systems with short mean free paths, should give valuable insights into the interaction between superfluid  $^3\text{He}$  and aerogel.*

A model for liquid  $^3\text{He}$  in aerogel based on a random distribution of short-ranged potentials acting on  $^3\text{He}$  quasiparticles has been shown to account semi-quantitatively for the reduction of the transition temperature,  $T_c$ , and the suppression of the superfluid density,  $\rho_s(T)$ .<sup>1</sup> Although the order parameter for  $^3\text{He}$  in aerogel is not firmly identified, measurements of the magnetization indicate that the low pressure phase is an ESP state.<sup>2</sup> A transverse NMR shift is observed and is roughly consistent with that of an axial state even though the B-phase is stable in pure  $^3\text{He}$  at these pressures. The addition of a small concentration of  $^4\text{He}$ , which coats the aerogel strands, induces a suppression of the magnetization for  $T < T_c^{\text{aerogel}}$ , indicative of a non-ESP state like the BW phase.<sup>2</sup> Thus, the stability of the superfluid state of  $^3\text{He}$  in aerogel is quite sensitive to the detailed interaction between  $^3\text{He}$  and the aerogel strands. Calculations of the free energy for  $^3\text{He}$  in aerogel which include anisotropic and magnetic scattering, and the effects of orientational correlations of the aerogel strands, confirm the sensitivity of the superfluid phases to the interaction between  $^3\text{He}$  and the aerogel.<sup>1,3</sup>

The high porosity of the aerogel implies that the silica structure does not significantly modify the bulk properties of normal  $^3\text{He}$ . The dominant effect of the aerogel structure is to scatter  $^3\text{He}$  quasiparticles moving at the

bulk Fermi velocity. If the coherence length,  $\xi_0 = \hbar v_f / 2\pi T_{c0}$ , is sufficiently long compared to the average distance between scattering centers then a reasonable starting point is to treat the aerogel as a homogenous scattering medium (HSM) described by a mean-free path  $\ell$ .<sup>1</sup> In this article we discuss the effects of a short mean free path on some of the transport properties of liquid  $^3\text{He}$ . The calculations presented below for sound propagation assume the BW phase is stable; however, the propagation and damping of low-frequency sound are expected to be qualitatively similar for other phases.




Fig. 1 -  $p$ - $T$  phase diagram vs.  $\ell$ . The geometric mean-free path is estimated to be  $1750 \text{ \AA}$ . The data is from Ref. 4.

The superfluid transition temperature for  $^3\text{He}$  in aerogel is suppressed by quasiparticle scattering off the aerogel structure. In the HSM model the suppression of  $T_c$  is given by the Abrikosov-Gorkov formula,

$$\ln(T_{c0}/T_c) = \Psi\left(\frac{1}{2} + \frac{1}{2} \frac{\xi_0 T_{c0}}{\ell T_c}\right) - \Psi\left(\frac{1}{2}\right), \quad (1)$$

with the pair-breaking parameter given by  $\xi_0/\ell$ .<sup>1</sup> Figure 1 shows the calculated aerogel transition temperature for several values of the mean-free path. The pressure dependence of the calculated phase diagram is determined by the pressure dependence of the bulk transition temperature and the Fermi velocity via  $\xi_0 = \hbar v_f / 2\pi T_{c0}$ . The experimental data for  $T_c(p)$  is from Matsumoto, et al.<sup>4</sup> for an aerogel with  $\approx 98\%$  open volume. For this porosity the typical diameter of the aerogel strands is  $d \simeq 30 \text{ \AA}$  and the mean distance between strands is  $D \simeq 325 \text{ \AA}$ , which should be compared with the bulk coherence length  $\xi_0 \simeq 700 \text{ \AA}$  at  $p = 1$  bar. The geometric mean-free path of the

aerogel determined from the surface to volume ratio,  $S \simeq 2.6 \times 10^5 \text{ cm}^{-1}$ , is  $\ell_{\text{geom}} = \frac{3\pi}{2}/S \simeq 1750 \text{ \AA}$ . The calculations show a mean-free-path of this order gives good agreement with the phase diagram at low temperatures and low pressures, i.e. for  $\xi(p) = \hbar v_f/2\pi k_B T_c(p, \ell) > D \simeq 325 \text{ \AA}$ , which corresponds to pressures  $p \lesssim 15 \text{ bar}$ . Also note that the mean-field phase diagram shows a zero-temperature phase transition as a function of pressure determined by  $\xi_0(p_c) = 0.28 \ell$ .

The transport properties of  $^3\text{He}$  should also be strongly affected by scattering from the aerogel. In the normal state the quasiparticle distribution function,  $\phi(\hat{\mathbf{p}}, \mathbf{R}; \epsilon, t)$ , satisfies the Boltzmann-Landau transport equation,

$$\frac{\partial \phi}{\partial t} + \mathbf{v}_f \cdot \nabla \phi + \left( \frac{\partial \phi_0}{\partial \epsilon} \right) \frac{\partial \mathcal{E}}{\partial t} = I[\phi], \quad (2)$$

where  $\mathcal{E}$  is the effective potential acting on the quasiparticles and  $I[\phi]$  is the collision integral. The effective potential consists of the external driving potentials and the internal potentials resulting from quasiparticle interactions. In pure liquid  $^3\text{He}$  the quasiparticle lifetime is determined by inelastic collisions between quasiparticles and is of order  $\tau_{\text{in}} \simeq 1 \mu\text{s} \cdot \text{mK}^2/T^2$  at  $p = 15 \text{ bar}$ . However, in aerogel the mean time between scatterings by the aerogel is  $\tau_{\text{el}} = \ell/v_f \simeq 4 \text{ ns}$  at the same pressure. Thus, for temperatures below  $T_* \simeq 16 \text{ mK}$  the collision rate is dominated by quasiparticle scattering by the aerogel:  $I_{\text{el}} = -\frac{1}{\tau_{\text{el}}} (\phi - \langle \phi \rangle_{\text{FS}})$ . This leads to strong reduction in the thermal conductivity and viscosity, and to an increase in the damping of hydrodynamic sound. The static transport coefficients exhibit cross-over behavior dictated by the scattering rate; the thermal conductivity and shear viscosity scale as

$$\kappa = \frac{1}{3} C_v v_f^2 \tau \sim \begin{cases} 1/T & , \\ T & \end{cases}, \quad \eta = \frac{1}{15} N_f v_f^2 p_f^2 \tau \sim \begin{cases} 1/T^2 & , \\ \text{const} & , \end{cases} \quad T > T_* \quad , \quad T < T_* \quad , \quad (3)$$

above and below  $T_*$ . Recent speculations that the low-temperature phase of  $^3\text{He}$  in aerogel for  $p < p_{cr}$  is not described by Fermi-liquid theory<sup>4,5</sup> can be tested by measuring these transport coefficients.

At low frequencies a compressible aerogel will move nearly in phase with the  $^3\text{He}$  density and longitudinal current mode; sound propagates but it is damped by the viscous coupling of the  $^3\text{He}$  to the aerogel,  $\alpha_1 = \frac{\omega^2}{\rho c_1^3} \eta$ , where  $c_1$  is the hydrodynamic sound velocity and  $\rho$  is the mass density of  $^3\text{He}$ . The viscous damping of hydrodynamic sound saturates for  $T < T_*$  at  $\alpha_1/q \simeq \frac{2}{5} (\omega \tau_{\text{el}}) / (1 + F_0^s)$ . At higher frequencies the impedance mismatch between  $^3\text{He}$  and the aerogel sound mode leads to an increasingly out-of-phase motion of  $^3\text{He}$  excitations and aerogel. Hydrodynamic sound may become

overdamped and reemerge as a diffusive mode. The frequency at which the cross-over from damped hydrodynamic sound to an overdamped diffusive mode occurs depends on the elastic compliance of the aerogel and the microscopic details of the coupling between  $^3\text{He}$  and the aerogel strands. Here we assume the frequency is above this cross-over, in which case the hydrodynamic mode is an overdamped diffusive mode,  $\omega = -i\mathcal{D}_s q^2$ , where  $\mathcal{D}_s = c_1^2 \tau$  is the acoustic diffusion constant with  $1/\tau = (1 + F_1^s/3)/\tau_{el}$ . At still higher frequencies,  $\omega\tau \gg 1$ , zero sound can propagate, albeit with relatively high attenuation,  $\alpha_0 \simeq 1/c_0\tau \sim 10^4 \text{ cm}^{-1}$  at  $p = 10 \text{ bar}$ . The regime for propagating zero sound is also pushed to higher frequencies,  $\omega/2\pi > v_f/2\pi\ell \simeq 42 \text{ MHz}$  at  $p = 10 \text{ bar}$ .




Fig. 2 - Suppression of  $\rho_s(T)$  by scattering at  $p = 10 \text{ bar}$  with  $F_1^s = 8.6$ . Results in Born approximation are shown for bulk  $^3\text{He-B}$  and mean-free paths  $\ell = 1600$  to  $3000 \text{ \AA}$ . The inset shows the 4<sup>th</sup> sound velocity for the same values of the mean-free path.

The collective mode spectrum of superfluid  $^3\text{He}$  in aerogel is also expected to show significant changes compared to bulk  $^3\text{He}$ .<sup>6,7,8</sup> A typical example is low frequency sound in superfluid  $^3\text{He}$  ( $\omega \ll \Delta/\hbar$ ). In bulk  $^3\text{He}$  one has collisionless zero sound and hydrodynamic first sound with nearly the same velocities,  $c_0^2 \approx c_1^2 = \frac{1}{3}(1 + F_0^s)(1 + \frac{1}{3}F_1^s)v_f^2$ , and small damping by quasiparticle-quasiparticle scattering (for reviews on sound and collective modes in  $^3\text{He}$  see e.g. Refs. 9, 10, 11). In aerogel, on the other hand, one expects a behavior which is similar to sound in  $^3\text{He}$  confined to small channels.<sup>12</sup> Sound will be weakly damped for  $q\ell \gg 1$ , ceases to be a well

defined mode for  $q\ell \approx 1$ , and reappears for  $q\ell \ll 1$  as fourth sound with a temperature dependent velocity,  $c_4 \simeq \sqrt{\rho_s(T)/\rho} c_1$ . Fig. 2 shows the effects of elastic scattering on the superfluid density.<sup>13</sup> Note the reduction in  $\rho_s/\rho$  at  $T = 0$ . For a mean-free path of  $\ell = 1800 \text{ \AA}$  less than 50% of the  $^3\text{He}$  mass density contributes to  $\rho_s(T = 0)$ . The inset shows the fourth sound velocity neglecting damping.




Fig. 3 - Spectral function of the density response  $\chi(\mathbf{q}, \omega)$  for  $q\ell = 0.001 \dots 10$ . The results are for  $T/T_c = 0.8$ ,  $1/\tau_{el}T_c = 0.01$ ,  $F_0^s = 50$  and  $F_1^s = 8.6$ ; the corresponding zero sound mode is at  $\omega/v_f q = c_0/v_f = 8.1$ .

Measurements of sound propagation in  $^3\text{He}$ -aerogel should provide a sensitive test of the HSM model. We study the spectrum of longitudinal sound in this model by calculating the linear response of the  $^3\text{He}$  density,  $\rho(\mathbf{q}; \omega)$ , to a driving force of wavevector  $\mathbf{q}$  and frequency  $\omega$ . Our driving force will be a scalar field described by a potential,  $\delta U_{ext}(\mathbf{q}; \omega)$ , which couples to the  $^3\text{He}$  density. More realistic “experimental driving forces” require more elaborate calculations, which are not called for given the present experimental status. The calculation follows the quasiclassical version<sup>14</sup> of the method of Betbeder-Matibet and Nozières.<sup>15</sup> In the low frequency limit one has to solve Boltzmann-Landau transport equations for the branches of particle-like and hole-like excitations with distribution functions  $\delta\phi_{B1, B2}(\hat{\mathbf{p}}, \mathbf{R}; \epsilon, t)$ . We keep the dominant Landau parameters,  $F_0^s$  and  $F_1^s$ , and obtain an effective scalar potential,  $\delta\tilde{u}(\mathbf{q}; \omega)$ , and longitudinal vector potential,  $\delta\tilde{a}(\mathbf{q}; \omega) = \mathbf{v}_f \cdot \delta\mathbf{A}(\mathbf{q}; \omega)$ . The collision terms in the

HMS model have the form,  $I_{B1,B2} = -\frac{1}{\tau(\epsilon)} \left( \delta\phi_{B1,B2} - \langle \delta\phi_{B1,B2} \rangle_{FS} \right)$ , where  $\tau(\epsilon) = \ell/v(\epsilon)$ , and  $v(\epsilon) = v_f \sqrt{\epsilon^2 - |\Delta|^2}/\epsilon$  is the energy dependent quasiparticle velocity in the superfluid state. In the low frequency limit the transport equations have to be supplemented by Landau's self-consistency equations for the effective potentials,  $\delta\tilde{u}$  and  $\delta\tilde{a}$ , and the particle conservation law,  $\rho + \nabla \cdot \mathbf{j} = 0$ . In this limit we can ignore the less important self-consistency equation for the amplitude,  $\delta|\Delta|$ , of the order parameter. The five coupled linear equations for the distribution functions, the effective potentials, and the phase,  $\delta\Psi(\mathbf{q};\omega)$ , of the order parameter can be solved, and will be described elsewhere.




Fig. 4 - Temperature dependence of the imaginary and real (inset) parts of the spectral function  $\chi(\mathbf{q},\omega)$  for  $T/T_c = 0.25 \dots 0.85$ . The results are shown for  $q\ell = 0.1$ ,  $qv_f = 0.001T_c$ ,  $F_0^s = 50$  and  $F_1^s = 8.6$ .

The calculated crossover from weakly damped zero sound to weakly damped fourth sound is shown in Fig. 3. We display the frequency dependent spectral function,  $-\mathcal{I}\text{m}\chi(q,\omega)$ , for various wavelengths at fixed temperature and elastic scattering time. One can see clearly the transition from zero sound at  $v_f q \gg 1/\tau_{el}$  to fourth sound at  $v_f q \ll 1/\tau_{el}$ . Fig. 4 shows the real and imaginary parts of the response function,  $\chi(q,\omega)$ , at various temperatures and fixed  $\tau_{el}$ . Because of the short mean free path,  $q\ell = 0.1$ , the zero sound resonance with wavevector  $\mathbf{q}$  is overdamped in the normal state and just below  $T_c$ . The damping decreases exponentially in the superfluid state for  $T \ll \Delta$  because of the freezing out of thermally excited quasiparticles; and, as can be seen from Fig. 4, a well defined zero sound mode develops.

We thank the Alexander von Humboldt-Stiftung, the Deutsche Forschungsgemeinschaft (SFB279) and the STC for Superconductivity (NSF 91-20000) for their support.

## REFERENCES

1. E. V. Thuneberg, S.-K. Yip, M. Fogelström, and J. A. Sauls, Phys. Rev. Lett. submitted (1996); <http://xxx@lanl.gov/cond-mat/abs/9601148>.
2. D. Sprague *et al.*, Phys. Rev. Lett. **75**, 661 (1995); **77**, 4568 (1996).
3. J. A. Sauls, E. V. Thuneberg, and S.-K. Yip, preprint (1996).
4. K. Matsumoto *et al.*, Phys. Rev. Lett. **79**, 253 (1997).
5. J. M. Parpia, QFS97 invited talk (1997).
6. M. J. McKenna *et al.*, Rev. Lett. **66**, 1878 (1991).
7. N. Mulders *et al.*, Phys. Rev. Lett. **67**, 695 (1991).
8. A. Kopf and H. Brand, J. Low Temp. Phys. (1997).
9. P. Wölfle, in *Progress in Low Temperature Physics*, Vol. VIIA, ed. D. F. Brewer (North Holland, Amsterdam), 191 (1978).
10. W. P. Halperin and E. Varoquaux, in *Helium Three*, eds. W. P. Halperin and L. P. Pitaevskii (Elsevier, Amsterdam), 353 (1990).
11. R. H. McKenzie and J. A. Sauls, in *Helium Three*, eds. W. P. Halperin and L. P. Pitaevskii (Elsevier, Amsterdam), 255 (1990).
12. H. Kojima, *et al.*, Phys. Rev. Lett. **32**, 141 (1974).
13. C. H. Choi and P. Muzikar, Phys. Rev. B **37**, 5947 (1988).
14. J. W. Serene and D. Rainer, Phys. Rep. **101**, 221 (1983).
15. O. Betbeder-Matibet and P. Nozières, Ann. Phys. **51**, 392 (1969).