

Diffusion process in a deterministically forced flow

Piotr Garbaczewski

Institute of Theoretical Physics, University of Wrocław, pl. M. Borna 9,
PL-50 204 Wrocław, Poland

December 31, 2021

PACS numbers: 02.50.-r, 05.20.+j, 03.65.-w, 47.27.-i

Abstract

We analyze circumstances under which the microscopic dynamics of particles which are driven by a forced, gradient-type flow can be consistently interpreted as a Markovian diffusion process. Special attention is paid to discriminating between forces that are presumed to act selectively upon diffusing particles, while leaving the random medium statistically at rest (Smoluchowski diffusion processes), and those perturbing the random medium itself and thus creating the nontrivial flows. We focus on the deterministic "stirring" scenarios.

To analyze random perturbations that are either superimposed upon or are intrinsic to a driving deterministic motion, quite typically a configuration space equation

$$\dot{\mathbf{x}} = \mathbf{v}(\mathbf{x}; t) \quad (1)$$

is invoked, which is next replaced by a formal infinitesimal representation of an Itô diffusion process

$$d\mathbf{X}(t) = \mathbf{b}(\mathbf{X}(t); t)dt + \sqrt{2D} d\mathbf{W}(t) \quad (2)$$

Here, $\mathbf{W}(t)$ stands for the normalised Wiener noise, and D for a diffusion constant.

Presented by P. Garbaczewski at the X Symposium on Statistical Physics, Zakopane, September 1-10, 1997

The dynamical meaning of $\mathfrak{b}(\mathfrak{x};t)$, and thus reasons for making a substantial difference between the forward drift of the process and the driving velocity field (1), relies on a specific diffusion input and its possible phase-space (e.g. Langevin or that coming from deterministic dynamical systems) implementation, that entail a detailed functional relationship of $\mathfrak{v}(\mathfrak{x};t)$ and $\mathfrak{b}(\mathfrak{x};t)$, and justify such notions like: diffusion in an external force field, diffusion under various strains, diffusion along, against or across the driving deterministic flow, [1]. We shall not touch upon an important issue of diffusion under shear, [2], when nontrivial vortices may arise, by assuming from the very beginning that only the gradient velocity fields and deterministic forces are of interest for us in the present paper.

The pertinent mathematical formalism corroborates both the Brownian motion of a single particle in flows of various origin and the diffusive transport of neutrally buoyant components in flows of the hydrodynamic type. However, our major issue is a probabilistic interpretation of various linear and nonlinear partial differential equations of physical relevance, hence with a slightly abstract flavour put against the generally favoured practical reasoning. Expressing that in more physical terms, we address an old-fashioned problem of "how much nonlinear", "how much time-dependent", and generally "how much arbitrary" can be the driving velocity field to yield a consistent stochastic diffusion process (or the Langevin-type dynamics). Another issue is to get hints about a possible non-deterministic origin of such fields, [3].

Clearly, in random media that are statistically at rest, diffusion of single tracers or dispersion of pollutants are well described by the Fickian outcome of the molecular agitation, also in the presence of external force fields (then, in terms of Smoluchowski diffusions). On the other hand, it is of fundamental importance to understand how flows in a random medium (fluid, as example) affect dispersion. Such velocity fields are normally postulated as a priori given agents in the formalism and their (molecular or else) origin is disregarded. Moreover, usually the force exerted upon tracers is viewed independently from the forcing ("stirring") that might possibly perturb the random medium itself and create nontrivial (driving) flows.

Except for suitable continuity and growth restrictions, necessary to guarantee the existence of the process $X(t)$ governed by the Itô stochastic differential equation, the choice of the driving velocity field $\mathfrak{v}(\mathfrak{x};t)$ and hence of the related drift $\mathfrak{b}(\mathfrak{x};t)$ is normally (in typical physical problems) regarded to be arbitrary (except for being "not too nonlinear", see however at van Kampen's discussion of that issue in Ref. [1]).

The situation looks deceptively simple, [2], if we are (for example) interested in a diffusion process interpretation of passive tracers dynamics in the a priori given flow whose velocity field is a solution of the nonlinear partial differential equation, be it

Euler, Navier-Stokes, Burgers or the like. An implicit assumption, that passively buoyant tracers in a fluid have a negligible effect on the flow, looks acceptable (basically, in case when the concentration of a passive component in a flow is small). Then, one is tempted to view directly the fluid velocity field $\mathbf{v}(\mathbf{x};t)$ as the forward drift $\mathbf{b}(\mathbf{x};t)$ of the process, with the contaminant being diffusively dispersed along the stream lines.

Here, apparent problems arise: irrespectively of a specific physical context and the phenomenology (like e.g. the Boltzmann equation with its, as yet, not well understood Brownian motion approximation) standing behind the involved partial differential equations, some stringent mathematical criterions must be met to justify the diffusion process scenario, be it merely a crude approximation of reality.

Namely, in general, the assumed nonlinear evolution rule for $\mathbf{v}(\mathbf{x};t)$ must be checked against the dynamics that is allowed to govern the space-time dependence of the forward drift field $\mathbf{b}(\mathbf{x};t)$ of the pertinent process, [4], which is not at all arbitrary. The latter is ruled by standard consistency conditions that are respected by any Markovian diffusion process, and additionally by the rules of the forward and backward Itô calculus, [1, 4], the mathematical input that is frequently ignored in the physical literature.

Normally, the pragmatically oriented authors do not pay any attention to such problems and feel free to use any (deterministically or not) motivated velocity fields as forward drifts. In that case, serious troubles follow.

Indeed, the closely related issue we have analyzed before, [5], where as a by-product of the discussion, the forced Burgers dynamics

$$\partial_t \mathbf{v}_B + (\mathbf{v}_B \cdot \nabla) \mathbf{v}_B = D \nabla^2 \mathbf{v}_B + \tilde{\mathbf{f}} \quad (3)$$

and the diffusion-convection equation

$$\partial_t c + (\mathbf{v}_B \cdot \nabla) c = D \nabla^2 c \quad (4)$$

(originally, for the concentration $c(\mathbf{x};t)$ of a passive component in a flow), in case of gradient velocity fields, were found to be generic to a Markovian diffusion process input and as generically incompatible with the standard continuity equation in the compressible regime. In that case, the dynamics of concentration (in general this notion does not coincide with the probability density !) results from the stochastic process whose density $\mathcal{P}(\mathbf{x};t)$ evolves according to the standard Fokker-Planck equation

$$\partial_t \mathcal{P} = D \nabla^2 \mathcal{P} - \nabla \cdot (\tilde{\mathbf{f}} \mathcal{P}) ; \quad (5)$$

the forward drift solves an evolution equation:

$$\partial_t \mathbf{b} + (\mathbf{b} \cdot \nabla) \mathbf{b} = D \nabla^2 \mathbf{b} + \tilde{\mathbf{f}} ; \quad (6)$$

and there holds

$$\dot{\mathfrak{b}} = \mathfrak{v}_B + 2D \tilde{r} \ln : \quad (7)$$

By combining intuitions which underly the self-diffusion description, [6], with those appropriate for probabilistic solutions of the so-called Schrodinger boundary-data and next-interpolation problem, [5, 7, 8], the above argument can be generalized to arbitrary conservatively forced diffusion processes, quite irrespectively of a physical context in which their usage can be justified.

Namely, let us consider a density $\rho(\mathbf{x};t); t \geq 0$ of a stochastic diffusion process, solving the Fokker-Planck equation (5). For drifts that are gradient fields, the potential in Eqs. (3) and (6) (whatever its functional form is), must allow for a representation formula, reminiscent of the probabilistic Cameron-Martin-Girsanov transformation:

$$\rho(\mathbf{x};t) = 2D \left[\partial_t + \frac{1}{2} \left(\frac{\mathfrak{b}^2}{2D} + \tilde{r} \cdot \mathfrak{b} \right) \right] ; \quad (8)$$

where $\mathfrak{b}(\mathbf{x};t) = 2D \tilde{r}(\mathbf{x};t)$. The formula (8) is a trivial identity, if we take for granted that all drifts are known from the beginning, like in case of typical Smoluchowski diffusions. Nonetheless, we always end up with a concrete space-time dependent function $\rho(\mathbf{x};t)$ which enters the partial differential equation (6). If we take Eq. (6) as a starting point with ρ a priori given, its solutions may be then sought for in turn (to yield the previous a priori given drifts, if the procedure is consistent).

Also, the functional properties of $\rho(\mathbf{x};t)$ are not an innocent feature of the formalism, since for the existence of the Markovian diffusion process with the forward drift $\mathfrak{b}(\mathbf{x};t)$, we must resort to potentials $\rho(\mathbf{x};t)$ that are not completely arbitrary functions. Technically, [7], the minimal requirement is that the admissible potential is a bounded from below continuous function. This restriction will have profound consequences for our further discussion of diffusion in a flow, although nothing serious happens if ρ is bounded and, for example, is the periodic space-time function.

Remark 1: If we set $\rho = \rho_1 + \rho_2$, and demand that $\rho_1 \in \mathcal{C}$ solves the Fokker-Planck equation with the very same drift $\mathfrak{b}(\mathbf{x};t)$ as ρ does, then as a necessary consequence of the general formalism, [5, 7], the concentration $c(\mathbf{x};t) = \frac{\rho_2(\mathbf{x};t)}{\rho(\mathbf{x};t)}$ solves an associated diffusion-convection equation $\partial_t c + (\mathfrak{v}_B - \tilde{r})c = D \Delta c$. Here, the flow velocity $\mathfrak{v}_B(\mathbf{x};t)$ coincides with the backward drift of the generic diffusion process with the density $\rho(\mathbf{x};t)$.

We should clearly discriminate between forces whose effect is a "stirring" of the random medium and those acting selectively on diffusing particles, with a negligible effect on the medium itself. For example, the traditional Smoluchowski diffusion processes in conservative force fields are considered in random media that are sta-

tistically at rest. Following the standard (phase-space, Langevin) methodology, let us set $\dot{\mathbf{x}} = -\frac{1}{\gamma} \mathbf{K}(\mathbf{x})$, where γ is a (large) friction coefficient and \mathbf{K} represents an external Newtonian force per unit of mass (e.g. an acceleration) that is of gradient form, $\mathbf{K} = -\nabla U$. Then, the effective potential reads:

$$U_{\text{eff}} = \frac{\gamma^2}{2} + \frac{D}{\gamma} \nabla \cdot \mathbf{K} \quad (9)$$

and the only distinction between the attractive or repulsive cases can be read out from the term $\nabla \cdot \mathbf{K}$. For example, the harmonic attraction/repulsion $\mathbf{K} = -\mathbf{x}$; $\gamma > 0$ would give rise to a harmonic repulsion, if interpreted in terms of $\nabla \cdot \mathbf{K}$, in view of $\nabla \cdot (-\mathbf{x}) = -3$. The situation would not change under the incompressibility condition (cf. also the probabilistic approaches to the Euler, Navier-Stokes and Boltzmann equations, [9]).

Notice that by formally changing a sign of γ we would arrive at the attractive variant of the problem, which is however incompatible with the diffusion process scenario in view of the unboundedness of U_{eff} from below.

We have thus arrived at the major point of our discussion: a priori, there is no way to incorporate the attractive forces which affect (drive) the flow and nonetheless generate a consistent diffusion-in-a-flow transport.

Clearly, there is no reason to exclude the attractive variants of the potential from considerations, since the deterministic motion is consistent with them.

Concluding, if the diffusion is to be involved we need to save the situation somehow, and this can be made only by incorporating the hitherto not considered "pressure" term effects. That is suggested by the general form of the compressible Euler ($\mathbf{F} = -\nabla V$ stands for external volume forces and ρ for the fluid density that itself undergoes a stochastic diffusion process):

$$\partial_t \mathbf{v}_E + (\mathbf{v}_E \cdot \nabla) \mathbf{v}_E = \mathbf{F} - \frac{1}{\rho} \nabla P \quad (10)$$

or the incompressible, [9], Navier-Stokes equation:

$$\partial_t \mathbf{v}_{NS} + (\mathbf{v}_{NS} \cdot \nabla) \mathbf{v}_{NS} = -\frac{1}{\rho} \nabla P + \nu \nabla^2 \mathbf{v}_{NS} ; \quad (11)$$

both to be compared with the equations (1) and (4), that set dynamical constraints for respectively backward and forward drifts of a Markovian diffusion process?

Let us stress again that the acceleration term \mathbf{F} in equations (10) and (11) normally is regarded as arbitrary, while the corresponding term $\nabla \cdot \mathbf{K}$ in (3), (6) and (8) involves a bounded from below function.

Since, in case of gradient velocity fields, the dissipation term in the incompressible Navier-Stokes equation (11) identically vanishes, we should concentrate on analyzing the possible "forward drift of the Markovian process" meaning of the Euler flow with the velocity field \mathbf{v}_E , (10).

At this point it is useful, at least on the formal grounds, to invoke the standard phase-space argument that is valid for a Markovian diffusion process taking place in a given flow $\mathbf{v}(\mathbf{x};t)$ with as yet unspecified dynamics nor physical origin. We account for an explicit force exerted upon diffusing particles, while not necessarily directly affecting the driving flow itself. Namely, [2, 4], let us set for infinitesimal increments of phase space random variables:

$$\begin{aligned} d\mathbf{X}(t) &= \mathbf{V}(t)dt \\ d\mathbf{V}(t) &= [\mathbf{v}(\mathbf{x};t) - \mathbf{V}(t)]dt + \sqrt{\frac{P}{2D}} d\mathbf{W}(t) : \end{aligned} \quad (12)$$

Following the leading idea of the Smoluchowski approximation, we assume that β is large, and consider the process for times significantly exceeding β^{-1} . Then, an appropriate choice of the velocity field $\mathbf{v}(\mathbf{x};t)$ (boundedness and growth restrictions are involved) may in principle guarantee, [4], the convergence of the spatial part $\mathbf{X}(t)$ of the process (12) to the Itô diffusion process with infinitesimal increments (where the force effects can be safely ignored if we are interested mostly in the driving motion):

$$d\mathbf{X}(t) = \mathbf{v}(\mathbf{x};t)dt + \sqrt{\frac{P}{2D}} d\mathbf{W}(t) : \quad (13)$$

However, one cannot blindly insert in the place of the forward drift $\mathbf{v}(\mathbf{x};t)$ any of the previously considered bulk velocity fields, without going into apparent contradictions. Specially, the equation (4) with $\mathbf{v}(\mathbf{x};t) \propto \mathbf{b}(\mathbf{x};t)$ must be valid.

By resorting to velocity fields $\mathbf{v}(\mathbf{x};t)$ which obey $\nabla \cdot \mathbf{v}(\mathbf{x};t) = 0$, we may pass from (6) to an equation of the Euler form, (10), provided (8) holds true and then the right-hand-side of (6) involves a bounded from below effective potential.

An additional requirement is that

$$\mathbf{F} - \frac{1}{\beta} \nabla P \stackrel{!}{=} \mathbf{f} : \quad (14)$$

Clearly, in case of a constant pressure we are left with the dynamical constraint $(\mathbf{b} \propto \mathbf{v}_E)$:

$$\partial_t \mathbf{b} + (\mathbf{b} \cdot \nabla) \mathbf{b} = \mathbf{F} = \mathbf{f} \quad (15)$$

combining simultaneously the Eulerian fluid and the Markov diffusion process inputs, if and only if \mathbf{F} is repulsive, e.g. $\nabla \cdot \mathbf{V}(\mathbf{x};t)$ is bounded from below. Quite

analogously, by setting $F = 0$, we would get a constraint on the admissible pressure term, in view of:

$$\partial_t \tilde{b} + (\tilde{b} \cdot \tilde{r}) \tilde{b} = -\frac{1}{\tilde{r}} \tilde{r} P = \tilde{r} \quad : \quad (16)$$

Both, in cases (15), (16) the effective potential must respect the functional dependence (on a forward drift and its potential) prescription (8). In addition, the Fokker-Planck equation (5) with the forward drift $\mathbf{v}_E(\mathbf{x};t) = \tilde{b}(\mathbf{x};t)$ must be valid for the density $\tilde{b}(\mathbf{x};t)$.

To our knowledge, in the literature there is only one known specific class of Markovian diffusion processes that would render the right-hand-side of Eq. (10) repulsive but nevertheless account for the troublesome Newtonian accelerations, e.g. those of the form $\tilde{r} V$, with $+V$ bounded from below. Such processes have forward drifts that for each suitable, bounded from below function $V(\mathbf{x})$ solve the nonlinear partial differential equation:

$$\partial_t \tilde{b} + (\tilde{b} \cdot \tilde{r}) \tilde{b} = -D \Delta \tilde{b} + \tilde{r} (2Q - V) \quad (17)$$

with the compensating pressure term :

$$Q = -2D^2 \frac{\Delta \tilde{b}}{\tilde{b}} = -\frac{1}{2} \tilde{u}^2 + D \tilde{r} \cdot \tilde{u} \quad (18)$$

$$\tilde{u}(\mathbf{x};t) = D \tilde{r} \ln \tilde{b}(\mathbf{x};t)$$

Their discussion can be found in Refs. [4, 5, 7, 8].

Clearly, we have:

$$F = \tilde{r} V ; \tilde{r} 2Q = -\frac{1}{\tilde{r}} \tilde{r} P \quad (19)$$

where:

$$P(\mathbf{x};t) = -2D^2 \tilde{b}(\mathbf{x};t) \ln \tilde{b}(\mathbf{x};t) \quad (20)$$

Effectively, P is here defined up to a time-dependent constant. Another admissible form of the pressure term reads (summation convention is implicit):

$$-\frac{1}{\tilde{r}} \tilde{r}_k [(2D^2 \partial_j \partial_k) \ln \tilde{b}] = \tilde{r}_j (2Q) \quad (21)$$

.

If we consider a subclass of processes for which the dissipation term identically vanishes (a number of examples can be found in Refs. [7]):

$$\Delta \tilde{b}(\mathbf{x};t) = 0 \quad (22)$$

the equation (17) takes a conspicuous Euler form (10), $\nabla_E \cdot \mathbf{v} = 0$.

Let us notice that (20), (21) provide for a generalisation of the more familiar, thermodynamically motivated and suited for ideal gases and fluids, equation of state $P = P(\rho)$. In case of density fields for which $\rho \propto \exp(-\phi)$, the standard relationship between the pressure and the density is reproduced. In case of density fields obeying $\rho \propto \exp(-\phi) = 0$, we are left with at most purely time dependent or a constant pressure. Pressure profiles may be highly complex for arbitrarily chosen initial density and/or the flow velocity fields.

To conclude the present discussion let us invoke Refs. [9, 6, 7]. The problem of a diffusion process interpretation of various partial differential equations is known to extend beyond the original parabolic equations setting, to general nonlinear velocity field equations. On the other hand, the nonlinear Markov jump processes associated with the Boltzmann equation, in the hydrodynamic limit, are believed to imply either an ordinary deterministic dynamics with the velocity field solving the Euler equation, or a diffusion process whose drift is a solution of the incompressible Navier-Stokes equation (in general, without our $\nabla \cdot \mathbf{v} = 0$ restriction), [6, 9]. The case of arbitrary external forcing has never been satisfactorily solved.

Our reasoning went otherwise. We asked for the admissible space-time dependence of general velocity fields that are to play the rôle of forward drifts of Markovian diffusion processes, and at the same time can be met in physically significant contexts. Therefore various forms of the Fokker-Planck equation for tracers driven by familiar compressible velocity fields were discussed.

Our finding is that solutions of the compressible Euler equation are appropriate for the description of the general non-deterministic (e.g. random and Markovian) dynamics running under the influence of both attractive and repulsive stirring forces, and refer to a class of Markovian diffusion processes originally introduced by E. Nelson, [4, 7, 3]. That involves only the gradient velocity fields (a couple of issues concerning the $\nabla \cdot \mathbf{v} \neq 0$ velocity fields and their nonconservative forcing have been raised in Refs. [5]).

Remark 2: Let us stress that a standard justification of the hydrodynamic limit for a tracer particle invokes a Brownian particle in an equilibrium fluid. An issue of how much the tracer particle disturbs the fluid (random medium) locally and how far away from the tracer particle the thermal equilibrium conditions regain their validity, [6], normally is disregarded. Moreover, in the standard derivation of local conservation laws from the Boltzmann equation, the forcing term on the right-hand-side of the Euler or Navier-Stokes equation up to scalings does coincide with the force acting on each single particle comprising the system. Thus, in this framework, there is no room for any discrimination between forces acting upon tagged particles and those perturbing the spatial flows (once on the level of local averages).

Quite on the contrary, the force term in the Kramers equation and this appearing in the related local conservation law for the forward drift or for the current velocity of the diffusion process are known not to coincide in general. Typically, the action of an external force is confined to diffusing (tagged) particles with no global or local effect on the surrounding random medium, cf. standard derivations of the Smoluchowski equation. This feature underlies problems with the diffusion process interpretation of general partial differential equations governing physically relevant velocity fields. Specifically, any external intervention (forcing) upon a stochastically evolving (in the diffusion process approximation) system gives rise to a perturbation of local flows, which seldom can be analyzed as forcing of any definite type on the molecular level. The Smoluchowski theory is here a notable exception, but there one has no room for genuine flows and velocity field profiles which are generated in the random medium.

Remark 3: It seems worthwhile to mention a close connection of the considered framework with the general issue of executing small random perturbations on the level of the classical Hamilton-Jacobi dynamics, [8], with the related issue of an optimal control of stochastic processes and with that of the so-called "viscosity solutions" of the Hamilton-Jacobi equation, [10]. In fact, our (Feynman-Kac, see [7]) potentials (8), (9) were introduced on the basis of probabilistic arguments via the Girsanov or Cameron-Martin theorems about transformations of drifts of the diffusion process. However, an implicit assumption that drifts are defined in terms of gradients of suitable logarithmic functions: $\mathbf{b} = 2D \tilde{\mathbf{r}} \log$ and $\mathbf{v}_b = \mathbf{b} = 2D \tilde{\mathbf{r}} \log$ (here, we employ the notation of our previous publications, [7], where $\tilde{\mathbf{r}}$ is a bounded solution of the forward generalized diffusion equation, while $\tilde{\mathbf{r}}$ that of its time adjoint) implies that the compatibility condition (8) can be rewritten in two equivalent forms, both involving the modified Hamilton-Jacobi equations. Namely, let us set $2D \log =$ and $2D \log =$. Then, we have: $= \partial_t + \frac{1}{2} \tilde{\mathbf{r}} \cdot \mathbf{j}^2 + D \Delta$ and at the same time $= \partial_t + \frac{1}{2} \tilde{\mathbf{r}} \cdot \mathbf{j}^2 - D \Delta$. The latter one is identified as the so-called Hamilton-Jacobi-Bellman programming equation in the optimal control of stochastic diffusion processes, [10, 8], and via the Hopf-Cole logarithmic transformation (take the gradient) is linked to the Burgers equation (3). An issue of viscosity solutions of the standard Hamilton-Jacobi equation has been extensively studied in the literature as the $D \rightarrow 0$ limit of solutions of the modified (e.g. Bellman) equation. It is thus clear, on the basis of our previous discussion, that an apparent obstacle is hidden in the assumption that a diffusion process is involved. Then, suitable restrictions upon $\tilde{\mathbf{r}}$ must be respected, and the attractive versus repulsive potential problem appears.

References

- [1] W . Horsthem ke, R . Lefever, "Noise-Induced Transitions", Springer-Verlag, Berlin, 1984; Z . Schuss, "Theory and Applications of Stochastic Differential Equations", Wiley, NY, 1980; N . G . van K am pen, "Stochastic Processes in Physics and Chemistry", North-Holland, Amsterdam, 1987
- [2] F . Sagues, W . Horsthem ke, Phys. Rev. A 34, (1986), 4136; Y . Katayama, R . Terauti, Eur. J. Phys. 17, (1996), 136; R . T . Foerster, T . G . M . van den Ven, J. Fluid Mech. 96, (1980), 105; A . S . Monin, A . M . Yaglom, "Statistical Fluid Mechanics", MIT Press, Cambridge, 1973
- [3] P . Garbaczewski, J . P . Vigier, Phys. Rev. A 46, (1992), 4634; P . Garbaczewski, pp. 169 in: "Coherent States: Past, Present, and Future", eds. D . H . Feng et al, World Scientific, Singapore, 1994
- [4] A . I . Ilin, R . Z . Khasinski, Theory Prob. Appl., 9, (1964), 466; E . Nelson, "Dynamical Theories of the Brownian Motion", Princeton, University Press, Princeton, 1967; E . Nelson, "Quantum Fluctuations", Princeton University, Princeton, 1985
- [5] P . Garbaczewski, G . Kondrat, R . Olkiewicz, Phys. Rev. E 55, (1997), 1401
- [6] H . Spohn, "Large Scale Dynamics of Interacting Particles", Springer, Berlin, 1992; J . L . Lebowitz, H . Spohn, J. Stat. Phys. 28, (1982), 539
- [7] Ph . Blanchard, P . Garbaczewski, Phys. Rev. E 49, (1994), 3815; P . Garbaczewski, R . Olkiewicz, J. Math. Phys. 37, (1996), 730;
- [8] J . C . Zambrini, J. Math. Phys. 27, (1986), 3207; J . C . Zambrini, pp. 393, in: "Chaos-The Interplay Between Stochastic and Deterministic Behaviour", eds. P . Garbaczewski, M . Wolf, A . Weron, Lecture Notes in Physics vol. 457, Springer-Verlag, Berlin, 1995
- [9] C . Constantini, R . Marra, J. Stat. Phys. 67, (1992), 229; C . Beck, G . Roepstor, Physica A 165, (1990), 270; R . Marra, pp. 564 in: "Stochastic Processes, Physics and geometry", eds. S . Albeverio et al, World Scientific, Singapore, 1989; C . Marchioro, M . Pulvirenti, "Vortex Methods in Two-Dimensional Fluid Dynamics", Lecture Notes in Physics vol. 203, Springer-Verlag, Berlin, 1984
- [10] W . H . Fleming, H . M . Soner, "Controlled Markov Processes and Viscosity Solutions", Springer-Verlag, Berlin, 1993