C ritical P roperties of Spectral Functions for the 1D Anisotropic t-J M odels with an Energy G ap

Tatsuya Fujii, Yasum asa Tsukam oto and Norio Kawakam i Department of Applied Physics, Osaka University, Suita, Osaka 565, Japan (June 7, 1997)

We exactly calculate the momentum-dependent critical exponents for spectral functions in the one-dimensional anisotropic t-J m odels with a gap either in the spin or charge excitation spectrum. Our approach is based on the Bethe ansatz technique combined with nite-size scaling techniques in conform al eld theory. It is found that the spectral functions show a power-law singularity, which occurs at frequencies determ ined by the dispersion of a massive spin (or charge) excitation. We discuss how the nontrivial contribution of a m assive excitation controls the singular behavior in optical response functions.

PACS: 75.10.Jm, 05.30.-d, 03.65.Sq

The dynam ical spectral properties of one-dim ensional (1D) highly correlated electron systems have been intensively studied. For example, there are a number of recent experiments on angular resolved photoem ission, [1 {5] which have revealed som e striking properties inherent in 1D systems, such as the spin-charge separation. [5] A lso, extensive theoretical studies on spectral functions have been done num erically [6{11] and analytically. [12{14]

A remarkable point in the recent photoem ission experim ent [2] for the insulator com pound SrC uO 2 is that it has successfully explored characteristic properties of a single hole doped in the 1D Mott insulator. In such insulating systems, only the spin sector is massless (Tom onaga-Luttinger (TL) liquid), while the charge sector is massive. This type of 1D electron system is sometimes referred to as the Luther-Em ery class, for which either spin or charge excitation is massive. It was naively believed that the one-particle G reen function for such system s with m assive excitations would show exponential decay instead of the power-law singularity. However, Sorella and Parola, [15,16] and Voit [17] theoretically predicted that the dynam ical spectral functions for these systems still show the power-law singularity at threshold energies, characterizing the critical properties of a doped hole in the 1D Mott insulator. In particular, Sorella and Parola [16] have demonstrated the above point by using the exact solution of the supersymmetric t-J model and conform al eld theory (CFT).

M otivated by the above investigations, we study the spectral properties of 1D highly correlated electron systems with a gap either in the spin or charge excitation spectrum. For this purpose, we consider two types of instructive models for 1D correlated electron systems: the rst model is the integrable an isotropic t-J model [18] away from half lling, for which the charge (spin) sector is massless (massive). The other one is the holedoped XXZ Heisenberg spin chain, which is realized by the anisotropic t-J m odel in the lim it of half lling with the condition J! 0. We explore this model as a typical system which has massless spin (massive charge) excitations. We exactly calculate the critical exponents of spectral functions for both of these models, by exploiting nite-size scaling ideas in CFT.W e discuss the critical

properties of the optical response functions in terms of the mom entum -dependent critical exponents obtained in this paper.

Let us start with the 1D anisotropic t-J m odel, which possesses a gap for spin excitations. This model is a deformed supersymmetric t-J model which was exactly solved by Barriev, [18]

$$H_{1} = \begin{pmatrix} X \\ P (C_{i;}^{y} C_{i+1}; + C_{i+1}^{y}, C_{i};)P \\ + \begin{pmatrix} X \\ C_{i;\#}^{i} C_{i;"} C_{i+1;"}^{y} C_{i+1;\#} + C_{i;"}^{y} C_{i;\#} C_{i+1;\#}^{y} C_{i+1;"} \\ & e n_{i;\#} n_{i+1;"} e n_{i;"} n_{i+1;\#}); \quad (1)$$

where the operator $P = \prod_{j=1}^{N} (1 n_{j,"} n_{j,\#})$ projects out the Hilbert space without double occupancy at all sites, is a deformation parameter which parametrizes and the anisotropy of the interaction. In the isotropic lim it (! 0), the Ham iltonian (1) reduces to the conventional supersymmetric t-J model which was exactly solved, [19{22] and its conform alproperties were clari ed. [23]

It is known that the spin excitation for (1) is always massive, while the charge excitation is massless away from half lling. [18,24] Hence, this model allows us to discuss the spectral properties of the system with the m assless charge and m assive spin excitations. We wish to emphasize here that although the following analysis is performed for the above speci c m odel, the obtained results can be generally applied to 1D correlated electron m odels with the spin gap, such as the attractive H ubbard m odel.

In the following, we derive the one-particle G reen function in the low-energy regime and the corresponding critical exponents. To this end, we derive the relevant Bethe equations which specify excitations when one electron is removed from the system . These equations can be straightforwardly deduced from those obtained by

Bariev. [18] The result reads

$$\frac{\sin((+ i))}{\sin((i))}^{N} = \frac{\sin((+ i = 2))}{\sin((i = 2))}$$

$$\frac{\Psi}{\sin((+ i))}^{N}; \quad (2)$$

where N (M) is the number of sites (down-spin electrons), and (= 1;2;:::;M) are charge rapidities which classify collective charge excitations. Note that a spin rapidity also appears in the above equation, which characterizes a massive spin excitation induced by the injected hole. A lthough massive spin excitations do not directly enter in the low-energy physics, they control the critical behavior of massless charge excitations via the nontrivial coupling between charge and spin sectors represented by (2), as will be shown momentarily.

In order to apply the sophisticated m ethods developed in CFT, [25] we now analyze the nite-size corrections to the ground-state energy and excitation energy including the e ect of the massive spin sector. [26,27] This manipulation enables us to exactly calculate the critical exponents for spectral functions in the low energy regime. The calculation is straightforwardly performed in a standard manner, [28(30] to yield the following expressions, which are typical for TL liquids classi ed by c = 1 CFT,

$$E_{0} = N_{0} - \frac{v_{c}}{6N};$$

$$E_{0} = s_{s} + c_{c} = e_{s}() + \frac{2 v_{c}}{N}x_{c};$$
(3)

where N $_0$ is the ground-state energy in the therm odynamic limit. The N-independent term $_s = e_s$ (socalled surface energy) gives an excitation energy for the massive spin sector. It is clearly seen that the spin excitation $_s$ is decoupled from the charge excitation $_c$. A ccording to the nite-size scaling idea in CFT, the scaling dimension for the charge sector is read from the universall=N corrections to the excitation energy $_c$, which is given as

$$x_{c} = \frac{1}{4 \frac{2}{c}} (M_{c} + n^{(s)})^{2} + \frac{2}{c} (D_{c} + d^{(s)})^{2};$$
 (4)

where D $_{\rm c}$ and M $_{\rm c}$ are the quantum numbers that label current excitations and charged excitations, which are subjected to the selection rule for fermions, D $_{\rm c}$ =

M $_{\rm c}{=}2$. The quantity $_{\rm c}{=}$ $_{\rm c}$ (v_0), often referred to as the dressed charge, is given as [18]

$${}_{c}(v) = 1 \qquad \qquad \begin{array}{c} Z & {}_{v_{0}} & Z \\ & + & \\ & {}_{v_{0}} & \end{array} \frac{1}{2} \quad {}^{\circ}(v \quad v_{j}^{\circ}) \ {}_{c}(v^{\circ}) dv^{\circ};$$
 (5)

where

$$(v;) = 2 \tan^{-1} (\operatorname{coth} () \tan (v=2));$$
 (6)

Note that the dressed charge features the U (1) critical line of c = 1 CFT when we change the strength of interaction or the density of electrons. Here, we stress that the scaling dimension x_c for the charge sector com – pletely determines the critical behavior of the spectral functions. Note, however, that two key quantities n^(s) and d^(s), which contain the e ect of the massive spin sector, are introduced in (4)

$$n^{(s)} = \begin{array}{ccc} Z & & Z \\ & & & \\ n^{(s)} = \begin{array}{c} + & & \\ & & \\ \end{array} \\ d^{(s)} = \begin{array}{c} \frac{1}{2} \left(z_{s} \left(\right) + z_{s} \left(\right) \right) \\ & & \\ & + \begin{array}{c} \frac{1}{2} \end{array} \\ & & \\ \end{array} \\ & & \\ \end{array} \\ \begin{array}{c} v_{0} \\ & \\ \end{array} \\ & & \\ \end{array} \\ \begin{array}{c} v_{0} \\ & \\ \end{array} \\ & & \\ \end{array} \\ (v) dv; \\ \end{array}$$

where

$$z_{s}(v) = (v_{Z} 2; =2) + (v v';)_{s}(v')dv';$$

$$s(v) = \frac{1}{2} v_{v_{0}}(v_{2}; =2) + \frac{1}{2} v_{v_{0}}(v_{v_{0}} 2; =2) + \frac{1}{2} v'(v_{v_{0}}, v';)_{s}(v')dv';$$

These quantities are alternatively represented in terms of the phase shifts $_L$ and $_R$ at the left and right Ferm i points in them assless charge sector: $n^{(s)} = _R + _L d^{(s)} =$

 $_{\rm R}$ $_{\rm L}$. A lthough the scaling dimension $x_{\rm c}$ reects c = 1 CFT for the massless sector, the massive spin sector also contributes to the scaling dimension via n $^{\rm (s)}$ and d $^{\rm (s)}$, as seen in (4). Note that the eq.(4) for scaling dimensions is typical for shifted c = 1 CFT, whose xed point is di erent from that of the static in purity problem, as pointed out by Sorella and Parola. [16,31]

W e are now ready to investigate the critical properties of the one-particle G reen function G (x;t). A coording to the spin-charge separation in elementary excitations for 1D system s, it is given by,

$$G (x;t) = i X (t)e^{i st i p_{s}x} < {}_{0}j^{\gamma}(0)j > < j^{(0)}j_{0} > e^{i ct i p_{c}x} = G_{s}(x;t) G_{c}(x;t):$$
(8)

As is well known, the G reen function G_s for the m assive spin sector gives the exponential decay, e.g., in the long-distance behavior of the equal-time G reen function G (x;t = 0). However, this is not true for the dynamical G reen function G (k;!) around ! = !s, for which the m assive spin excitation w_s determ ines the shift of the energy, whereas the m assless charge excitation still causes the infrared singularity, resulting in the power-law behavior around ! = !s. In order to clarify how such a

power-law singularity appears, we rst concentrate on the low-energy behavior of the G reen function G $_{\rm c}$ in the m assless charge sector. Exploiting nite-size scaling techniques in CFT, we can write down its asymptotic form as

$$G_{c}(x;t) = \frac{e^{i\mathcal{Q}_{c}x}}{(x - i\mathcal{U}_{c}t)^{2} + (x + i\mathcal{V}_{c}t)^{2}};$$
 (9)

where v_c is the velocity of charge excitations and $Q_c = 2Q_F (D_c + d^{(s)})$. Here are conformal dimensions which are related to the scaling dimension x_c ,

$$x_c = + + :$$
 (10)

To get the critical exponent for the one-particle G reen function, we have to set M $_{\rm c}$ = 1 and D $_{\rm c}$ = 1=2 in (4).

The spectral function A(k;!) is now given via the Fourier transform ation as

$$A(k;!) = \frac{1}{2} \operatorname{Im} G(k;!) \quad (! \quad !_{s}(k \quad Q_{c}))^{X(k)}; \quad (11)$$

with the critical exponent X $(k) = 2x_c$ 1. It is seen from this formula that the most relevant singularity in the spectral function occurs at frequencies determined by the massive spin spectrum $!_s (k \quad Q_c)$, and the corresponding critical exponent X (k) is dependent on the momentum k of an injected hole. Note that the quantities which give rise to the momentum dependence of the critical exponent are the phase shifts n^(s) and d^(s).

The momentum dependence of a critical exponent X (k) is shown in Fig. 1. Here the momentum is plotted in the unit of inverse lattice spacing 1=a. It is clear that X (k) has values with either a positive or a negative sign, which respectively results in convergence or divergence power-law behavior around the threshold energy Q_c). Therefore, we can say that the spectral in-!₅(k tensity around the threshold would be strongly momentum dependent. In particular, for the region exhibiting convergence power-law behavior, i.e. for X (k) > 0, the edge singularity is completely sm eared, and m ay not be observed clearly in experiments. We note that the plateaulike structures in part of the momentum region in Fig. 1 for the case of small (close to isotropic case) re ect the fact that n^(s) is weakly dependent on the momentum for the case of 0.

W e now proceed to another interesting an isotropic t-J m odel,

$$H_{2} = tPf (C_{i;}^{y} C_{i+1;} + C_{i+1;}^{y} C_{i;}) gP$$

$$X \stackrel{i;}{J} f_{i}^{x} \stackrel{x}{_{i+1}} + \frac{y}{_{i}} \stackrel{y}{_{i+1}} + \cos(2) (\stackrel{z}{_{i}} \stackrel{z}{_{i+1}} 1)$$

$$+ H (\stackrel{i}{_{i}} 1)g;$$

where are Paulim atrices, is a non-negative quantity which param etrizes the anisotropy, and H is an applied m agnetic eld. This model is realized by doping holes into the integrable 1D XXZ spin model. [32] If the condition J=t 1 is satisfied, we can perform the exact evaluation of the critical exponents based on the idea of CFT. In the following, we wish to deal with the case where a single hole is doped into the above insulating XXZ spin chain, for which the spin excitation still remains massless. Note that the present system belongs to the same universality class as the repulsive H ubbard model at half lling, and the results obtained below should be complementary to those of Sorella and P arola for the supersymmetric case. [16]

We now investigate the critical behavior of the spectral function at half lling in the lim it $J \\ 0$. Repeating the same calculation as for the previous model (1), we immediately end up with the following scaling dimension for spin excitations,

$$x_{s} = \frac{1}{4 \frac{2}{s}} (M_{s} + n^{(c)})^{2} + \frac{2}{s} (D_{s} + d^{(c)})^{2}; \quad (13)$$

where the quantum numbers have been chosen as M $_{\rm s}$ =

1 and D $_{s}$ = 1=2 to satisfy the ferm ion selection rule D $_{s}$ = M $_{s}$ =2. The dressed charge $_{s}$ = $_{s}$ ($_{0}$), is given by

$$_{s}() = 1 + \int_{0}^{Z} \frac{1}{2} K (;)_{s}() d;$$
(14)

where

$$K(i;) = \frac{\sin 4}{\sinh(i + i2)\sinh(i - i2)};$$
 (15)

The charge contribution to the massless spin sector is explicitly written down in terms of two phase shifts

$$n^{(c)} = \frac{1}{2} (m + 1); \quad d^{(c)} = 0;$$
 (16)

where m is the magnetization of the system in the unit of $g_B = 1$. Note that the spin and charge sectors in the present model exchange their role compared with the t-J model (1). It turns out that in the lim it J! 0, n^(c) and d^(c) do not depend on the momentum of an injected hole, in contrast to the model (1). The critical behavior of the spectral function A (k;!) is determined by the above scaling dimension,

$$A(k;!) = \frac{1}{2} \operatorname{Im} G(k;!) \quad (! \quad !_{c} (k \quad Q_{s}))^{X}; \quad (17)$$

where $Q_s = 2Q_F$ ($D_s + d^{(c)}$) and $!_c(k) = 2\cos(k)$ is the excitation spectrum for the massive charge sector.

(12) The computed critical exponents X are shown in Fig. 2 as a function of the magnetization. For the completely polarized case (m = 1), the dressed charge _s equals to unity as in the noninteracting case. Therefore, the

critical exponent takes the same constant value, 1=2, even if the anisotropy parameter is changed. In the lim it ! =4 the XXZ m odel reduces to the XY m odel, which is nothing but a free ferm ion m odel. A lthough it is naively expected that the critical exponent for such a free system is trivial, the presence of two phase shifts n^(c) and d^(c) result in the nontrivial exponents, as is the case for the ordinary X-ray singularity problem. In the opposite lim it, ! =2, the XXZ m odel reduces to the XXX m odel. The resulting critical exponent 1=2 at m = 0 exactly coincides with the one obtained by Sorella and P arola, [15] and Voit. [17]

In sum m ary, we have studied the critical properties of spectral functions for the 1D anisotropic t-J m odels com – bining the Bethe ansatz technique with nite-size scaling m ethods in CFT. In particular, we have focused on the m odels possessing an excitation gap either in the spin or charge spectrum by exploiting the two anisotropic t-J m odels. By investigating critical properties exactly, we have seen that spectral functions show power-law behavior around the threshold energies, as found by Sorella and Parola for the isotropic supersym m etric m odel. The singularity in spectral functions occurs at frequencies determ ined by the dispersion of a m assive excitation, re-

ecting the nontrivial interaction between two elementary excitations. It is interesting to extend the present analysis to the case with orbital degeneracy, which would provide useful information about optical response functions for som e M n oxides, which have been investigated intensively in recent years.

W e would like to thank S.Fujim oto, T.Fukui and S. K um ada for their valuable discussions. This work was partly supported by a G rant-in-A id from the M inistry of E ducation, Science, Sport and Culture, Japan.

- [L] B. Dardel, D. Malterre, M. Grioni, P. Weibel, Y. Bear and F. Levy: Phys. Rev. Lett. 67 (1991) 3144.
- [2] Y. Hwu, P. Almeras, M. Marsi, H. Berger, F. Levy, M. Grioni, D. Malterre and G. Margaritondo: Phys. Rev. B 46 (1992) 13624.
- [3] C. Coluzza, H. Berger, P. Almeras, F. Gozzo, G. Margaritondo, L. Forro and Y. Hwu: Phys. Rev. B 47 (1993) 6625.
- [4] M. Nakamura, A. Sekiyama, H. Namatame, A. Fujimori,
 H. Yoshihara, T. Ohtani, A. M isu and M. Takano: Phys.
 Rev. B 49 (1994) 16191.
- [5] C.Kim, A.Y.Matsuura, Z.-X. Shen, N.Motoyama, H. Eisaki, S.Uchida, T.Tohyama and S.Maekawa: Phys. Rev.Lett. 77 (1996) 4054.
- [6] R. Preuss, A. Muram atsu, W. von der Linden, P. Dieterich, F. F. A ssaad and W. Hanke: Phys. Rev. Lett. 73 (1994) 732.

- [7] S.Haas and E.Daggoto: Phys.Rev.B 52 (1995) 14396.
- [8] T. Tohyam a and S. Maekawa: J. Phys. Soc. Jpn. 65 (1990) 1902.
- [9] J. Lorenzana and R. Eder: preprint cond-m at/9608035.
- [10] J. Favand, S. Haas, K. Penc, F. M ila and E. D agotto: preprint cond-m at/9611223.
- [11] K. Penc, K Hallberg, F. M ila and H. Shiba: preprint cond-m at/9701051; The spin-charge factorized wave function in U ! 1 used in this paper was originally introduced in M.Ogata and H.Shiba: Phys.Rev.B 41 (1990) 2326; M.Ogata, T.Sugiyam a and H.Shiba: ibid. 43 (1991) 8401.
- [12] V.M eden and K.Shonham m er: Phys.Rev.B 46 (1992) 15753.
- [13] J.Voit: Phys.Rev.B 47 (1993) 6740.
- [14] H.Suzuura and N.Nagaosa: preprint cond-m at/9702212.
- [15] S. Sollera and Parola: J. Phys.: Condens. Matter 4 (1992) 3589.
- [16] S.Sollera and Parola: Phys.Rev.Lett.76 (1996) 4604.
- [17] J.Voit: preprint cond-m at/9602087.
- [18] R.Z.Bariev: Phys.Rev.B 49 (1994) 1474.
- [19] B.Sutherland: Phys. Rev. B 12 (1975) 3795.
- [20] P.Schlottm ann: Phys. Rev. B 36 (1987) 5177.
- [21] P.Bares and G.Blatter: Phys.Rev.Lett.64 (1990) 2567.
- [22] P. Bares, G. Blatter and M. Ogata: Phys. Rev. B 44 (1991) 130.
- [23] N.Kawakam iand S.K.Yang: Phys.Rev.Lett.65 (1990) 2309; J.Phys.: Condens. M atter 3 (1991) 5983.
- [24] J. Shiraishi, Y. M orita and M. Kohmoto: J. Phys. A 30 (1997) 831; R. Sato: J. Phys.: Condens. M atter 8 (1996) 8363.
- [25] A.A.Belavin, A.M. Polyakov and A.B.Zam olodchikov: Nucl.Phys.B 241 (1984) 333.
- [26] J.L.Cardy: Nucl. Phys. B 270 (1986) 186.
- [27] I.A eck: Phys. Rev. Lett. 56 (1986) 746.
- [28] F.W oynarovich: J.Phys.A 22 (1989) 4243.
- [29] N.Kawakam iand S.K.Yang: Phys.Lett.A 148 (1990) 359.
- [30] H.Frahm and V.Korepin: Phys. Rev. B 42 (1990) 10553.
- [31] Y.T sukam oto, T.Fu jii and N.K aw akam i: in preparation
- [32] N.M. Bogoliubov, A.G. Izergin and V.E.K orepin: Nucl. Phys. B 275 (1986) 687.

FIG.1. M om entum -dependent critical exponents X (k) for the spectral function in the 1D anisotropic t-J m odel(1) for which the spin sector is massive and the charge sector is massless.

FIG.2. Critical exponents X for the spectral function as a function of the magnetization m. The model is the hole-doped H eisenberg X X Z spin chain in the lim it of J=t 1, for which the spin sector is massless and the charge sector is massive.