

Superfluid ^3He , Particle Physics and Cosmology

G.E. Volovik

Low Temperature Laboratory, Helsinki University of Technology, Box 2200, FIN-02015 HUT, Finland
and

Landau Institute for Theoretical Physics, Moscow, Russia

(December 31, 2021)

Superfluid ^3He -A and high-temperature superconductors both have gapless fermionic quasiparticles with the "relativistic" spectrum close to the gap nodes. The interaction of the "relativistic" fermions with bosonic collective modes is described by the quantum field theory, which results in a close connection with particle physics. Many phenomena in high-energy physics and cosmology can thus be simulated in superfluid phases of ^3He and in unconventional superconductors. This includes axial anomaly, vacuum polarization, zero-charge effect, fermionic charge of the vacuum, baryogenesis, event horizon, vacuum instability, Hawking radiation, etc. Analogs of some of these phenomena, which are related to the axial anomaly, have been experimentally simulated in superfluid ^3He . This includes the baryogenesis by textures (Manchester), the baryogenesis by cosmic strings (Manchester) and the generation of the primordial magnetic field via the axial anomaly (Helsinki).

It is now well realized that the Universe and its broken symmetry ground state – the physical vacuum – may behave like a condensed matter system, in which the complicated degenerate ground state is developed due to the symmetry breaking. The condensed matter counterparts of Universe are represented by superconductors and by superfluid phases of ^3He : both systems contain quantum Bose and Fermi fields, describing the interaction of the elementary quasiparticles with each other and with the degenerate vacuum [1,2]. This analogy allows us to simulate in condensed matter many properties of the physical vacuum, while the direct experiments with the physical vacuum are still far from the realization.

The important property of the physical vacuum is its high anisotropy (Fig. 1).

The physical vacuum (superfluid ground state) of superfluid ^3He -A resembles the electroweak vacuum. The fermionic excitations of the ^3He -A in the vicinity of the gap nodes are "relativistic" chiral fermions (Fig. 2). They are the left-handed near the north pole, where they have a linear momentum \mathbf{p} close to $p_F \hat{\mathbf{l}}$ ($\hat{\mathbf{l}}$ is the direction of spontaneous angular momentum in ^3He -A) and they are right-handed near the south pole where $\mathbf{p} = -p_F \hat{\mathbf{l}}$. Another important similarity is that the symmetry breaking pattern $SU(2) \times U(1) \rightarrow U(1)$ in electro-weak interactions is the same as the symmetry breaking pattern $SO(3) \times U(1) \rightarrow U(1)$ in ^3He -A (Fig. 3).

That is why the physical effects are actually the same in the vacuum of the high energy physics and in superfluid ^3He -A. Thus the ^3He -A is the right condensed matter for simulations of the vacuum properties (Fig. 4). The main difference is in the terminology, so we need the dictionary for the translation from the particles physics language to that of ^3He -A (Fig. 5).

Here we discuss 3 experiments in superfluid ^3He which simulate the processes in the early Universe. The first two are related to the cosmological problem why the Universe contains much more matter than antimatter. The




FIG. 1. (top): The spectrum of fermionic excitations of the physical vacuum contain the branch of the chiral particles: left-handed neutrino. The right-handed neutrino is absent: it is the remarkable manifestation of the violation of the left-right symmetry in the vacuum. Another symmetry, which is broken in the present Universe, is the $SU(2)$ symmetry of weak interactions. In the unbroken symmetry state of the early Universe, the left leptons (neutrino and left electron) formed the $SU(2)$ doublet, while the right electron was the $SU(2)$ singlet. (bottom): During the cool down of the Universe the phase transition (or the crossover) occurred, at which the $SU(2)$ symmetry was broken. As a result the left and right electrons were hybridized forming the present electronic spectrum with the gap $\Delta = m_e c^2$. The electric properties of the vacuum thus exhibited the metal-insulator transition: The metallic state of the vacuum with the Fermi point in the electronic spectrum was transformed to the insulating state with the gap.




FIG. 2. Gap nodes and chiral fermions in ³He-A. Close to the gap nodes the gapless quasiparticle represent the massless fermions moving in the electromagnetic and gravity fields and obeying the relativistic equation $g^{\mu\nu}(p_\mu - eA_\mu)(p_\nu - eA_\nu) = 0$. Here $e = \pm$ is the "electric charge" and simultaneously the chirality of the quasiparticles. The righthanded particles have positive charge $e = +1$ and are in the vicinity of the north pole, while the left particles with $e = -1$ are in the vicinity of the south pole. $p_\mu = (E, \mathbf{p})$ is the 4-momentum; $A_0 = \mu_R$ plays the part of the chemical potential for the righthanded particles. The texture of $\hat{\mathbf{l}}$ field provides the vector potential $\mathbf{A} = k_F \hat{\mathbf{l}}$ and also the metric tensor of the gravitational field: $g^{ik} = c_\perp^2 (\delta^{ik} - \hat{l}^i \hat{l}^k) + c_\parallel^2 \hat{l}^i \hat{l}^k$, $g^{00} = -1$, the quantities $c_\parallel = p_F/m$ and $c_\perp = \Delta_0/p_F$ correspond to velocities of "light" propagating along and transverse to $\hat{\mathbf{l}}$, here m is the mass of the ³He atom and Δ_0 is the amplitude of the gap.

³He experiments allow to test different scenaria in which the nonconservation of the baryons is caused by the so called axial anomaly, which governs the nucleation of the fermionic charge from the vacuum. The essence of the axial anomaly is presented in Fig. 6. The chiral anomaly equation [3,4] is obtained using the picture of Landau levels for the right-handed particle in the applied magnetic field. This anomaly equation which describes the nucleation of the fermionic charges in the presence of magnetic and electric fields describes both the production of the baryons in the electroweak vacuum (baryogenesis) and the production of the linear momentum in the superfluid ³He-A (momentogenesis) (Fig. 7). In ³He-A the effective magnetic and electric fields are generated by the moving continuous vortex texture. The anomalous production of the linear momentum leads to the additional force acting on the continuous vortex in ³He-A (Fig. 8), which was measured in the Manchester experiment [5]. The




FIG. 3. The symmetry breaking in the electroweak transitions is similar to that in the transition from the normal ³He to ³He-A.

- Why ³He can help**
- We need broken symmetry **condensed matter** to simulate **quantum vacuum**.
 - **liquid crystals**: but they are not quantum
 - **superfluid ⁴He**: but it does not contain Fermi quasiparticles
 - **superconductors**:
Bogoliubov fermionic quasiparticles vs quarks
Nambu - Jona-Lasinio model of quark condensate.
but the limited number of degrees of freedom
 - **superfluid phases of ³He** are right systems
 - Superfluid ³He represent **interacting Fermi and Bose quantum fields**
analogs of
leptons, gauge fields, Higgs fields, gravitons, W-bosons, ...
 - Symmetry breaking is similar to that in physical vacuum
³He-A = electroweak vacuum
 - Various topological objects
3 vortices + vortex sheet in ³He-A and 4 vortices in ³He-B
(observed in Helsinki)
³He-A vortices = Z- and W- cosmic strings
1/2-vortex = Alice string
spin-mass vortex = cosmic string terminating on cosmic wall
nonaxisymmetric vortex = Witten superconducting string
 - Probing of quantum vacuum
flow instability = vacuum instability in strong fields
supercritical motion of texture = event horizon in black hole
 - Anomalies in quantum field theory
Anomalous force on vortex (Manchester experiments) = **baryogenesis**
 - Modelling of processes in hot Universe (Helsinki)
Nucleation of vortices by neutrons = strings formation in Big-Bang

FIG. 4. Superfluid ³He is the right condensed matter for simulation of the properties of the physical vacuum.

anomaly equation has thus been verified in the ³He-A experiments with the moving vortices. The axial anomaly is also responsible for the dynamics of singular vortices and for the baryogenesis by the singular cosmic string. In this case the spectral flow from the vacuum is realized by

Dictionary	
Superfluid ^3He	Universe
Superfluid ground state (no excitations)	Quantum vacuum (no elementary particles)
Bogoliubov quasiparticles	Quarks and leptons
Collective modes	Gauge and Higgs bosons
Texture	Gauge field
$k_F \mathbf{l}$	Vector potential \mathbf{A}
$k_F d\mathbf{l}/dt$	Electric field \mathbf{E}
$k_F \nabla \times \mathbf{l}$	Magnetic field $\mathbf{B} = \nabla \times \mathbf{A}$
Vortex	Cosmic string
Bound states in the core	Fermion zero modes
Exchange between superfluid and normal components	Axial anomaly: nucleation of charge from vacuum
Momentogenesis by moving vortex	Baryogenesis by cosmic string







FIG. 5. Dictionary

the bound states of quasiparticles in the core of vortices and strings (Figs. 9,10). The baryoproduction in the core of the cosmic string was simulated in the Manchester experiment on dynamics of singular vortices in $^3\text{He-B}$ [5]. There is also an opposite effect: the anomalous transformation of the chiral particles into the magnetic field. In the Universe this mechanism can be responsible for the formation of the primordial magnetic field [6], while in the $^3\text{He-A}$ the similar anomaly equations describes the collapse of the excitation momenta towards the formation of the textures – counterpart of the magnetic field in cosmology (Figs. 11,12). Such collapse was recently observed in Helsinki rotating cryostat [7].

We discussed only 3 experiments in superfluid ^3He related to the properites of the electroweak vacuum. In all of them the chiral anomaly is an important mechanism. It regulates the nucleation of the fermionic charge from the vacuum as was observed in [5] and the inverse process of the nucleation of the effective magnetic field from the fermion current as was observed in [7]. There are many other connections between the superfluid ^3He and other branches of physics which should be exploited (Fig. 13).



$$\dot{n} = (1 / 4\pi^2) [(e_R)^2 - (e_L)^2] \mathbf{E} \cdot \mathbf{B}$$

FIG. 6. Chiral anomaly: the spectral flow from the occupied energy levels caused by the applied electric field leads to the creation of the fermionic charge from the vacuum if the left-right symmetry is violated.

- [1] G.E. Volovik and T. Vachaspati, Int. J. Mod. Phys. **B 10**, 471 (1996).
- [2] G.E. Volovik, Exotic properties of superfluid ^3He , World Scientific, Singapore-New Jersey-London-Hong Kong, 1992.
- [3] S. Adler, Phys. Rev. **177**, 2426 (1969).
- [4] J.S. Bell and R. Jackiw, Nuovo Cim. **A60**, 47 (1969).
- [5] T.D.C. Bevan, A.J. Manninen, J.B. Cook, J.R. Hook, H.E. Hall, T. Vachaspati and G.E. Volovik, Nature **386**, 689 (1997).
- [6] M. Joyce, M. Shaposhnikov, Phys. Rev. Lett., **79**, 1193 (1997).
- [7] V.M.H. Ruutu, J. Kopu, M. Krusius, U. Parts, B. Placais, E.V. Thuneberg, and W. Xu, to appear in Phys. Rev. Lett.

Anomalous nonconservation of baryonic charge and ^3He momentum

Baryon nonconservation in Electroweak Theory

$$\dot{\mathbf{B}} = (1/4\pi^2) \mathbf{B}_Y \cdot \mathbf{E}_Y \sum_q [B_{qR} (Y_{qR})^2 - B_{qL} (Y_{qL})^2] + (1/4\pi^2) \mathbf{B}_W \cdot \mathbf{E}_W \sum_q [B_{qR} (W_{qR})^2 - B_{qL} (W_{qL})^2]$$

B -- baryonic charge of matter
 \mathbf{B}_Y and \mathbf{E}_Y -- magnetic and electric fields of hypercharge
 \mathbf{B}_W and \mathbf{E}_W -- weak magnetic and electric fields
 B_{qR} and B_{qL} -- baryonic charge of right and left quark
 Y_{qR} and Y_{qL} -- hypercharge of right and left quark
 W_{qR} and W_{qL} -- weak charge of right and left quark

Left-Right asymmetry gives net product of baryonic charge

$$\dot{\mathbf{B}} = (3/8\pi^2) [\mathbf{B}_Y \cdot \mathbf{E}_Y - \mathbf{B}_W \cdot \mathbf{E}_W]$$

Momentum transfer in Superfluid $^3\text{He-A}$

$$\dot{\mathbf{P}} = (1/4\pi^2) (\mathbf{B} \cdot \mathbf{E}) [\mathbf{P}_R - \mathbf{P}_L]$$

\mathbf{P} -- total momentum of quasiparticles
 $\mathbf{B} = \mathbf{k}_F \nabla \times \mathbf{l}$ and $\mathbf{E} = \mathbf{k}_F d\mathbf{l}/dt$ -- "magnetic" and "electric" fields
 \mathbf{P}_R and \mathbf{P}_L -- momenta of created right and left quasiparticle
Left-Right asymmetry, $\mathbf{P}_R = p_F \mathbf{l} = -\mathbf{P}_L$, gives net product of momentum

$$\dot{\mathbf{P}} = (1/2\pi^2) (\mathbf{B} \cdot \mathbf{E}) p_F \mathbf{l}$$





FIG. 7. The same anomaly equation leads to production of the baryonic charge B in high energy physics and to production of the linear momentum \mathbf{P} in superfluid $^3\text{He-A}$.

Momentogenesis by continuous $N=2$ vortex



Momentum transfer from the vacuum (superfluid) to the heat bath (matter) gives extra force on $N=2$ vortex

$$\begin{aligned} \mathbf{F} &= \int d^3r \dot{\mathbf{P}} = (1/2\pi^2) \int d^3r (\mathbf{B} \cdot \mathbf{E}) p_F \mathbf{l} \\ &= (1/2\pi^2) \hbar k_F^3 \int d^3r (\nabla \times \mathbf{l} \cdot d\mathbf{l}/dt) \mathbf{l} \\ &= \pi \hbar N (1/3\pi^2) k_F^3 \hat{\mathbf{z}} \times (\mathbf{v}_n - \mathbf{v}_L) \end{aligned}$$


This force effectively cancels the Magnus force

$$\mathbf{F}_M = \pi \hbar N n \mathbf{z} \times (\mathbf{v}_s - \mathbf{v}_L)$$

The particle density n deviates from $(1/3\pi^2) k_F^3$ due to small asymmetry between particles and holes

FIG. 8. The order parameter texture (analog of magnetic field) in the core of continuous vortex is converted into the excess of the fermionic charge (linear momentum) due to the axial anomaly in the moving continuous vortex. The moving vortex generates the time dependence of the order parameter, which is equivalent to electric field.


Cosmic string vs Singular vortex



core radius r_c
 $=$ coherence length ξ in condensed matter vortex
 $= 1/m_{\text{Higgs}}$ in electroweak string (m_{Higgs} - mass of Higgs-boson)
 $\lambda =$ penetration length of magnetic field in superconductor
 $= 1/m_Z$ in electroweak Z-string (m_Z - mass of Z-boson)
 $= \infty$ in electrically neutral superfluid ^3He

FIG. 9. Cosmic string is the counterpart of the Abrikosov vortex in superconductor. It also contains the bound states of the fermions, which are important for the baryoproduction.

Anomalous nonconservation of linear momentum in superconductors and in $^3\text{He-B}$ vortex



When vortex moves along x the angular momentum changes:

$$\dot{\mathbf{L}}_z = \dot{\mathbf{x}} p_y = v_x p_y$$

This gives the flow of levels from Fermi sea:

$$\dot{\mathbf{n}} = \dot{\mathbf{L}}_z / \hbar = v_x p_y / \hbar$$

The flow of linear momentum (transverse force on vortex):

$$\begin{aligned} \mathbf{F}_y &= \langle \dot{p}_y \rangle = \langle p_y \dot{\mathbf{n}} \rangle = \langle p_y^2 \rangle v_x / \hbar = \\ &= \pi \hbar (1/3\pi^2) k_F^3 v_x \end{aligned}$$

FIG. 10. The spectral flow of the momentum in the core of the moving singular vortex leads to the experimentally observed reactive force on a vortex in superfluid $^3\text{He-B}$.

Counterflow vs chemical potential for chiral fermions





FIG. 11. Instability of the counterflow. The counterflow generated by the rotation of the cryostat is equivalent to the nonzero chemical potential for the right electrons. When the velocity of the counterflow $\mathbf{v}_n - \mathbf{v}_s$ in the \hat{l} direction (corresponding to the chemical potential μ_R of the chiral electrons) exceeds some critical value, the instability occurs and the container becomes filled with the \hat{l} -texture which corresponds to the hypermagnetic field.

Flow instability vs instability of state with excess of chiral charge



Particle physics ${}^3\text{He-A}$

Chern-Simons term leading to helical instability:
 $(1/2\pi^2) \mu \mathbf{A}(\nabla \times \mathbf{A})$

Chemical potential $\mu = \rho_F l_0(\mathbf{v}_s - \mathbf{v}_n)$
 Vector potential $\mathbf{A} = \rho_F(l - l_0)$
 Magnetic field = helical texture
 $F_{ik} = \nabla_i A_k - \nabla_k A_i = k_F(\nabla_i l - \nabla_k l)$
 Magnetic field energy = Energy of texture
 $(1/12\pi^2) \ln(\Lambda/T) \sqrt{-g} g^{ij} g^{kl} F_{ik} F_{jl}$

Metric tensor $g^{ij} = c_{||}^2 |i\rangle\langle i| + c_{\perp}^2 (\delta^{ij} - |i\rangle\langle i|)$
 Planck energy $\Lambda = \text{gap amplitude } \Delta_0$

FIG. 12. The excess of the chiral right-handed electrons in early Universe can be effectively converted to the hypermagnetic field due to the mechanism of chiral anomaly. This is described essentially by the same equations as the counterflow instability observed in ${}^3\text{He-A}$.




FIG. 13. Connections of superfluid ${}^3\text{He}$ to other branches of physics.