

Quantum Phase Transitions in Josephson-Junction Chains

M.-S. Choi¹, J. Y. Ji², M. Y. Choi², J. Choi³, and S.-I. Lee¹

¹ Department of Physics, Pohang University of Science and Technology, Pohang 790-784, Korea

² Department of Physics and Center for Theoretical Physics, Seoul National University, Seoul 151-742, Korea

³ Department of Physics, Keimyung University, Taegu 704-701, Korea

(Oct. 20 1997)

We investigate the quantum phase transition in a one-dimensional chain of ultra-small superconducting grains, considering both the self- and junction capacitances. At zero temperature, the system is transformed into a two-dimensional system of classical vortices, where the junction capacitance introduces anisotropy in the interaction between vortices. This leads to the superconductor-insulator transition of the Berezinskii-Kosterlitz-Thouless type, as the ratios of the Josephson coupling energy to the charging energies are varied. It is found that the junction capacitance plays a role similar to that of dissipation and tends to suppress quantum fluctuations; nevertheless the insulator region survives even for arbitrarily large values of the junction capacitance.

PACS Numbers: 74.50.+r, 67.40.Db

Quantum phase transitions, which are induced by quantum fluctuations at zero temperature, are distinguished from classical phase transitions in several important respects; this has attracted much attention in recent years [1]. In particular, advances in fabrication techniques have made available arrays of ultra-small superconducting grains, where the charging energy dominates the Josephson coupling energy and accordingly, quantum fluctuation effects are of paramount importance. Such Josephson-junction arrays have become a prototype system displaying quantum phase transitions between the superconducting and the insulating phases. In the vicinity of the superconductor-insulator transition, the fluctuation effects depend crucially on the dimensionality of the system. In the case of two-dimensional (2D) arrays, rich effects of quantum fluctuations and resulting phase transitions have been examined for a rather general form of the capacitance matrix, although there still exist unsettled issues in the quantum regime, such as low-temperature re-entrance [2,4]. On the other hand, one-dimensional (1D) chains of Josephson junctions, where quantum fluctuations should be more important, have been studied mainly in the two limiting cases: the self-charging model and the nearest-neighbor model where only nearest neighboring charges interact [5,6]. In the 1D system with only self-capacitance, the role of dissipation on the quantum phase transition [7] as well as the persistence current and voltage [8] has also been considered.

This paper investigates the quantum phase transitions in general Josephson-junction chains with both the self- and junction capacitances. At zero temperature, the system is transformed into a two-dimensional system of classical vortices, where the junction capacitance introduces anisotropy in the interaction between vortices. This leads to the superconductor-insulator transition of the Berezinskii-Kosterlitz-Thouless (BKT) type [9,10], as the ratios of the Josephson coupling energy to the charging energies are varied. Interestingly, the junction capaci-

tance here plays a role similar to that of dissipation and tends to suppress quantum fluctuations, enhancing superconductivity. However, the suppression is not strong enough, and the insulator region still remains even for arbitrarily large values of the junction capacitance.

We consider a one-dimensional array of N ultra-small superconducting grains, each of which is coupled to nearest-neighboring grains via Josephson junctions. The system is characterized by three energy scales, Josephson coupling energy E_J and charging energies $E_0 = e^2/2C_0$ and $E_1 = e^2/2C_1$, where C_0 is the self-capacitance of each grain and C_1 is the junction capacitance between nearest-neighboring grains. It is described by the Hamiltonian

$$H = \frac{1}{2K_0} \sum_{i,j=1}^N n_i C_{ij}^{-1} n_j - K_0 \sum_{i=1}^N \cos(\phi_{i+1} - \phi_i); \quad (1)$$

where the number n_i of Cooper pairs and the phase ϕ_i of the superconducting order parameter at site i are quantum-mechanically conjugate variables: $[\phi_i; n_j] = i \delta_{ij}$, and the energy has been rescaled in units of the Josephson plasma frequency $\frac{\hbar!}{p} \frac{E_J}{8E_0E_J}$. Here we have introduced $K_0 = \frac{E_J}{8E_0}$ to describe the case $C_0 \neq 0$, and written the capacitance matrix in the form

$$C_{ij} = \delta_{ij} + \frac{C_1}{C_0} (\delta_{i,j-1} + \delta_{i,j+1});$$

where $\frac{C_1}{C_0} = 1$. On the other hand, when $C_0 = 0$, it is convenient to introduce $K_1 = \frac{E_J}{8E_1}$. Note that K_0 can also be written in the form: $K_0 = K_1 = K$.

In the imaginary-time path-integral representation, the partition function of the system reads

$$Z = \int \prod_{j; n_j} \mathcal{D} \phi_j \exp \left(- \sum_{j=1}^N \frac{d\phi_j}{2} \right) \exp \left(- \sum_{i=1}^N L[\phi; n] \right) \quad (2)$$

with the action

$$L[\mathbf{n};] = i \sum_j \mathbf{n}_j \cdot \nabla_{\mathbf{x}_j} + \frac{1}{2K_0} \sum_{i,j} \mathbf{n}_i \cdot C_{ij}^{-1} \mathbf{n}_j - K_0 \sum_j \cos \theta_{\mathbf{x}_j}; \quad (3)$$

where the temperature has been rescaled according to $\hbar v_p$. In Eq. (3), $\nabla_{\mathbf{x}}$ and $\nabla_{\mathbf{t}}$ denote the difference operator with respect to the position \mathbf{x} and to the imaginary time τ , respectively: $\nabla_{\mathbf{x}_j} = \nabla_{\mathbf{x}_{j+1}} - \nabla_{\mathbf{x}_j}$ and $\nabla_{\mathbf{t}_j} = \nabla_{\mathbf{t}_{j+1}} - \nabla_{\mathbf{t}_j}$, and the (imaginary) time slice in the interval $[0; \beta]$ has been chosen to be unity (in units of $\hbar v_p$) [11]. Our main concern here is the quantum phase transition at zero temperature, and accordingly, the limit $\beta \rightarrow \infty$ as well as the thermodynamic limit $N \rightarrow \infty$ is to be taken. We then apply the Villain approximation [12] and integrate out $\mathbf{f}_j; g_j$, to obtain the partition function in terms of an additional set of integer variables $\mathbf{m}_j; g_j$ (as well as $\mathbf{f}_j; g_j$). The corresponding action is given by

$$L[\mathbf{n}; \mathbf{m}] = \frac{1}{2K_0} \sum_{i,j} \mathbf{n}_i \cdot C_{ij}^{-1} \mathbf{n}_j + \frac{1}{2K_0} \sum_j \mathbf{m}_j^2; \quad (4)$$

where the two sets of integer variables satisfy the constraint $\nabla_{\mathbf{x}} \mathbf{m}_j + \nabla_{\mathbf{t}} \mathbf{n}_j = 0$. This constraint is conveniently taken into account by introducing an integer field $A_{j;\sim}$ defined on the space-time dual lattice $(\tilde{j}; \sim)$ ($j+1=2; \tau+1=2$) in such a way that $\mathbf{m}_j = \nabla_{\mathbf{x}} A_{j;\sim}$ and $\mathbf{n}_j = \nabla_{\mathbf{t}} A_{j;\sim}$. Henceforth we will work on the dual lattice, and drop for simplicity the tilde sign over site indices. The partition function is thus written in terms of the unconstrained summation of $\exp L[\mathbf{A}]$ over \mathbf{A}_j 's, with the action

$$L[\mathbf{A}] = \frac{1}{2K_0} \sum_{i,j} (\nabla_{\mathbf{x}} A_j) \cdot C_{ij}^{-1} (\nabla_{\mathbf{x}} A_j) + \frac{1}{2K_0} \sum_j (\nabla_{\mathbf{t}} A_j)^2; \quad (5)$$

Finally, the Poisson summation formula, which decomposes A_j into a real-valued field and a new integer field v_j , allows us to integrate out the real-valued field A_j . This leads, apart from the spin-wave part, to the 2D system of classical vortices described by the Hamiltonian

$$H_v = 2^2 K_0 \sum_{i,j} v_i \cdot U(i;j) \cdot v_j; \quad (6)$$

where the vortex interaction is given by

$$U(\mathbf{x};) = \int_0^{2\pi} \frac{dq}{2\pi} \int_0^{2\pi} \frac{d!}{2\pi} e^{i\mathbf{q}\cdot\mathbf{x} - i!} \mathcal{U}(\mathbf{q};!);$$

$$\mathcal{U}(\mathbf{q};!) = \frac{C(\mathbf{q})}{\mathcal{C}(\mathbf{q})} + (!); \quad (7)$$

with the Fourier transforms of the lattice Laplacian $(z) = 2(1 - \cos z)$ and of the capacitance $\mathcal{C}(\mathbf{q}) = 1 + \sum_{\mathbf{r}} C(\mathbf{q})$. Note that unless $C_0 = 0$, the diagonal piece $U_0 = U(\mathbf{x}=0; \tau=0)$ becomes arbitrarily large as N and β are increased to infinity. Therefore the diagonal term $2^2 K_0 U_0 (\sum_j v_j)^2$ should vanish in Eq. (6), leaving the vorticity neutrality condition: $\sum_j v_j = 0$, and the Hamiltonian (6) may be written in the form

$$H_v = K_0 \sum_{i,j} v_i \cdot \mathcal{U}(i;j) \cdot v_j; \quad (8)$$

with $\mathcal{U}(\mathbf{x};) = 2 [\sum_{\mathbf{r}} U(\mathbf{x};)]$. The behavior of the (reduced) vortex interaction $\mathcal{U}(\mathbf{x};)$ for several values of

is displayed in Figs. 1 and 2, which manifests the logarithmic behavior at large length scales. In particular, the short-range anisotropy prominent for large values of K_0 decreases rapidly with the distance. According to the renormalization group (RG) theory of the 2D Coulomb gas, such short-range anisotropy should not affect the universality class. Thus in the spirit of the RG theory, the system is expected to exhibit qualitatively the same critical behavior as the 2D Coulomb gas [13].

We now investigate the quantum phase transitions displayed by the Hamiltonian in Eq. (6). First we consider the simplest case of $C_1 = 0$, which has been studied in Ref. [5]. In this self-charging limit, the interaction in Eq. (7) takes the simple form $\mathcal{U}(\mathbf{q};!) = [C(\mathbf{q}) + (!)]^{-1}$ or $\mathcal{U}(\mathbf{x};) = \frac{1}{\ln(x^2 + \tau^2) + \frac{3}{2} \ln 2 + E_E}$, where E_E is the Euler number. Thus the Hamiltonian (8) describes precisely the isotropic 2D Coulomb gas, and the system undergoes a BKT transition [9,10,12] from the insulating phase to the superconducting one as K_0 is increased [5].

We next consider the opposite limit $C_0 = 0$, where the relevant (dimensionless) coupling constant is K_p instead of K_0 . Accordingly, it is proper to use $\hbar v_p$ instead of $\hbar v$ in rescaling the energy and the imaginary time, which in turn gives $\mathcal{C}(\mathbf{q}) = \sum_{\mathbf{r}} C(\mathbf{q})$ and $\mathcal{U}(\mathbf{q};!) = [1 + (!)]^{-1}$. As a result, we obtain the Hamiltonian in the form

$$H_v = 2^2 K_1 \sum_{i,j} v_i \cdot U(i;j) \cdot v_j; \quad (9)$$

where, in sharp contrast to the previous self-charging case, the vortex interaction $\Phi(x; \nu) = \sum_j U(x; \nu)$ is short-ranged:

$$\Phi(x; \nu) = \sum_{j \neq 0} \frac{h}{2\pi} e^{i \nu \phi_j} \quad (10)$$

for $j \neq 0$. It is further of particular importance that U_0 in general does not diverge, which implies that the diagonal term in the Hamiltonian (9) does not give the vorticity neutrality condition. Thus (unbound) free vortices become pervasive, and the system remains insulating for any nonzero value of K_1 . This can also be understood in the charge representation, where the partition function reads

$$Z = \sum_{\{n_j\}} \sum_{\{q_j\}} \exp \left[-\frac{1}{N} \sum_{q_j} G(q_j) \right] \quad (11)$$

with $G(q; \nu) = \nu [(\nu) + 1]^{-1}$. It follows from the analytic continuation of $G(q; \nu)$ that the charge excitation has a gap $E_g = h\nu$. Thus the long-range interaction C_{ij}^{-1} between charges gives rise to the gap, resulting in an insulator. It is also of interest to compare this case with the "nearest-neighbor model" in Ref. [5]: In the latter, only the nearest-neighbor charges interact, which allows only bound vortices. Consequently, the system is superconducting for all finite values of the interaction strength.

We now turn to the more realistic case that ν is small but nonzero, i.e., $C_0 = C_1$. In this case, the vortex

interaction Φ is, to the order of ν^2 , isotropic and has the asymptotic behavior: $\Phi(x; \nu) \sim \ln(x^2 + \nu^2) + \nu^2$, where ν^2 is the vortex pair creation energy (per vortex) given by $\nu^2 = (3/2) \ln 2 + E + (\nu)^2 + O(\nu^4)$. Hence the system again reduces to the 2D Coulomb gas, with the vortex fugacity $y(\nu) = \exp[-2K_0 \nu^2]$ diminished by the factor $\exp[-2K_0(\nu)^2]$. The standard RG theory of the BKT transition [10,12] then predicts the transition point $K_0^c(\nu)$ slightly decreased from $K_0^c(0)$ by the amount

$$\frac{K_0^c(\nu)}{K_0^c(0)} = \left(1 - \frac{2K_0^c(0)}{2K_0^c(0)} \frac{4}{3} \nu^2 + O(\nu^4) \right) \quad (12)$$

In this limit ($\nu \rightarrow 0$), the Hamiltonian in Eq. (1) can also be represented in terms of phase variables, which yields a 2D XY model with an additional interaction for nonzero ν . To see this, we neglect $O(\nu^4)$, and write the inverse capacitance matrix in the form

$$C_{ij}^{-1} = (1 - \nu^2) \delta_{ij} + \nu^2 (\delta_{i,j+1} + \delta_{i,j-1}) \quad (13)$$

Note that without the off-diagonal term in C_{ij}^{-1} , the first and the second terms in the action (3) would just be the Villain form of the cosine action along the x direction [5,8]. To examine the effects of the off-diagonal term, we use the identity

$$\exp \left[-\frac{\nu^2}{4K_0} (\partial_x n_j)^2 \right] = \int_{-\pi}^{\pi} dz_j \exp \left[i z_j \partial_x n_j - \frac{1}{K_0} z_j^2 \right]$$

which allows to separate the charge variables at different sites. The resulting action then takes the Villain form of the cosine action with the argument

$\theta_j + i \frac{\nu^2}{2K_0} \partial_x z_j$ along the x direction. To the order of ν^2 , the Gaussian integration over z_j leads to the effective phase Hamiltonian

$$H_e[\theta] = K_0 \sum_j \left[\frac{1}{2} (\partial_x \theta_j)^2 + \frac{1}{4K_0} (\partial_x \theta_j)^2 \right] + \sum_j \left[\cos(\theta_j) + \cos(\theta_j + \nu^2 \partial_x z_j) \right] + \sum_j \left[\cos(\theta_j) + \cos(\theta_j + \nu^2 \partial_x z_j) \right] \quad (14)$$

where the anisotropy has been removed by rescaling again the x axis by the factor $(1 - \nu^2)^{-1/2} = 1 + \nu^2/2 = 4K_0$. Note that Eq. (14) reduces, for $\nu = 0$, to the standard 2D XY Hamiltonian. Here the junction capacitance not only enhances the effective coupling of the XY model but also

introduces an additional interaction given by the second term in Eq. (14). Interestingly, the latter is very similar in form to the dissipation term in the effective action, which is known to suppress quantum fluctuations [3,7]. Therefore both effects contribute to the enhancement of

phase coherence, and it is concluded that the junction capacitance in general tends to suppress quantum fluctuations induced by the self-capacitance, thus reducing the insulator region.

For larger values of ϵ , Figs. 1 and 2 (b) show that the

$$\langle \mathcal{U} \rangle(0; \epsilon) = \frac{2}{\epsilon} + \ln \epsilon + E + 2 \ln 2 - \frac{1}{2} \ln 5 - \frac{127}{150} + O(\epsilon^{-1}; \epsilon^{-2}) \quad (15)$$


Although the anisotropy in the short-range behavior of $\langle \mathcal{U} \rangle$ may slightly alter the details of the RG flow, the qualitative features at large length scales are expected unaffected. Thus it is concluded that for any value of finite ϵ the system undergoes the superconductor-insulator transition of the BKT universality class.

Figure 3 displays the schematic phase diagram on the K_0 - K_1 plane. Although the precise phase boundary, i.e., the detailed behavior of $K_0^c(\epsilon)$, in general depends on the microscopic length scales such as the (imaginary) time slice τ , the universal (large-length scale) behavior should not be affected. Thus the phase boundary between the superconducting and the insulating phases is concluded to belong to the BKT universality class. As ϵ ($K_1=K_0$) is increased, the vortex creation energy also grows monotonically, resulting in the decrease of $K_0^c(\epsilon)$ toward the limiting value 2π . (Note that the value 2π corresponds to the zero vortex fugacity or the infinite vortex creation energy.) This lowering of the transition point and the resulting enhancement of superconductivity reflects the role of the junction capacitance: It tends to suppress the quantum fluctuations induced by the self-capacitance, as manifested in the phase representation given by Eq. (14). There has been few experimental studies of quantum phase transitions in 1D Josephson junction arrays. Recent interest in 1D tunnel junction arrays [14] and advances in submicron fabrication techniques strongly motivate experiments in this field.

MSC is grateful to W. G. Choe, G. S. Jeon, and K. H. Wagenblast for helpful discussions. This work was supported in part by the Basic Science Research Institute Program, Ministry of Education of Korea and in part by the Korea Science and Engineering Foundation through the SRC Program.

vortex creation energy $\langle \mathcal{U} \rangle$ and the short-range behavior of the interaction $\langle \mathcal{U} \rangle(x; \epsilon)$ are highly anisotropic. Still at large length/time scales, $\langle \mathcal{U} \rangle(x; \epsilon)$ becomes isotropic and logarithmic, which can also be confirmed by the asymptotic expansion for large ϵ :

[2] R. Fazio and G. Schon, Phys. Rev. B 43, 5307 (1991); A. van Otterlo, K. H. Wagenblast, R. Fazio, and G. Schon, *ibid.* 48, 3316 (1993).
 [3] B. J. Kim and M. Y. Choi, Phys. Rev. B 52 (5), 3624 (1995); *ibid.* 56, 395 (1997).
 [4] C. Rojas and J. V. Jose, Phys. Rev. B 54, 12361 (1996).
 [5] R. M. Bradley and S. Doniach, Phys. Rev. B 30, 1138 (1984).
 [6] It should be noted that the second case (the nearest-neighbor model in Ref. [5]) is rather unrealistic and in particular does not correspond to the case $C_0 = 0$ and $C_1 \neq 0$ shown schematically in Fig. 2 of Ref. [5]. In the latter case, the charge interaction C_{ij}^{-1} is long-ranged rather than between nearest neighbors.
 [7] P. A. Bobbert, R. Fazio, G. Schon, and G. T. Zimanyi, Phys. Rev. B 41, 4009 (1990); P. A. Bobbert, R. Fazio, G. Schon, and A. D. Zaikin, *ibid.* 45, 2294 (1992).
 [8] M. Y. Choi, Phys. Rev. B 48, 15920 (1993).
 [9] V. L. Berezinskii, Zh. Eksp. Teor. Fiz. 59, 907 (1970) [Sov. Phys. JETP 32, 493 (1971)]; J. M. Kosterlitz and D. J. Thouless, J. Phys. C 6, 1181 (1973).
 [10] J. M. Kosterlitz, J. Phys. C 7, 1047 (1974).
 [11] The critical behavior of the system should not be affected by the choice of τ (see Ref. [1]). Here all the dynamics of the system occurs over the time scale $1/\tau$, making it proper to choose \hbar/τ .
 [12] J. V. Jose, L. P. Kadano, S. Kirkpatrick, and D. R. Nelson, Phys. Rev. B 16, 1217 (1977).
 [13] P. Minnhagen, Rev. Mod. Phys. 59, 1001 (1987).
 [14] See, e.g., P. Delsing, D. B. Haviland, and P. D. Avidsson, Czech. J. Phys. 46, 2359 (1996), and reference therein.



[1] See, e.g., S. L. Sondhi, S. M. Girvin, J. P. Carini, and D. Shahar, Rev. Mod. Phys. 69, 315 (1997) and reference therein.




FIG .1. Vortex interaction potential $\hat{\psi}$ along (a) the x-axis and (b) the τ -axis for $\beta = 0; 0.5; 1; 5$, and 50 (from the bottom to the top).





FIG .2. Plot of the vortex interaction $\hat{\psi}(x; \tau)$ for (a) $\beta = 0.1$ and (b) $\beta = 30$.




FIG .3. Schematic phase diagram of a Josephson-junction chain at zero temperature. As β gets increased, K_0^c decreases toward the limiting value $2/\pi$.