

Hopf Term for a Two-Dimensional Electron Gas:
Comment to Reply to Comment

In our Comment [1] to the Letter [2] we questioned the derivation of the Hopf term in the hydrodynamic action for the Skyrme ion dynamics in QHE. Our main argument was that the description in terms of Euler angles used in [2] was potentially dangerous since these angles were ill-defined. The Reply [3] to our Comment confirmed our apprehension. In their Reply [3] Apel and Bychkov used their hydrodynamic action to derive the spin of the Skyrme ion from the adiabatic rotation of the Skyrme ion. They obtained the value of spin $s = (1/2)Q$ where

$$Q = \frac{1}{4} \int d^2r f(\theta_x, \theta_y) \theta_y [\cos \theta_x] \theta_x [\cos \theta_y] g; \quad (1)$$

They claimed that Q is the topological charge of Skyrme ion. Calculating Q for the configuration of the Euler angles

$$\cos \theta_x(r=1) = 1; \cos \theta_x(r=0) = -1; \theta_y = \arctan(y=x) \quad (2)$$

which corresponds to the Skyrme ion with the topological charge 1, they obtained $Q = 1$ and thus suggested that their hydrodynamic action does contain the Hopf term responsible for the spin $1/2$ of the Skyrme ion.

Thus the controversy is reduced to the estimation of the integral in Eq.(1), which is an easy task. Let us check the statement that the Eq.(1) gives the value $Q = 1$ for the configuration in Eq.(2). Though the angle θ is ill-defined, its derivatives are well defined. Thus the integrand in the Eq.(1) is also well defined in the whole 2D plane including the origin $r = 0$, though it diverges there. That is why one can easily estimate the integral either by direct calculations or by converting it to a contour integral.

Let us calculate the integral in Eq.(1) directly. It can be expressed as a sum of two terms:

$$Q = Q_1 + Q_2; \quad (3)$$

where

$$Q_1 = \frac{1}{4} \int d^2r f(\theta_x \cos \theta_y) (\theta_y) (\theta_y \cos \theta_x) (\theta_x) g; \quad (4)$$

and

$$Q_2 = \frac{1}{4} \int d^2r f \cos(\theta_x \theta_y - \theta_y \theta_x) g; \quad (5)$$

The first integral gives $Q_1 = 1$. In the second integral one has

$$(\theta_x \theta_y - \theta_y \theta_x) = 2 \theta(r); \quad (6)$$

Using the configuration in Eq.(2) one obtains $Q_2 = -1/2$. Altogether this gives

$$Q = Q_1 + Q_2 = 1 - 1/2 = 1/2; \quad (7)$$

instead of $Q = 1$.

This mistake in calculation of the integral in Eq. (1) (Eq.(4) of [3]) is rather common, since the θ -function contribution is often forgotten, and it serves as a nice illustration of the potential danger of using the Euler angles for the description of the Hopf term. As was observed by Apel and Bychkov [3] the same result $Q = 1/2$ can be obtained from Eq.(1) by converting the 2D integral to a contour integral around $r = 1$. But they disregarded this as a wrong way of calculation of Eq.(1).

Instead Apel and Bychkov suggested an alternative way of the estimation of the Eq.(1): they introduced the hole in the 2D space, where the Euler angles have singularity, and considered the 2D space with removed point. In this case the volume integral in Eq.(1) can be transformed into the contour integrals at infinity and around the infinitely thin cylindrical surface embracing the hole. Such description in terms of the holes gives the correct value for the topological invariant $Q = 1/2 + 1/2 = 1$. But if one introduces holes for the calculation of topological invariant Q , one should also introduce the holes in the 3D space-time (the space-time with removed lines) for the calculation of the Hopf invariant in action: one cannot use different approaches for the calculation of Q and of Hopf invariant! In this case the problem becomes complicated, since one must estimate the contribution of these dynamical holes into the hydrodynamic action. Apparently this was not made in [2]: the hole contribution has not been considered there.

Conclusion:

(i) If the holes in 2D space are introduced, the calculations of the Hopf term, made by Apel and Bychkov [2], should be reconsidered.

(ii) If the holes are not introduced, then the Eq.(1) should be valid in the whole 2D space and thus the Hopf-like action calculated in [2] gives the wrong result $Q = 1/2$ for Skyrme ion with topological charge $Q = 1$.

In both cases the derivation of the Hopf term in [2] remains questionable.

G.E.Volovik

Helsinki University of Technology, Low Temperature Laboratory, P.O.Box 2200, FIN-02015 HUT, Finland

Landau Institute for Theoretical Physics, 117334 Moscow, Russia

November 10, 1997

PACS numbers: 73.20.Dx, 71.35.Ji, 73.20.Mf, 75.30.Et

[1] G.E.Volovik and V.M.Yakovenko, Phys.Rev.Lett. 79, 3791 (1997).

[2] W.Apel and Yu.A.Bychkov, Phys.Rev.Lett. 78, 2188 (1997).

[3] W.Apel and Yu.A.Bychkov, Phys.Rev.Lett. 79, 3792 (1997).