

Comment on "Shell Filling and Spin Effects in a Few Electron Quantum Dot"

The experimental observation of shell and spin effects in the ground-state addition spectrum of a semiconductor quantum dot has been reported in a recent Letter by Tarucha *et al.* [1]. By applying a gate voltage, the chemical potential of the dot is varied and one observes Coulomb oscillations in the current vs gate voltage. Current peaks indicate the tunneling of the $N + 1$ st electron into the N -electron dot when the chemical potential $\mu_N = E(N + 1) - E(N)$ gets into resonance with the source and drain potentials.

The authors show the addition spectrum vs magnetic field for $N = 0$ to 23. They explain the observed structures by a model of non-interacting electrons in a disc with parabolic lateral confinement $V(r) = \frac{1}{2}m^* \omega_0^2 r^2$ ($m^* = 0.067m_0$ in GaAs). A confinement energy of $\hbar\omega_0 = 3$ meV fits well to the measured minimum of the 7th-electron addition energy at $B = 1.3$ T, which they attribute to the crossing of the 6th and 7th single-particle energy levels. Spin effects are interpreted in the framework of a constant-interaction model with fitting parameters including the exchange energy of parallel spins.

Using current- and spin-density functional theory (CSDFT) in local-density approximation (LDA) [2,3], we have calculated the addition-energy spectrum of a parabolic circular quantum dot in a perpendicular magnetic field with up to 20 electrons. Our calculations give a satisfactory agreement with the experiment (see Fig. 1). This is remarkable, since we use a mean-field approach and calculate the exchange-energy by employing the local spin-density approximation. Thus no further fitting parameters appear besides $\hbar\omega_0$ for the bare confinement (in contrast with the model in Ref. [1]). The latter turns out to be larger (5 meV) than what the experimentalists assumed for their fit. The reason for this difference is that the bare confinement is screened by the interacting electrons. It is also interesting to regard our calculations as a test of the LDA for a small number of electrons in the dot.

Due to interaction, the shell structure of the non-interacting model (connected with "magic numbers" $N = 2, 6, 12$ etc.) at zero magnetic field is partially broken. In the Kohn-Sham (KS) scheme of CSDFT, only the single-particle angular-momentum states with opposite signs are still exactly degenerate. Nevertheless, we obtain a spectrum with increased addition energies for the 7th and 13th electron as seen in experiment at $B = 0$.

We confirm the observed Hund's rule for small magnetic fields which favors spin alignment. In Fig. 1 this leads to a reduction of the addition energy if the electron is placed into the same spin state as its predecessor (e.g. for $N = 4; 8$ or 14) and vice versa ($N = 5; 9$ or 15).

Quasi-degenerate spin states at intermediate magnetic fields give rise to pairs of parallel addition-energy traces (indicating spin saturation for even N) [1]. Deviations from this behavior occur when the spin-saturated system is near a level crossing of different angular-

momentum states at the Fermi energy. Then again, Hund's rule applies since it becomes energetically favorable to occupy different angular-momentum states and align spins in order to gain exchange energy. As a result, the addition of, e.g., the 7th electron around $B = 1.2$ T [Fig. 1(a)] is especially costly where the exchange gap of the 6-electron dot is the largest [cf. Fig. 1(b)]. Correspondingly, the addition-energies for the 5th and 6th electron (now spin-aligned) are no longer parallel. This effect has not been discussed in Ref. [1] although it is (weakly) visible in Fig. 2 of Ref. [1] for $N = 6 \rightarrow 7$ at $B = 1.3$ T and occasionally more obvious for larger N .




FIG. 1. (a) Addition spectrum vs magnetic field for a circular dot with $\hbar\omega_0 = 5$ meV and $N = 2::20$. This is to be compared with Fig. 2 in Ref. [1]. The encircled region is discussed in the text. (b) KS energy levels for $N = 6$ showing the exchange gap around $B = 1.2$ T. Arrows indicate spin states.

Oliver Steffens and Ulrich Rössler
Institut für Theoretische Physik, Universität Regensburg,
D-93040 Regensburg, Germany

January 11, 2022

PACS number(s): 73.20.Dx, 71.15.Mb, 73.23.-b

- [1] S. Tarucha, D.G. Austing, T. Honda, R.J. van der Hage, and L.P. Kouwenhoven, Phys. Rev. Lett. **77**, 3613 (1996).
- [2] G. Vignale and M. Rasolt, Phys. Rev. B **37**, 10685 (1988).
- [3] M. Ferconi and G. Vignale, Phys. Rev. B **50**, 14722 (1994).