

Overscreened multi-channel $SU(N)$ Kondo model: large- N solution and conformal field theory

Olivier Parcollet, Antoine Georges

Laboratoire de Physique Théorique de l'École Normale Supérieure, 24 rue Lhomond, 75231 Paris Cedex 05, France

Gabriel Kotliar, Anirvan Sengupta

Serin Physics Laboratory, Rutgers University, Piscataway, NJ 08854, USA
(March 23, 2024)

Abstract

The multichannel Kondo model with $SU(N)$ spin symmetry and $SU(K)$ channel symmetry is considered. The impurity spin is chosen to transform as an antisymmetric representation of $SU(N)$, corresponding to a fixed number of Abrikosov fermions $\sum_P f_P^\dagger f_P = Q$. For more than one channel ($K > 1$), and all values of N and Q , the model displays non-Fermi behaviour associated with the overscreening of the impurity spin. Universal low-temperature thermodynamic and transport properties of this non-Fermi liquid state are computed using conformal field theory methods. A large- N limit of the model is then considered, in which $K=N$ and $Q=N$ are held fixed. Spectral densities satisfy coupled integral equations in this limit, corresponding to a (time-dependent) saddle-point. A low-frequency, low-temperature analysis of these equations reveals universal scaling properties in the variable ω/T , in agreement with conformal invariance. The universal scaling form is obtained analytically and used to compute the low-temperature universal properties of the model in the large- N limit, such as the $T=0$ residual entropy and residual resistivity, and the critical exponents associated with the specific heat and susceptibility. The connections with the "non-crossing approximation" and the previous work of Cox and Ruckenstein are discussed.

PACS numbers: 75.20.Hr, 75.30.Mb, 71.10.Hf. Preprint LPTENS : 97/55

I. INTRODUCTION AND MODEL

Multichannel Kondo impurity models [1,2] have recently attracted considerable attention, for several reasons. First, in the overscreened case, they provide an explicit example of a non-Fermi liquid ground-state. Second, these models can be studied by a variety of controlled techniques, and provide invaluable testing grounds for theoretical methods dealing with correlated electron systems. One of the most recent and fruitful development in this respect has been the conformal field-theory approach developed by Afeck and Ludwig [4-6]. Finally, multichannel models have experimental relevance to tunneling phenomena in quantum dots and two level systems [3], and possibly also to some heavy-fermion compounds [2].

In this paper, we consider a generalisation of the multi-channel Kondo model, in which the spin symmetry group is extended from $SU(2)$ to $SU(N)$. In addition, the model has a $SU(K)$ symmetry among the K "channels" (flavours) of conduction electrons. We focus here on spin representations which are such that the model is in the non-Fermi liquid overscreened regime. We shall derive in this paper several universal properties of this $SU(N) \times SU(K)$ Kondo model in the low-temperature regime. Specifically, we obtain: the zero-temperature residual entropy, the zero-temperature impurity resistivity and T -matrix, and the critical exponents governing the leading low-temperature behaviour of the impurity specific heat, susceptibility and resistivity.

These results will be obtained using two different approaches. It is one of the main motivations of this paper to compare these two approaches in some detail. First (Sec.III), we apply CFT methods to study the model for general values of N and K . Then, we study the limit of large N and K , with $K/N = \text{fixed}$. This limit was previously considered by Cox and Ruckenstein [7], in connection with the "non-crossing approximation" (NCA). There is a crucial difference between our work and that of Ref. [7] however, which is that we keep track of the quantum number specifying the spin representation of the impurity by imposing a constraint (on the Abrikosov fermions representing the impurity) which also scales proportionally to N . As a result, the solution of the model at large N follows from a true saddle-point principle, with controllable fluctuations in $1/N$. Hence, a detailed quantitative comparison of the large- N limit to the CFT results can be made. The saddle-point equations are coupled integral equations similar in structure to those of the NCA, except for the different handling of the constraint. The $T = 0$ impurity entropy and residual resistivity are obtained for the first time in analytical form in this paper from a low-energy analysis of these coupled integral equations and shown to agree with the large- N limit of the CFT results. We also demonstrate that the spectral functions resulting from these equations take a universal scaling form in the limit $T \rightarrow 0$, which is precisely that expected from the conformal invariance of the problem.

The Hamiltonian of the model considered in this paper reads:

$$H = \sum_{\mathbf{p}} \sum_{i=1}^K \sum_{\alpha=1}^N \epsilon_{\mathbf{p}} c_{\mathbf{p}i\alpha}^\dagger c_{\mathbf{p}i\alpha} + J_K \sum_{A=1}^{N_K-1} S^A \sum_{\mathbf{p}\mathbf{p}'} c_{\mathbf{p}i}^\dagger t^A c_{\mathbf{p}'i} \quad (1)$$

In this expression, $c_{\mathbf{p}i}^\dagger$ creates an electron in the conduction band, with momentum \mathbf{p} , channel (flavour) index $i = 1, \dots, K$ and $SU(N)$ spin index $\alpha = 1, \dots, N$. The conduction electrons transform under the fundamental representation of the $SU(N)$ group, with generators t^A ($A = 1, \dots, N^2 - 1$). They interact with a localised spin degree of freedom placed

at the origin, $S = fS^A; A = 1; \dots; N$ lg which is assumed to transform under a given irreducible representation R of the $SU(N)$ group.

In the one-channel case ($K = 1$), and when R is taken to be the fundamental representation, this is the Coqblin-Schrieffer model of a conduction gas interacting with a localised atomic level with angular momentum j ($N = 2j + 1$) [9]. In this article, we are interested in the possible non-Fermi liquid behaviour associated with the multi-channel generalisation ($K > 1$) [1] [2]. We shall mostly focus on the case where R corresponds to antisymmetric tensors of Q indices, i.e the Young tableau associated with R is made of a single column of Q indices. In that case, it is convenient to use an explicit representation of the localised spin in terms of N species of auxiliary fermions f_i ($i = 1; \dots; N$), constrained to obey:

$$\sum_{i=1}^N f_i^\dagger f_i = Q \quad (2)$$

so that the $N^2 - 1$ (traceless) components of S can be represented as: $S = f^\dagger f \frac{Q}{N}$. For these choices of R , the Hamiltonian can be written as (after a reshuffling of indices using a Fierz identity):

$$H = \sum_{p, i=1}^X \sum_{i=1}^K \sum_{i=1}^N (c_{pi}^\dagger c_{pi}) + J_K \sum_{p, p^0 i}^X (f_i^\dagger f_{p^0 i} \frac{Q}{N}) c_{pi}^\dagger c_{p^0 i} \quad (3)$$

In a recent paper [10], the case of a symmetric representation of the impurity spin (corresponding to a Young tableau made of a single line of P boxes) has been considered by two of us. In that case, a transition from overscreening to underscreening is found as a function of the "size" P of the impurity spin. In contrast, the antisymmetric representations considered in the present paper always lead to overscreening (except for $K = 1$ which is exactly screened), as shown below. As long as only the overscreened regime is considered, the analysis of the present paper applies to symmetric representations as well, up to some straightforward replacements.

II. STRONG COUPLING ANALYSIS

It is easily checked that a weak antiferromagnetic coupling ($J_K > 0$) grows under renormalisation for all K and N , and all representations R of the local spin. What is needed is a physical argument in order to determine whether the RG flow takes J_K all the way to strong coupling (underscreened or exactly screened cases), or whether an intermediate non-Fermi liquid fixed point exists (overscreened cases).

Following the Nozières and Blandin [1] analysis of the multichannel $SU(2)$ model, we consider the strong-coupling fixed point $J_K = +1$. In this limit, the impurity spin binds a certain number of conduction electrons (at most N/K because of the Pauli principle). The resulting bound-state corresponds to a new spin representation R_{sc} which is dictated by the minimisation of the Kondo energy. For the specific choices of R above, we have proven that:

In the one-channel case ($K = 1$) and for arbitrary N and Q , R_{sc} is the "singlet" representation (of dimension $d(R_{sc}) = 1$). It is obtained by binding $N - Q$ conduction electrons to the Q pseudo-fermions f . The impurity spin is thus exactly screened at strong-coupling.

For all multi-channel cases ($K \geq 2$, arbitrary N and Q), the ground-state at the strong-coupling fixed point is the representation R_{sc} characterised by a rectangular Young tableau with $N - Q$ lines and $K - 1$ columns. Its dimension $d(R_{sc})$ (i.e the degeneracy of the strong-coupling bound-state) is larger than the degeneracy at zero coupling, given by $d(R) = \frac{N!}{Q!(N - Q)!}$.

The Young-tableau associated with the strong-coupling state in both cases is depicted in Fig.1. The detailed proof of these statements and the explicit construction of R_{sc} are given in Appendix A. These properties are sufficient to establish that :

In the one-channel case, the strong coupling fixed point is stable under R.G., and hence the impurity spin is exactly screened by the Kondo effect.

In the multi-channel case, a direct R.G. flow from weak to strong-coupling is impossible, thereby suggesting the existence of an intermediate coupling fixed-point ("overscreening").

The connection between these statements and the above results on the nature and degeneracy of the strong-coupling bound-state is clear on physical grounds. Indeed, it is not possible to flow under renormalisation from a fixed point with a lower ground-state degeneracy to a fixed-point with a higher one, because the effective number of degrees of freedom can only decrease under R.G. Hence, no flow away from the strong-coupling fixed point is possible in the one-channel ($K = 1$) case since the strong-coupling state is nondegenerate. Also, no direct flow from weak to strong coupling is possible for $K \geq 2$ since $d(R_{sc}) > d(R)$. These statements can be made more rigorous [6] by considering the impurity entropy defined as:

$$S_{imp} = \lim_{T \rightarrow 0} \lim_{V \rightarrow \infty} [S(T) - S_{bulk}(T)] \quad (4)$$

Where S_{bulk} denotes the contribution to the entropy which is proportional to the volume V (and is simply the contribution of the conduction electron gas), and care has been taken in specifying the order of the infinite-volume and zero-temperature limits. At the weak coupling fixed point $S_{imp}(J_K = 0) = \ln d(R)$, while $S_{imp}(J_K = \infty) = \ln d(R_{sc})$ at strong-coupling. S_{imp} must decrease under renormalisation [6], a property which is the analogue for boundary critical phenomena to Zamolodchikov's "c-theorem" in the bulk. This suggests a R.G. flow of the kind indicated above. This conclusion can of course be confirmed by a perturbative calculation (in the hopping amplitude) around the strong-coupling fixed-point [1].

The value of S_{imp} will be calculated below at the intermediate fixed-point in the overscreened case, and found to be non-integer (as in the $N = 2$ case [6]).

III. CONFORMAL FIELD THEORY APPROACH

Having established the existence of an intermediate fixed-point for $K \geq 2$, we sketch some of its properties that can be obtained from conformal field-theory (CFT) methods. This is a straightforward extension to the $SU(N)$ case of Aeck and Ludwig's approach for $SU(2)$ [4,5]. The aim of this section is not to present a complete conformal field-theory solution, but

simply to derive those properties which will be compared with the large- N explicit solution given below.

In the CFT approach, the model (1) is first mapped at low-energy onto a $1+1$ -dimensional model of NK chiral fermions. At a fixed point, this model has a local conformal symmetry based on the Kac-Moody algebra $\mathfrak{SU}_K(N)_s \oplus \mathfrak{SU}_N(K)_f \oplus \mathfrak{u}(1)_c$ corresponding to the spin-flavour-charge decomposition of the degrees of freedom. The free-fermion spectrum at the weak-coupling fixed point can be organised in multiplets of this symmetry algebra: to each level corresponds a primary operator in the spin, flavour and charge sectors. A major insight [4] is then that the spectrum at the infra-red stable, intermediate coupling fixed point can be obtained from a "fusion principle". Specifically, the spectrum is obtained by acting, in the spin sector, on the primary operator associated with a given free-fermion state, with the primary operator of the $\mathfrak{SU}_K(N)_s$ algebra corresponding to the representation R of the impurity spin (leaving unchanged the flavour and charge sectors). The "fusion rules" of the algebra determine the new operators associated with each energy level at the intermediate fixed point.

This fusion principle also relates the impurity entropy S_{imp} as defined above to the "modular S-matrix" S_0^R of the $\mathfrak{SU}_K(N)$ algebra (this is the matrix which specifies the action of a modular transformation on the irreducible characters of the algebra corresponding to a given irreducible representation R). Specifically, denoting by $R = 0$ the trivial (identity) representation:

$$S_{\text{imp}} = \ln \frac{S_0^R}{S_0^0} \quad (5)$$

The expression of the modular S-matrix for $\mathfrak{SU}_K(N)$ can be found in the literature [11]. We have found particularly useful to make use of an elegant formulation introduced by Douglas [12], which is briefly explained in Appendices A and B. For the representations R associated to a single column n of length Q (corresponding to (2)), one finds using this representation:

$$S_{\text{imp}} = \ln \prod_{n=1}^Q \frac{\sin \frac{(N+1-n)}{N+K}}{\sin \frac{n}{N+K}} \quad (6)$$

It is easily checked that indeed $S_{\text{imp}} < \ln d(R)$ for all values of N , Q and K . Note also that this expression correctly yields $S_{\text{imp}} = 0$ in the exactly screened case $K = 1$ (for arbitrary $N; Q$).

The low-temperature behaviour of various physical quantities can also be obtained from the CFT approach. At the intermediate coupling fixed point, the local impurity spin acquires the scaling dimension of the primary operator of the $\mathfrak{SU}_K(N)_s$ algebra associated with the $(N^2 - 1)$ dimensional adjoint representation of $SU(N)$. Its conformal dimension s (such that $\langle S(0)S(t) \rangle \sim t^{-2s}$ at $T = 0$) reads:

$$s = \frac{N}{N+K} \quad (7)$$

Integrating this correlation function, this implies that the local susceptibility $\chi_{\text{loc}} / T^{R_{1=T=0}} < S(0)S(\infty) < \infty$ (corresponding to the coupling of an external field to the impurity spin only) diverges at low temperature when $K \geq N$, while it remains finite for $K < N$:

$$\begin{aligned}
K > N &: \chi_{loc} = \frac{1}{T} \quad (K - N) = (K + N) \\
K = N &: \chi_{loc} = \ln T \\
K < N &: \chi_{loc} = \text{const.}
\end{aligned} \tag{8}$$

Exactly at the fixed point, the singular contributions to the specific heat and impurity susceptibility $\chi_{imp} = \chi_{bulk}$ (defined by coupling a magnetic field to the total spin density) vanish [4,13]. Indeed, the singular behaviour is controlled by the leading irrelevant operator compatible with all symmetries that can be generated. An obvious candidate for this operator is the spin, flavour and charge singlet obtained by contracting the spin current with the adjoint primary operator above, $J_1 \sim$. It has dimension $1 + \frac{N}{N+K} = 1 + \chi_s$. This leads to a singular contribution to the impurity susceptibility (arising from perturbation theory at second order in the irrelevant operator [4,13]) of the same nature than for χ_{loc} given in (8):

$$\chi_{imp} = \chi_{loc} \cdot$$

Another irrelevant operator can be constructed in the flavour sector in an analogous manner, namely $J_1^f \sim$. This operator has dimension $K = (N + K)$, which is thus lower than the above operator in the spin sector when $K < N$. This operator could a priori contribute to the low-temperature behaviour of the specific heat, which would lead to a divergent specific heat coefficient $C = T$ for both $K > N$ and $K < N$. On the basis of the explicit large- N calculation presented below, we believe however that this operator is not generated for model (1) when the conduction band d.o.s is taken to be perfectly flat and the cutoff is taken to infinity (conformal limit), so that the specific heat ratio has the same behaviour than the susceptibilities above: $C = T \chi_{loc} = \chi_{imp}$. If the model is extended to an impurity spin with internal flavour degrees of freedom, this operator will however show up (leading to $C = T \chi_{loc} = T \chi_{imp} = T \chi_s$ for $K < N$). It also appears (see below) if an Anderson model generalisation of model (1) is considered away from particle-hole symmetry.

IV. SADDLE-POINT EQUATIONS IN THE LARGE- N LIMIT

We now turn to the analysis of the large- N limit of this model. This will be done by setting:

$$K = N \quad ; \quad J_K = \frac{J}{N} \tag{9}$$

and taking the limit $N \rightarrow \infty$ for fixed values of β , and J , so that the number of channels is also taken to be large. In Ref. [7] (see also [2]), Cox and Ruckenstein considered this limit while holding Q fixed ($Q = 1$). They obtained in this limit identical results to those of the "non-crossing approximation" (NCA) [8]. Here, we shall proceed in a different manner by taking Q to be large as well:

$$Q = q_0 N \tag{10}$$

This insures that a true saddle-point exists, with controllable fluctuations order by order in $1/N$. It will also allow us to study the dependence on the representation R of the local spin, parametrised by q_0 [14]. The approach of Ref. [7] is recovered in the limit $q_0 \rightarrow 0$ (or 1). The action corresponding to the functional integral formulation of model (3) reads:

$$\begin{aligned}
S = & \sum_0^R d_0^P \sum_0^R d_0^{0P} c_1^Y(\cdot) G_0^{-1}(\cdot) c_1(\cdot) \\
& + \sum_0^R d_0^P f^Y(\cdot) @ f(\cdot) + \sum_0^R d_i(\cdot) (f^Y(\cdot) f(\cdot)) \otimes N \\
& + \frac{J}{N} \sum_0^R d_0^P \sum_i c_1^Y(\cdot) c_1(\cdot) f^Y(\cdot) f(\cdot) \otimes
\end{aligned} \quad (11)$$

In this expression, the conduction electrons have been integrated out in the bulk, keeping only degrees of freedom at the impurity site. $G_0(i!_n) = (i!_n - \mu)^{-1}$ is the on-site Green's function associated with the conduction electron bath. In order to decouple the Kondo interaction, an auxiliary bosonic field $B_i(\cdot)$ is introduced in each channel, and the conduction electrons can be integrated out, leaving us with the effective action:

$$\begin{aligned}
S_{\text{eff}} = & \sum_0^R d_0^P f^Y(\cdot) @ f(\cdot) + \sum_0^R d_i(\cdot) (f^Y(\cdot) f(\cdot)) \otimes N + \frac{1}{J} \sum_0^R d_0^P \sum_i B_i^Y B_i \\
& + \frac{1}{N} \sum_0^R d_0^P \sum_i B_i(\cdot) f^Y(\cdot) G_0(\cdot) B_i^Y(\cdot) f(\cdot)
\end{aligned} \quad (12)$$

The quartic term in this expression can be decoupled formally using two bi-local fields $Q(\cdot; \cdot)$ and $\bar{Q}(\cdot; \cdot)$ conjugate to $\sum_i B_i^Y(\cdot) B_i(\cdot)$ and $\sum_i f^Y(\cdot) f(\cdot)$ respectively, leading to the action:

$$\begin{aligned}
S = & \sum_{0,ZZ}^Z d_0^X f^Y(\cdot) @ f(\cdot) + \frac{1}{J} \sum_{0,i}^Z d_0^X B_i^Y B_i + \sum_{0,ZZ}^Z d_i(\cdot) (f^Y(\cdot) f(\cdot)) \otimes N \\
& N \sum_{0,ZZ}^Z d_0^0 \bar{Q}(\cdot; \cdot) G_0^{-1}(\cdot) Q(\cdot; \cdot) + \sum_{0,ZZ}^Z d_0^0 Q(\cdot; \cdot) \sum_i B_i^+(\cdot) B_i(\cdot) \\
& \sum_{0,ZZ}^Z d_0^0 \bar{Q}(\cdot; \cdot) \sum_i f^+(\cdot) f(\cdot)
\end{aligned} \quad (13)$$

The B and f fields can now be integrated out to yield:

$$\begin{aligned}
S = & N \sum_{0,ZZ}^Z d_0^0 \bar{Q}(\cdot; \cdot) G_0^{-1}(\cdot) Q(\cdot; \cdot) + N q \sum_{0,ZZ}^Z d_i(\cdot) \\
& N \text{Tr} \ln[(\otimes \sum_i(\cdot))(\cdot) + \bar{Q}(\cdot; \cdot)] + K \text{Tr} \ln \left[\frac{1}{J} (\cdot) Q(\cdot; \cdot) \right]
\end{aligned} \quad (14)$$

This final form of the action involves only the three fields Q , \bar{Q} and \cdot , and scales globally as N thanks to the scalings $K = N$ and $Q = qN$. Hence, it can be solved by the saddle-point method in the large- N limit. At the saddle-point, $\sum_{sp}(\cdot) = \sum_i$ becomes static and purely imaginary, while $Q_{sp} = Q(\cdot; \cdot)$ and $\bar{Q}_{sp} = \bar{Q}(\cdot; \cdot)$ retain time-dependence but depend only on the time difference \cdot (they identify with the bosonic and fermionic self-energies, respectively).

The final form of the coupled saddle-point equations for the fermionic and bosonic Green's functions $G_f(\cdot) < T f(\cdot) f(0) >$, $G_B(\cdot) < T B(\cdot) B(0) >$ and for the Lagrange multiplier field read:

$$f(\cdot) = G_0(\cdot) G_B(\cdot); \quad B(\cdot) = G_0(\cdot) G_f(\cdot) \quad (15)$$

where the self-energies Σ_f and Σ_B are defined by:

$$G_f^{-1}(i!_n) = i!_n + \Sigma_f(i!_n); \quad G_B^{-1}(i_n) = \frac{1}{J} - \Sigma_B(i_n) \quad (16)$$

In these expressions $\omega_n = (2n + 1)\pi$ and $\omega_n = 2n\pi$ denote fermionic and bosonic Matsubara frequencies. Let us note that the field $B(\omega)$ is simply a commuting auxiliary field rather than a true boson (its equal-time commutator vanishes). As a result $G_B(\omega)$ is a periodic function, but does not share the usual other properties of a bosonic Green function (in particular at high frequency).

Finally, ρ is determined by the third saddle point equation :

$$G_f(\omega = 0) - \frac{1}{N} \sum_n G_f(i\omega_n) e^{i\omega_n 0^+} = \rho \quad (17)$$

These equations are identical in structure to the usual NCA equations, except for the last equation (17) which implements the constraint and allows us to keep track of the choice of representation for the impurity spin.

V. SCALING ANALYSIS AT LOW FREQUENCY AND TEMPERATURE

A. General considerations

The analysis of the NCA equations in Ref. [15] can be applied in order to find the behaviour of the Green's functions in the low temperature, long time regime defined by: T_K^{-1}

$\gg 1$ (where T_K is the Kondo temperature). In this regime, a power-law decay of the Green's functions is found:

$$G_f(\omega) \sim \frac{A_f}{\omega^{2_f}} ; G_B(\omega) \sim \frac{A_B}{\omega^{2_B}} ; (T_K^{-1} \gg 1) \quad (18)$$

The scaling dimensions 2_f and 2_B can be determined explicitly by inserting this form into the above saddle-point equations and making a low-frequency analysis, as explained in Appendix C. This yields:

$$2_f = \frac{1}{1 + \dots} ; 2_B = \frac{1}{1 + \dots} \quad (19)$$

The overall consistency of (15,16) at large time also constrains the product of amplitudes $A_f A_B$ (Eq.(C8) in Appendix C) and dictates the behaviour of the self-energies (denoting by $\rho_0 = \text{Im } G_0(i0^+) =$ the conduction bath density of states at the Fermi level) :

$$\rho_f(\omega) \sim \rho_0 A_B \frac{1}{\omega^{2_B + 1}} ; \rho_B(\omega) \sim A_f \rho_0 \frac{1}{\omega^{2_f + 1}} \quad (20)$$

together with the sum rule:

$$\rho_B(\omega = 0; \omega = 1) = \frac{1}{J} \quad (21)$$

The expression (19) of the scaling dimensions 2_f and 2_B is in complete agreement with the CFT result. Indeed, the fermionic field transforms as the fundamental representation of the $SU(N)_K$ spin algebra, while the auxiliary bosonic field transforms as the fundamental representation of the $SU(K)_N$ flavour algebra, leading to: $2_f = \frac{N^2 - 1}{N(N + K)}$ and $2_B =$

$\frac{K^2 - 1}{K(N + K)}$, which agrees with (19) in the large- N limit. Also, the local impurity spin correlation function, given in the large- N limit by $\langle S_i S_{i+1} \rangle / G_f(\omega) G_f(\omega)$ is found to decay as $1 = 2^{-s}$, with $s = 2 - \epsilon = 1 + \frac{1}{2}$ in agreement with the CFT result (7).

As will be shown below however, these asymptotic behaviours at $T = 0$ do not provide enough information to allow for the computation of the low-temperature behaviour of the impurity free-energy and to determine the $T = 0$ impurity entropy (4). One actually needs to determine the Green's functions in the limit $\epsilon \rightarrow 1$, but for an arbitrary value of the ratio $\epsilon = \frac{1}{2}$ (i.e. to analyze the low-temperature, low-frequency behaviour of (15,16) keeping the ratio $\epsilon = T$ fixed [17]). It is easily seen that in this limit the Green's functions and their associated spectral densities $\rho_{f,B}(\omega) = \frac{1}{\pi} \text{Im} G_{f,B}(\omega + i0^+)$ obey a scaling behaviour:

$$G_{f,B}(\omega) = A_{f,B} T^{2-\epsilon} g_{f,B}(\frac{\omega}{T}) \quad ; \quad \epsilon = \text{arbitrary} \quad (22a)$$

$$\rho_f(\omega) = A_f T^{2-\epsilon} f(\frac{\omega}{T}) \quad ; \quad \rho_B(\omega) = A_B T^{2-\epsilon} B(\frac{\omega}{T}) \quad (22b)$$

In these expressions, $g_{f,B}$ and $f_{f,B}$ are universal scaling functions which depend only on ϵ and q_0 and not on the specific shape of the conduction band or the cutoff. These scaling functions will now be found in explicit form.

B. The particle-hole symmetric representation $q_0 = 1=2$

We shall first discuss the case where the representation R has $Q = N=2$ boxes ($q_0 = 1=2$), for which there is a particle-hole symmetry among pseudo-fermions under $f \leftrightarrow f^\dagger$. The expression of the scaling functions $g_{f,B}$ in that case can be easily guessed from general principles of conformal invariance. The idea is that, in the limit $T_K \rightarrow 0$ with ϵ fixed, the finite-temperature Green's function can be obtained from the $T = 0$ Green's function by applying the conformal transformation $z = \exp(i2\pi \frac{\omega}{T_K})$ [16]. Applying this to the $T = 0$ power-law decay given by (18), one obtains the well-known result for the scaling functions ($\sim \omega^{-\epsilon} = \omega^{-1}$):

$$g_f(\omega; q_0 = 1=2) = \frac{1}{\sin \omega} \quad ; \quad g_B(\omega; q_0 = 1=2) = \frac{1}{\sin \omega} \quad (23)$$

with the periodicity requirements $g_f(\omega + 1) = g_f(\omega)$, $g_B(\omega + 1) = g_B(\omega)$. Note that these functions satisfy the additional symmetry $g_{f,B}(1 - \omega) = g_{f,B}(\omega)$ indicating that they can only apply to the particle-hole symmetric case $q_0 = 1=2$. The corresponding form of the scaling functions associated with the spectral densities (22b) reads, with $\omega = T$ (after a calculation detailed in Appendix C):

$$\begin{aligned} f(\frac{\omega}{T}; q_0 = 1=2) &= \frac{1}{2} (2 - \epsilon)^{2-\epsilon} \cosh \frac{\frac{\omega}{T}}{2} \frac{(\frac{\omega}{T} + i\frac{1}{2}) (\frac{\omega}{T} - i\frac{1}{2})}{(2 - \epsilon)} \\ B(\frac{\omega}{T}; q_0 = 1=2) &= \frac{1}{2} (2 - \epsilon)^{2-\epsilon} \sinh \frac{\frac{\omega}{T}}{2} \frac{(\frac{\omega}{T} + i\frac{1}{2}) (\frac{\omega}{T} - i\frac{1}{2})}{(2 - \epsilon)} \end{aligned} \quad (24)$$

Note that the $\omega \rightarrow 0$ singularity of the $T = 0$ case is now recovered for $\omega \rightarrow 0$ at $T = 0$:

$$g_{f,B}(\omega; q_0 = 1/2) = \frac{1}{\omega! + 1} \frac{1}{(2 - \omega_{f,B})} \frac{1}{\omega!} \quad (25)$$

It is an interesting calculation, performed in detail in Appendix C, to check that indeed these scaling functions do solve the large-N equations (15,16) at finite temperature in the scaling regime. That NCA-like integral equations obey the finite-temperature scaling properties dictated by conformal invariance has not, to our knowledge, been pointed out in the previous literature.

C. General values of q_0

1. Spectral asymmetry

Let us move to the general case of representations with $q_0 \neq \frac{1}{2}$ in which the particle-hole symmetry between pseudo-fermions is broken. The exponent of the power-law singularity in the $T = 0$ spectral densities is not affected by this asymmetry. It does induce however an asymmetry of the prefactors associated with positive and negative frequencies as $\omega \rightarrow 0$. We introduce an angle θ to parametrize this asymmetry, defined such that:

$$g_f(\omega \rightarrow 0^+) = h(\omega; \theta) \frac{\sin(\frac{\omega}{2})}{\omega!} ; \quad g_f(\omega \rightarrow 0^-) = h(\omega; \theta) \frac{\sin(\frac{\omega}{2})}{(\omega!)^2} \quad (26)$$

where $h(\omega; \theta)$ is a constant prefactor. The explicit dependence of θ on q will be derived below. This corresponds to the following analytic behaviour of the Green's function in the complex frequency plane, as $z \rightarrow 0$:

$$G_f^R(z) = h(\omega; \theta) \frac{e^{i\frac{\omega}{2}}}{z^{1-\frac{\omega}{2}}} \quad \text{Im } z > 0 \quad (27)$$

Equivalently, this means that the symmetry $G_f(\omega) = G_f(-\omega)$ is broken, and that the scaling function $g_f(\omega)$ must satisfy (from the behaviour of its Fourier transform):

$$\frac{g_f(0^+)}{g_f(1)} = \frac{\sin(\frac{\omega}{2})}{\sin(\frac{\omega}{2})} \quad (28)$$

We have found, by an explicit analysis of the saddle-point and constraint equations in the scaling regime, which is detailed in Appendix C, that the full scaling functions for this asymmetric case are very simply related to the symmetric ones at $q_0 = 1/2$, through:

$$g_{f,B}(\omega; q) = \frac{e^{(\omega - \frac{1}{2})}}{\cosh \frac{\omega}{2}} g_{f,B}(\omega; q_0 = 1/2) = \frac{e^{(\omega - \frac{1}{2})}}{\cosh \frac{\omega}{2}} \frac{1}{\sin \omega} \quad (29)$$

where the parameter θ is simply related to q so as to obey (28):

$$\theta = \ln \frac{\sin \frac{\omega}{2(1+q)}}{\sin \frac{\omega}{2(1-q)}} \quad (30)$$

Fourier transforming, this leads to the scaling functions for the spectral densities:

$$\begin{aligned} \rho_f(t; q_0) &= \frac{\cosh \frac{t}{2}}{\cosh \frac{t+}{2} \cosh \frac{t-}{2}} \rho_f(t+; q_0 = 1/2) \\ \rho_B(t; q_0) &= \frac{\sinh \frac{t}{2}}{\sinh \frac{t+}{2} \cosh \frac{t-}{2}} \rho_B(t+; q_0 = 1/2) \end{aligned} \quad (31)$$

The thermal scaling function for the fermionic spectral density ρ_f and the bosonic one ρ_B are plotted in Fig.(2) for various values of the asymmetry parameter α . We also note the expression for the maximally asymmetric case $\alpha \rightarrow 1$ (corresponding, as shown below, to $q_0 \rightarrow 0$, i.e. to the limit $Q \rightarrow N$ as in the usual NCA).

$$\rho_f(t) \xrightarrow{\alpha \rightarrow 1} e^{\frac{t}{2}} \frac{(2)^{2-\alpha-1} \rho_f + \frac{t}{2}}{(2-\alpha)} \quad (32)$$

We also note, for further use, the expressions of the full Green's functions in the complex frequency plane (defined by $g_{f=B}(z; \alpha) = \int_{-1}^{R+1} dt \frac{f=B(t)}{z-t}$), in the scaling regime for $\text{Im } z > 0$:

$$g_f(z; \alpha) = \frac{2i(2)^{2-\alpha-1}}{\cosh \frac{z}{2} (2-\alpha) \sin 2-\alpha} \frac{z+}{2} \cos \frac{z+}{2} \sin \frac{z+}{2} \quad (33)$$

$$g_B(z; \alpha) = \frac{2(2)^{2-\alpha-1}}{\cosh \frac{z}{2} (2-\alpha) \sin 2-\alpha} \frac{z+}{2} \sin \frac{z+}{2} \sin \frac{z+}{2} \quad (34)$$

The reader interested in the details of these calculations is directed to Appendix C.

At this stage, the point which remains to be clarified is the explicit relation between the asymmetry parameter α and the parameter q_0 specifying the representation. This is the subject of the next section.

Before turning to this point, we briefly comment on the CFT interpretation of the asymmetry parameter α (or α) associated with the particle-hole asymmetry of the fermionic fields. The form (23) of the correlation functions at finite temperature in the scaling limit can be viewed as those of the exponential of a compact bosonic field with periodic boundary conditions. The asymmetric generalization (29) corresponds to a shifted boundary condition on the boson (i.e. to a twisted boundary condition for its exponential).

2. Relation between q_0 and α

Let us clarify the relation between the spectral asymmetry parameter α , and the parameter q_0 specifying the spin representation. That such a relation exists in universal form is a remarkable fact: indeed α is a low-energy parameter associated with the low-frequency behaviour of the spectral density, while q_0 is the total pseudo-fermion number related by the constraint equation (17) to an integral of the spectral density over all frequencies. The

situation is similar to that of the Friedel sum rule in impurity models, or to Luttinger theorem in a Fermi liquid, and indeed the derivation of the relation between q_0 and χ follows similar lines [19]. It is in a sense a Friedel sum rule for the quasiparticles carrying the spin degrees of freedom (namely, the pseudofermions f).

We start from the constraint equation (17) written at zero-temperature as:

$$q_0 = i \lim_{t \rightarrow 0^+} \int_0^Z \frac{d\omega}{2} G_f(\omega) e^{i\omega t} \quad (35)$$

In this expression, and below in this section, $G_f(\omega)$ and $G_B(\omega)$ denote the (Feynman) $T = 0$ Green's functions while the retarded Green's functions are denoted by G^R . Using analytic continuation of (16), we have:

$$\frac{\partial \ln G_f(\omega)}{\partial \omega} G_f(\omega) \frac{\partial G_f(\omega)}{\partial \omega} = G_f(\omega) \quad (36)$$

$$\frac{\partial \ln G_B(\omega)}{\partial \omega} G_B(\omega) \frac{\partial G_B(\omega)}{\partial \omega} = 0 \quad (37)$$

so that (35) can be rewritten as:

$$q_0 = i \lim_{t \rightarrow 0^+} \int_0^Z \frac{d\omega}{2} \left[\frac{\partial \ln G_f(\omega)}{\partial \omega} G_f(\omega) \frac{\partial G_f(\omega)}{\partial \omega} - \frac{\partial \ln G_B(\omega)}{\partial \omega} G_B(\omega) \frac{\partial G_B(\omega)}{\partial \omega} \right] e^{i\omega t} \quad (38)$$

In this expression, the bosonic part (which vanishes altogether) has been included in order to transform further the terms involving derivatives of the self energy, using analyticity. This transformation is only possible if both fermionic and bosonic terms are considered. This is because the Luttinger-Ward functional [19] of this model involves both Green's functions. It has a simple explicit expression which reads:

$$\Gamma_{LW}(G_f; G_B) = \sum_{ji} \int_0^Z dt G_0(t) G_{fj}(-t) G_{Bi}(t) \quad (39)$$

such that the saddle-point equations (15) are recovered by derivation:

$$G_{fj}(t) = \frac{\Gamma_{LW}}{G_{fj}(-t)}; \quad G_{Bi}(t) = \frac{\Gamma_{LW}}{G_{Bi}(-t)} \quad (40)$$

From the existence of Γ_{LW} , we obtain the sum rule:

$$\int_{-1}^Z d\omega \left[G_f(\omega) \frac{\partial G_f(\omega)}{\partial \omega} - G_B(\omega) \frac{\partial G_B(\omega)}{\partial \omega} \right] = 0 \quad (41)$$

(Note that there is no logarithmic divergence in this integral). After integrating by parts and using the asymptotic behaviour of the two Green's functions in order to eliminate the boundary terms, we get:

$$q_0 = i \int_{-1}^Z \frac{d\omega}{2} \left[\frac{\partial \ln G_f(\omega)}{\partial \omega} - \frac{\partial \ln G_B(\omega)}{\partial \omega} \right] e^{i\omega t} \quad (42)$$

Since $G(!) = G^R(!)$ for $! > 0$ and $G(!) = \overline{G^R(!)}$ for $! < 0$ (with G^R the retarded Green's function), this can be transformed using:

$$\int_{-1}^1 \frac{d!}{2} \frac{\partial \ln G_{fB}(!)}{\partial !} e^{i!0^+} = \int_{-1}^1 \frac{d!}{2} \frac{\partial \ln G_{fB}^R(!)}{\partial !} e^{i!0^+} + \int_{-1}^0 \frac{d!}{2} \frac{\partial}{\partial !} \ln \left(\frac{G_{fB}^R(!)}{G_{fB}^R(!)^A} \right) e^{i!0^+} \quad (43)$$

The first integrals in the right hand side can be deformed in the upper plane and their sum vanishes [19]. Thus we obtain (denoting by $\arg G$ the argument of G):

$$q_f = \arg G_f^R(0) - \arg G_f^R(-1) - \arg G_B^R(0) - \arg G_B^R(-1) \quad (44)$$

$\arg G_f^R(0)$ directly follows from the parametrisation (27) defining ϕ . It can also be read off from the behaviour of the scaling function $g_f(z)$ for $z \rightarrow 1$. Thus, from (33) we can also read off $\arg G_B^R(0)$:

$$\arg G_f^R(0) = -\phi_f; \quad \arg G_B^R(0) = -\phi_f \quad (45)$$

Taking into account that $G_f^R(!) \rightarrow 1$ as $! \rightarrow 1$ and that $\text{Im } G_f^R < 0$ we have $\arg G_f^R(-1) = \phi_f$. Similarly, we have $\arg G_B^R(-1) = 0$. Inserting these expressions into (44), we finally obtain the desired relation between q_f and ϕ (or ϕ_f):

$$\frac{(1 + \phi_f)}{2} = \frac{1}{2} \phi_f; \quad \phi_f = \ln \frac{\sin \frac{\phi_0}{1+\phi_f}}{\sin \frac{(1-\phi_0)}{1+\phi_f}} \quad (46)$$

This, together with (33), fully determines the universal scaling form of the spectral functions in the low-frequency, low temperature limit.

VI. PHYSICAL QUANTITIES AND COMPARISON WITH THE CFT APPROACH

A. Impurity residual entropy at $T = 0$

The impurity contribution to the free energy (per colour of spin) $f_{\text{imp}} = (F - F_{\text{bulk}})/N$ reads, at the saddle point:

$$f_{\text{imp}} = q_0 + T \sum_n^X \ln G_f(i!_n) - T \sum_n^X \ln G_B(i!_n) - \int_0^Z d\phi_f(\phi) G_f(\phi) \quad (47)$$

This expression can be derived either directly from the saddle-point effective action (in which case the last term arises from the quadratic term in Q and \bar{Q}), or from the relation between the free energy and the Luttinger-Ward functional. $F_{\text{bulk}} = N^2 T \text{Tr} \ln G_0$ is the free energy of the conduction electrons. In (47) the formulas $\text{Tr} \ln G$ are ambiguous. We must precisely define which regularisation of these sums we consider: the actual value of the sums depends on the precise definition of the functional integral. For the fermionic field,

the standard procedure of adding and subtracting the contribution of a free local fermion, and introducing an oscillating term to regularize the Matsubara sum holds:

$$\text{Tr} \ln G_f = -T \ln 2 + T \sum_n \ln (i!_n G_f(i!_n)) e^{i!_n 0^+} \quad (48)$$

The situation is somewhat less familiar for the bosonic field. As pointed out above, the latter is merely a commuting auxiliary field (rather than a true boson). We have found that the correct regularisation to be used is:

$$\text{Tr} \ln G_B = T \lim_{N \rightarrow \infty} \sum_{n=-N}^{N-1} \ln (J G_B(i!_n)) \quad (49)$$

The factor of J takes into account the determinant introduced by the decoupling with B , and a symmetric definition of the (convergent) Matsubara sum has been used. Some details and justifications about these regularisations are given in Appendix (D).

We shall perform a low-temperature expansion of Eq.(47), considering successively the particle-hole symmetric ($q_0 = 1/2$) and asymmetric ($q_0 \neq 1/2$) cases, which require rather different treatments.

1. The particle-hole symmetric point $q_0 = \frac{1}{2}$

In this case $\epsilon = 0$, so that the first term in Eq.(47) does not contribute. Let us consider the last term in (47). Using the spectral representation of G_f and the definition of f_ϵ we obtain easily (for $\epsilon = 0$):

$$\sum_0^Z d!_{-f}(\epsilon) G_f(i!_n) = \sum_1^{Z+1} d!_{-f}(\epsilon) \frac{f_\epsilon(i!_n)}{1 + e^{i!_n}} \quad (50)$$

We subtract the value at $T = 0$:

$$(T) - (T = 0) = \sum_0^{Z+1} d!_{-f}(\epsilon) (f_\epsilon(i!_n; T) - f_\epsilon(i!_n; T = 0)) + 2 \sum_0^{Z+1} d!_{-f}(\epsilon) \frac{f_\epsilon(i!_n)}{1 + e^{i!_n}} \quad (51)$$

In the second term, we can replace f_ϵ by its scaling limit. So this term is of order $O(T^{2-\epsilon+1})$. We know the asymptotics of f_ϵ : $f_\epsilon(x) \sim x!_1 C_1 x^{2-\epsilon-1} + C_2 x^{2-\epsilon-3}$. (The term $x^{2-\epsilon-2}$ cancels due to the particle-hole symmetry). Thus, the first term in (51) is of the form: $T^{2-\epsilon+1} \int_0^1 dx x (f_\epsilon(x) - C_1 x^{2-\epsilon-1})$ (the integral is convergent). We conclude that $(T) = (0) + O(T^{2-\epsilon+1})$, so that the last term in (47) does not contribute to the zero-temperature entropy in the particle-hole symmetric case.

Let us express the remaining terms in (47) as integrals over real frequencies, using the regularisations introduced above. As detailed in Appendix D, this leads to the following expression, involving the argument of the (finite-temperature) retarded Green's functions:

$$f_{\text{imp}} = -\frac{1}{2} \sum_1^{Z+1} d!_{-f}(\epsilon) \arctan \frac{G_f^0(i!_n)}{G_f^0(i!_n)} - \frac{1}{2} \sum_1^{Z+1} d!_{-f}(\epsilon) \arctan \frac{G_B^0(i!_n)}{G_B^0(i!_n)} \quad (52)$$

In these expressions n_F (resp. n_B) are the Fermi (resp. Bose) factor.

At this point, it would seem that in order to perform a low-temperature expansion of the free-energy, one has to make a Sommerfeld expansion of the Fermi and Bose factors. This is not the case however, for two reasons: i) the argument of the Green's functions appearing in (52) are not continuous at $t = 0$, so that a linear term in T does appear (as expected from the non-zero value of S_{imp}) ii) the Green's functions have an intrinsic temperature dependence, and the full scaling functions computed above must be used in (52). More precisely, when computing the difference $f_{\text{imp}}(T) - f_{\text{imp}}(T = 0)$, the leading term is obtained by replacing the Green's function by their scaling form (33).

These considerations lead to the following expression of the impurity entropy (per spin colour) $s_{\text{imp}} = S_{\text{imp}}/N$ at zero-temperature for $q_0 = \frac{1}{2}$:

$$s_{\text{imp}} = \frac{1}{2} \int_0^1 dt (a_f(t) - a_f(t=1)) - \frac{1}{2} \int_0^1 dt \frac{1}{e^{tj} + 1} a_f(t) \text{sgn}(t) - \frac{1}{2} \int_0^1 dt (a_b(t) - a_b(t=1)) + \frac{1}{2} \int_0^1 dt a_b(t) \text{sgn}(t) \frac{1}{e^{tj} - 1} \quad (53)$$

In this expression, $a_{f,b}$ denotes the arguments of the scaling functions, obtained from (33):

$$a_f(t) = \arctan \frac{g_f^0(t)}{g_f^{\infty}(t)} = \arctan \cot(\theta_f) \tanh \frac{t}{2} \quad (54)$$

$$a_b(t) = \arctan \frac{g_b^0(t)}{g_b^{\infty}(t)} = \arctan \tan(\theta_f) \tanh \frac{t}{2} \quad (55)$$

From (53), we obtain with $t = \tan \theta_f$

$$\frac{s_{\text{imp}}}{+1} = \frac{1}{2} \int_0^1 du \frac{2 \arctan t}{(1+u^2)} u \arctan \frac{u}{t} + \frac{\arctan(ut)}{u} \frac{\arctan(ut)}{u(1+u^2)} \quad (56)$$

To perform the integration, we note that $\frac{\partial}{\partial t} \frac{\partial}{\partial t} (1+t^2)^{\frac{s_{\text{imp}}}{1+}} = \frac{2t}{(1+t^2)}$ and we obtain finally the simple expression

$$s_{\text{imp}}(q_0 = \frac{1}{2}) = \ln 2 + \frac{1}{2} \int_0^1 \frac{\ln(1+u^2)}{(1+u^2)} du \quad (57)$$

This can also be rewritten, after a change of integration variable, as:

$$s_{\text{imp}}(q_0 = 1/2) = \frac{1}{N} S_{\text{imp}} = \frac{1}{2} \int_0^1 \frac{1}{1+u^2} du + \frac{1}{2} \int_0^1 \frac{1}{1+u^2} du \quad (58)$$

with

$$f(x) = \int_0^x \ln \sin(u) du$$

This coincides with the large- N limit of the CFT result, Eq.(6), in the particle-hole symmetric case.

2. The general case $q_0 \neq 1=2$

For $q_0 \neq \frac{1}{2}$, the first term in Eq.(47) also contributes to the entropy. Indeed, as shown below, the Lagrange multiplier $\beta(T)$ at the saddle-point has a term which is linear in temperature. This stems from a very general thermodynamic relation, which is derived by taking the derivative of Z with respect to q_0 in the functional integral, leading to:

$$\frac{1}{Z} \frac{\partial Z}{\partial q_0} = \frac{\partial F}{\partial q_0} \quad (59)$$

where the average is to be understood with the action (11). At the saddle-point, we thus have: $\beta = \frac{\partial F_{\text{imp}}}{\partial q_0}$ and in particular:

$$\frac{\partial \beta}{\partial T} \Big|_{T=0} = \frac{\partial S_{\text{imp}}}{\partial q_0} \quad (60)$$

We shall directly use this equation in order to compute the residual entropy, by calculating the linear correction in T to β , and then integrating over q_0 . This method shortcuts the full low-temperature expansion of the free-energy (as done in the previous section), which actually turns out to be quite a difficult task to perform correctly for $q_0 \neq 1=2$ [22]. In order to calculate this linear correction, we shall relate β to the high-frequency behaviour of the fermion Green's function. As $G_f(i!_n) \rightarrow 0$ when $!_n \rightarrow \infty$ we have:

$$G_f(i!_n) = \frac{1}{i!_n} - \frac{1}{(i!_n)^2} + o\left(\frac{1}{(i!_n)^2}\right) \quad (61)$$

This shows that β is the discontinuity of the derivative of $G_f(\beta)$ with respect to β at $\beta = 0$:

$$\partial G_f(0^+) + \partial G_f(\beta) = \sum_{!_n=1}^{\infty} d!_n G_f(!_n) \quad (62)$$

Let us define $g(\beta)$ by:

$$G_f(\beta) = \frac{e^{-\beta/2}}{\cosh \frac{\beta}{2}} g(\beta) \quad (63)$$

where β is the spectral asymmetry parameter in Eq.(46) (at this stage we emphasize that the full finite temperature, finite cutoff, Green's function G_f is considered). Equation (62) can be rewritten as:

$$\beta = T + \tanh \frac{\beta}{2} \left(\partial g(0^+) - \partial g(\beta) \right) = \partial g(0^+) + \partial g(\beta) \quad (64)$$

where we have used that $G_f(0^+) + G_f(\beta) = 1$. Denoting by $g(!)$ the spectral function associated with g , we have:

$$\partial g(0^+) - \partial g(\beta) = \sum_{!_n=1}^{\infty} d!_n \frac{g(!_n) - g(\beta)}{1 + e^{-\beta}} \quad (65)$$

In the scaling limit, the spectral function g must become particle-hole symmetric (since the effect of the particle-hole asymmetry in this limit is entirely captured by γ in (63)), and must coincide with $A_f T^{2-\epsilon+1} f(\omega/T; q_0 = 1/2)$. Hence, following the same reasoning than for above, the term in (65) is of order $\text{const.} + O(T^{2-\epsilon+1})$. Thus we have :

$$\frac{\partial}{\partial T} \Big|_{T=0} = \frac{\partial A}{\partial T} \Big|_{T=0} \quad (66)$$

where $A = \partial g(0^+) + \partial g(0^-)$ is the discontinuity of the derivative ∂g . A reflects the particle-hole asymmetry of g and thus vanishes in the scaling limit. Actually the derivative $\frac{\partial A}{\partial T}$ also vanishes as $T \rightarrow 0$ as we now show. Consider first sending the bare cutoff to infinity (along with J) so as to keep the Kondo temperature fixed. In this limit A takes the form : $A = T f \frac{T}{T_K}$. The low energy scaling limit, in which Eq. (29) holds, can be reached by fixing T and sending T_K to infinity. Since g must become particle-hole symmetric in this limit, this implies that $f(x)$ vanishes at small x . Hence, taking a derivative with respect to temperature, of $A = T f \frac{T}{T_K}$ we find that $\frac{\partial A}{\partial T} \Big|_{T=0} = 0$. Thus we finally obtain :

$$\frac{\partial S_{\text{imp}}}{\partial q_0} = \frac{\partial}{\partial T} \Big|_{T=0} = \quad (67)$$

where (q_0) is given in Eq.(46). Integrating this equation over q_0 (taking into account as a boundary condition the value of $S_{\text{imp}}(q_0 = 1/2)$ obtained above), we finally derive the expression of the entropy :

$$S_{\text{imp}} - \frac{1}{N} S_{\text{imp}} = \frac{1}{N} \int_0^1 f \frac{1}{1+u} du - \int_0^1 f \frac{1}{1+u} (1-q_0) du - \int_0^1 f \frac{1}{1+u} q_0 du \quad (68)$$

with, as above: $f(x) = \int_0^x \ln \sin(u) du$. The expression (68) coincides precisely with the large N limit of the CFT result (6) [23]. A plot of the residual entropy and of the asymmetry parameter γ as a function of q_0 is displayed on Fig.3. S_{imp} is maximal at $q_0 = 1/2$ and vanishes as $q_0 \rightarrow 0$ as expected.

In this section, we have discovered that the spectral asymmetry parameter (γ) shares a fairly simple relation with the residual entropy, given by Eq.(67). These are two universal quantities, characteristic of the fixed-point. Remarkably, γ also coincides with the term proportional to T in S (while S itself is non-universal, its linear term in T is). It is tempting to speculate that a deeper interpretation of these facts is still to be found.

B. Internal Energy and Specific Heat

The low-temperature behaviour of the internal energy in the large- N limit can be obtained by two different methods. We shall briefly describe both since they emphasize different and complementary aspects of the physics.

In the first method, we use the effective action in the form (11), before the decoupling with the auxiliary bosonic field $B_i(\omega)$ is made. We thus have a quartic interaction vertex between the conduction electrons at the origin and the Abrikosov fermions representing the quasiparticles in the spin sector, which reads:

$$\frac{J}{N} \sum_{i=1}^N \sum_{\alpha} (f^{\alpha} f^{\alpha}) = \frac{Q}{N} \sum_{i=1}^N c_i^{\alpha} c_i^{\alpha} \quad (69)$$

One can then perform a skeleton expansion of the free-energy functional in terms of the interacting Green's functions for the pseudo-fermions and the conduction electrons, $G_f(\omega)$ and $G_c(\omega)$. The first-order (Hartree) contribution to this functional vanishes because the spin operator in (69) is written in a traceless manner. The next contribution, at second order, yields the most singular contribution at low temperature and reads:

$$E / J^2 \int_0^{\infty} d\omega G_c(\omega)^2 G_f(\omega) \quad (70)$$

At the saddle-point, the interacting conduction electron Green's function is $G_c(\omega) / G_f(\omega) G_B(\omega)$, and hence its dominant long-time behaviour is: $G_c(\omega) \sim 1/\omega$. Inserting this, together with $G_f(\omega) \sim 1/\omega^{2-\epsilon}$ in (70), we see that the leading low-temperature behaviour to the energy reads $E / c_1 T^{4-\epsilon+1} + c_2 T^2 + \dots$, and hence to the specific heat coefficient:

$$\begin{aligned} > 1 : C = T^{-\epsilon+1} \sim \frac{1}{T^{1-\epsilon}} \\ = 1 : C = T^{-\epsilon} \ln 1/T \\ < 1 : C = T^{-\epsilon} \text{ const.} \end{aligned} \quad (71)$$

which agrees with the CFT result described above. We note that there is a quite precise connection between this calculation and the CFT approach: the operator appearing in the Kondo interaction (69) acquires conformal dimension $2(\epsilon_c + \epsilon_f) = 1 + 2\epsilon_f$ and has the appropriate structure of the scalar product of a spin current with the operator S (transforming as the adjoint). Therefore, it is the large- N version of the leading irrelevant operator associated with the spin sector, as described in Sec.III. It is satisfying that the leading low- T behaviour comes from the second-order contribution of this operator in this formalism as well. We note that for the simple Kondo model (1), in the scaling limit, the analogous irrelevant operator in the flavour sector does not show up in the calculation of the energy in the large- N solution. We shall comment further on this point below.

The second method to investigate the internal energy is to push the low-temperature expansion of the free-energy to higher orders. To this end, we need to compute higher-order terms in the expansion (22a) of the Green's functions in the scaling regime. This computation is detailed in Appendix C.3, and leads to:

$$G_f(\omega) = A_f \left[g_f - \frac{1}{\omega^2} + \frac{1}{\omega^4} g_f^{(2)} - \frac{1}{\omega^6} + \frac{1}{\omega^8} g_f^{(3)} - \frac{1}{\omega^{10}} + \dots \right] \quad (72)$$

$$G_B(\omega) = A_B \left[g_B - \frac{1}{\omega^2} + \frac{1}{\omega^4} g_B^{(2)} - \frac{1}{\omega^6} + \frac{1}{\omega^8} g_B^{(3)} - \frac{1}{\omega^{10}} + \dots \right] \quad (73)$$

Let us emphasize that the exponents appearing in this expansion are not symmetric between the bosonic and fermionic degrees of freedom. This is because we are dealing with the Kondo model for which the auxiliary field (bosonic) propagator has no frequency dependence in the non-interacting theory. Also, the expansion given in (72) assumes a perfectly flat conduction

band in the limit of an infinite bandwidth (conformal limit). Using this expansion into the expression (52) of the free-energy leads to a specific heat coefficient: $C = T^{-\alpha} [c_0 T^{2-\alpha-1} + c_1 T^{4-\alpha-1} + c_2 + \dots]$. The coefficient actually vanishes, so that the behavior in (71) is recovered. The vanishing of c_0 was clear in the first approach, where it followed from the absence of Hartree terms. In the CFT approach, it is associated with the fact that the leading irrelevant operator does not contribute to the free energy at first order. The vanishing of c_0 implies non-trivial sum rules relating the scaling functions $g_{f,B}$ and $g_{f,B}^{(2)}$ (which we have not attempted to check explicitly).

We also note that this behaviour of the specific heat is modified when an Anderson model version of the present model is considered (as in Ref. [7]). Because the non-interacting slave boson propagator has a frequency dependence, the exponents of the second-order terms as written in (72) are only correct for $\alpha > 1$ for the Anderson model. For $\alpha < 1$, the term $T^{4-\alpha} g_f^{(2)}(\omega)$ is replaced by $T^{1-\alpha} g_f^{(2)}(\omega)$, while $T^{1-\alpha} g_B^{(2)}(\omega)$ is replaced by $T^{4-\alpha} g_B^{(2)}(\omega)$. As a result, one finds a diverging specific heat coefficient in both cases, with $C = T^{-\alpha} T^{-(1-\alpha)} = T^{1-\alpha}$ for $\alpha > 1$ and $C = T^{-\alpha} T^{-(4-\alpha)} = T^{1-\alpha}$ for $\alpha < 1$. The behaviour for $\alpha < 1$ is due to the leading irrelevant operator in the flavour sector. Similarly, for $\alpha < 1$ in the Anderson model, the susceptibility associated with the flavour (channel) sector χ_f is found to diverge [7], so that a finite Wilson ratio can still be defined as $T^{-1} \chi_f = C$ for $\alpha < 1$.

C. Resistivity and T-matrix

In order to discuss transport properties, we define a scattering T-matrix for the conduction electrons in the usual manner (for a single impurity):

$$G(\mathbf{k}; \mathbf{k}^0; ! + i0^+) = G_0(\mathbf{k}; ! + i0^+) \delta_{\mathbf{k}, \mathbf{k}^0} + G_0(\mathbf{k}; ! + i0^+) T(!) G_0(\mathbf{k}^0; ! + i0^+) \quad (74)$$

where G and G_0 denote the interacting and non-interacting conduction-electron Green's functions, respectively. Taking a flat particle-hole symmetric band for the conduction electron and denoting by ρ_0 the local non-interacting density of states, this yields the local conduction electron Green's function in the form:

$$G(! + i0^+) = \sum_{\mathbf{k}, \mathbf{k}^0} G(\mathbf{k}; \mathbf{k}^0) = \frac{1}{i \rho_0 (1 - i \rho_0 T(!))} \quad (75)$$

Following Ref. [5], we parametrize the zero-frequency limit of the T-matrix in terms of a scattering amplitude S^1 as:

$$T(! = 0) = \frac{i}{2 \rho_0} (1 - S^1) \quad (76)$$

so that, the zero frequency electron Green's function reads:

$$G(i0^+) = \frac{1}{i \rho_0} \frac{1 + S^1}{2} \quad (77)$$

$S^1 = 1$ corresponds to no scattering at all, while $S^1 = -1$ corresponds to maximal unitary scattering (i.e. π -phase shift and vanishing conduction electron density of states at the

in purity site). In the overscreened case, as noted in [5], S^1 is in general such that $|S^1| < 1$, reflecting the non-Fermi liquid nature of the model, and the fact that the actual quasiparticles bear no resemblance to the original electrons. In addition here, we shall find a new feature: namely that S^1 is in fact a complex number for non particle-hole symmetric spin representations (i.e. $q_0 \neq 1/2$).

We first derive an expression for S^1 for arbitrary N, K and spin representation $Q = N q_0$ by generalizing to $SU(N)$ the CFT approach of Ref. [5]. There, it was shown that S^1 can be expressed as a ratio of elements of the modular S -matrix S_{ij} of the $SU(N)_K$ algebra. Denoting by 0 the identity representation, by F the fundamental representation (corresponding to a Young tableau with a single box) and by R the representation in which the impurity lives (Young tableau with a single column of Q boxes), one has [4,5]:

$$S^1 = \frac{S_{FR} = S_{0R}}{S_{F0} = S_{00}} \quad (78)$$

The evaluation of these elements of the modular S -matrix can be done along the same lines than the conformal field theory calculation of the entropy, described above. Some details are given in Appendix B. The result is:

$$S^1 = \frac{\sin \frac{(N+1)}{N+K} \exp \frac{i(1-2q_0)}{N+K} \sin \frac{N}{N+K} \exp \frac{i(N+1)(1-2q_0)}{N+K}}{\sin \frac{N}{N+K}} \quad (79)$$

Notice that S^1 has both real and imaginary parts in the absence of particle hole symmetry $q_0 \neq \frac{1}{2}$.

Let us take the large- N limit of this expression, with $K = N$ fixed. This reads, to first non-trivial order:

$$S^1 = 1 + \frac{2}{N(1+q_0)} \cot \frac{1}{1+q_0} \frac{\cos \frac{(1-2q_0)}{1+q_0}}{\sin \frac{1}{1+q_0}} - \frac{i}{N(1+q_0)} \frac{1}{2q_0} \frac{\sin \frac{(1-2q_0)}{1+q_0}}{\sin \frac{1}{1+q_0}} \quad (80)$$

We now show how to recover this expression from an analysis of the integral equations of the direct large- N solution. Coupling an external source to the conduction electrons in the functional integral formulation of the model, it is easily seen that the conduction electron T -matrix is given, in the large- N limit, by:

$$T(\omega) = \frac{1}{N} G(\omega + i0^+) \quad (81)$$

where G denotes the convolution of the fermion and auxiliary boson Green's function:

$$G(\omega) = G_f(\omega) G_B(\omega) \quad (82)$$

Hence, we have at $T = 0$:

$$S^1 = 1 - \frac{2i}{N} G(i0^+) \quad (83)$$

We first make use of the scaling limit of the two Green's functions, given by (29), and obtain the scaling form of G :

$$G(\omega) = G_f(\omega) G_B(\omega) = \frac{A_f A_B}{\cosh^2 \frac{\omega}{2} \sin \frac{\omega}{2}} + \quad (84)$$

We note that, in this scaling limit, the particle-hole asymmetry of the impurity Green's function has been lost altogether: ϵ has cancelled completely in the dependence of G in this limit and is only present in the prefactor. Thus, only $\text{Re} S^1$ can be extracted from the scaling limit, while $\text{Im} S^1$ requires a more sophisticated analysis. Eq. (84) implies $\text{Im} G = A_f A_B$, and hence $\text{Re} S^1 - 1 = 2^2 A_f A_B \omega = (N \cosh^2 \frac{\omega}{2})$. We make use of the expression (C8) derived in Appendix C for the product of amplitudes $A_f A_B$ and obtain:

$$\text{Re} S^1 - 1 = \frac{1}{(1 + \epsilon) N} \text{Re} \tan \epsilon \frac{i}{2} \quad (85)$$

After expressing ϵ in terms of q using (46) as:

$$\tanh \frac{\omega}{2} = \cot(\epsilon) \tan \frac{(1 - 2q)}{2(1 + \epsilon)} \quad (86)$$

$\text{Re} S^1$ coincides with the real part of (80).

We now consider $\text{Im} S^1$, for which we need to go beyond the scaling limit and use global properties of the Green's functions. First expressing ϵ as a convolution on the imaginary axis and using $\partial_i G_0(i!) = 2i \omega_0(i!)$ in the limit of a flat particle-hole symmetric band we obtain:

$$i \omega_0 - 2 G(i) = A + B(\epsilon) \quad (87)$$

with the definitions:

$$A = \int_{-\infty}^{\infty} d! G_f(i!) \partial_i \epsilon_f(i!) \quad (88)$$

$$B(\epsilon) = \int_{-\infty}^{\infty} d! [G_f(i! + i) \partial_i \epsilon_f(i!) - G_f(i!) \partial_i \epsilon_f(i!)] \quad (89)$$

In the limit of vanishing ϵ , $B(\epsilon)$ can be calculated from the scaling limit of G_f . We obtain:

$$B(\epsilon \rightarrow 0^+) = \frac{1}{1 + \epsilon} \left(e^{\frac{i(1 - 2q)}{1 + \epsilon}} + e^{2i\epsilon} \right) \frac{1}{\sin 2\epsilon} + i \quad (90)$$

On the other hand, A contains high frequency information that is lost in the scaling limit. We find:

$$A = \frac{i}{1 + \epsilon} (1 - 2q) \quad (91)$$

The details of these calculations are provided in Appendix E.

Combining (91,90,87,83) we find agreement with the large N limit of S_1 in Eq. (80).

For a dilute array of impurities (of concentration n_{imp}), the conduction electron-self energy is given by $(i + i0^+)^{-1} n_{\text{imp}} T(i)$, to lowest order in n_{imp} . As shown above, $T(i)$ is given in the large- N approach by the Fourier transform of $G_f(\omega) G_B(\omega)$. The expansion (72) yields the long-time behaviour $G_f(\omega) = A_f^{(2)} \omega^2 + A_f^{(4)} \omega^4 + \dots$ and $G_B(\omega) = A_B^{(2)} \omega^2 + A_B^{(4)} \omega^4 + \dots$.

From the fact that $2_{\text{f}} + 2_{\text{B}} = 1$, this implies: $G_{\text{f}}(\omega) G_{\text{B}}(\omega) = 1 = 1 + 1 = 2_{\text{f}} + 2_{\text{B}}$. Hence the resistivity behaves as:

$$\rho(T) = \rho_{\text{imp}} + \frac{1}{2} \frac{\text{Re} S^1}{cT^{2_{\text{f}}}} + \dots \quad (92)$$

Where ρ_{u} is the impurity resistivity in the unitary limit. For the same reasons as above, the Anderson model result would lead to an exponent 2_{B} in the regime $\omega < 1$ [7].

D. The limit of a large number of channels ($N \rightarrow \infty$)

We finally emphasize that all the expressions derived above greatly simplify in the limit of a large number of channels $N \rightarrow \infty$. This is expected, since in this limit the non-Fermi liquid intermediate coupling fixed point becomes perturbatively accessible from the weak-coupling one [1,24]. The physics of the fixed point can be viewed as an almost free spin of "size" $Q = N q_0$ weakly coupled to the conduction electrons. Indeed the large- N expansion of the entropy (53), the Green function G_{f} (33) and the twist (46) are:

$$\begin{aligned} S_{\text{imp}} &= - (q_0 \ln q_0 + (1 - q_0) \ln(1 - q_0)) - \frac{2 q_0 (1 - q_0)}{6 N^2} + \dots \\ g_{\text{f}}(\omega) &= \frac{e^{i(\omega - \frac{1}{2})}}{\cosh \frac{\omega}{2}} = 1 + \frac{1}{2} \ln \frac{1}{\sin \frac{\omega}{2}} + \dots \\ \rho_{\text{f}}(\omega) &= \frac{1}{T} \left(\frac{1}{T} + \dots \right) \\ &= \ln \frac{q_0}{1 - q_0} + \dots \end{aligned} \quad (93)$$

and the leading terms in these expansions are given by the corresponding quantities for a free spin of "size" $N q_0$. Moreover the scattering matrix S^1 and resistivity have the following expansion:

$$\begin{aligned} \text{Re} S^1 &= 1 - \frac{2 q_0 (1 - q_0)}{N^2} + \dots; \quad \text{Im} S^1 = 0 - \frac{1}{3} \frac{1}{N^2} + \dots \\ \rho(T = 0) = \rho_{\text{imp}} &= \frac{1}{N} q_0 (1 - q_0) \frac{1}{2} + \dots \end{aligned} \quad (94)$$

while the anomalous dimensions read $\gamma_{\text{S}} = 2_{\text{f}} = \frac{1}{2} - \frac{1}{2} + \dots, 2_{\text{B}} = 1 - \frac{1}{2} + \frac{1}{2} + \dots$.

VII. CONCLUSION

In this paper, we have focused on the non-Fermi liquid overscreened regime of the $SU(N)$ $SU(K)$ multichannel Kondo model. This model has actually a wider range of possible behaviour, which become apparent when other kinds of representations of the impurity spin are considered. In a recent short paper [10], two of us have studied fully symmetric representations corresponding to Young tableaux with a single line of P boxes. (This amounts

to consider Schwinger bosons in place of the Abrikosov fermions used in the present work). It was demonstrated that, in that case, a transition occurs as a function of the "size" P of the impurity spin, from overscreening (for $P < K$) to underscreening (for $P > K$), with an exactly screened point in between ($P = K$). The large- N analysis of the overscreened regime $P < K$ is essentially identical to that presented in the present paper for antisymmetric representations.

Obviously, an interesting open problem is to understand the physics of the model for more general impurity spin representations, involving both "bosonic" and "fermionic" degrees of freedom (corresponding respectively to the horizontal and vertical directions in the associated Young tableau). CFT methods are a precious guide in achieving this goal. In particular, the formulas and rules given in Appendices A and B allow for an easy derivation of the impurity $T = 0$ residual entropy and zero-frequency T -matrix, using Alicki and Ludwig's fusion principle and the identification of these quantities in terms of modular S -matrices.

An open question which certainly deserves further studies is to identify which of these more general spin representations are such that a direct large- N solution of the model can be found. This question has obvious potential applications to the multi-impurity problem and Kondo lattice models.

During the course of this study, we learned of a work by A. Jerez, N. Andrei and G. Zarand on the same model using the Bethe Ansatz method. Our results and conclusions agree when a comparison is possible (in particular for the impurity residual entropy and low-temperature behaviour of physical quantities).

ACKNOWLEDGMENTS

We are most grateful to N. Andrei for numerous discussions on the connections between the various approaches. We also acknowledge discussions with S. Sachdev at an early stage of this work, about the importance of the thermal scaling functions. This work has been partly supported by a CNRS-NSF collaborative research grant NSF-INT-14273C O O P. Work at Rutgers was also supported by grant NSF 9529138.

APPENDIX A : THE STRONG COUPLING STATE

We now describe in more details the proof of the statements in Sec. (II) about the nature and degeneracy of the strong-coupling state R_{sc} . For a general reference on the group theory material used in this appendix, the reader is referred e.g. to Ref. [18]. Let us note N_Y the number of electrons brought on the impurity site and by Y the Young tableau with N_Y boxes associated with the representation in which the conduction electrons on the impurity site combine. Because of the Pauli principle, the length of any of its lines must be smaller than K (and hence N_Y must be smaller than NK). Indeed, we must antisymmetrize the wave function separately for each flavour.

The Kondo energy is given by :

$$E = J_K \sum_{\mathbf{S}} \mathbf{S} \cdot \mathbf{S}^0 \quad (\text{A1})$$

with

$$S = f^\dagger f \quad \frac{Q}{N} \quad S^0 = c^\dagger c \quad (A 2)$$

in which f denotes the pseudo-fermion and c the conduction electrons at the impurity site. We can introduce the linear combinations :

$$T = \frac{S + S^0}{2} \quad > \quad T = \frac{S - S^0}{2} \quad < \quad T = S = \quad (A 3)$$

such that:

$$X_{S S} = X_{T^2} \quad (A 4)$$

This leads to the following expression of the Kondo energy:

$$\frac{2E}{J_K} = C_2(R_{sc}) - C_2(Y) - C_2(R) \quad (A 5)$$

in which $C_2(Z)$ denotes the quadratic Casimir operator of the representation Z . The representation R_{sc} is the specific component of $Y \otimes R$ associated with the bound-state formed by the impurity spin and the conduction electrons at strong-coupling. We recall that R is a column of length Q in this paper.

We have to minimize E over all possible choices of Y and of R_{sc} .

First let us recall that for a general representation Y , C_2 is given by :

$$C_2 = \frac{1}{N} \text{tr} A^{-1} \left(\frac{n}{2} + 1 \right) \quad (A 6)$$

where n_i ($1 \leq i \leq N-1$) is the number of columns with length i in the Young tableau Y and A is the Cartan matrix of the $SU(N)$ group [18]. Let us denote by f_j ($1 \leq j \leq N$) the length of the line j in the tableau. Then we have

$$C_2 = \frac{1}{N} \left(\frac{1}{2} \sum_{j=1}^N (f_j - j + N)^2 - \frac{1}{2N} N_Y^2 + \frac{N}{2} \frac{1}{N_Y} - \frac{N(N-1)(2N-1)}{12} \frac{1}{A} \right) \quad (A 7)$$

with $N_Y = \sum_{j=1}^N f_j$ is the number of boxes of Y . Note that with this definition, all f_j 's can be shifted by the same constant without changing the representation (this is because a column of length N can be removed without changing the representation). (A 7) can be given a simple interpretation in terms of N "particles" occupying a set of fermionic levels. This interpretation was introduced in a slightly different form by Douglas [12]. Let $p_j = f_j - j + N$ be the position of the particle j . Because Y is a Young tableau, the particles are ordered and cannot be on the same level. Fig.(4) gives an example of the construction of the diagram associated with a simple Young tableau.

A simple construction of all allowed Young tableaux appearing in the tensor product $Y \otimes R$ [18] can be given in this fermionic language. Starting with the diagram associated with Y , we choose Q particles and raise each of them by one level beginning with the one in

the highest level. (We note that, in the fermionic interpretation, adding a box to line i corresponds to raising the i -th particle up by one level). An example is given in Fig.(5). Let us denote by p_i the positions of the N "particles" in Y , and by p_i^0 the new positions in a given allowed component of $Y \rightarrow R$. The Kondo energy is given by:

$$E = \frac{1}{4N} \sum_{j=1}^N (p_i^0 - p_i^2) + \frac{N_Y}{2N} - \frac{2N-1}{8} C_2(R)^A \quad (A 8)$$

The last two terms are constant (R is held fixed) and can be dropped in the minimisation process.

The p_i 's can be decomposed in two sets: those for which $p_i^0 = p_i$ (we have $N - Q$ of them) and those for which $p_i^0 = p_i + 1$ (Q of them). Let us denote by P the sum of the latter ones. We have:

$$E = \frac{1}{4N} (Q + 2P) + \frac{N_Y}{2N} - \frac{2N-1}{8} C_2(R) \quad (A 9)$$

Thus the lowest energy is achieved for the smallest possible value of P . Since a given shift $p \rightarrow p+1$ can only appear once in the sum (because double occupancies are forbidden and a given particle cannot be raised twice), the absolute minimum is obtained when we sum on all the lowest Q shifts. This implies that the diagram associated with Y has Q particles on the Q lowest levels (from 0 to $Q-1$) and none on the Q -th level.

The upper part of the diagram (above level Q) is then determined by the maximisation of N_Y . Going back to the language of Young tableaux, the minimum is thus achieved when Y is a rectangle of height $N - Q$ and width K , and R_{sc} is given by the same tableau with the first column removed.

Two cases must thus be distinguished:

For ($K = 1$) and for arbitrary N and Q , R_{sc} is the trivial (singlet) representation (of dimension $d(R_{sc}) = 1$).

For ($K \geq 2$, arbitrary N and Q) the dimension $d(R_{sc})$ is larger than the dimension of R . Indeed, denoting by $d_K(R_{sc})$ the dimension R_{sc} for K -channels, we have the recursion relation (from the "hook law" [20]):

$$\frac{d_{K+1}}{d_K} = \frac{(N+K)(N+K-1)}{(Q+K)(Q+K-1)} \cdot \frac{(N+K-Q+1)}{(Q+K+1-Q)} \quad (A 10)$$

(because $Q < N$). It increases with K . The $K = 2$ case is just a column of length $N - Q$ which has the dimension of R . Moreover the inequality is strict for $K > 2$.

APPENDIX B: CONSTRUCTION OF MODULAR S-MATRICES

If two representations R and R^0 correspond to fermion configurations with positions fp_1, \dots, fp_N and fp_1^0, \dots, fp_N^0 (See Appendix A) respectively, then the modular S-matrix element is :

$$S_{R;R^0} = C_{N,K} e^{\frac{2iN}{N+K} \sum_{j=1}^N p_j^0} \det [e^{\frac{2ip_j}{N+K}}] \quad (B1)$$

with $p = \sum_{i=1}^N p_i = N$, $p^0 = \sum_{j=1}^N p_j^0 = N$ and $C_{N,K}$ is a constant which depends only on N and K . Since for the trivial representation 0, the p 's are the consequent integers $0; 1; \dots; N-2; N-1$, $S_{0;R}$ involves a determinant of the form

$$\begin{vmatrix} 1 & 1 & & 1 & 1 \\ z_1 & z_2 & & z_N & z_N \\ z_1^2 & z_2^2 & & z_N^2 & z_N^2 \\ \vdots & \vdots & & \vdots & \vdots \\ z_1^{N-2} & z_2^{N-2} & & z_N^{N-2} & z_N^{N-2} \\ z_1^{N-1} & z_2^{N-1} & & z_N^{N-1} & z_N^{N-1} \end{vmatrix}$$

where $z_j = e^{\frac{2ip_j}{N+K}}$, $p_1; \dots; p_N$ being the positions of fermions corresponding to the representation R . This is just the Van Der Monde determinant $(z) = \prod_{i < j} (z_i - z_j)$. To calculate the T matrix, we also need to know the S -matrix element between the fundamental representation F and an arbitrary representation R . For F , the positions of the fermions are $0; 1; 2; \dots; N-2; N-1$. Therefore $S_{F;R}$ involves the determinant

$$\begin{vmatrix} 1 & 1 & & 1 & 1 \\ z_1 & z_2 & & z_N & z_N \\ z_1^2 & z_2^2 & & z_N^2 & z_N^2 \\ \vdots & \vdots & & \vdots & \vdots \\ z_1^{N-2} & z_2^{N-2} & & z_N^{N-2} & z_N^{N-2} \\ z_1^{N-1} & z_2^{N-1} & & z_N^{N-1} & z_N^{N-1} \end{vmatrix}$$

This determinant has the same antisymmetry property in z_i 's as the Van Der Monde determinant. However, the present determinant is one order higher than (z) as a homogeneous polynomial in z 's. A little reflection shows it is $(\prod_{i=1}^N z_i) (z)$.

Finally we find :

$$\frac{S_{F;R}}{S_{0;R}} = e^{\frac{2iN}{N+K} \sum_{j=1}^N p_j^0} e^{\frac{2ip_j}{N+K}} \quad (B2)$$

Using this formula we deduce (79) from (78).

APPENDIX C : SOLUTION OF THE SADDLE POINT EQUATIONS IN THE SCALING REGIME

In this appendix we solve (15) in the scaling regime as explained in section (V A) and obtain the scaled spectral densities and Green functions.

1. Scaling functions

First we show that (29) is solution of the saddle point equations in the scaling regime. We deal with an arbitrary q_0 . Let us denote by f_{fB} the scaling function of the fermionic and bosonic self-energies :

$$f_{fB}(\omega) = A_{B,f} \omega^{2B_f-1} f_{fB}(-\omega) \quad (C1)$$

G_0 is the local Green function for the conduction electron. Its density of states does not depend on T . So its scaling form is :

$$G_0(\omega) = \frac{1}{\sin \omega} \quad (C2)$$

with $\omega_0 = \frac{1}{2} \text{Im} G_0(i=0)$. Using this formula and (15), f_{fB} are related to g_{fB} . We insert the scaling form (22a) into (16). Matching the power in ω leads to $2_f + 2_B = 1$ and :

$$\begin{aligned} A_f^{-1} g_f^{-1}(i!_n) &= (f(i!_0))^{2_B} A_B (f(i!_n) - f(i!_0)) ; \\ A_B^{-1} g_B^{-1}(i_n) &= \frac{1}{J} B(0)^{2_f} A_f (B(i_n) - B(0)) \end{aligned} \quad (C3)$$

with $i!_n = i(2n+1)$ and $i_n = i2n$. The term $i!_n$ in (16) vanishes in this scaling limit because $f_{fB} < 1$. We assume that at zero-temperature

$$f(0) = \frac{1}{J} \quad B(0) = 0 \quad (C4)$$

so disappears of these equations at lower-order.

We insert our Ansatz into (C3) with the following Fourier transform formulas (which follow from [21] 3.631) :

$$g_f(i!_n) = \frac{(2)^{2_f} T^{2_f-1} i(1)^{n+1} (1 - 2_f)}{\cosh \frac{1}{2} \left(\frac{1}{2} + \frac{i_n}{2} \right)} \quad (C5a)$$

$$g_B(i_n) = \frac{(2)^{2_B} T^{2_B-1} (1)^{n+1} (1 - 2_B)}{\cosh \frac{1}{2} \left(\frac{n}{2} + \frac{i}{2} \right)} \quad (C5b)$$

$$f(i!_n) - f(i!_0) = \frac{i_0 (2)^{2_B+1} T^{2_B} (1)^{n+1} (1 - 2_B)}{\cosh \frac{1}{2} \left(\frac{1}{2} + \frac{i_n}{2} \right)} \quad (n=0) \quad (C5c)$$

$$B(i_n) - B(0) = \frac{0 (2)^{2_f+1} T^{2_f} (1)^{n+1} (1 - 2_f)}{\cosh \frac{1}{2} \left(\frac{n}{2} + \frac{i}{2} \right)} \quad (n=0) \quad (C5d)$$

We see that (29) is solution of (C3) provided that :

The precise form of the cancellation (C 4) at finite temperature is (at leading order in T) :

$$\frac{1}{J_K} \mathcal{Z}_B(0) = \frac{A_f (2)^{2_f+1} T^{2_f} (2_f)}{\cosh \frac{1}{2} \left(\frac{1}{2} \right)^{2_f+1} T^{2_f} (2_f)} \quad (C 6)$$

$$\mathcal{Z}_f(i!_0) = \frac{i A_B (2)^{2_B+1} T^{2_B} (2_B)}{\cosh \frac{1}{2} \left(\frac{1}{2} \right)^{2_B+1} T^{2_B} (2_B)} \quad (C 7)$$

$$(19) \text{ is obeyed : } 2_f = \frac{1}{1+} ; 2_B = \frac{1}{1+}$$

We have the relation between amplitudes :

$$2_B = 2_A A_B (1 - 2_B) (2_B) \frac{\sin \frac{1}{2}}{\cosh^2 \frac{1}{2}} \quad (C 8)$$

In our scaling forms, is the same for the fermionic and the bosonic function. One can check easily that for a more general Ansatz with f and B the saddle-point equations imply $f = B$.

2. Spectral densities

We calculate now the scaled spectral density from the above scaling function. Denoting $= 1$ for fermions and $= 1$ for bosons we have the general formula.

$$G_{fB}(\omega) = \int_0^{\omega} \frac{e^{-\omega''}}{1 - e^{-\omega''}} \mathcal{Z}_{fB}(\omega'') d\omega'' \quad (C 9)$$

In the scaling regime, we have to solve :

$$\frac{e^{(x - \frac{1}{2})}}{\cosh \frac{1}{2}} \frac{1}{\sin(x)} = \int_0^{\omega} \frac{e^{-xu}}{1 - e^{-u}} \mathcal{Z}_{fB}(u) du \quad 0 < x < 1 \quad (C 10)$$

Setting $t = i(x - \frac{1}{2})$ we see it is sufficient to solve :

$$\frac{e^{-it}}{\cosh \frac{1}{2}} \frac{1}{\cosh(t)} = \int_0^{\omega} \frac{e^{-itu}}{e^{\frac{u}{2}} - e^{-\frac{u}{2}}} \mathcal{Z}_{fB}(u) du \quad |\text{Im } t| < \frac{1}{2} \quad (C 11)$$

Due to the properties of Fourier transformation, we can just solve for $\omega = 0$, and obtain the solution for arbitrary ω with (31). With

$$\int_0^{\omega} dt \frac{1}{\cosh(t)} e^{-itu} = (2)^{-1} \frac{\frac{1}{2} + \frac{iu}{2}}{\left(\frac{1}{2} \right)^2 - \left(\frac{iu}{2} \right)^2} \quad 0 < \omega < 1 \quad u \text{ real} \quad (C 12)$$

(see formula 3.313.2 of [21]), we find the result given in the text (24) :

$$\begin{aligned} f_f(t; q_0 = 1/2) &= \frac{1}{2} (2 - t)^2 f_f^{-1} \cosh \frac{t}{2} \frac{(-f_f + i\frac{t}{2})(-f_f - i\frac{t}{2})}{(2 - f_f)} \\ f_B(t; q_0 = 1/2) &= \frac{1}{2} (2 - t)^2 f_B^{-1} \sinh \frac{t}{2} \frac{(-f_B + i\frac{t}{2})(-f_B - i\frac{t}{2})}{(2 - f_B)} \end{aligned} \quad (C13)$$

The asymptotic behaviour follows from formula 8.328 of [21].

We then derive the full Green function by taking a Hilbert transform :

$$g(z) = \frac{1}{2\pi i} \int_{-1}^{Z+1} dx \frac{f(x)}{z - x} \quad (C14)$$

We end (33) using :

The representation

$$\frac{1}{z - u} = \frac{1}{i} \int_0^{Z+1} e^{i(z-u)d} \quad \text{Im } z > 0 \quad (C15)$$

The Fourier formula which inverts (C12)

The formula

$$\int_0^{Z+1} dx \frac{e^{izx}}{\sinh \frac{x}{2}} = 2 \int_0^{Z+1} dx \frac{e^{\frac{izx}{2}} (1 - e^{-x})}{1 - e^{-\frac{x}{2}}} \quad 0 < \text{Im } z < 1 \quad z \text{ real} \quad (C16)$$

which results from formula 3.112.1 of [21].

Finally, we comment on the treatment of the constraint equation (17) in our derivation of the scaling functions. The relation between f_f and q_0 has been derived from a Luttinger sum-rule, which holds at zero-temperature. So one may worry whether the scaling form does satisfy the leading low-temperature corrections to the $T = 0$ constraint equation. We show now that this is actually the case. Starting from Eq.(17) written as:

$$\int_{-1}^{Z+1} dx f_f(x) = q_0 \quad (C17)$$

we subtract the relation at $T = 0$ and take into account the asymptotic behaviour of f_f given by Eq. (25) to obtain :

$$\int_{-1}^{Z+1} dx f_f(x) - \int_{-1}^{Z+1} dx f_f(x) \Big|_{T=0} = \int_{-1}^{Z+1} dx \frac{e^{-x} f_f(x)}{\cosh \frac{x}{2} (2 - f_f)} + \int_{-1}^{Z+1} dx \frac{\text{sgn} x f_f(x)}{e^{x/2} + 1} = 0 \quad (C18)$$

It is a rather strong constraint on the scaling function f_f that this equation should hold, and it is satisfying that the explicit form obtained for f_f does satisfy (C18). This proves that f_f is really a solution of the full system (15,16,17) in the scaling regime at fixed q_0 .

3. Higher-order terms in the scaling expansion

Here, we give some indications on the derivation the expansion in (72). Let us start from the long-time expansion for T_K^{-1} :

$$G_f(\epsilon) = \frac{A_f}{2} \epsilon^{\alpha_f} + \frac{A_f^{(2)}}{2} \epsilon^{\alpha_f+1} \quad G_B(\epsilon) = \frac{A_B}{2} \epsilon^{\alpha_B} + \frac{A_B^{(2)}}{2} \epsilon^{\alpha_B+1} \quad (C19)$$

in which α_f and α_B are exponents to be determined below. Then we have :

$$G_f(\epsilon) = A_f C_{2-f-1} \epsilon^{2-f-1} + A_f^{(2)} C_{-1} \epsilon^{-1} \quad (C20)$$

with $C = \int_0^R dt \frac{e^{-it}}{t+1}$, and a similar expression for G_B . We can then deduce the expansions of α_f and α_B , and insert them into the saddle point equation. We find :

$$\begin{aligned} \frac{A_f C_{2-f-1} \epsilon^{2-f-1}}{A_f C_{2-f-1}} &= \frac{A_f^{(2)} C_{-1} \epsilon^{-1}}{A_f^2 C_{2-f-1}^2} \epsilon^{+1-4-f} = \frac{C_{2-B} A_B \epsilon^0}{A_f^2 C_{2-f-1}^2} \epsilon^{2-B} \frac{C A_B^{(2)} \epsilon^0}{A_f^2 C_{2-f-1}^2} \epsilon^0 \\ \frac{A_B C_{2-B-1} \epsilon^{2-B-1}}{A_B C_{2-B-1}} &= \frac{A_B^{(2)} C_{-1} \epsilon^{-1}}{A_B^2 C_{2-B-1}^2} \epsilon^{+1-4-B} = \frac{C_{2-f} A_f \epsilon^0}{A_B^2 C_{2-B-1}^2} \epsilon^{2-f} \frac{C A_f^{(2)} \epsilon^0}{A_B^2 C_{2-B-1}^2} \epsilon^0 \end{aligned} \quad (C21)$$

The first order yields:

$$= 0 A_f A_B C_{2-f} C_{2-B-1} = 0 A_f A_B C_{2-B} C_{2-f-1} \quad (C22)$$

which, using:

$$C_{-1} / C \quad (C23)$$

gives (19) again. The second equation leads to $\alpha_f = \alpha_B + 1 - 4 - f$.

First suppose $\alpha_f < 1$: in this case we must drop the ϵ^{-1} term but we have

$$\frac{C_{-1} C_{-1}}{C_{2-B-1} C_{2-f-1}} = \frac{C C}{C_{2-B} C_{2-f}} \quad (C24)$$

which implies $\alpha_f = 2 - f$ or $\alpha_f = 2 - f - 1$ (taking C23 into account). So this possibility must be rejected. Finally we are lead to $\alpha_f = 1$ and $\alpha_B = 4 - f$.

The higher order corrections can be dealt with in a similar manner. Restoring the scaling functions, this leads to (72).

APPENDIX D : CALCULATION OF THE RESIDUAL ENTROPY

1. The formula of the free energy

We first give a few more details on the regularisation in (47). We will check that (49) is the right formula for the pseudo-boson.

In the following we will denote by Tr the regularisation with $e^{i\epsilon n^0}$ and by Tr_{sym} the regularisation defined in (49). We note that $\text{Tr}_{\text{sym}} = (\text{Tr}_+ + \text{Tr}_-)/2$ as can be checked explicitly using a spectral representation of the function to be summed.

Let us introduce the following notation for any quantity A (function of ϵ): $A = A^+ - A^-$. As the free energy is particle-hole symmetric, we have :

$$= (\text{Tr} \ln G_f) - (\text{Tr} \ln G_B) \quad (D1)$$

Let us consider

$$(i!_n) = \ln \frac{i!_n + \frac{1}{i!_n}}{i!_n} \quad (D2)$$

such as

$$(\text{Tr} \ln G_f) = (\epsilon = 0) \quad (D3)$$

As ϵ is particle-hole symmetric, we have $(\epsilon = 0^+) = (\epsilon = 0^-)$. As its asymptotic behaviour is $(i!_n) \sim \frac{2}{i!_n}$, its discontinuity is $(\epsilon = 0^+) - (\epsilon = 0^-) = 2$. We obtain :

$$(\text{Tr} \ln G_f) = \quad (D4)$$

This implies that the bosonic term does not contribute to (D1). But there is an analogous relation for the boson : we first calculate the discontinuity of G_B from the saddle-point equations, use an analogous function and obtain $(\text{Tr} \ln G_B) = (1 - 2g)J=2$. So we find

$$(\text{Tr}_{\text{sym}} \ln G_B) = 0 \quad (D5)$$

So we have checked that (49) is the right regularisation for the bosonic term.

2. Derivation of (52)

We consider first the fermionic term. Let $G_0(i!_n) = \frac{1}{i!_n}$ be the Green function of free electrons. We have :

$$\begin{aligned} \text{Tr} \ln G_f &= \text{Tr} \ln 2 - \frac{1}{2} \int_{-R}^R d! (\text{Im} \ln G_f - \text{Im} \ln G_0) n_F(!) \\ &= \text{Tr} \ln 2 + \frac{1}{2} \int_{-R}^R d! \arctan \frac{G_f^0(x)}{G_f^{\text{eff}}(x)} + \frac{1}{2} \int_{-R}^R d! \arctan \frac{G_f^0(x)}{G_f^{\text{eff}}(x)} n_F(!) \end{aligned} \quad (D6)$$

$$= \frac{1}{2} \int_{-1}^1 d! \arctan \frac{G_f^0(x)}{G_f^{\text{eff}}(x)} - \frac{1}{2} n_F(!) \quad (D7)$$

The bosonic term is obtained by an analogous calculation. In the particle hole symmetric case considered in the text the three regularisations for the bosonic term are equivalent. We have :

$$\begin{aligned} \text{Tr}_{\text{sym}} \ln G_B &= \frac{1}{2} \int_{-1}^1 d! \text{Im} \ln (JG(!)) n_B(!) \\ &= \frac{1}{2} \int_{-1}^1 d! \arctan \frac{G_B^{\text{eff}}(x)}{G_B^0(x)} n_B(!) \end{aligned} \quad (D8)$$

Finally we find the formula quoted in the text Eq. (52).

APPENDIX E: SOME DETAILS OF THE T-MATRIX CALCULATION

In this appendix, we calculate A and $B(0^+)$.

1. Computation of A

Using the definition of \mathcal{G}_f and introducing a oscillating term to regulate the 2 integrals, we have :

$$A = \int_0^Z d! G_f(i!) \mathcal{G}_f(i!) \quad (E1)$$

$$= i \int_0^Z d! G_f(i!) e^{i!0^+} + \int_0^Z d! \mathcal{G}_f \ln G_f(i!) e^{i!0^+} \quad (E2)$$

The first term is $2i \phi_f$. Using (46) and

$$\int_0^Z d! \frac{e^{i!0^+}}{i!} = i \quad (E3)$$

(the integral is to be understood as a principal part), we have with $\mathcal{G}_f(z) = \ln(zG_f(z))$

$$A = 2i(1 + \frac{1}{2}) + \int_0^Z dz \mathcal{G}_f(z) \quad (E4)$$

$$= 2i(1 + \frac{1}{2}) - \lim_{\epsilon \rightarrow 0} 2i \operatorname{Im} \left(\frac{1}{i!} \right) (i!) \quad (E5)$$

$$= 2i \quad (E6)$$

We used $\mathcal{G}_f(i!) = 0$ and $\mathcal{G}_f(i!) = \ln(A^{2-\epsilon}) - i$ with A a real constant. Finally we find (91).

2. Computation of $B(0^+)$

$$B(\epsilon) = i \int_0^Z d! (G_f(i! + i) - G_f(i!)) \int_0^Z d! (G_f(i! + i) - G_f(i!)) \mathcal{G}_f^{-1}(i!) \quad (E7)$$

We replace G_f by the scaling function g_f . The second term is of order 1 whereas the first is $O(T^{2-\epsilon})$ and can be neglected. We have then

$$B(\epsilon) = \int_0^Z dx (g_f(i(x+1)\sim) - g_f(ix\sim)) \mathcal{G}_f^{-1}(ix\sim) \quad (E8)$$

with $\sim = \frac{\epsilon}{T}$. We want $B(\epsilon = 0^+; T = 0)$ which is obtained by taking the limit $\sim \rightarrow 0$ in the previous scaling limit of B . To perform this limit we use $g_f(z) = \overline{g_f}(z)$ and the following expansion for g :

$$g_f(ix) = x! + 1 - cA x^{2-\epsilon-1} \quad \text{with} \quad A = i \cosh \frac{\epsilon}{2} + i \epsilon_f \quad (E9)$$

where c is a real constant (Eq. (E9) is obtained directly from (33)). We find :

$$B(0^+) = (1 - 2\epsilon) \int_1^{\infty} dx \frac{e^{ix0^+}}{x^2 - \epsilon(x+1)^{1-2\epsilon}} - \frac{A}{A} \int_1^{\infty} dx \frac{e^{ix0^+}}{x^2 - \epsilon(x+1)^{1-2\epsilon}} + \int_0^{\infty} dx \frac{e^{ix0^+}}{x^2 - \epsilon(x+1)^{1-2\epsilon}} - \int_1^{\infty} dx \frac{e^{ix0^+}}{x} \quad (\text{E10})$$

The last term is a principal part and is given by (E3). We then use the following identity :

$$\begin{aligned} 0 &= \int_{R+i0^+}^{\infty} dz \frac{e^{iz0^+}}{z^2 - \epsilon(z+1)^{1-2\epsilon}} \\ &= \int_1^{\infty} dx \frac{e^{ix0^+}}{x^2 - \epsilon(x+1)^{1-2\epsilon}} + e^{2i\epsilon} \int_1^{\infty} dx \frac{e^{ix0^+}}{x^2 - \epsilon(x+1)^{1-2\epsilon}} \\ &\quad + \int_0^{\infty} dx \frac{e^{ix0^+}}{x^2 - \epsilon(x+1)^{1-2\epsilon}} \end{aligned} \quad (\text{E11})$$

We find

$$\begin{aligned} B(0^+) &= (1 - 2\epsilon) \left[\frac{A}{A} + e^{2i\epsilon} \int_1^{\infty} dx \frac{e^{ix0^+}}{x^2 - \epsilon(x+1)^{1-2\epsilon}} + i \right] \\ &= \frac{A}{1+A} + e^{2i\epsilon} \frac{A}{\sin 2\epsilon} + i \end{aligned} \quad (\text{E12})$$

A simple calculation with (E9) shows :

$$\frac{A}{A} = e^{\frac{i}{1+\epsilon}(1-2\epsilon_0)} \quad (\text{E13})$$

and we find (90).

REFERENCES

- Unité propre du CNRS (UP 701) associée à l'ENS et à l'Université Paris-Sud.
- [1] P. Nozières and A. Blandin, J. Phys. (Paris) 41, 193 (1980).
 - [2] For a recent review on non-Fermi liquid fixed points in Kondo models, see D. L. Cox and A. Zawadowski, preprint cond-mat/9704103, to appear in Advances in Physics.
 - [3] See e.g. "Correlated fermions and transport in mesoscopic systems", T Martin, G Montambaux and J. Tran Thanh Van eds, (Frontieres, 1996), and references therein.
 - [4] IA eck and A W W . Ludwig, Nucl. Phys. B 352, 849 (1991); Nucl. Phys. B 360, 641 (1991).
 - [5] IA eck and A W W . Ludwig, Phys. Rev. B 48, 7297 (1993).
 - [6] IA eck and A W W . Ludwig, Phys. Rev. Lett. 67, 161 (1991).
 - [7] D. L. Cox and A. E. Ruckenstein, Phys. Rev. Lett. 71, 1613 (1993).
 - [8] For a review, see N. E. Bickers, Rev. Mod. Phys. 59, 845 (1987).
 - [9] B. Coqblin and J. R. Schrieffer, Phys. Rev. 185, 847 (1969).
 - [10] O. Parcollet and A. Georges, "Transition from overscreening to underscreening in the multichannel Kondo model: exact solution at large-N" preprint cond-mat/9707337 to appear in Phys. Rev. Lett.
 - [11] V. Kac and D. Peterson, Adv. Math. 53, 125 (1984); V. Kac and M. Wakimoto, Adv. Math. 70, 156 (1988); see also: E. J. Mlawer, S. G. Naculich, H. A. Riggs and H. J. Schnitzer, Nucl. Phys., B 352, 863 (1991).
 - [12] M. R. Douglas preprint hep-th 9403119.
 - [13] V. J. Emery and S. Kivelson, Phys. Rev. B 46, 10812 (1992); A. Sengupta and A. Georges, Phys. Rev. B 49, 10020 (1994); D. G. Clarke, T. Giamarchi and B. Shraiman Phys. Rev. B 48, 7070 (1993).
 - [14] In the exactly screened one-channel case, the q_0 dependence has been investigated both in large N and by the Bethe Ansatz method by P. Coleman and N. Andrei, J. Phys. C 19, 3211, 1986. In that case, Bose condensation of the auxiliary field $B(\cdot)$ takes place in the large-N limit.
 - [15] E. Müller-Hartmann, Z. Phys. B 57, 281 (1984).
 - [16] See e.g. chapter 24 in A. M. Tsvelik "Quantum Field Theory in Condensed Matter Physics", Cambridge University Press, 1995.
 - [17] See also the recent work by S. Sachdev, cond-mat/9705206 and cond-mat/9705266.
 - [18] J. F. Cornwell Group Theory in Physics, vol. 2, Academic Press.
 - [19] Our proof is closest to that of Luttinger's theorem in A. A. Abrikosov, L. P. Gor'kov and I. E. Dzialoshinski, Methods of Quantum Field Theory in Statistical Physics, revised edition by R. A. Silverman, Dover (New York), 1963.
 - [20] H. Georgi Lie Algebras in Particle Physics, Frontiers in Physics, Addison-Wesley.
 - [21] I. S. Gradshteyn and I. M. Ryzhik Table of integrals, series and products, Academic Press, 1980.
 - [22] More precisely, we have found that inserting the scaling functions in $f_{\text{imp}}(T) - f_{\text{imp}}(0)$ and expanding to linear order in T leads, for $q_0 \notin 1=2$, to an incorrect result. Higher order terms apparently cannot be ignored in this expansion! The situation is somewhat similar to $\text{Im } S^1$ in Sec. V C.
 - [23] A numerical calculation of the impurity entropy within the standard NCA approach,

(which differs from ours) has recently appeared in T.-S. Kim and D. L. Cox, Phys. Rev. B 55, 12594 (1997).

[24] J. Gan, N. Andrei and P. Coleman Phys. Rev. Lett 70, 686 (1993)

FIGURES




FIG .1. Young tableau corresponding to the strong-coupling state





FIG. 2. Plot of ϕ_f and ϕ_b as a function of $\tilde{\omega}-\alpha$ for different values of the asymmetry parameter α : $\alpha = 0; -2; -5; -1$ ($\phi_f = \phi_b$).




FIG .3. Residual entropy s_{imp} and α vs. q_0 for $\beta = 1.5$




FIG .4. An example of an $SU(N)$ Young tableau (for $N = 5$) and its associated fermionic representation




FIG. 5. An example of the general composition rule explained in the text. We show the fermionic diagram associated with Y , the resulting fermionic diagrams and their transcription in terms of Young tableaux. (N is arbitrary in this example as evidenced by the ...)