

The exact realisation of the Lanczos Method for a quantum Many-Body System

N.S. Witte

Research Centre for High Energy Physics,
School of Physics, University of Melbourne,
Parkville, Victoria 3052, AUSTRALIA.

April 15, 2024

Abstract

The Lanczos process has been analytically and exactly carried out for the spin 1/2 isotropic XY chain in the thermodynamic limit, yielding a form for the Lanczos coefficient $c_n^2(s)$. This coefficient has a monotonic variation for real positive s and confirms a general theorem on the ground state properties of extensive Many-body Systems. The Taylor expansion of the coefficient about $s = 0$ has a finite radius of convergence, and ground state estimates based on a finite truncation of this are shown to be asymptotic.

PACS: 05.30.-d, 11.15.Tk, 71.10.Pm, 75.10.Jm

The Lanczos Algorithm is one of the few reliable general methods for computing ground state and some excited state properties of interacting quantum Many-body Systems. It has been traditionally employed as a numerical technique on small finite systems, with attendant round-off error problems, although the main obstacle to its further development is the rapid growth of the number of basis states with system size. We wish to demonstrate that the Lanczos Method can be used exactly and analytically for quantum Many-body Systems, and in addition where the thermodynamic limit can also be treated exactly. We demonstrate this for a member of an exactly solvable class of such systems, those falling into the universality class of the two-dimensional Ising Model, namely the isotropic spin 1/2 XY Model. In fact we have provided another notion of solvability, both in the mechanics of constructing the solution and in terms of its general applicability. In addition we state some general theorems arising from the Analytic Lanczos Method Formalism, which apply to both integrable and non-integrable models, where in the latter case a truncation in an expansion necessarily takes place. Given the exact results for our model the effects of this truncation can be examined in minute detail, and we present such a convergence study.

E-mail: nswphysics.unimelb.edu.au

In our convention the Lanczos process generates an orthonormal basis $|j_n\rangle$ from a suitable trial state $|j_0\rangle$ with the recurrence [1, 2]

$$H|j_n\rangle = c_n|j_{n-1}\rangle + c_{n+1}|j_{n+1}\rangle \quad (1)$$

All of our attention focuses on the Lanczos coefficients so defined, c_n and c_{n+1} , rather than the basis states and we perform these steps exactly. It was found recently [3, 4] that the coefficients arising from an extensive Many-body System satisfy a convergence property $c_n(N) \sim N^{-1/2}$ (s) and $c_{n+1}(N) \sim N^{-1/2}$ (s) as $n; N \rightarrow \infty$ where the scaled Lanczos iteration number is $s = n/N$. With such a description, and employing theorems on the extremal zeros of Orthogonal Polynomials, a number of exact general theorems [5] for the ground state energy [3], ground state averages [6] and gaps [7, 4] of an arbitrary extensive Many-body System were found. For example the ground state energy density is given by

$$E_0 = \inf_{s>0} [c(s)^2 - c(s)^2] \quad (2)$$

For the N site chain, our model Hamiltonian is taken to be

$$H = \sum_{i=1}^N S_i^x S_{i+1}^x + S_i^y S_{i+1}^y \quad ; \quad (3)$$

where the label i indexes the sites, and where we only consider the c -cyclic problem, as the thermodynamic limit will be taken. This model, being at the isotropic point, is at a critical point of an anisotropic family of models, and consequently has vanishing excited state gaps. The trial state was taken to be the classical Neel state even though this state is in fact, as far from the true ground state as one could possibly be but its simplicity allows for ease of moment computation, certainly compared to alternatives. So as a "natural" choice we proceed by only considering this choice. In the spin picture the Neel state $|N\rangle$ will be represented by $|j \uparrow \uparrow \uparrow \dots \uparrow \uparrow \uparrow\rangle$, where the quantisation axis is in the z -direction. If we denote the vacuum state for the quasiparticles by $|j_0\rangle$ then the Neel

state can be expressed in terms of the quasiparticle operators ψ_q by

$$|N\rangle = \prod_{q=1}^N \frac{1 + \frac{\psi_q^\dagger \psi_q}{2}}{2} |0\rangle; \quad (4)$$

Here the spectrum of allowed momentum values have been divided up into the set $Z = [0; \pi/2) \cup (\pi/2; \pi)$ and its complement Z^c , and there is a bijection from Z to Z^c defined by $q \rightarrow q + \pi/2$ if $q \in Z$ and $q \rightarrow q - \pi/2$ if $q \in Z^c$. Basically the Neel state is created from the quasiparticles vacuum state by pairs of excitations whose momenta sum to $\pi/2$.

One first constructs the cumulant generating function from its Taylor series expansion about $t = 0$,

$$\ln \langle e^{t \sum_{i=1}^N \psi_i^\dagger \psi_i} \rangle = \sum_{n=1}^N \frac{t^n}{n!} e^{F(t)}; \quad (5)$$

and the exact generating function $F(t)$ was found to be [8]

$$F(t) = \frac{N}{2} \int_0^{\pi/2} dq \ln \cosh(t \epsilon_q); \quad (6)$$

where the quasiparticle energy is $\epsilon_q = j \cos(q)$. So from the Taylor series expansion about $t = 0$ we have the cumulants

$$2n = \frac{2^{2n}}{4n} \frac{1}{(n!)^2} B_{2n} N \quad n \geq 1; \quad (7)$$

where B_{2n} are the standard Bernoulli numbers.

Our moment generating function is also essentially the characteristic function of the underlying measure defining the Lanczos Process, so a Fourier inversion integral yields the weight function $w(\lambda)$ in which the only parameter is the system size N . In what follows we will only need the leading order contribution as $N \rightarrow \infty$ so we perform a stationary phase approximation yielding

$$\ln w(\lambda) = N \left[\frac{1}{2} \int_0^{\pi/2} dq \ln \cosh(\lambda \epsilon_q) \right] + O(1); \quad (8)$$

where λ is the stationary point determined from

$$\lambda = \frac{1}{2} \int_0^{\pi/2} dq \epsilon_q \tanh(\lambda \epsilon_q); \quad (9)$$

and $\epsilon = E/N$ the energy density.

A key link to complete our analysis is a very recent result from the theory of random Matrix Ensembles [9] in which the Lanczos coefficients are determined from the weight function to leading order in $n; N \rightarrow \infty$ by the supplementary condition

$$0 = \frac{1}{2} \int_0^{\pi/2} dq \frac{u^0(\lambda)}{4 \epsilon_q^2 (\epsilon_q)^2}; \quad (10)$$

and the normalisation condition

$$1 = \frac{1}{2} \int_0^{\pi/2} dq \frac{u^0(\lambda)}{4 \epsilon_q^2 (\epsilon_q)^2}; \quad (11)$$

where $u(\lambda) = \ln w(\lambda)$. This pair of equations arises from the continuum Coulomb gas approach to the eigenvalue distribution of random Matrix Ensembles, in the solution for the density of eigenvalues $\rho(\lambda)$, where the $u(\lambda)$ plays the role of a confining one-body potential. The explicit solution for this charge density can be constructed, but we do not require this here. An equivalent derivation [10] of the above results, which does not involve Matrix Ensembles, employs a continuum Coulomb gas approach to the distribution of stationary points arising from the leading order stationary phase approximation to the Selberg integral expression for the Hankel determinants.

In our model $\epsilon(s) = 0$ and our fundamental result then is the following pair of implicit equations for $\lambda^2(s)$

$$\begin{aligned} s &= \frac{1}{2} \int_0^{\pi/2} dq \frac{1}{4 \epsilon_q^2} \\ &= \frac{1}{2} \int_0^{\pi/2} dq \cos(q) \tanh(\lambda \cos(q)); \end{aligned} \quad (12)$$

In Figure 1 we have plotted the dependence of λ on s in a form which illustrates the ground state theorem, Eq.(2), ($\lambda_0 = 1$). Clearly $\lambda^2(s)$ approaches λ_0 as $s \rightarrow 1$, i.e. there is no minimum but an infimum, and the monotonicity of this Lanczos coefficient can be easily shown from the defining equations

$$\frac{ds}{d\lambda^2} = \frac{1}{2} \int_0^{\pi/2} dq \frac{d}{d\lambda^2} \frac{1}{4 \epsilon_q^2}; \quad (13)$$

with $\lambda^2 = \lambda_0^2$. The asymptotic behaviour of this Lanczos coefficient as $s \rightarrow 1$ is also demonstrated in Figure 1, which can be analytically shown to be

$$\lambda^2(s) \sim 1 - \exp(-\frac{1}{4} \frac{1}{3} s); \quad (14)$$

where the pure number $1/4 \approx 0.25$. This is found using the leading order solution for λ as $s \rightarrow 1$

$$\lambda = \frac{1}{2} \int_0^{\pi/2} dq \frac{1}{4 \epsilon_q^2} = \frac{1}{2} \int_0^{\pi/2} dq \frac{1}{4 \cos^2(q)}; \quad (15)$$

Of great interest is the Taylor expansion of $\lambda^2(s)$ about $s = 0$ [11, 12], the "Planchette Expansion", as this forms the basis of the application of the general theorems to non-integrable models. Using the exact defining equations one can perform the two reversions of series developments, which has the initial term s

$$\begin{aligned} \lambda^2(s) &= \frac{1}{2^2} s + \frac{3}{2^2} s^2 + \frac{1}{3} s^3 + \frac{5}{3^2} s^4 + \frac{17}{3 \cdot 5} s^5 \\ &+ \frac{2 \cdot 7 \cdot 127}{3^3 \cdot 5^2} s^6 + \frac{2^2 \cdot 61 \cdot 367}{3^4 \cdot 5^2 \cdot 7} s^7 + \frac{5 \cdot 137}{7^2} s^8 \\ &+ \frac{2 \cdot 17 \cdot 119417}{3^6 \cdot 5 \cdot 7^2} s^9 + \frac{11 \cdot 103 \cdot 299539}{3^8 \cdot 5^3 \cdot 7^2} s^{10} + \dots \end{aligned} \quad (16)$$

This coincides exactly with an early calculation [13] and an independent one using the cumulants directly, although we have calculated many more terms. To indicate the asymptotic behaviour of the coefficients in this Taylor expansion, and the convergence of the series we have plotted the n th root of the coefficients in Figure 2 versus the index. There is a clear limiting value indicating a finite radius of convergence for the expansion, and an interesting quasi-periodic cusp structure superimposed. Another indication of this convergence is given in Figure 3 where the complex zeros of the truncated $d^2(s)$ are plotted in the complex s -plane for a sequence of truncation orders. Clearly here the distribution of zeros approaches a limiting curve, with a simple shape about $s = 0$ and at a finite distance from the origin. As the Lanczos Process terminates when $n! = 0$ at some nonzero n , this curve gives the breakdown of the Plaque Expansion (the exact $d^2(s)$ for reals > 0 is nonzero). One can in fact prove this exactly from considerations of the analytic behaviour of $d(s)$ as a function of complex s . This transformation of the s -plane is analytic except on the imaginary axis $j = j > = 2$ with a sequence of branch points at $s_k = i(k + \frac{1}{2})$. The real part of d has a discontinuity across these cuts (the defined value on the cut is zero), and both parts have inverse square-root singularities when approaching the branch points from one direction along the imaginary axis. The closest zero of $d(s)$ to the origin is relevant in determining the radius of convergence and this occurs as $s \rightarrow 0^+$ (in fact all the zeros lie on the imaginary axis, some at s_k) and therefore the value of s_0 is given by

$$s_0 = \frac{1}{2} \lim_{n \rightarrow \infty} \sqrt[n]{d_n} = \frac{1}{2} \lim_{n \rightarrow \infty} \sqrt[n]{d_n} ; \quad (17)$$

where $d_n = d^{(n)}(0)$. The integrand has the following series development

$$d(s) = \frac{k_0 + 1}{2} + \frac{1}{2} \sum_{k=1}^{\infty} \frac{(k + \frac{1}{2})}{(k + \frac{1}{2})^2 - s^2} ; \quad (18)$$

where $k_0 = b = \frac{1}{2}c$. Using this it is easy to show that

$$s_0 = \frac{1}{2} \lim_{n \rightarrow \infty} \sqrt[n]{d_n} = \frac{1}{2} \lim_{n \rightarrow \infty} \sqrt[n]{d_n} = \frac{1}{2} \lim_{n \rightarrow \infty} \sqrt[n]{d_n} ; \quad (19)$$

Another relevant set of zeros are those of $d^2(s) = ds$ and these are plotted in Figure 4 for the same sequence of truncations. At finite truncations a finite minimum or a complex conjugate pair of stationary points (i.e. an inflection point) of $d^2(s)$ always exists (this pair lie close to the positive real axis), and these can be seen to lie just inside the curve where the Lanczos process breaks down. Finally we plot the relative error in the ground state energy estimates in Figure 5 using the truncated

series directly in the theorem, Eq. (2), versus truncation order. After an initial reduction in the error to the one percent level, the error just continues to fluctuate about this level. The reason why no further improvement is possible with a direct application of the ground state theorem is that the minimum (or inflection point) is pinned just inside the radius of convergence, whereas it should tend to infinity in order to converge.

We have found the exact solution to the Lanczos process for a quantum Many-body System in the thermodynamic limit using results from the theory of random Matrix Ensembles and Orthogonal Polynomial Systems. Our exact method applies to any model where the cumulant generating function can be found, where it should be noted that in this case it was not necessary to fully diagonalise the problem in order to find this. Our exact solution provides an illustration of some general theorems applying to arbitrary extensive Many-body Systems and also demonstrate that these theorems continue to work as an approximate technique even at a critical point in the model, albeit to only a semi-quantitative level. The analytic Lanczos method has recently been applied to a related model with a quantum phase transition, the Ising model in a transverse field [14, 15], to elucidate the dependence of the Lanczos Process away from, near and at criticality. This work was supported by the Australian Research Council.

References

- [1] Y. Saad, Numerical Methods for Large Eigenvalue Problems, Lecture Notes, 1991.
- [2] B. N. Parlett, The Symmetric Eigenvalue Problem, Prentice-Hall, Englewood Cliffs, 1980.
- [3] L. C. L. Hollenberg and N. S. Witte, Phys. Rev. B 54, 16309 (1996).
- [4] N. S. Witte, Many-body orthogonal polynomial systems, in Special Functions and Differential Equations, edited by K. S. Rao, R. Jagannathan, G. Vandenberghe, and J. Van der Jeugt, pages 326-335, New Delhi, 1997, Allied Publishers.
- [5] N. S. Witte, The analytic Lanczos method, in Recent Progress in Many-Body Theories, edited by D. Neilson, Singapore, 1998, World Scientific.
- [6] N. S. Witte, L. C. L. Hollenberg, and Z. Weihong, Phys. Rev. B 55, 10412 (1997).
- [7] L. C. L. Hollenberg, M. P. Wilson, and N. S. Witte, Phys. Lett. B 361, 81 (1995).
- [8] N. S. Witte, Int. J. Mod. Phys. B 11, 1503 (1997).
- [9] Y. Chen and M. E. H. Ismail, J. Phys. A: Math. Gen. 30, 6633 (1997).

- [10] N. S. Witte, The lanczos algorithm for the extensive many-body system in the thermodynamic limit, 1997.
- [11] B. G. Lindsay, Ann. Stat. 17, 711 (1989).
- [12] L. C. L. Hollenberg, Phys. Rev. D 47, 1640 (1993).
- [13] L. C. L. Hollenberg, Phys. Lett. A 182, 238 (1993).
- [14] N. S. Witte and R. Shankar, Moment formalism applied to a solvable model of a quantum phase transition. I. the t -expansion method, 1997.
- [15] N. S. Witte and L. C. L. Hollenberg, Moment formalism applied to a solvable model of a quantum phase transition. II. the lanczos method, 1997.

List of Figures

- | | | |
|---|---|---|
| 1 | Dependence of the Lanczos coefficient b_n on the scaled Lanczos iteration number $s = n/N$ | 4 |
| 2 | The n th root of the Taylor series coefficient b_n defined through the Lanczos coefficient $\beta^2(s) = \sum_{n=1}^{\infty} b_n s^n$ versus index n , indicating the radius of convergence of this series. | 5 |
| 3 | The complex zeros of the truncated Taylor series expansion for $\beta^2(s)$ in the complex s -plane, for truncation orders $r = 25, 50, 75, 100$, and 125 | 5 |
| 4 | The complex zeros of the truncated Taylor series expansion for $d\beta^2(s)/ds$ in the complex s -plane, for truncation orders $r = 25, 50, 75, 100$, and 125 | 6 |
| 5 | The relative error in estimates of the ground state energy using the exact ground state energy theorem and the truncated Taylor series expansion for $\beta^2(s)$, versus the truncation order r | 6 |




Figure 1:




Figure 2:




Figure 3:




Figure 4:




Figure 5: