

Fixed Points in Self-Similar Analysis of Time Series

S. Gluzman¹ and V. I. Yukalov²

¹International Centre of Condensed Matter Physics
University of Brasilia, Brasilia, DF 70919-970, Brazil
and

²Centre for Interdisciplinary Studies in Chemical Physics
University of Western Ontario, London, Ontario N6A 3K7, Canada

Abstract

Two possible definitions of fixed points in the self-similar analysis of time series are considered. One definition is based on the minimal difference condition and another, on a simple averaging. From studying stock market time series, one may conclude that these two definitions are practically equivalent. A forecast is made for the stock market indices for the end of March 1998.

Time series analysis and forecasting have a long history and abundant literature, to mention just a few Refs. [1-3]. When analysing time series, one usually aims at constructing a particular model that could represent the available historical data and, after such a model is constructed, one could use it for predicting future. This kind of approach has been found rather reasonable for describing sufficiently smooth evolution, but it fails in treating large fluctuations, like those happening in stock markets. This failure is caused by quite irregular evolution of markets whose calm development is often interrupted by sudden strong deviations producing booms and crashes. Such deviations are not regular cyclic oscillations

but rather are chaotic events, alike heterophase fluctuations in statistical systems [4]. Similarly to the latter, strong market fluctuations are also of coherent nature, having their origin in the collective nonlinear interactions of many trading agents. The coherent collective behaviour of traders is often termed the crowd or herd behaviour [5-7]. Strong nonlinearity and nonequilibrium of stock markets make them one of the most complex systems existing in nature, comparable with human brain.

A novel approach to analysing and forecasting time series has been recently suggested [8,9]. Being based on the self-similar approximation theory [10-17], this technique can be called the self-similar analysis of time series. In this approach, instead of trying to construct a particular model imitating the dynamical system generating time series, we assume that the evolution of the system is self-similar. This is almost the same as to say that the dynamics of the considered system is governed by some laws. Since the observed time series data are the product of a self-similar evolution, the information on some kind of self-similarity is to be hidden in these data. The role of the self-similar analysis is to extract this hidden information.

We applied the self-similar analysis to stock market time series in Refs. [8,9], where we used two definitions of fixed points resulting in two possible forecasts, $f_n(n+1)$ and $f_n(n+1)$. Here and in what follows we use the notation of Ref. [9]. The aim of the present letter is to pay a special attention to comparing these two ways of defining fixed points. We consider stock market indices for the cases when the answer is known and also make predictions for the end of March 1998.

NYSE Composite. Let us try to make a forecast for the end of February 1998. The following data are available in the period of time from August 31, 1997 till January 31, 1998, taken with one month resolution:

470:48 (Aug:31); 497:23; 481:14; 499:10; 511:19; 510:63 (Jan:31):

The self-similar exponential approximations and corresponding multipliers can be obtained:

$$f_2(3) = 531:074; \quad f_3(4) = 563:269; \quad f_4(5) = 449:993; \quad f_5(6) = 550:72;$$

$$f_2(3) = 538:364; \quad f_3(4) = 566:264; \quad f_4(5) = 438:115; \quad f_5(6) = 539:475;$$

$$M_2(3) = 0:38; \quad M_3(4) = 1:022; \quad M_4(5) = 0:215; \quad M_5(6) = 0:013;$$

$$M_2(3) = 0.495; \quad M_3(4) = 1.115; \quad M_4(5) = 0.347; \quad M_5(6) = 0.005:$$

By the end of February the index was 544.26, which should be compared with $f_5(6)$. Let us make a forecast for the end of March, 1998:

$$f_2(3) = 492.69; \quad f_3(4) = 518.884; \quad f_4(5) = 8042; \quad f_5(6) = 425.835;$$

$$f_2(3) = 484.587; \quad f_3(4) = 522.012; \quad f_4(5) = 2.127 \cdot 10^{14}; \quad f_5(6) = 420.252;$$

$$M_2(3) = 0.184; \quad M_3(4) = 0.042; \quad M_4(5) = 385; \quad M_5(6) = 0.605;$$

$$M_2(3) = 0.492; \quad M_3(4) = 0.1; \quad M_4(5) = 1.12 \cdot 10^{15}; \quad M_5(6) = 0.726:$$

The optimal forecast is $f_3(4)$.

S&P. Let us try to make a forecast for the end of February 1998. The following data are available in the period of time from August 31, 1997 till January 31, 1998, taken with one month resolution:

$$900 \text{ (Aug:31)}; \quad 950; \quad 915; \quad 955; \quad 970; \quad 980 \text{ (Jan:31)}:$$

The self-similar exponential approximations and corresponding multipliers can be obtained:

$$f_2(3) = 994.161; \quad f_3(4) = 1004; \quad f_4(5) = 832.896; \quad f_5(6) = 1058;$$

$$f_2(3) = 999.82; \quad f_3(4) = 995.585; \quad f_4(5) = 798.7576; \quad f_5(6) = 1038;$$

$$M_2(3) = 0.607; \quad M_3(4) = 0.055; \quad M_4(5) = 0.286; \quad M_5(6) = 0.0115;$$

$$M_2(3) = 0.683; \quad M_3(4) = 0.076; \quad M_4(5) = 0.455; \quad M_5(6) = 0.0025:$$

By the end of February the index was 1049, which should be compared with $f_5(6)$. Let us make a forecast for the end of March, 1998:

$$f_2(3) = 953.742; \quad f_3(4) = 979.487; \quad f_4(5) = 915; \quad f_5(6) = 862.455;$$

$$f_2(3) = 941.302; \quad f_3(4) = 985.725; \quad f_4(5) = 915; \quad f_5(6) = 874.712;$$

$$M_2(3) = 0.106; \quad M_3(4) = 0.035; \quad M_4(5) = 0; \quad M_5(6) = 0.104;$$

$$M_2(3) = 0.48; \quad M_3(4) = 0.169; \quad M_4(5) = 0; \quad M_5(6) = 0.105;$$

The optimal forecasts are $f_4(5) = f_4(5)$.

Dow Jones. Let us try to make a forecast for the end of February 1998. The following data are available in the period of time from August 31, 1997 till January 31, 1998, taken with one month resolution:

$$7670 \text{ (Aug:31)}; \quad 8000; \quad 7442; \quad 7823; \quad 7908; \quad 7907 \text{ (Jan:31)};$$

The self-similar exponential approximations and corresponding multipliers can be obtained:

$$f_2(3) = 8043; \quad f_3(4) = 8235; \quad f_4(5) = 6347; \quad f_5(6) = 9069;$$

$$f_2(3) = 8091; \quad f_3(4) = 8152; \quad f_4(5) = 5965; \quad f_5(6) = 8898;$$

$$M_2(3) = 0.396; \quad M_3(4) = 0.015; \quad M_4(5) = 0.436; \quad M_5(6) = 0.007;$$

$$M_2(3) = 0.499; \quad M_3(4) = 0.038; \quad M_4(5) = 0.605; \quad M_5(6) = 0.0015;$$

By the end of February the index was 8546, which should be compared with $f_5(6)$. Let us make a forecast for the end of March, 1998:

$$f_2(3) = 7578; \quad f_3(4) = 7968; \quad f_4(5) = 7442; \quad f_5(6) = 6721;$$

$$f_2(3) = 7431; \quad f_3(4) = 8032; \quad f_4(5) = 7442; \quad f_5(6) = 6890;$$

$$M_2(3) = 0.18; \quad M_3(4) = 0.049; \quad M_4(5) = 0; \quad M_5(6) = 0.15;$$

$$M_2(3) = 0.444; \quad M_3(4) = 0.127; \quad M_4(5) = 0; \quad M_5(6) = 0.168;$$

The optimal forecasts are $f_4(5) = f_4(5)$.

Nasdaq Composite. Let us try to make a forecast for the end of February 1998. The following data are available in the period of time from August 31, 1997 till January 31, 1998, taken with one month resolution:

$$1600 \text{ (Aug:31)}; \quad 1690; \quad 1594; \quad 1601; \quad 1570; \quad 1619 \text{ (Jan:31)};$$

The self-similar exponential approximations and corresponding multipliers can be obtained:

$$f_2(3) = 1506; \quad f_3(4) = 1612; \quad f_4(5) = 1391; \quad f_5(6) = 1878;$$

$$f_2(3) = 1482; \quad f_3(4) = 1618; \quad f_4(5) = 1323; \quad f_5(6) = 1843;$$

$$M_2(3) = 0.302; \quad M_3(4) = 0.0535; \quad M_4(5) = 0.433; \quad M_5(6) = 0.011;$$

$$M_2(3) = 0.429; \quad M_3(4) = 0.064; \quad M_4(5) = 0.567; \quad M_5(6) = 0.0025;$$

By the end of February the index was 1771, which should be compared with $f_5(6)$. Let us make a forecast for the end of March, 1998:

$$f_2(3) = 1569.6; \quad f_3(4) = 1560; \quad f_4(5) = 1663; \quad f_5(6) = 1477;$$

$$f_2(3) = 1566; \quad f_3(4) = 1575; \quad f_4(5) = 1693; \quad f_5(6) = 1509;$$

$$M_2(3) = 0.025; \quad M_3(4) = 0.151; \quad M_4(5) = 0.137; \quad M_5(6) = 0.188;$$

$$M_2(3) = 0.5; \quad M_3(4) = 0.175; \quad M_4(5) = 0.257; \quad M_5(6) = 0.105;$$

The optimal forecast is $f_2(3)$.

As follows from the analysis of forecasts for February, two ways of defining fixed points and leading to $f_n(n+1)$ or $f_{n+1}(n+1)$, respectively, are practically equivalent, the corresponding optimal forecasts being close to each other. How the forecast for March works, we shall check in a month.

References

- [1] D.C. Montgomery and L.A. Johnson, *Forecasting and Time Series Analysis* (McGraw-Hill, New York, 1976).
- [2] R.C. Pindyck and D.L. Rubinfeld, *Econometric Models and Economic Forecasts* (McGraw-Hill, New York, 1991).

- [3] C.W.J. Granger and P. Newbold, *Forecasting Economic Time Series* (Academic, Orlando, 1986).
- [4] V.I. Yukalov, *Phys. Rep.* 208, 395 (1991).
- [5] D. Sornette and A. Johansen, *Physica A* 245, 411 (1997).
- [6] R. Cont and J.P. Bouchaud, preprint cond-m at/9712318 (1997).
- [7] J.P. Bouchaud and R. Cont, preprint cond-m at/9801279 (1998).
- [8] S. Gluzman and V.I. Yukalov, preprint cond-m at/9708143 (1997).
- [9] S. Gluzman and V.I. Yukalov, preprint cond-m at/9710336 (1997).
- [10] V.I. Yukalov, *Physica A* 167, 833 (1990).
- [11] V.I. Yukalov, *J. Math. Phys.* 32, 1235 (1991).
- [12] V.I. Yukalov, *J. Math. Phys.* 33, 3994 (1992).
- [13] V.I. Yukalov and E.P. Yukalova, *Physica A* 206, 553 (1994).
- [14] V.I. Yukalov and E.P. Yukalova, *Physica A* 225, 336 (1996).
- [15] V.I. Yukalov and S. Gluzman, *Phys. Rev. Lett.* 79, 333 (1997).
- [16] S. Gluzman and V.I. Yukalov, *Phys. Rev. E* 55, 3983 (1997).
- [17] V.I. Yukalov and S. Gluzman, *Phys. Rev. E* 55, 6552 (1997).