

Coulomb drag at $\nu = \frac{1}{2}$: Composite fermion pairing fluctuations

Iddo Ussishkin and Ady Stern

Department of Condensed Matter Physics, The Weizmann Institute of Science, Rehovot 76100, Israel
(August 2, 1998)

We consider the Coulomb drag between two two-dimensional electron layers at filling factor $\nu = \frac{1}{2}$ each, using a strong coupling approach within the composite fermion picture. Due to an attractive interlayer interaction, composite fermions are expected to form a paired state below a critical temperature T_c . We find that above T_c pairing fluctuations make the longitudinal transresistivity ρ_{xx} increase with decreasing temperature. The pairing mechanism we study is very sensitive to density variations in the two layers, and to an applied current. We discuss possible relation to an experiment by Lilly et al. [Phys. Rev. Lett. 80, 1714 (1998)].

Double layer systems in the quantum Hall regime exhibit a rich variety of physical phenomena, including transitions between compressible and incompressible states, as well as interlayer coherence [1]. A powerful probe to such systems is a Coulomb drag experiment, in which a current I_1 is driven through one layer, and the voltage V_2 is measured on the other layer. The transresistivity is defined as $\rho_{xy} = V_2/I_1$ for a square sample. A Coulomb drag measurement probes the interlayer interaction, the single layer response, and interlayer correlations.

We concentrate here on the Coulomb drag between identical layers at filling factor $\nu = \frac{1}{2}$ each, which has been recently measured by Lilly et al. [2]. The weak coupling limit of this problem was considered using a composite fermion picture [3,4] and an electronic picture [5], leading to three main results. First, ρ_{xy} is expected to be much larger (typically by 3(4 orders of magnitude) than at zero magnetic field, due to the slow decay of density fluctuations [5]. Second, the low temperature dependence is $\rho_{xy} \sim T^{-4=3}$ [5]. Finally, within a weak coupling theory ρ_{xy} vanishes at $T = 0$. The observed low temperature behavior [2] seems to contradict the last conclusion, and indicates the need to go beyond weak coupling theory.

In this paper we consider the Coulomb drag using a strong coupling approach within the composite fermion picture. Part of the interlayer interaction between composite fermions is attractive [6], and as a result a composite fermion paired state is expected to form below a critical temperature T_c [7]. In electronic terms, such a state is an incompressible quantum Hall state of the double layer system, and is expected to have a large transresistivity, similar to that discussed in Ref. [8]. Experimentally this state is not observed, and ρ_{xy} is always much smaller than the single layer resistivity ρ_{xx} [2]. Motivated by this experimental finding, we consider here the contribution of fluctuations of the paired state above T_c . Our central result is that by including the contribution of pairing fluctuations the transresistivity becomes

$$\rho_{xy} = \frac{2}{4} \frac{e^2}{\rho_{xx}} \frac{T_c}{h T} + \rho_{xy}^{(0)}; \quad (1)$$

where $0 \leq \rho_{xy}^{(0)}$, and $\rho_{xy}^{(0)}$ is the weak coupling con-

tribution [5]. This expression is valid provided that $T > T_c$, but $\rho_{xx} > \rho_{xx}^{(0)}$. Unlike $\rho_{xy}^{(0)}$, the first term in (1) increases with decreasing temperature. While a similar term affects ρ_{xx} as well, ρ_{xy} is much more sensitive to it, being typically three orders of magnitude smaller than ρ_{xx} . Within the mechanism we study, the drag voltage is always parallel to the current, i.e., there is no Hall drag.

In the composite fermion approach [9], the system of electrons is transformed by a suitable Chern-Simons transformation to a system of composite fermions in an average zero field, carrying flux lines attached to them. In the double layer system, if the interlayer spacing d is large, a single-layer transformation is to be performed on each layer separately, leading to the Hamiltonian, in units where $\hbar = c = 1$,

$$H = \frac{1}{2m_b} \sum_i \int d^2r \psi_i^\dagger(r) [i \nabla - \mathbf{a}(r)]^2 \psi_i(r) + \frac{1}{2} \sum_{ij} \int d^2r d^2r^0 : \psi_i^\dagger(r) v_{ij}(r, r^0) : \psi_j(r^0); \quad (2)$$

Here, $\psi_i^\dagger(r)$ is the creation operator for a composite fermion at point r in layer i , m_b is the electron's band mass, $\psi_i(r) = \psi_i^\dagger(r) \psi_i(r) - n$ is the density (with the average value n subtracted), $\mathbf{a}_i(r)$ is the Chern-Simons gauge field (with the external vector potential subtracted), and the colons stand for normal order. The density and the gauge field are related by $\nabla \cdot \mathbf{a}_i(r) = r \mathbf{a}_i(r)$. The Coulomb interaction is given by $v_{ij}(r) = e^2 / (r^2 + d^2(1 - \epsilon_{ij}))$, where ϵ is the bulk dielectric constant. Since the Chern-Simons transformation is carried out separately on each layer, the electronic transresistivity (both longitudinal and Hall components) is identical to that of the composite fermions. At $\nu = \frac{1}{2}$, composite fermions experience no magnetic field on average. Consequently, their Hall transresistivity vanishes, and so does the electronic one. We find this result to hold also around $\nu = \frac{1}{2}$.

The composite fermions interact through the Coulomb interaction and through an intralayer Chern-Simons interaction introduced by the transformation (2). Within

the random phase approximation, the most singular element of the gauge propagator is between anti-symmetric transverse components. This element mediates an interlayer interaction between transverse currents [6],

$$U_{11} = \frac{k_1 \hat{Q}}{m} \frac{k_2 \hat{Q}}{m} \frac{Q=k_F}{T_0 (Q=k_F)^3} : \quad (3)$$

Here, composite fermions of momentum k_1, k_2 are scattered to $k_1 + Q, k_2 - Q$, the energy transfer is $\hbar\omega$, the Fermi momentum is $k_F = \frac{1}{4} n$, and the limits $q \ll k_F = m$ and $q \ll k_F$ are used. The energy scale T_0 is, in the limit of strong screening, $T_0 = e^2 n d$ [5]. The ratio of T_0 and $E_F = \frac{1}{2} k_F$ is proportional to $k_F d$. We assume $T_0 \ll E_F = T$. We use a dressed interaction vertex, inversely proportional to the composite fermion effective mass m^* . The contribution of all other components of the gauge propagator is neglected. The strong singularity of (3) in the limit $Q \rightarrow 0$, which results from the slow dynamics of charge relaxation in the $\nu = 1/2$ state [5], poses the main difficulty in the present calculation.

From Eq. (3) it follows that the interlayer interaction between fermions moving in opposite directions ($k_2 = -k_1$) is attractive [6]. The only bare interlayer interaction is the electrostatic one. Thus, this attractive interaction may be understood in terms of the electronic Coulomb interaction: a composite fermion in momentum state k carries a current j/k . In electronic terms, such a current is accompanied by a chemical potential gradient $(2\hbar = e^2) \nabla j$. The $\nu = \frac{1}{2}$ state is incompressible, so that a chemical potential gradient leads to a density gradient, which is proportional to $\nabla j/k$. The attraction between a composite fermion of momentum k in the first layer and a composite fermion of momentum $-k$ in the second layer results, then, from the Coulomb attraction between the density gradients accompanying the two composite fermions, which are oppositely directed.

The composite fermion attraction is expected to lead to interlayer Cooper pairing below T_c [7]. Above T_c , pairing fluctuations enhance the composite fermion conductivity tensor, and hence σ_D . As in conventional superconductivity [10], there are Aslamazov-Larkin [11] and Maki-Thompson [12] contributions. Pair breaking mechanisms are significant in this problem (see [7] and below). Consequently, the Aslamazov-Larkin term dominates close to T_c . Physically, it corresponds to the enhancement of the conductivity due to the formation of composite fermion Cooper pairs by thermal fluctuations.

We now outline the calculation leading to Eq. (1). A detailed presentation will appear elsewhere. We use the finite temperature Matsubara Green function method. To treat the divergences resulting from the singular interaction (3) at finite temperatures, we separate the interaction, depending on the bosonic Matsubara energy transfer ($\omega_m = 2\pi m T$), into a static part, U^{stat} , when $\omega_m = 0$, and a dynamic part, U^{dyn} , when $\omega_m \neq 0$. The

infrared divergences result from the static part, which may be viewed as analogous to a singular disorder potential that is identical in both layers. These divergences are exactly canceled in the calculation, a result which is a manifestation of Anderson's theorem.

We use the following notation: Calligraphic letters (e.g. U) denote matrices, with indices corresponding to both momentum and Matsubara energy. Matrix multiplication is defined with the accompanying factors,

$$(A B)_{\mathbf{k} \omega; \mathbf{k}' \omega'} = \sum_{\mathbf{m}} \sum_{\mathbf{m}'} \frac{d^2 \mathbf{K}}{(2\pi)^2} A_{\mathbf{k} \mathbf{m}; \mathbf{k}' \mathbf{m}'} B_{\mathbf{k}' \mathbf{m}'; \mathbf{k} \omega'} : \quad (4)$$

The trace over a matrix of this type (denoted by tr) and a multiplication with a vector, which is marked with an arrow, are similarly defined.




FIG. 1. Diagrams for (a) The Aslamazov-Larkin contribution to the composite-fermion transconductivity $\sim \chi^{\text{AL}}$ (b) The particle-particle ladder K (c) The static ladder C . Here, U^{stat} is marked with a dashed line, and U^{dyn} with a dashed wiggly line. The Green function is denoted by a bold line. The vertex function qJ in (a) is derived from Eq. (18).

The diagrams involved in the calculation of the Aslamazov-Larkin contribution are given in Fig. 1. The composite-fermion transconductivity is given by

$$\chi^{\text{AL}} = \lim_{\omega \rightarrow 0} \frac{ie^2}{\omega} \sum_{\mathbf{m}} \sum_{\mathbf{m}'} \frac{d^2 \mathbf{q}}{(2\pi)^2} \mathbf{q}^2 \text{tr} \left[\mathbf{J}(\mathbf{q}; \mathbf{m}) \mathbf{J}(\mathbf{q}; \mathbf{m}') \right] : \quad (5)$$

Here, $K(\mathbf{q}; \mathbf{m})$ is the particle-particle ladder [Fig. 1 (b) and (c)], with total incoming momentum \mathbf{q} and total incoming (bosonic) Matsubara energy ω_m . The vertex function is given by qJ , where J , unlike K , is regular in the long-wavelength low-energy limit at $T = T_c$. The external frequency is ω_m . The calculation is carried out to leading order in $T - T_c$.

Because of the separation of the interaction into a static and dynamic part, the equation for the particle-particle ladder K is also separated into two parts. The integralequation for K [see Fig. 1 (b)] is given by

$$K(q; !_m) = U^{\text{dyn}}(q) + U^{\text{dyn}}(q) C(q; !_m) K(q; !_m) : \quad (6)$$

where the static ladder C is the solution of the integral equation [see Fig. 1(c)]

$$C(q; !_m) = G(q; !_m) + G(q; !_m) U^{\text{stat}}(q) C(q; !_m) : \quad (7)$$

Here,

$$G_{k_m, k^0_m}(q; !_m) = \frac{(2!)^2}{T} m m^0^2 (k - k^0) G(k + q; m + !_m) G(k; m) ; \quad (8)$$

where $m = (2m + 1) T$ is a fermionic Matsubara energy. The static ladder is analogous to the Cooperon in disordered systems. For the Green function, $G(k; m) = [i_m - k + (k; m)]^{-1}$, we use the self-consistent Born approximation for the static interaction,

$$(k; m) = T \int \frac{d^2 k^0}{(2!)^2} U_{k_m, k^0_m}^{\text{stat}}(q=0) G(k^0; m) : \quad (9)$$

Eq. (9) results from the observation that the most singular term in the intralayer interaction is U_{11} . The factor T appeared from the separation of the Matsubara sum. The contribution of U^{dyn} to the self-energy is non-divergent. Thus, its inclusion here is not essential. The angular averaging of U^{stat} is, for small $|k - k^0|$,

$$\int \frac{d}{2} U_{k_m, k^0_m}^{\text{stat}}(q=0) = \frac{k_F^3}{4 m^2 T_0 |k - k^0|} m m^0 ; \quad (10)$$

where k and k^0 are assumed be close to k_F , and θ is the angle between k and k^0 . The divergence due to the interaction is thus logarithmic. To regularize the divergence, the singular term $1 = |k - k^0|$ may be replaced with $\tanh[(k - k^0)/(k - k^0)] = (k - k^0)$, and the limit $|k - k^0| \rightarrow 1$ taken at the end of the calculation. In our case the terms involving U^{stat} are canceled exactly.

We first describe the solution of the particle-particle ladder [Eqs. (6)-(9)] at $q = !_m = 0$, where the pairing instability emerges at the critical temperature T_c . The static ladder C has a singularity at $q = 0$ (after analytic continuation $i_m \rightarrow +i$), where resonant scattering occurs. The terms contributing to the singular part C^s are those with multiple scattering lines. As a result, C^s is separable in the momentum variables:

$$C_{k_m, k^0_m}^s = \frac{m m^0}{T} \frac{1}{N_0} \frac{\text{Im} G^+(k; 0) \text{Im} G^+(k^0; 0)}{j_m j} ; \quad (11)$$

where N_0 is the density of states at the Fermi energy and the energy is measured from the Fermi surface.

The particle-particle ladder $K(q = !_m = 0)$ is singular at $T = T_c$. To calculate the singular part K^s [using

Eq. (6)], an angular averaging over the Fermi surface of $U^{\text{dyn}}(q = 0)$ is needed [cf. (10)]. For $m \ll m^0$ it is

$$U_{m, m^0}^{\text{dyn}} = \frac{2}{3} \frac{k_F^2}{3 m^2 T_0^{2=3} j_m j} : \quad (12)$$

This result assumes $T_0 (|k - k^0| = k_F)^3 = j_m j$ an assumption justified a posteriori by Eq. (15). After averaging, the dynamic interaction is momentum independent, and thus K^s is momentum independent and separable in the energy variables. We therefore write

$$K_{k_m, k^0_m}^s = \frac{1}{N_0} \frac{T_c}{T} m m^0 : \quad (13)$$

Using Eq. (6), at $T = T_c$ the vector χ , whose km component is χ_m , satisfies $\chi = U^{\text{dyn}} C \chi$, i.e.,

$$\chi_m = \frac{X}{T} \int_{m^0 \in m} U_{m, m^0}^{\text{dyn}} \frac{N_0}{j_m j} \chi_{m^0} : \quad (14)$$

The highest temperature for which (14) has a solution is T_c . Note that Eq. (14) is independent of U^{stat} . Solving (14) numerically, we find

$$T_c = 0.32 \frac{E_F^3}{T_0^2} ; \quad (15)$$

where we used $N_0 = m^2$ and $E_F = k_F^2 = 2m$. Eq. (15) is in agreement with Ref. [7]. As pointed out by Bonesteel et al. [7], T_c is reduced by symmetric density fluctuations, which the composite fermions view as a fluctuating pair-breaking magnetic field. Impurities, which exert a different disordered potential in each of the two layers, further suppress T_c . Additional controllable mechanisms for T_c suppression are discussed below.

An expansion of the results for small q and $!$ is needed for the denominator of K^s in (13), while the numerator is regular in the long-wavelength low-energy limit at $T = T_c$. At finite q and $!_m > 0$, Eq. (14) is written with U^{dyn} and the denominator of C^s taken at q and $!_m$, and the sum limited to $m^0 = 0$ and $m^0 < m$. The temperature at which Eq. (14) has a solution, which is T_c at $q = !_m = 0$, is then expanded for small q and $!$ (after analytic continuation $i!_m \rightarrow +i$). We find,

$$K_{k_m, k^0_m}^s = \frac{1}{N_0} \frac{T_c}{T} \frac{1}{i! + q} m m^0 : \quad (16)$$

Numerically we find 0.12 . The calculation of χ , and of the vertex function is difficult since both depend on U^{stat} . However, they are related by

$$\chi_y = J \chi = \frac{2N_0}{T_c} : \quad (17)$$

Here, the expansion of the vertex function to first order in q at $!_m = \tilde{m} = 0$ is qJ . This equality is a consequence of a Ward identity. It is shown by writing J as

$$J = \frac{1}{2} \sum_{x,y} \left(\partial_q^2 C^s + 2\partial_q C^s \right) \psi^{dyn} \partial_q C^s \quad (18)$$

$$+ 2\partial_q C^s \partial_q U^{dyn} \tilde{C} + 2C^s \partial_q U^{dyn} \partial_q C^s ;$$

and comparing with the expansion for ρ , described above. We note that diagrams with ψ_a^2 vertices appear in (18) when a derivative with respect to q is applied to a transverse current vertex of U [see Eq. (3)]. Eq. (17) leads to a cancelation of both J and ρ from ρ .

Finally, ρ may be calculated. The calculation of the composite-fermion transconductivity $\sim \rho^{A1}$ (5), using Eqs. (16) and (17), is similar to the Aalam azov-Larkin calculation [11]. The conductivity tensor is then inverted to obtain the composite-fermion transresistivity, which is equal to the (physical) electronic transresistivity ρ . Assuming ρ_{xx} , we finally obtain Eq. (1). We now comment on some aspects of that result.

First, we note that pairing fluctuations lead to an increase of the transresistivity as the temperature is decreased, in sharp contrast to the temperature dependence of the weak coupling contribution.

Second, the pairing contribution to ρ is much more sensitive to a small density difference n between the layers than the weak coupling one. Such a density difference amounts to a difference $\mu = (\partial \mu / \partial n) n$ between the chemical potentials of the two layers, where $\partial \mu / \partial n = \epsilon^2 = \kappa$. Pairing between the layers is gradually suppressed by μ , in a way similar to the suppression of conventional Cooper pairing by spin polarization. Complete suppression of the pairing contribution to ρ happens when $\mu = T_c$, i.e., when $n = n(T_c = E_F)$. Similarly, the pairing mechanism is also sensitive to a symmetric deviation of the density in both layers to a filling factor $\nu = \frac{1}{2} + \delta$. In that case both layers are subject to a uniform Chern-Simons magnetic field B / ν , which suppresses pairing. In realistic samples we expect both dependences, on symmetric and asymmetric density deviation from $\nu = 1/2$, to be mostly determined by the inevitable disorder-induced fluctuations in the density.

Third, we note that the pairing contribution is also easily suppressed by the current j_1 flowing in one of the layers. Qualitatively, such a current shifts the composite fermion Fermi sphere in the current carrying layer relative to the one in the other layer by κ / j_1 , and thus suppresses the critical temperature to $T_c \sim (\kappa / j_1)^2$. In a very naive estimate $\kappa = T_c$ and $\kappa \sim j_1 = ne$.

Experimentally, Lilly et al. [2] observed a non-monotonic temperature dependence of ρ over some narrow ranges of magnetic field in the vicinity of $\nu = \frac{1}{2}$. The precise position of these ranges seems to be sample specific, and the non-monotonic temperature dependence is observed only at very low currents. The analysis above suggests that this phenomenon is the result of pairing fluctuations in those parts of the sample at which both layers are at filling factor very close to $\frac{1}{2}$. The precise

values of magnetic field at which such parts exist depend on the specific disorder configuration.

A quantitative comparison between theory and experiment is hard to achieve at this stage, since the one parameter in Eq. (1), T_c , is hard to calculate, and there is not enough experimental data to allow for a fit.

It was recently suggested [13] that the low temperature behavior of ρ observed by Lilly et al. was due to the interlayer distance being close to the critical value where the two layers form an incompressible state of the (1;1;1) type. The mechanism we suggest is different from the one leading to a (1;1;1) state. In particular, the Hall transresistivity is zero in our mechanism, in contrast to the finite value predicted in [13]. A measurement which separates the longitudinal and Hall components of the transresistivity may distinguish between the two mechanisms.

In summary, we have calculated the contribution of pairing fluctuations to the transresistivity in a system of two layers at $\nu = \frac{1}{2}$ each. We find that above the critical temperature, pairing fluctuations, introduced by the interlayer attractive interaction between composite-fermions, enhance the transresistivity. In particular, the transresistivity does not vanish as the low temperature limit is approached.

We thank J.P. Eisenstein, B.I. Halperin, A.K. Kamenev, D.E. Khmelnitskii, M.P. Lilly, and A.H. MacDonald for stimulating discussions. This research was supported by the U.S.-Israel Binational Science Foundation (95/250) and the Minerva Foundation (Munich, Germany).

Note added: A related preprint (F. Zhou and Y.B. Kim, cond-mat/9807321), with which some of our results disagree, has been very recently posted on the electronic archive.

[1] For reviews, see Perspectives in Quantum Hall Effects, edited by S. Das Sarma and A. Pinczuk (Wiley, New York, 1997), pp. 37(70 and 161(224).

[2] M.P. Lilly, J.P. Eisenstein, L.N. Pfeiffer, and K.W. West, Phys. Rev. Lett. 80, 1714 (1998).

[3] Y.B. Kim and A.J. Millis, preprint cond-mat/9611125.

[4] S. Sakhi, Phys. Rev. B 56, 4098 (1997).

[5] I.Ussishkin and A. Stern, Phys. Rev. B 56, 4013 (1997); Physica E 1, 176 (1997).

[6] N.E. Bonesteel, Phys. Rev. B 48, 11484 (1993).

[7] N.E. Bonesteel, I.A. MacDonald, and C. Nayak, Phys. Rev. Lett. 77, 3009 (1996).

[8] G. Vignale and A.H. MacDonald, Phys. Rev. Lett. 76, 2786 (1996).

[9] B.I. Halperin, P.A. Lee, and N. Read, Phys. Rev. B 47, 7312 (1993).

[10] W.J. Skocpol and M. Tinkham, Rep. Prog. Phys. 38, 1049 (1975).

[11] L.G. Aalam azov and A.I. Larkin, Fizika Tverdogo Tela 10, 1104 (1968) [Sov. Phys. Solid State 10, 875 (1968)].

[12] R.S. Thompson, Phys. Rev. B 1, 327 (1970).

[13] K. Yang, preprint cond-mat/9806153.