

Solidity of Viscous Liquids

Jeppe C. Dyre

Department of Mathematics and Physics (IMFUFA), Roskilde University, P.O. Box 260, DK-4000 Roskilde, Denmark
(March 23, 2024)

Recent NMR experiments on supercooled toluene and glycerol by Hinze and Bohmer show that small rotation angles dominate with only few large molecular rotations. These results are here interpreted by assuming that viscous liquids are solid-like on short length scales. A characteristic length, the "solidity length", separates solid-like behavior from liquid-like behavior.

The viscosity of a liquid approaching the glass transition [12] is typically a factor 10^{15} larger than the viscosity of ordinary liquids like room-temperature water or ethanol. Although viscosity is just a parameter entering the Navier-Stokes equation believed to describe all liquids close to equilibrium, this enormous difference raises the question: Are viscous liquids qualitatively different from ordinary liquids or is the difference just quantitative? Below, it is argued that the former is the case. The idea is that viscous liquids behave like solids on short length scales. It is shown that this leads to a prediction consistent with the results of recent NMR experiments by Hinze and Bohmer [13,14].

In many phenomenological models of viscous liquids [3,15,24] flow proceeds via sudden reorientations of molecules, "flow events", which are rare because of the large energy barriers to be overcome [1,17,18,21]. Kauzmann referred to flow events as "jumps of molecular units of flow between different positions of equilibrium in the liquid's quasicrystalline lattice" [1]. It is this point of view that is explored here: Most molecular motion is purely vibrational and in the time between two flow events a viscous liquid is in a state of elastic equilibrium, just like a solid. However, elastic equilibrium only persists on a certain length scale beyond which the liquid does not display solid-like behavior (this point is returned to below).

Recently, Hinze and Bohmer studied reorientation of toluene and glycerol molecules by means of two-dimensional time-domain NMR spectroscopy [13,14]. The rotation angle distribution is dominated by small angles with a small, but significant fraction of larger rotation angles. These findings were interpreted as follows [14]. The large-angle rotations are those required to cross a local energy barrier. Upon barrier crossing local strains are created. These strains are relaxed through small positional and angular adjustments, not only by the molecules in the immediate vicinity but also by those further away. Briefly, large-angle rotations are "causes"

and small-angle rotations are "effects". Accepting this picture, we now proceed to show that the rotation angle distribution for small angles may be derived from the fact that viscous liquids have slow density fluctuations, assuming these liquids are solid-like on short length scales.

Consider first density fluctuations. Viscous liquids have long average relaxation times (roughly proportional to viscosity according to the Maxwell relation). These long relaxation times are basically the time between two flow events involving the same molecule. Not only enthalpy or shear stress relaxes on this time scale, but so does density. This has been known for many years from the fact that glass has smaller compressibility than corresponding equilibrium viscous liquid. More recently, measurements of the frequency-dependent bulk modulus of viscous liquids [25] revealed a loss peak around the inverse of the Maxwell relaxation time; via the fluctuation-dissipation theorem this shows directly that there are slow density fluctuations. Slow density fluctuations in viscous liquids lead to "dynamic heterogeneities", a major research topic in the 1990's [26]. Dynamic heterogeneities have been observed, e.g., in light scattering experiments [27,28], NMR experiments [29], time resolved optical spectroscopy [30], and computer simulations [31].

As mentioned, slow density fluctuations take place on a time-scale basically determined by the rate of flow events. A flow event is a rapid reorientation of molecules, probably lasting just a few picoseconds. After a flow event the molecules involved have different relative orientations. Generally, density changes somewhat at the place of a flow event (this is the cause of slow density fluctuations). As a simple model, assume isotropic flow events involving molecules confined to a sphere of radius r_0 before the flow event. We now proceed to calculate the rotation angle probability distribution for small angles. The induced movement of the surroundings is calculated by means of solid elasticity theory. If the change of radius is r , the displacement of the surroundings is given [32,34] by $u_r = r(r-r_0)^2$, where r is the distance to the flow event. The average rotation angle is proportional to the strain tensor, which in turn is formed from first order derivatives of u_r . Consequently, $\langle \theta \rangle \propto r^3$ (a detailed calculation gives $\langle \theta^2 \rangle = (6/5)r_0^4 \langle r^2 \rangle r^{-6}$). The rotation angle probability distribution is given by $P(\theta) = P(r) |dr/d\theta|$, where $P(r) \propto r^2$ from geometry. Since $|dr/d\theta| \propto r^4$ we thus find $P(\theta) \propto r^6$. Thus, [for small rotation angles] P is given by

$$P(\theta) \propto \theta^{-2} \quad (1)$$

Presently, it is not possible to determine $P(\theta)$ accurately

from experiments. It should be noted, though, that since $\sin(\theta) \approx \theta$ for small θ , Eq. (1) is consistent with the rotation angle distribution tentatively inferred from NMR experiments on glycerol, $P(\theta)/1 = \sin^2(\theta)$ [14].

The rotation angle distribution Eq. (1) is not normalizable, reflecting the fact that in the above derivation all molecules of the liquid rotate slightly following a single ω event. This, however, is not realistic; there is a "solidity length" l , beyond which ω events effectively do not induce molecular rotations. To estimate l note that elastic displacements propagate with the velocity of sound, c . Consider a sphere with radius R . Within this sphere there are $N = (R/r_0)^3$ possible locations for ω events. A molecule at the center of the sphere only "feels" the effects from any ω event in the sphere if the following condition is obeyed: The displacement deriving from such a ω event must propagate throughout the sphere and elastic equilibrium be reestablished before the next ω event occurs. If τ is the average relaxation time, the average time between two ω events within the sphere is $\tau/N = \tau/(R/r_0)^3$. This time must be longer than or equal to R/c . To estimate the solidity length l we use equality for $R = l$ and note that c is, of course, the sound velocity of the glassy state, c_{glass} . This leads to

$$l^4 = r_0^3 \tau c_{\text{glass}} \quad (2)$$

The solidity length diverges slowly as $\tau \rightarrow \infty$. To get a feeling of the order of magnitude of l , consider the case where $\tau = 1$ s. Assuming $r_0 = 5$ Å and $c_{\text{glass}} = 10^3$ m/s one finds $l \approx 6,000$ Å.

To conclude, below the glass transition, of course, viscous liquids are solid for all practical purposes. However, it has been argued here that even above the glass transition viscous liquids in certain respects behave more like solids than like less-viscous liquids. The solid-like behavior takes place on length scales below the solidity length l . Note that l diverges when τ diverges; however l is probably unrelated to the Adam-Gibbs characteristic length that also diverges with τ [17]. Rather, the solidity length l is similar to the length scale "related to solid-like behavior" recently discussed by Ahluwalia and Das within ideal mode-coupling theory [35], a length scale representative of the distance over which the liquid has enough structure to sustain propagating shear waves. However, the exact relation between the length discussed by Ahluwalia and Das and the solidity length introduced here remains to be determined.

ACKNOWLEDGMENTS

This work was supported by the Danish Natural Science Research Council.

- [1] W. Kauzmann, *Chem. Rev.* **43**, 219 (1948).
- [2] G. Harrison, *The Dynamic Properties of Supercooled Liquids* (Academic Press, New York, 1976).
- [3] S. Brawer, *Relaxation in Viscous Liquids and Glasses* (American Ceramic Society, Columbus, Ohio, 1985).
- [4] G. P. Johari, in *Relaxations in Complex Systems*, Eds. K. L. Ngai and G. B. Wright (U. S. Government Printing Office, Washington, DC, 1985), p. 17.
- [5] J. Jackle, *Rep. Progr. Phys.* **49**, 171 (1986).
- [6] C. A. Angell, *J. Phys. Chem. Solids* **49**, 863 (1988).
- [7] G. W. Scherer, *J. Non-Cryst. Solids* **123**, 75 (1990).
- [8] C. A. Angell, *J. Non-Cryst. Solids* **131-133**, 13 (1991).
- [9] A. Hunt, *J. Non-Cryst. Solids* **160**, 183 (1993).
- [10] C. A. Angell, *Science* **267**, 1924 (1995).
- [11] U. Mohanty, *Adv. Chem. Phys.* **89**, 89 (1995).
- [12] M. D. Ediger, C. A. Angell, and S. R. Nagel, *J. Phys. Chem.* **100**, 13200 (1996).
- [13] G. Hinze, *Phys. Rev. E* **57**, 2010 (1998).
- [14] R. Bohmer and G. Hinze, *J. Chem. Phys.* **109**, 241 (1998).
- [15] J. H. Gibbs and E. A. DiMarzio, *J. Chem. Phys.* **28**, 373 (1958).
- [16] M. H. Cohen and D. Turnbull, *J. Chem. Phys.* **31**, 1164 (1959).
- [17] G. Adam and J. H. Gibbs, *J. Chem. Phys.* **43**, 139 (1965).
- [18] M. Goldstein, *J. Chem. Phys.* **51**, 3728 (1969).
- [19] G. Williams, in: *Dielectric and Related Molecular Processes*, Specialist Periodical Report, Vol. 2, Ed. M. Davies (Chem. Soc., London, 1975), p. 151.
- [20] E. Donth, *Glasübergang* (Akademie-Verlag, Berlin, 1981).
- [21] F. H. Stillinger, *J. Chem. Phys.* **89**, 6461 (1988).
- [22] R. V. Chamberlin, *Phys. Rev. B* **48**, 15638 (1993).
- [23] J. C. Dyre, *Phys. Rev. B* **51**, 12276 (1995).
- [24] D. Kivelson, G. Tarjus, X. Zhao, and S. A. Kivelson, *Phys. Rev. E* **53**, 751 (1996).
- [25] T. Christensen and N. B. Olsen, *Phys. Rev. B* **49**, 15396 (1994).
- [26] D. N. Perera and P. Harrowell, *J. Chem. Phys.* **104**, 2369 (1996).
- [27] C. T. Moynihan and J. Schroeder, *J. Non-Cryst. Solids* **160**, 52 (1993).
- [28] C. H. Wang and E. W. Fischer, *J. Chem. Phys.* **105**, 7316 (1996).
- [29] R. Bohmer, G. Hinze, G. Diezemann, B. Geil, and H. Sillescu, *Europhys. Lett.* **36**, 55 (1996).
- [30] M. T. Cicerone, F. R. Blackburn, and M. D. Ediger, *J. Chem. Phys.* **102**, 471 (1995).
- [31] M. M. Hurley and P. Harrowell, *Phys. Rev. E* **52**, 1694 (1995).
- [32] L. D. Landau and E. M. Lifshitz, *Theory of Elasticity* (Pergamon Press, Oxford, 1970).
- [33] G. Wylie, *Phys. Rep.* **61**, 327 (1980).
- [34] J. C. Dyre, *J. Non-Cryst. Solids*, to be published (1998).
- [35] R. Ahluwalia and S. P. Das, *Phys. Rev. E* **57**, 5771 (1998).