C L.K ane D epartm ent of P hysics, U niversity of P ennsylvania P hiladelphia, P ennsylvania 19104 M arch 1, 2022)

We study the problem of resonant tunneling between a Ferm i liquid and the edge of a = 1=3 fractional quantum H all state. In the lim it of weak coupling, the system is adequately described within the sequential tunneling approximation. At low temperatures, however, the system crosses over to a strong coupling phase, which we analyze using renorm alization group techniques. We nd that at low temperatures a `perfect" can be achieved by tuning two parameters. This resonance has a peak conductance $G = e^2=2h$ and is a realization of the weak backscattering lim it of the g = 1=2 Luttinger liquid, as well as the non Ferm i liquid xed point of the two channel K ondo problem. We discuss several regimes which may be experimentally accessible as well as the implications for resonances recently observed in cleaved edge overgrow th structures.

PACS: 73.20Dx, 73.40Hm, 73.40Gk, 72.10Fx

I. IN TRODUCTION

The rich structure of the fractional quantum Hall effect o ers a controlled laboratory for studying the physics of strong correlation. As rst argued by W ert, the low energy excitations at the edge of a quantum Hall state form a novel quantum uid known as a chiral Luttinger liquid. This theory has led to a number of predictions regarding the e ects of tunneling and resonant tunneling between quantum Hall states¹³. Experiments by Milliken Um bach and Webb⁴ probed tunneling and resonant tunneling at a quantum point contact. They observed a striking di erence in behavior at lling factors = 1 = 1=3, which was consistent with these predicand tions. M ore recent experim ents on a cleaved edge overgrowth structure by Chang, P fei er and W est observed the predicted power law scaling of the tunneling current with voltage and temperature at = 1=3 over several decades of current. W hile questions rem ain about other

lling factors, the agreem ent between experiment and theory has been quite encouraging for the Laughlin states at = 1 and 1=3.

In this paper we analyze the problem of resonant tunneling between two di erent quantum Hall states, = 1 and 1/3 - or equivalently between a Ferm i liquid and = 1=3. Our motivation is twofold. First, in their recent cleaved edge overgrow th experiments, G rayson et al. have observed interesting resonances in tunneling between a metal and = 1=3⁷. Second, this problem is interesting from a theoretical point of view because, as we explain below, a \perfect" resonance in this problem constitutes a realization of the perfectly transmitting xed point of a g = 1=2 Luttinger liquid, which is equivalent to the non Ferm i liquid xed point of the two channel K ondo problem.

O ur analysis is closely related to that of resonant tunneling in a = 1=3 quantum point contact, which is closely related to the problem of tunneling through a barrier in a g = 1=3 Luttinger liquid⁸. The lim it of strong pinch o corresponds to a barrier with very sm all transm ission, whereas the lim it weak pinch o corresponds to a small barrier which backscatters only weakly. As argued in Ref. 3 a perfect resonance is controlled by the perfectly transm itting xed point. Such a resonance has a peak conductance of e^2 =3h and a universal lineshape which is determ ined by the crossover between the perfectly transm itting and perfectly re ecting xed points.

The problem of tunneling between = 1 and = 1=3 can be mapped onto that of tunneling through a barrier in a g = 1=2 Luttinger liquid. This has led to the interesting suggestion that the interface between = 1 and = 1=3 m ay allow for the realization of the perfectly transm itting limit of the g = 1=2 Luttinger liquid^{9;10}. Such a state would have interesting properties. Since the \perfect" two term inal conductance $e^2=2h$ is larger than the conductance of the quantum H all state, $e^2=3h$ it would act as a DC step-up transform er^{10} . In addition, the weak backscattering in this state would in principle be detected in a shot noise experiment¹¹.

However, realizing this state presents a subtle problem. Unlike the = 1=3 point contact, where one can physically separate the counterm oving edge states, there is no direct way of \engineering" the g = 1=2 perfectly transm itting xed point. In principle it could be achieved using the adiabatic contact, suggested by Chklovskii and Halperin¹⁰, in which the quantum Hall uids are connected sm oothly by a quantum wire, in which the Luttinger parameter g can vary continuously. Recently, Sandler et al.¹¹ have argued that the perfectly transm itting

xed point may be approached by scaling to su ciently high energies. However, since there are many irrelevant operators at this xed point, it is unlikely that the system would ow near the xed point without tuning several parameters.

In this paper, we show that the perfectly transm itting lim it can be reached by resonantly tunneling through an impurity state. A \perfect" resonance can be reached at zero tem perature by tuning two parameters. Such a resonance has a peak two term inal conductance of $e^2=2h$ and has a universal temperature dependent lineshape which has been computed in the context of a g = 1=2 luttinger liquid⁸. The remainder of the paper is organized as follows. A fler describing the model, we discuss the resonances in the weak tunneling limit, when the tunneling is sequential. Then using a renormalization group analysis we study the crossover to the perfect resonance, F inally we discuss the experimental implications of our results and the connection with the recent observations by G rayson et al.⁷

II. THE LUTTINGER LIQUID MODEL

We begin with the chiral Luttinger liquid model describing the coupling between the = 1=3 and = 1 edge states and the impurity state. The Ham iltonian may be written, $H = H_0^1 + H_0^3 + H_{imp} + H_T^1 + H_T^3$. The edge states are described by

$$H_0^m = dx u_m (\theta_{x m})^2$$
(1)

where the elds m satisfy the Kac Moody algebra

$$[m (x^{0}); 0_{x m} (x)] = (2 i=m) (x x);$$
 (2)

and u_m determ ines the velocity of propagation. We consider a single impurity state with energy " $_0$,

$$H_{imp} = "_0 d^Y d; \qquad (3)$$

where d^y creates an electron in that state. The tunneling of electrons between the impurity state and the edges is described by

$$H_{T}^{m} = t_{m} c^{1} d^{y} e^{im} + hc:$$
(4)

where t_1 and t_3 are dimensionless tunneling amplitudes and $\ _{\rm c}$ is a short time cuto $\ .$

A way from a resonance, when " $_{0} \in 0$, electrons may tunnel virtually through the impurity. This leads to a nonresonant conductance which vanishes as T² due to the power law tunneling density of states of the = 1=3 edge. However when " $_{0}$ is tuned through zero, the energy denom inator for the virtual state becomes small, and there can be resonant transmission through the im purity state. We will analyze the resonances rst in the perturbative sequential tunneling limit, and then perform a renorm alization group analysis, which can describe the perfect resonance.

III. SEQUENTIAL TUNNELING LIM IT

In the limit, t_m 1, tunneling is so infrequent that successive hops onto and o of the inpurity will be uncorrelated. This \sequential tunneling" regime has been studied in detail in the context of Luttinger liquid theory by Cham on and W en¹² and by Furusaki and N agaosa¹³. The result of this analysis is that the tunneling conductance is given by

$$G = \frac{e^2}{2 h} \frac{3 ("_0; T) 1 ("_0; T)}{3 ("_0; T) + 1 ("_0; T)} \frac{1}{T} \operatorname{sech}^2 \frac{"_0}{2T}:$$
(5)

m (";T) is the inverse lifetime for tunneling into the = 1=m lead from the impurity, which rejects the power law dependence of the tunneling density of states,

$$(") / "^2 m^0 1$$
: (6)

Here m_{m}^{0} is the bare scaling dimension of the electron tunneling operator, given by

In the following section we will see that m can be renormalized. We therefore compute the lifetimes for general m,

$$m (";T) = 2 t_{m c}^{2} (2 T_{c})^{2 m 1} F_{m} (\frac{"}{2 T}); \quad (8)$$

with

$$F_{m}(x) = 2\cosh x \frac{j(m + ix)\hat{j}}{(2m)}$$
: (9)

In the perturbative regime, in which m are given by (7), this form ula reduces to

where c_1 and c_3 are dimensionless constants of order unity. It is useful to distinguish the following limiting cases:

(1) W hen $_1$ $_3$, the conductance is limited by the tunneling into the = 1 edge. In this regime,

$$G = \frac{e^2}{2 h} \frac{1}{T} \operatorname{sech}^2 \frac{"_0}{2T}; \qquad (11)$$

and the resonances are Ferm i liquid like, with a peak conductance scaling as 1=T and a lineshape given by the derivative of the Ferm i function.

(2) W hen $_{3}$ 1 the conductance is limited by tunneling into the = 1=3 edge. In this case,

$$G / \frac{\mathbf{u}_0^2 + {}^2 T^2}{T} \operatorname{sech}^2 \frac{\mathbf{u}_0}{2T}$$
: (12)

Thus, the peak conductance decreases as T as the tem – perature is lowered in contrast to the T 2 behavior o resonance. The lineshape is slightly modil ed from the derivative of the Ferm i function.

IV.RENORMALIZATION GROUP ANALYSIS

It is important to emphasize that the sequential tunneling approximation is valid only when tunneling is uncorrelated. This will be true provided $_1$; $_3$ T. To go beyond this approximation it is useful to develop a perturbative renormalization group analysis. We will now



FIG.1. Renorm alization group ow diagram as a function of t_3 and $_3$ for $t_1 = 0$. For $t_3 < t_3 t_3$ ows to zero, and the impurity state decouples from the = 1=3 state. For $t_3 > t_3$ the impurity state merges with the = 1=3 state.

focus on the resonance peaks, when $"_0 = 0$. To leading order, the scaling equations for t_1 and t_3 are given by

$$\frac{dt_m}{d'} = (1 \qquad _m)t_m : \qquad (13)$$

To low est order in $t_m = m$ is simply given by (7). Thus, to leading order, t_3 is irrelevant, while t_1 is relevant. The scaling behavior of m in (8) follows directly,

$$m (f_{m}^{R})^{2}T / T^{2 m 1}:$$
 (14)

where t_m^R is the renormalized value of t_m at an energy scale of order T. The perturbative analysis is then valid provided t_m^R 1, or equivalently m T.

For nite $\frac{1}{2}$, m is renormalized. This e ect has been worked out in detail by the author in Appendix B of Ref.8 in the context of resonant tunneling in a Luttinger liquid. Here we generalize those results to the problem of tunneling between = 1 and = 1=3. The resulting ow equations for m are

$$\frac{d_{1}}{d'} = 4_{1}t_{1}^{2} - 2(t_{1} + t_{3} - 2)t_{3}^{2}$$
(15)

$$\frac{d_{3}}{d'} = 4_{3}t_{3}^{2} - 2(t_{1} + t_{3} - 2)t_{1}^{2}$$
(16)

It is instructive to analyze the renormalization group ows in limiting cases.

(1) For $t_3 = 0$, the resonant state is coupled only to the = 1 edge. In this case, t_1 is relevant and the system ow s to a strongly coupled phase with t_2 ! 1, 1 = 0 and 3 = 2. Since 3 = 2, t_3 is an irrelevant perturbation in this phase. The tunneling conductance can then be found perturbatively to vary as $G / t_3^2 T^2$. This is the same as non resonant tunneling between = 1 and = 1=3. In this phase the impurity has e ectively m erged with the = 1 quantum Hall uid.

(2) For $t_1 = 0$, the resonant state is coupled only to the = 1=3 edge. To leading order t_3 is irrelevant. How ever,

 $_3$ is renorm alized dow nw ard from its initial value of 3=2. Thus, as indicated in Fig. 1, when t_3 reaches a critical



FIG.2. Zero temperature phase diagram for an inpurity resonantly coupled to = 1 and = 1=3 states as a function of the couplings t_1 and t_3 . For large t_1 the inpurity merges with the = 1 state, while for large t_3 the inpurity merges with the = 1=3 state. At the transition between the two phases is the perfect resonance, which is described by the perfectly transmitting xed point of a Luttinger liquid at q = 1=2.

value t_3 0.22 the system ow s to a K osterlitz-T houless like xed point at $\xi = 0$, $_3 = 1$. For $t_3 > t_3$ the system ow s to a strongly coupled phase with $\xi ! 1$, $_3 = 0$ and $_1 = 2$. Here, the in purity has merged with the = 1=3 quantum Hall uid. Treating ξ perturbatively again leads to a tunneling conductance varying as T^2 . On the other hand, for $t_3 < t_3$, t_3 ow s to zero with a modi ed exponent: $1 < _3 < 3=2$. In this case $_3$ (T) / $T^{2_3 2}$. Since t_1 remains relevant, how ever, this state is unstable for nite ξ and at low tem peratures the in purity merges with the = 1 quantum Hall uid.

W hen both t_1 and t_3 are nite it is clear that there are two phases: the in purity merges with either the = 1 or the = 1=3 quantum Hall uids. For xed₁tas t_3 is increased the system undergoes a transition between these two phases, as indicated in the phase diagram Fig. 2. Integration of the ow equations shows that precisely at this transition, t_1 and t_3 are equal and grow to in nity, with $_1 = _3 = 1=2$. We will analyze this case in the following section.

V.THE PERFECT RESONANCE

The xed point describing the phase transition is not accessible within this perturbative analysis. However, when $_1 = _3$ and $t_1 = t_3$, this problem is equivalent to resonant tunneling between two g = 1=2 Luttinger liquids in the lim it in which the two tunneling barriers are symmetric. As argued in Ref. 8, this is precisely the condition for a \perfect resonance", where the system ow s to the perfectly transmitting xed point.

Equivalently, this model is can be mapped to the anisotropic two channel K ondo problem 14,15 . In this

mapping, the occupation of the impurity state corresponds to the state of the K ondo spin, and the two leads correspond to the two channels. The tunneling matrix elements t_m play the role of the transverse couplings J_2^i for the two channels, i. The scaling dimensions $_m$ are related to J_z^i . W hen $_1 = _3 = 1=2$, the K ondo m odel is in the Toulouse limit.

Under this mapping, the two phases in which the in purity state merges with either = 1 or = 1=3 can easily be identied. When there is channel anisotropy in the two channel K ondo problem the K ondo spin tends to form a singlet with one or the other channels. As the couplings are varied, there is a transition in which the K ondo spin switches its allegiance. Precisely at this transition, which occurs when the channels are symmetric, the system ows to the non Fermiliquid xed point in which the spin is shared between the two channels.

Having established that the resonance xed point is equivalent to those of both the perfectly transmitting g = 1=2 Luttinger liquid and the two channel K ondo problem, we may now describe its properties. Precisely on resonance, the conductance is $G = e^2=2h$. Slightly away from the resonance, the lineshape is controlled by the single relevant operator at the resonant xed point. This operator describes equivalently $2k_F$ backscattering in the Luttinger liquid model, or channel anisotropy in the K ondo m odel. This leads to a resonance with a tem – perature dependent width proportional to T¹⁼² and a universal lineshape⁸.

$$G(T;) = G(=T^{1-2})$$
 (17)

where is the tuning parameter for the resonance and

$$G'(X) = \frac{e^2}{2h} \int_{-1}^{2} dy \frac{e^y}{(e^y + 1)^2} \frac{y^2}{y^2 + X^4} :$$
(18)

VI.EXPERIMENTAL IMPLICATIONS

The resonances depend critically on temperature as well as the coupling between the impurity state and the two leads. Here we outline the various possible regimes, depending on these parameters. It is most useful to characterize the coupling to the leads by the inverse lifetimes $_1$ and $_3$, though one must keep in m ind that they can depend on temperature. In the limit of weak coupling, $_1$ is temperature independent, while $_3$ / T². For stronger coupling the temperature dependence is altered.

If T $_1$; $_3$, then tunneling is sequential, and the resonances should be accurately described by equation (5). There are three distinct cases depending on $_1$ and $_3$:

(1) Ferm i Liquid dom inated sequential tunneling If 1 3, then the transm ission is limited by the contact to the Ferm i liquid. The resonances are then insensitive to the Luttinger liquid correlations, and have the Ferm i liquid form (11).

(2) = 1=3 dom inated sequential tunneling Since $_3$ decreases as the temperature is lowered, while $_1$ remains temperature independent, it is possible that the

two could cross at a temperature above $_1$. In this case the system would remain in the sequential tunneling regime, but the tunneling would be limited by $_3$. Provideded the coupling to = 1=3 is su ciently weak that $_3$ has not renormalized signi cantly then the resonances peaks will vary as T and the lineshape will be given by (12). This will be the case provided t_3 1, or equivalently $_3$ (T) $_c$ T², where $_c$ ¹ is the high energy cuto .

(3) Renormalized sequential tunneling For $_{c}T^{2}$

3 T, 1 T, one is no longer in the weak coupling limit. Nonetheless, as shown in Fig. 1, for $t_3 < t_3$ and $t_1 = 0$ the system ows to a xed point with $t_3 = 0$. In this limit, 3 T, so tunneling is sequential, how ever, the exponents 1 and 3 are renormalized: $1 < _3 < 3=2$, and $1=2 < _1 < 2$ 3. In this regime, the resonances can be described by combining (5), (8) and (9). The peak heights will vary as T. If $_1 < _3$ then 1 < < 0. If $_1 > _3$ then 0 < < 1.

(4) Strong Coupling Regime W hen $T < _1$ or $_3$, the system is in a regime where the impurity is strongly coupled to the leads. This regime will be characterize by the zero tem perature phase diagram shown in Fig. 2. Generically, the impurity will be strongly coupled to either the Fermi liquid or the = 1=3 lead. In that case, the conductance will have the \circ resonance" T^2 tem perature dependence. However, if the coupling t_3 is su ciently strong, and the couplings are such that the system is precisely on the phase boundary, then the system will be at a perfect resonance. A chieving the perfect resonance requires two param eters to be tuned: the resonance energy $"_0$ and the coupling to = 1=3, t₃. At the resonance the peak conductance should increase as the tem perature is low ered, approaching a peak value of $e^2 = 2h$, while the width becom es narrower, ultim ately having the universal lineshape given in (17) and (18).

Recently, in their cleaved edge overgrowth experiments, G rayson et al. have observed resonances in tunneling between a metal and the edge of a = 1=3 quantum Hall state as a function of magnetic eld. These resonances have a peak conductance which increases as approximately 1=T as the temperature is lowered and a line shape that is well t by the derivative of the Ferm i function. Since the peak height of the resonances is quite small (of order $005e^2$ =h at 25 mK), it is most likely that in this temperature range the system is in Ferm i liquid dom inated sequential tunneling regime. From the peak height at 25 mK, one can estimate 1 0:1mK, which also gives a lower bound on 3, so that 0:1mK < 3 < 25mK.

In these samples signatures of Luttinger liquid behavior in the resonances may not occur until much lower temperatures, and reaching the strong coupling regime may prove di cult. Nonetheless it would be interesting to probe such resonances either by going to lower tem – perature or by somehow increasing the coupling to the impurity state. A second tuning parameter, such as a gate voltage would allow additional exibility in controlling 1 and 3, and would allow for the possibility of tuning to the perfect resonance. The rst signature of strong coupling behavior would be a deviation from the 1=T dependence of the peak heights of the resonances.

ACKNOW LEDGMENTS

It is a pleasure to thank C.deC.Chamon, E.Fradkin and S.Sondhi for helpful discussions and A.Chang, M. Grayson and D.T sui for sharing their experimental results prior to publication. This work has been supported by the National Science Foundation under grant DMR 95-05425.

- ¹ X G.W en, Phys.Rev.B 43, 11025 (1991); Phys.Rev.Lett. 64, 2206 (1990); Phys.Rev.B 44 5708 (1991).
- ² C L. K ane and M P A. Fisher, Phys. Rev. Lett. 68, 1220 (1992).
- ³ K.Moon et al, Phys. Rev. Lett. 71, 4381 (1993).
- ⁴ F P. M illiken, C P. Umbach and R A. Webb, Solid State Commun. 97, 309 (1996).
- ⁵ A M .Chang,LN.Pfei erandKW .West,Phys.Rev.Lett. 77,2538 (1996).
- ⁶ M.Grayson, D.C.Tsui, L.N.Pfeier, K.W.West and A.M. Chang, Phys. Rev. Lett. 80, 1062 (1998).
- ⁷ M. Grayson, D.C. Tsui, L.N. Pfei er, K.W. West, and A.M. Chang, \Tunneling Resonance at the Edge of the 1/3 FQHE," to be published.
- ⁸ C L. Kane and M P A. Fisher, Phys. Rev. B 46, 15233 (1992).
- ⁹ C.de C.Cham on and E.Fradkin, Phys.Rev.B 56, 2012 (1997).
- ¹⁰ D B.Chklovskii and B.I.Halperin, Phys.Rev.B 57, 3871 (1998).
- ¹¹ N P.Sandler, C.de C.Cham on and E.Fradkin, \W hat do noise m easurem ents reveal about fractional charge in FQH liquids?", cond_m at/9806335.
- ¹² C.de C.Cham on and X G.W en, Phys. Rev. Lett. 70, 2605 (1993).
- ¹³ A. Furusaki and N. Nagoaosa, Phys. Rev. B 47, 3827 (1993).
- ¹⁴ A. Furusaki and K A. M atveev, Phys. Rev. Lett. 75, 709 (1995).
- ¹⁵ H.Yiand C.L.Kane, Phys. Rev. B 57, 5579 (1998).