₿T_EX

The equation of state and speci c heat of the electron gas on a one dimensional lattice

V ladan Celebonovic

Institute of Physics, Pregrevica 118,11080 Zem un-Beograd, Yugoslavia vladan@phybg.ac.yu

vælebonovid sezam pro.yu

to appear in PhysLow-D in Structures.,1/2,XXX (1999)

Abstract iJ sing previous results and general therm odynam ical form alism, an expression is obtained for the speci c heat perparticle under constant volum e of a degenerate non-relativistic electron gas on a 1D lattice. The result is a non-linear function of the tem perature, and it could have applications in studies of quasi one-dimensional organic metals.

Introduction

The purpose of this paper is to determ ine the equation of state and speci c heat of the electron gas on a one dimensional lattice. In experiments, the electronic speci c heat is determined indirectly, only after somehow estimating and substracting the phonon heat capacity from the measured heat capacity of the specimen Ω n the purely theoretical side, the electronic speci c heat is directly related to the exact single particle G reen's functions [la]. The calculations, to be reported in the following section, are motivated by recent theoretical and experimental results on transport processes in quasi one-dimensional (Q 1D) organic metals [1], [2]. A preliminary version of the calculation has already been published [3].

Q 1D organic m etals, later nam ed the Bechgaard salts, were discovered in 1980 [4], [5]. It was experim entally shown soon afterwards that their transport properties are widely di erent from those of ordinary m etals (for example [6] and references given there). For a general review of the eld of organic conductors see [7].

The calculations

It will be assumed in the calculations that the number of particles N in the system is a variable. The starting point is a well known therm odynam ical relation [8]

$$dG = S dT + V dP + dN$$
(1)

where G = N and all the symbols have their standard meanings. Inserting the de nition of G into eq.(1), it follows that

$$SdT + VdP + dN = dN + Nd$$
 (2)

D i erentiating eq.(2) with respect to T gives

$$V (1 + \frac{1}{V} \frac{\partial V}{\partial T}) \frac{\partial P}{\partial T} = S + \frac{\partial S}{\partial T} + \frac{\partial N}{\partial T} \frac{\partial}{\partial T} + N \frac{\partial}{\partial T}$$
(3)

In the last equation we have used $dP = \frac{\partial P}{\partial T} dT$ and similar relations for other variables in eq.(2).

Equation (3) is expressed in terms of the parameters of the system in bulk C hanging the variables to the number density n (N = nV) and the entropy per particle s (S = nsV) after some algebra one arrives at

$$V \left(1 + \frac{1}{V} \frac{\partial V}{\partial T}\right) \frac{\partial P}{\partial T} = ns\left(V + \frac{\partial V}{\partial T}\right) + V \frac{\partial n}{\partial T}\left(s + \frac{\partial}{\partial T}\right) + nV \frac{\partial}{\partial T}\left(s + \right) + n\frac{\partial}{\partial T} \frac{\partial V}{\partial T}$$
(4)

W e have thus obtained the EOS of any material in di erential form. In applications to Q 1D organic metals, it can be considerably simplied. Experiments on these materials are usually performed under constant volume [7]. This implies that all terms in eq.(4) containing derivatives of the volume can be disregarded A coordingly, the nalform of the EOS of Q 1D organic metals is

$$\frac{\partial P}{\partial T} = (n + \frac{\partial n}{\partial T})(s + \frac{\partial}{\partial T}) + n\frac{\partial s}{\partial T}$$
(5)

The speci c heat per particle is given by [8]

$$c_{V} = \frac{T}{n} \left(\frac{\theta^{2} P}{\theta T^{2}} \right)_{V} \qquad T \left(\frac{\theta^{2}}{\theta T^{2}} \right)_{V}$$
(6)

D i erentiating eq.(5), and inserting the result into eq.(6), gives the follow - ing nalexpression for the speci c heat per particle

$$c_{V} = \frac{T}{n} \left(s + \frac{\theta}{\theta T}\right) \left(\frac{\theta n}{\theta T} + \frac{\theta^{2} n}{\theta T^{2}}\right) + \frac{T}{n} \frac{\theta n}{\theta T} \left(2\frac{\theta s}{\theta T} + \frac{\theta^{2}}{\theta T^{2}}\right) + T \left(\frac{\theta s}{\theta T} + \frac{\theta^{2} s}{\theta T^{2}}\right)$$
(7)

D iscussion

Equation (7) is considerably more complex than the preliminary result already published in [3]. This di erence is due several simplifying assumptions made in [3], which were dropped in the present paper. Basically, it was assumed there that the entropy per particle is temperature independent, that the total number of particles in the system is a constant, and the term $\frac{\Theta n}{\Theta T} \frac{\Theta}{\Theta T}$ was neglected.

The aim of the calculation reported here was to obtain an expression for the di erential form of the EOS and the speci c heat ,but without all these simpli cations. The result for the EOS is given by eq.(4). It was derived using general therm odynam ical argum ents, and accordingly can be applied to any material. In the special case of Q1D organic metals, one can introduce the constant volume condition into eq.(4) and thus obtain eqs.(5) and (7).

In application to the degenerate electron gas on a 1D lattice, one has to introduce into eq.(7) appropriate expressions for the chem ical potential, num ber density and entropy perparticle. It has recently been shown that the chem ical potential of the electron gas on a 1D lattice is given by [9]

$$= \frac{(t)^{6} (af \ 1) jj}{1:1029 + :1694 (t)^{2} + :0654 (t)^{4}}$$
(8)

where a denotes the lattice constant, the hopping, f the band lling and the inverse temperature. The in uence of the lattice is contained in this expression through the presence of the lattice constant a:

The entropy per electron and the num ber density should also be expressed so as to take into account the existence of the lattice W ork on this problem is currently in progress, but as a rst approximation one could use existing results for the degenerate electron gas [10], [13]. The entropy per electron is given by

$$s = \frac{Q}{n_e} \frac{Q}{QT} F_{3=2} ()$$
(9)

where $F_{3=2}$ () is a special case of a Ferm i-D irac integral [10], n_e denotes the electronic number density, and Q is a combination of known constants (such as the electron m ass and P lanck's constant). The number density of a degenerate electron gas at low teem perature can be expressed in analytical form as [10]

$$n_e = AT^{15=2} [1 + BT^{3=2} +]$$
 (10)

A; B again denote combinations of known constants. Inserting expressions (8) - (10) in eq.(7) one would get the nal result for the speci c heat of the electron gas on a 1D lattice.

How can eq.(7) be applied in the analysis of experimental data on Q 1D organic metals? It can be used in analyzing data on the speci c heat and therm al conductivity of these materials, such as those discussed in [1].

The specic cheat of a solid can be decomposed into phonon and electronic parts. Their relative contributions to the "summary" specic heat of the specimen are functions of the parameters of the system. The most im – portant of them are the temperature and the density Expression (8) for the chemical potential was derived assuming that the Fermi liquid picture can be used for describing electrons in Q1D organic metals. This assumption is at present a subject of active discussion (see, for example, [11], [12]). Holding to this assumption, it then follows that eq.(7) can be applied to determining the phonon part of the specic cheat from the experimental data. On the other hand, starting from the measured values of the specic cheat, and the somehow theoretically estimated phonon contribution, one could obtain the electronic part. Comparing this function with the predictions of eq.(7), one could test the applicability of the Fermi liquid picture to electrons in Q1D organic metals.

The reasoning behind applications of eq.(7) to studies of the therm al conductivity of Q 1D organic m etals is essentially the same. It can be shown from elementary considerations [14] that the therm al conductivity is proportional to the speci c heat. Taking eq.(7) in account, this im plies that the therm al conductivity of Q 1D organic m etals is a nonlinear function of the tem perature, in line with recent experiments [1] A gain, this could be used to test theoretical models of the organic m etals. W ork aim ing at details of this com parison is currently in preparation.

A dknow ledgem ent

This paper is dedicated to the memory of the late Heinz J.Schulz from Laboratoire de Physique des Solides in Orsay (France) who introduced me to the eld.

References

[1] S Belin and K Behnia, PhysRevLett., 79, 2125 (1997). [1a] C M .Varm a, preprint cond-m at/9607105 v2 (1996). [2] V Celebonovic, PhysLow-D in Struct., 3/4, 65 (1997) and preprint cond-m at/9702147. BIV Celebonovic, Solid State Phenomena, 61-62, 135 (1998c). [4] K Bechgaard, C S Jacobsen, K M ortensen et al, Solid State Comm .,33,1119 (1980). [5] D Jerom e, A M azaud, M R ibault and K Bechgaard, C R A cad Sc Paris, B 290, 27 (1980); D Jerom e, A M azaud, M R ibault and K Bechgaard, J Physique (Paris), Lettres, 41, L95 (1980). [6] D Jerom e and F C reuzet, in: N ovel Superconductivity (ed by SAW olf and VZK resin), p.103, Plenum Press, New York (1987). [7] T. Ishiguro and K. Yam a ji, Organic Conductors, Springer Verlag, Berlin (1990). [8] H E Stanley, Introduction to phase transitions and critical phenom ena, Clarendon Press, Oxford (1971). [9] V Celebonovic, PhysLow-D in Struct., 11/12,25 (1996) and preprint cond-m at/9512016. [10] V Celebonovic, PublA stron ObsBelgrade, 60, 16 (1998a) and preprint astro-ph/9802279. [11] J.Voit, Rep Progr Phys., 58,975 (1995). [12] C Bourbonnais, preprint cond-m at/9611064 v2 (1996); A Schwartz M D ressel G G runner et al, Phys Rev., B 58, 1261 (1998); JM oser,M G abay,P A uban-Senzier,D Jerom e,K Bechgaard and JM Fabre, EurPhys. J., B1, 39 (1998) . [13] L D Landau and E M Lifchitz, Statisticheskaya zika, Vol.1, Nauka, Moscow (1976). [14] N W A shcroft and N D M erm in Solid State Physics, Holt, Rinehart and Winston, London (1976).