

Non-perturbative results for level correlations from the replica nonlinear model

I. V. Yurkevich and I. V. Lerner

School of Physics and Astronomy, The University of Birmingham, Edgbaston, Birmingham B15 2TT, UK
(August 26, 2021)

We show that for all the three standard symmetry classes (unitary, orthogonal and symplectic), the conventional replica nonlinear model gives the correct non-perturbative result for the two-level correlation functions $R_2(l)$ of electrons in disordered metals in the limit of large l . In this limit, non-perturbative oscillatory contributions arise from a degenerate saddle-point manifold within this model which corresponds to the replica-symmetry breaking. Moreover, we demonstrate that in the unitary case the very same results can be extracted from the wellknown exact integral representation for $R_2(l)$.

I. INTRODUCTION

Starting from the seminal papers of Wegner¹ and Efetov,² a field-theoretical description based on the nonlinear model (NL M) has become one of the main analytical approaches to various problems in disordered electronic systems. The ensemble averaging over all configurations of disorder is performed either using bosonic^{1,3} or fermionic⁴ n -replicated fields and taking the $n \rightarrow 0$ limit in the results, or using supersymmetric (\mathbb{Z}_2 -graded) fields;⁵

The very first application of this approach was a derivation^{3,4,2} of the renormalization-group (RG) equations of the scaling theory⁶ of Anderson localization. For such a perturbative derivation, generalized later for mesoscopic systems⁷, both the replica and the supersymmetric methods are equally well justified: they just ensure the cancellation of unphysical vacuum loops in a diagrammatic expansion. However, it soon became conventional wisdom that there existed two sets of problems for which only one of these methods was applicable.

On the one hand, the fermionic replica NL M has been generalized by Finkelstein⁸ to include interactions between electrons. The interest in this approach has been greatly enhanced by the recent discovery⁹ of an apparent metal-insulator transition in 2D disordered systems in zero magnetic field. Although it has been recently demonstrated¹⁰ that the Keldysh technique provides a viable alternative to the replica approach, the latter still remains one of the best available tools for consideration of interacting electrons in disordered systems. In the very least, it is clear that there is no simple way of applying the supersymmetry method to a many-particle fermionic system.

On the other hand, the viability of the replica approach was undermined by the existence of a set of problems which could apparently be solved only with the help of the supersymmetry method. The first, and possibly most famous, of such problems was solved by Efetov¹¹ who used the supersymmetric NL M to derive the two-level correlation function (TLCF) in the universal ergodic regime for electrons in disordered metallic grains. The results proved to be identical to those for eigenvalue correlations in random matrix theory,¹² as had much earlier been assumed by Gorkov and Eliashberg.¹³ This first microscopic derivation of essentially non-perturbative results has opened the way to numerous new results (mainly obtained during the last decade) for which using the supersymmetry method seemed to be absolutely essential (many of these results have been reviewed in a recent book by Efetov¹⁴).

In many ways, though, this first non-trivial "supersymmetric" result¹¹ seemed to be the best illustration of why the replica method could only be used within a perturbative approach. For the easiest case of the unitary symmetry the irreducible TLCF is given by

$$R_2(l) = \frac{\sin^2 l}{l^2}; \quad (1.1)$$

where l is the distance between two levels in units of Δ and Δ is the mean level spacing. This result is valid in the ergodic regime, i.e. for $l \gg g$ where $g \ll 1$ is the dimensionless conductance. For $l \ll 1$, the TLCF averaged over fast oscillations, could be readily obtained from the standard diagrammatic techniques¹⁵ with $l=1$ being the perturbation parameter (or from the perturbation theory in the framework of either the supersymmetric or replica NL M). However, the non-perturbative factor $\sin^2 l$ cannot be restored from the perturbation series. Since the replica trick is well justified only within the perturbative approach, it might seem rather hopeless to obtain the result (1.1) within the replica approach. And indeed, quite involved calculations by Verbaarschot and Zimbauer¹⁶ have shown that a direct application of the replica trick (using either the bosonic or fermionic NL M) has apparently not reproduced the TLCF given by Eq. (1.1).

Therefore, a very recent result of Kamenev and Mezard¹⁷ makes a real breakthrough in this area. Using the replica-symmetry breaking within an effective fermionic field theory corresponding to the large N limit of the Gaussian unitary ensemble of $N \times N$ random matrices, they have reproduced the non-perturbative result (1.1), albeit only in the region

! 1. This raises hope that one might eventually apply the replica approach for obtaining non-perturbative results for interacting electrons in disordered systems. However, the method used in Ref. 17 does not appear to be extendible to the standard NL M describing electrons in a random potential. Although the large- N field theory written in terms of the n replica matrix fields Q is, in principle, totally analogous to the standard NL M, this analogy has been considerably smeared by the very essence of the method applied. Namely, the standard $\exp[N \text{tr} \ln(-iQ)]$ term has been represented after a shift of variables as $(\det Q)^N$ and the resulting Itzykson-Zuber integral (similar to that introduced in the context of the supersymmetric method by Guhr¹⁸) was calculated taking into account replica-symmetry breaking.¹⁷

Although the method used in Ref. 17 could hardly be generalized directly, the message was very clear: the replica approach taken together with the symmetry-breaking, can reproduce the results of the supersymmetric approach.

The purpose of this paper is to demonstrate that non-perturbative results for all the classical symmetry ensembles (unitary, orthogonal and symplectic) can be derived in the framework of the standard fermionic NL M with replica-symmetry breaking. We believe that the present derivation is both more general and considerably easier than the original one¹⁷ and can be straightforwardly extended to include interactions. Independently, Alex Kamenev and Marc Mezard have presented in their second paper¹⁹ a derivation which is similar in spirit, although considerably different from that suggested here.

Before describing our derivation, we want to emphasize the following. It is the standard fermionic NL M which does contain correct (at least for large $!$) non-perturbative oscillatory contributions to the level correlation functions. The replica-symmetry breaking which involves a set of additional saddle-point submanifolds within the standard NL M is just a very convenient method of calculating the large- $!$ asymptotic behavior of the correlation functions including these oscillatory contributions. Similarly, within the supersymmetric NL M Andreev and Altshuler²⁰ have reproduced the large- $!$ limit of Eq. (1.1) by the saddle-point calculation of the supersymmetric integral that involved an additional saddle-point which breaks the supersymmetry. The very same integral has been exactly calculated by Efetov¹¹ without breaking the supersymmetry. We extend this analogy to the fermionic replica NL M by showing, with the help of a trick which resembles the replica-symmetry breaking, that the $n = 0$ limit of the exact integral representation of the TL CF obtained by Verbaarschot and Zimbauer¹⁶ does contain, at least for large $!$, the correct oscillatory behavior.

The paper is organised as follows. In the next section we recall the formulation of the standard replica fermionic NL M and introduce all relevant notations. In Section III we present in detail a calculation of the large $!$ non-perturbative contribution to the TL CF for the unitary ensemble, based on replica-symmetry breaking. In Section IV we outline similar calculations for the orthogonal and symplectic ensembles. In Section V we extract the very same results from the exact integral representation for R_2 obtained by Verbaarschot and Zimbauer¹⁶ for the unitary ensemble. Finally, in Section VI we summarize our results and discuss perspectives of the replica method.

II. FORMULATION OF THE PROBLEM

We consider the two-level correlation function (TL CF) defined by

$$R_2(!) = \frac{1}{2} h(!!) (!!) i = 1 : \quad (2.1)$$

Here $h::i$ stands for the ensemble averaging, $(!!)$ is the electronic density of states per unit volume defined in terms of the spectrum $f''g$ for a given sample as $(!!) = L^{-d} \sum_{\alpha} \delta(E - E_{\alpha})$, $h(!!) i = 1/L^d$, and L, d are the sample size and dimensionality. We will measure all energies in units of $=$, in which the TL CF is as written in Eq. (1.1). In these units we express R_2 via the product of the retarded and advanced Green's functions as follows:

$$R_2(!) = \frac{1}{2} [e S_2(!) - 1]; \quad S_2(!) = \int_0^D G^r(!! + \frac{i}{2}) G^a(!! - \frac{i}{2}) : \quad (2.2)$$

The function S_2 can be expressed in the standard way^{11;21} in terms of the replica NL M:

$$S_2(!) = \lim_{n \rightarrow 0} \frac{1}{n^2} \frac{\partial^2}{\partial !^2} Z_n(!); \quad Z_n(!) = \int D Q \exp(-F[Q; !]) \quad (2.3)$$

where $F[Q; !]$ is the non-linear sigma model functional

$$F[Q; !] = \frac{1}{L^d} \int d^d r \text{Tr} \left[\frac{1}{8} D (r Q)^2 - \frac{i!}{4} Q \right] : \quad (2.4)$$

This functional has been derived⁴ for the model of free electrons in a random potential represented in terms of the fermionic (anticommuting) replicas. As a result, the field $Q(r)$ is an Hermitian matrix of rank $2n$ whose elements are quaternions for the orthogonal symmetry ensemble, complex numbers for the unitary symmetry ensemble which arises when time-reversal symmetry in the model is broken, and real numbers for the symplectic symmetry ensemble, which arises when spin-rotation symmetry is broken. It satisfies the standard constraints

$$Q^2 = 1_{2n}; \quad \text{Tr} Q = 0; \quad (2.5)$$

These constraints are resolved by representing Q as follows:

$$Q = U^y U = T^y T; \quad \epsilon = \text{diag}(1_n; -1_n); \quad T = \exp \begin{pmatrix} 0_n & t \\ t^\dagger & 0_n \end{pmatrix}; \quad (2.6)$$

Here t is an arbitrary $n \times n$ matrix, and T is obtained by factorizing matrices $U \in S(2n)$ with respect to redundant matrices $R \in S(n) \times S(n)$ that commute with ϵ , i.e. $U = RT$, where $S(n)$ is the appropriate symmetry group. These conditions mean that Q belongs to the compact Grassmannian manifold²² (coset space), $S(2n)/S(n) \times S(n)$. For three different classes of symmetry, $S(n)$ becomes the symplectic group, $Sp(n)$, for the orthogonal class, the unitary group $U(n)$ for the unitary class, and the orthogonal group, $SO(n)$, for the symplectic class,

The ergodic regime corresponds to the level separations much smaller than the Thouless energy, $D = L^2$ (which in the chosen units coincides, up to a numerical factor, with the dimensionless conductance g). In this regime the gradient term in Eq. (2.4) may be neglected, and the NL M functional reduces to the zero-dimensional limit:¹¹

$$F[Q; !] = \frac{i!}{4} \text{Tr}[Q]; \quad (2.7)$$

with Q becoming a spatially homogeneous matrix. Here and in Eq. (2.4) $\epsilon = 1$ for the orthogonal class, and $\epsilon = 2$ for the unitary and symplectic classes. This factor arises because unitary and symplectic classes have been obtained from orthogonal⁴ by the suppression of massive modes corresponding to the time-reversal or spin-rotational symmetry breaking and a subsequent reduction of the Q matrix rank. The coefficient $i!$ also absorbs an extra factor in the symplectic case due to the redefinition of the mean level spacing in the chosen units.

III. UNITARY ENSEMBLE

We limit all further considerations to the ergodic regime only, and rewrite explicitly $Z(!)$ given by Eqs. (2.3) and (2.7):

$$Z_n(!) = \int DQ \exp \left[-\frac{i!}{2} \text{Tr} Q \right]; \quad (3.1)$$

where the measure is defined by

$$DQ = \prod_{i,j=1}^n d^{ra}_{ij} d^{ra}_{ij}; \quad d^{ra}_{ij} = dT_{ij} d\tau_{ij}; \quad (3.2)$$

Here T is the matrix parameterizing Q , Eq. (2.6), and r and a refer to the replica indices which originate from G^r and G^a , respectively. In the large- $!$ limit this integral is mainly contributed by the extrema of the functional which obey the standard condition $[Q] = 0$. This condition is satisfied by any matrix of the form $Q = \text{diag}(Q^r; Q^a)$, where Q^r and Q^a are the $n \times n$ Hermitian matrices whose eigenvalues are ± 1 and $\text{Tr}(Q^r + Q^a) = 0$. This defines a highly degenerate saddle-point manifold which consists of C_{2n}^n submanifolds specified by a particular distribution of n eigenvalues ± 1 and n eigenvalues ± 1 between Q^r and Q^a . These submanifolds can be divided into $n+1$ classes of equivalence, $Q_p = \text{diag}(Q_p^r; Q_p^a)$, labeled by $\text{Tr} Q_p^r = \text{Tr} Q_p^a = n - 2p$, with $p = 0, 1, \dots, n$. The p -th class has weight $(C_n^p)^2$, with $C_n^p = \frac{n!}{p!(n-p)!}$.

The matrix Q_p^r with $(Q_p^r)^2 = 1_n$ and $\text{Tr} Q_p^r = n - 2p$ can be parameterized by analogy with Eq. (2.6) as

$$Q_p^r = (T_p^r)^y T_p^r; \quad T_p^r = \text{diag}(1_p; -1_p); \quad T_p^r = \exp \begin{pmatrix} 0_{n-p} & t^r \\ (t^r)^y & 0_p \end{pmatrix}; \quad (3.3)$$

where t^r is an arbitrary $p \times (n-p)$ matrix. This defines the coset space $G_p = U(n)/U(n-p) \times U(p)$, i.e. Therefore, $Q_p = \text{diag}(Q_p^r; Q_p^a)$ belongs to the manifold $G_p \times G_p$ and can be parameterized as

$$Q_p = T_p^Y T_p; \quad T_p = \text{diag}(T_p^r; T_p^a); \quad p = \text{diag}(p; -p) \quad (3.4)$$

The integer p specifies the replica-symmetry breaking, as it describes the number of the -1 eigenvalues in each Q block (equal to the number of the $+1$ eigenvalues in each Q^a block): in the symmetry-unbroken case, $p = 0$, and hence retarded and advanced blocks, $Q^{r;a}$, contain only positive or negative eigenvalues, respectively. We want to emphasize that manifolds $G_p \subset G_0$ with $p \notin p^0$ cannot appear within the degenerate saddle-point manifold, $[Q] = 0$, of the functional (2.7). Indeed, the corresponding matrices have non-zero trace, and thus do not belong to the Q space defined by Eq. (2.6). Naturally, one could have derived the replica NL M with different numbers of elements corresponding to the retarded and advanced blocks, as in Ref. 1. In this case, the replica-symmetry breaking would still mean the redistribution of $+1$'s and -1 's eigenvalues between these blocks, keeping the value of $\text{Tr} Q$ the same (equal to the difference between the numbers of replicas in the retarded and advanced blocks in the symmetry-unbroken case, $p = 0$) in each p -th manifold.

Now we need to take into account contributions from 'massive' modes (with $m \neq 0$) in the vicinity of each manifold (3.4). In the large- n limit these contributions may be considered as independent and the partition function is then represented by the sum of all of them:

$$Z_n(!) = \sum_{p=0}^n \frac{X^n}{(C_n^p)^2} \int DQ \exp \left\{ -\frac{i}{2} \text{Tr}_p Q \right\}; \quad (3.5)$$

Here we have used the fact that the similarity transformation in the vicinity of p , i.e. $U_p U^Y$, covers the entire symmetric manifold of the NL M, Eq. (2.6), including all the massive modes. Having substituted this into the functional (2.7), we have reduced $\text{Tr} U_p U^Y$ to $\text{Tr}_p T^Y T = \text{Tr}_p Q$, where we have substituted $U = RT$, as defined after Eq. (2.6). Let us emphasize again that the above representation of Z as the sum over all p can be justified only as a perturbative (in $1/n$) procedure: a possible overlapping of massive modes originated from different manifolds is irrelevant in the large- n limit.

Each term in the sum (3.5) contains both massive and massless modes. Indeed, we have used above the factorization $U = RT$ with R being block-diagonal matrices commuting with p . The matrices T in $\text{Tr}_p Q = \text{Tr}_p T^Y T$ still contain the subset of matrices commuting with p that correspond to the massless modes. Therefore, we need to parameterize T in a way which enables us to factorize out these massless modes and perform the integration over the massive ones.

The most suitable parameterization of T , analogous to that used in Ref. 16 for the bosonic NL M, can be obtained by expanding the matrix exponent in Eq. (2.6). By introducing matrix $B = \frac{1}{2}(\frac{1}{2}T + \frac{1}{2}T^Y)$, we represent T and thus $Q = T^Y T$ as follows:

$$T = \frac{1_n}{B^Y} \frac{B B^Y}{B^Y} \frac{1_n}{B^Y B}; \quad Q = \frac{1_n}{B^Y} \frac{2B B^Y}{1_n B^Y} \frac{1_n}{(1_n - 2B^Y B)}; \quad (3.6)$$

The matrix B in this parameterization is not unconstrained, though. The $Q = Q^Y$ condition is fulfilled only when the matrices $\frac{1_n}{B^Y} B B^Y$ and $\frac{1_n}{B^Y B} B^Y B$ are Hermitian. This is so only when all the eigenvalues of $B B^Y$ and $B^Y B$ do not exceed unity. Only under this constraint does Q , parameterized as in Eq. 3.6, still belong to the coset space $U(2n)/U(n) \times U(n)$. Nevertheless, this parameterization is very convenient. First, the corresponding Jacobian is unity (see Appendix), so that the measure of integration (3.2) can be written simply as

$$DQ = \prod_{i,j} dB_{ij} dB_{ij}^Y \quad DB; \quad (3.7)$$

In addition, the representation of all the exponents in the sum (3.5) in terms of B is also very simple, $\text{Tr}_p Q = 2(n - 2p) - 2\text{Tr}_p (B B^Y + B^Y B)$, so that we obtain:

$$Z_n(!) = \sum_{p=0}^n \frac{X^n}{(C_n^p)^2} e^{i!(2p-n)} Z_n^p(!); \quad (3.8)$$

$$Z_n^p(!) = \int DB \exp \{ -i! \text{Tr}_p (B B^Y + B^Y B) \}; \quad (3.9)$$

The region of integration in (3.9) is restricted by the constraint described after Eq. (3.6). Last, but not least, the parameterization (3.6) allows us to separate out the massless modes, which obey the condition $[T; p] = 0$, in each

integral (3.9). Indeed, this condition is satisfied by all matrices T constructed from B which anticommute with ρ , i.e. have the off-diagonal block structure.

This means that in the representation of B in the block form reflecting the structure of $\rho = \text{diag}(1_{n-p}; 1_p)$,

$$B = \begin{pmatrix} B_1 & b_1 \\ b_1^Y & B_2 \end{pmatrix}; \quad (3.10)$$

the matrices $B_{1,2}$ represent the massive modes, and $b_{1,2}$ massless. When the massive modes are suppressed ($B_1 = 0$ and $B_2 = 0$), the T matrices in Eq. (3.6) constructed from p $(n-p)$ matrices b_2 only, parameterize the same degenerate p -th manifold, $G_p \subset G_p$ described in Eq. (3.4), as one expects.

By substituting the representation (3.10) into Eq. (3.9), we reduce Z_n^p to the product of integrals over the massive and massless modes:

$$Z_n^p(!) = \int_{\mathbb{R}^{2(n-p)}} \int_{\mathbb{R}^{2p}} \exp \left[-\frac{1}{2} \text{tr} (B_1 B_1^Y + B_2 B_2^Y) \right] \int_{\mathbb{R}^{2p}} D b_1 D b_2; \quad (3.11)$$

Here the region of integration over $b_{1,2}$ depends on $B_{1,2}$ due to the constraint on the eigenvalues of the matrices $B B^Y$ and $B^Y B$ in the representation (3.10). Since the integral over $B_{1,2}$ is contributed only by the region where both $\text{tr} B_1 B_1^Y$ and $\text{tr} B_2 B_2^Y < 1$, in the leading in 1 approximation we may put both $B_{1,2}$ to 0 in the constraint of the integration region over the massless modes $b_{1,2}$. In this approximation, as we have noticed after Eq. (3.10), matrices $b_{1,2}$ parameterize the p -th manifold (3.4) so that

$$\int_{\mathbb{R}^{2p}} D b_1 D b_2 = \int_{Q_p} dQ_p = \text{Vol}(G_p); \quad (3.12)$$

where the measure of integration over dQ_p is defined in terms of T_p in the same way that dQ is defined in terms of T , Eq. (3.2), and $\text{Vol}(G_p)$ is the volume of the compact coset space G_p . This volume is expressed via the well-known volumes of the unitary group, $\text{Vol}(U(n))$, as follows:

$$\text{Vol}(G_p) = \frac{\text{Vol}(U(n))}{\text{Vol}(U(n-p)) \text{Vol}(U(p))} = (2)^{\frac{1}{2} [n^2 - (n-p)^2 - p^2]} \prod_{j=1}^p \frac{(1+j)}{(n+2-j)}; \quad (3.13)$$

In the same large- 1 approximation, the variables $B_{1,2}$ parameterizing the massive modes are unconstrained. Then the Gaussian integral over the $2[(n-p)^2 + p^2]$ independent massive modes yields

$$Z_n^p(!) = \int_{\mathbb{R}^{2(n-p)}} \int_{\mathbb{R}^{2p}} \exp \left[-\frac{1}{2} \text{tr} (B_1 B_1^Y + B_2 B_2^Y) \right] = \frac{(n-p)^2}{i!} \frac{p^2}{i!} \quad (3.14)$$

Combining Eqs. (3.14) and (3.13) and omitting an irrelevant overall factor which goes to 1 when $n \rightarrow 0$, we arrive at the following expression which is essentially the same as that derived in Ref. 17 via the Itzykson-Zuber integral:

$$Z_n(!) = \sum_{p=0}^n F_n^p \int_{\mathbb{R}^{2p}} \frac{e^{i! (2p-n)}}{(2!)^{(n-p)^2 + p^2}}; \quad F_n^p = C_n^p \prod_{j=1}^p \frac{(1+j)}{(n+2-j)}; \quad (3.15)$$

Here the summation over p has been extended to 1 since $F_n^p = 0$ for all integer $n > p$. This allows one to take the replica limit, $n \rightarrow 0$, in each of the terms in Eq. (3.15). Due to the fact that $F_n^p / n^p \rightarrow 0$ as $n \rightarrow 0$, only the terms with $p = 0$ and $p = 1$ in Eq. (3.15) contribute to S_2 in Eq. (2.3). Omitting all the terms with $p \geq 2$, one obtains

$$Z_n(!) = \frac{e^{i! n}}{i! n^2} + n^2 \frac{e^{i! (2-n)}}{4! (n-1)^2 + 1}; \quad (3.16)$$

Substituting this into Eqs. (2.3) and (2.2) and keeping the leading in 1 terms only, one arrives at the expression (1.1) for the T L C F. Although this expression looks as being exact, it has been actually derived only in the large- 1 limit, as is the case in the Itzykson-Zuber calculation of Kamenev and Mezard¹⁷, and in the 'supersymmetry breaking' method of Andreev and Altshuler.²⁰

A generalization to both these symmetries from the unitary one is straightforward: elements of all matrices in the previous section were complex numbers, while now they become quaternion numbers^{4,22} in the orthogonal symmetry and real numbers in the symplectic symmetry. In other words, $S(n)$, $S(p)$, etc, should be substituted for $U(n)$, $U(p)$, etc, in all formulae of Section III, with S being the symplectic group Sp for the orthogonal symmetry and the orthogonal group SO for the symplectic symmetry (such an inversion is characteristic of the fermionic replica approach). One should also redefine the measure of integration, Eq. (3.2), in terms of independent matrix elements and substitute $2!$ by $1!$ as defined in Eq. (2.7).

After these redefinitions, the partition function Z_n is represented by the sum (3.8), with each term in this sum given by the product of integrals over the massive and massless modes, Eq. (3.11). The number of massive modes is $(4 - \epsilon)(n - p)^2 + p^2$, where $\epsilon = 1; 2, 4$ for the orthogonal, unitary and symplectic symmetry is the standard Dyson parameter. For orthogonal and symplectic symmetries, the Gaussian integral (3.14) over the massive modes yields

$$Z_n^p(1) = \begin{cases} (1!)^{2[(n-p)^2 + p^2]} & \text{orthogonal symmetry} \\ (2!)^{[(n-p)^2 + p^2]/2} & \text{symplectic symmetry} \end{cases} \quad (4.1)$$

The integrals over massless modes, Eq. (3.12), give the volumes, $2^{(G_p)}$ of the corresponding coset spaces. The results can be written as

$$(G_p) = \begin{cases} \prod_{j=1}^p (2)^{2(n-p)p} \frac{(1+2j)}{[1+2(n+1-j)]} & \text{orthogonal symmetry} \\ \prod_{j=1}^p (1)^{(n-p)p/2} \frac{(1+j/2)}{[1+(n+1-j)/2]} & \text{symplectic symmetry} \end{cases} \quad (4.2)$$

Combining Eqs. 4.1 and 4.2, we obtain the partition function for both the symmetries. It can be written in terms of ϵ , together with the unitary Z_n , Eq. (3.15), in universal form:

$$Z_n(1) = \sum_{p=0}^X F^2(n;p) \frac{e^{-i!(2p-n)}}{(2!)^{\frac{\epsilon}{2}[(n-p)^2 + p^2]}}; \quad F(n;p) = C_n^p \prod_{j=1}^p \frac{(1+\frac{\epsilon}{2}j)}{(1+\frac{\epsilon}{2}(n+1-j))} \quad (4.3)$$

In the $n \rightarrow 0$ limit only the terms with $p = 0; 1$ survive for $\epsilon = 1; 2$, while for $\epsilon = 4$ one needs to keep the $p = 2$ term as well:

$$Z_{n \rightarrow 0}(1) = 1 - \frac{i!n + n^2}{2} - \frac{!^2}{2} \ln ! + \frac{!^2(1+2\epsilon)}{(2!)^4} \frac{2!}{e!} + \frac{e^{4i!}}{!^4 2^8 !^4} \quad (4.4)$$

Substituting this into Eqs. (2.3) and (2.2), we obtain the T LCF as follows:

$$R_2(1) = \frac{1}{!^2} + \frac{!^2(1+2\epsilon)}{(2!)^4} \cos 2! + \frac{\cos 4!}{!^4 32 !^4} \quad (4.5)$$

This reproduces the correct large- $!$ asymptotic behavior of R_2 for all three ensembles, which includes the non-perturbative oscillatory factors.

As we have derived these results with a non-standard treatment of the standard fermionic NLM, they also should be contained in the exact integral representation obtained for this model by Verbaarschot and Zirnbauer.¹⁶ We will show now that this is, indeed, the case.

V. THE LARGE- $!$ LIMIT OF THE VERBAARSCHOT-ZIRNBAUER INTEGRAL

The 'zero-mode' partition function, Eq. (2.7), can be exactly represented in the following form

$$Z_{n+1}(1) = \frac{Z_{n+1}}{1} (U(n)) \prod_{i=1}^n \frac{Y_i}{2} e^{-i! - i d_i}; \quad (1) \prod_{i < j} (j - i) : \quad (5.1)$$

which is equivalent (with accuracy up to factors going to 1 in the $n \rightarrow 0$ limit) to the representation for S_2 given in Eq. (2.24) of the paper by Verbaarschot and Zimbauer.¹⁶

The leading in $1/n$ contributions to this highly oscillatory integral (which does not have stationary points inside integration region) come from the end points. To single out these contributions, we must take some ϵ 's close to $+1$ and the rest close to -1 , which imitates replica symmetry breaking. Let us choose $n-p$ of ϵ 's close to $+1$ and p of ϵ 's close to -1 . Then we can split up the Vandermonde determinant in the following way:

$$\Delta^2(\epsilon) = \prod_{i,j}^{Y^n} (j - \epsilon_i \epsilon_j) = 2^{p(n-p)} \prod_{i,j}^{Y^n} (j - \epsilon_i \epsilon_j)$$

where

$$\prod_{i,j=1}^{Y^p} (j - \epsilon_i \epsilon_j) = \prod_{i,j=n-p+1}^{Y^n} (j - \epsilon_i \epsilon_j) \quad (5.2)$$

Reducing the integral (5.1) to the sum of such contributions only, we represent it as

$$Z_n(!) = \sum_{p=1}^{X^n} (C_n^p)^2 2^{2p(n-p)} \int_1^{Z+1} \prod_{j=1}^{Y^p} e^{i! \epsilon_j} d\epsilon_j \int_1^{Z+1} \prod_{j=n-p+1}^{Y^n} e^{i! \epsilon_j} d\epsilon_j \quad (5.3)$$

Since in each of the integrals all the variables are close to one of the limits of integration, the second limit was extended to infinity. Now we make substitutions $\epsilon_i = 1 - x_i$ in the first integral, and $\epsilon_i = -1 + x_i$ in the second one, reducing the above sum to the form:

$$Z_n(!) = \sum_{p=1}^{X^n} (C_n^p)^2 2^{2p(n-p)} e^{i!(2p-n)} I_{n-p} I_p \quad (5.4)$$

where I_p are integrals of the Selberg's type:¹²

$$I_p = \int_0^Z \prod_{j=1}^{Y^p} (x - \epsilon_j) dx e^{i! x_j} \quad (5.5)$$

Substituting the known Selberg integrals and discarding an overall factor which goes to unity in the replica limit we arrive at

$$Z_n(!) = \sum_{p=0}^{X^n} F_n^p \int_0^Z \frac{e^{i!(2p-n)}}{2! (n-p)^2 + p^2}; \quad F_n^p = C_n^p \prod_{j=1}^{Y^p} \frac{(1+j)}{(n+2-j)} \quad (5.6)$$

This expression is exactly the same as Eq. 3.15 obtained in Section III with the help of the replica-symmetry breaking. Therefore, the exact representation (5.1) does contain the true oscillatory asymptotic behavior of the TLGF.

The authors of Ref. 16 have also drawn attention to the fact that there is an apparent contradiction between the $n \rightarrow 0$ limit for S_2 obtained from the replica trick and the exact supersymmetric result. Indeed, if $!$ is put to 0 in the expression for S_2 following from Eqs. (5.1) and (2.3) the $n \rightarrow 0$ limit is taken after that, one obtains $S_2(!=0) = -1$. This cannot be correct as $\langle e S_2(!=0) \rangle > 0$ as follows from the definition (2.2) and, moreover, it is known that $S_2(!=0) \neq -1$. What is interesting, however, is that if one separates the singular, $S_2^{\text{sing}}(!)$, and regular, $S_2^{\text{reg}}(!)$, parts of $S_2(!)$, then $S_2^{\text{reg}}(!=0) = -1$. Therefore, the replica method gives $S_2^{\text{reg}}(!=0)$ correctly, and it is just S_2^{sing} which is missing. However, the fact that $Z_n(!=0)$ is finite for any integer n does not necessarily imply that it is also finite (as a function of $!$) in the replica limit. For example, if the expansion of $S_2(!)$ before taking the $n \rightarrow 0$ limit contained a term proportional to $!^{n^2-1}$, it would be singular in the replica limit. In other words, if a non-trivial dependence on the order of limits $n \rightarrow 0$ and $! \rightarrow 0$ existed, in a spirit of the replica trick the $n \rightarrow 0$ limit should be taken first. At the moment, though, this is only a speculation as we have not yet succeeded in calculating the integral (5.1) for small $!$. However, the fact that the large- $!$ limit of this integral reproduces the correct results (1.1) makes it plausible that there are only technical difficulties rather than one of principle in the application of the replica method.

VI. CONCLUSIONS

The purpose of this paper was to demonstrate explicitly that non-perturbative oscillatory contributions to the T LCF of electrons in a random potential could be extracted from the standard N L M formulated in fermionic replicas by Efetov, Larkin and Khmelnitskii many years ago. To this end, all one needs is to parameterize all the non-trivial saddle-point manifolds corresponding to the broken replica-symmetry and describing ‘massless modes’ of the theory, and expand the action in the vicinity of these manifolds to include ‘massive modes’. The very similar approach has been used in the supersymmetric N L M : the non-perturbative oscillations have been extracted by the expansion around two extremal points one of which breaks the supersymmetry.²⁰ Since the exact supersymmetric calculation of the T LCF was well known,¹¹ it was clear that the supersymmetry breaking²⁰ was just a convenient method of extracting the large- l limit (and going beyond the universal ‘zero-mode’ approximation). It is not clear at the moment whether exact calculation is possible within the replica approach. However, as we have shown in the Section V, the exact integral representation of $R_2(!)$ does at least describe the correct behavior in the large- l limit.

On the other hand, there is a number of non-perturbative results where all the leading contributions come from the small- l limit (see, e.g., Ref. 23): the existence of a nontrivial spatially inhomogeneous saddle point in the non-compact sector of the supersymmetric N L M was the main source of these contributions. At the moment it is unclear whether such a saddle point could be found within the fermionic replica approach. Nevertheless, the fact that some of these results had earlier been obtained^{7,24} within the RG treatment of the fermionic replica N L M prompts that it could be technical rather than methodological difficulties involved.

Despite the fact that the replicas prove now to be a trusted tool for non-perturbative calculations, there is no doubt that the supersymmetric N L M is the best tool to obtain non-perturbative results for non-interacting electrons. But the further elaboration of the non-perturbative methods within the replica trick will be important for the problems where supersymmetry cannot be applied. This is the situation with the interacting electrons in a random potential. Such systems can be described by N L M with fermionic replicas only and we believe that the further development of methods similar to presented here will enable one to obtain at last non-perturbative results for disordered systems of interacting electrons.

ACKNOWLEDGMENTS

This work has been supported by the EPSRC grant GR/K 95505.

APPENDIX A : CALCULATION OF JACOBIAN

To calculate differentials of matrices entering the measure (3.2) we use the following rational substitution:

$$B = \frac{2}{1-y} \quad \Rightarrow \quad T = \begin{pmatrix} 0 & 1+y \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 2 & 1+y \end{pmatrix}^{-1} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad (A.1)$$

Then the differentials in the measure (3.2) are given by

$$d^{12}(T) = dT \cdot T^{-1} = 2 \frac{1}{1-y} (d \begin{pmatrix} 1 & 2 \\ 2 & 1+y \end{pmatrix}^{-1}) \frac{1}{1-y} : \quad (A.2)$$

On the other hand we can find the differential of B , entering the measure (3.7) as follows:

$$dB = 2 \frac{1}{1-y} (d \begin{pmatrix} 1 & 2 \\ 2 & 1+y \end{pmatrix}^{-1}) \frac{1}{1-y} : \quad (A.3)$$

Comparing Eqs. (A.2) and (A.3) one can see that the Jacobians of transitions $d^{12}(T) \rightarrow d$ and $dB \rightarrow d$ are exactly the same. From that fact one deduces that the Jacobian of the transition $d^{12}(T) \rightarrow dB$ is unity.

- ¹ F. Wegner, Z. Phys. B 35, 207 (1979).
- ² K. B. Efetov, Sov. Phys. JETP 55, 514 (1982).
- ³ L. Schafer and F. Wegner, Z. Phys. B 38, 113 (1980); A. J. McKane and M. Stone, Ann. Phys. (N.Y.) 131, 36 (1981).
- ⁴ K. B. Efetov, A. I. Larkin, and D. E. Khmel'nitskii, Sov. Phys. JETP 52, 568 (1980).
- ⁵ J. J. M. Verbaarschot, H. A. W. Eidenmüller, and M. R. Zimbauer, Phys. Rep. 129, 367 (1985).
- ⁶ E. Abraham, P. W. Anderson, D. C. Licciardello, and T. V. Ramakrishnan, Phys. Rev. Lett. 42, 673 (1979).
- ⁷ B. L. Altshuler, V. E. Kravtsov, and I. V. Lerner, Sov. Phys. JETP 64, 1352 (1986).
- ⁸ A. M. Finkelstein, Sov. Phys. JETP 57, 97 (1983); Z. Phys. B 56, 189 (1984).
- ⁹ S. V. Kravchenko et al., J. Phys. {Condens. Matter 7, L41 (1995); Phys. Rev. B 51, 7038 (1995).
- ¹⁰ A. Kamenev and A. Andreev, cond-mat/9810191; C. Chamon, A. W. W. Ludwig, and C. Nayak, cond-mat/9810282.
- ¹¹ K. B. Efetov, Sov. Phys. JETP 56, 467 (1982).
- ¹² M. L. Mehta, Random matrices, Academic Press, Boston (1991).
- ¹³ L. P. Gor'kov and G. M. Eliashberg, Sov. Phys. JETP 21, 940 (1965).
- ¹⁴ K. B. Efetov, Supersymmetry in Disorder and Chaos, Cambridge University Press, Cambridge (1997).
- ¹⁵ B. L. Altshuler and B. I. Shklovskii, Sov. Phys. JETP 64, 127 (1986).
- ¹⁶ J. J. M. Verbaarschot and M. R. Zimbauer, J. Phys. A 17, 1093 (1985).
- ¹⁷ A. Kamenev and M. Mezard, cond-mat/9901110.
- ¹⁸ T. Guhr, J. Math. Phys. 32, 336 (1991).
- ¹⁹ A. Kamenev and M. Mezard, cond-mat/9903001.
- ²⁰ A. V. Andreev and B. L. Altshuler, Phys. Rev. Lett. 75, 902 (1995).
- ²¹ R. A. Smith, I. V. Lerner, and B. L. Altshuler, Phys. Rev. B 58, 10343 (1998).
- ²² B. A. Dubrovin, A. T. Fomenko, and S. P. Novikov, Modern Geometry, Part II, Springer-Verlag, Berlin (1985).
- ²³ B. A. Muzikantskii and D. E. Khmel'nitskii, Phys. Rev. B 51, 5480 (1995); K. B. Efetov and V. I. Fal'ko, Europhys. Lett. 32, 627 (1995); A. D. Mirlin, Phys. Rev. B 53, 1186 (1996); V. E. Kravtsov and I. V. Yurkevich, Phys. Rev. Lett. 78, 3354 (1997).
- ²⁴ B. L. Altshuler, V. E. Kravtsov, and I. V. Lerner, in Mesoscopic Phenomena in Solids, edited by B. L. Altshuler, P. A. Lee, and R. A. Webb, North-Holland, Amsterdam, 449 (1991).