Evolutionary dynam ics in the Bak-Sneppen model on small-world networks

R.V.Kulkami, E.Alm aas and D.Stroud

Department of Physics, The Ohio State University, Columbus, OH 43210

(M arch 23, 2024)

We study the dynamics of the Bak-Sneppen model on small-world networks. For each site in the network, we de nea \connectance," which measures the distance to all other sites. We nd radically di erent patterns of activity for di erent sites, depending on their connectance and also on the topology of the network. For a given network, the site with the minimal connectance shows long periods of stasis interrupted by much smaller periods of activity. In contrast, the activity pattern for the maxim ally connected site appears uniform on the same time scale. We discuss the signi cance of these results for speciation events.

PACS num bers: 87.10.+ e, 87.23.-n, 05.40-a

The theory of punctuated equilibrium (PE) [1] states that (i) most evolutionary change associated with life on Earth occurs during speciation events, and (ii) the tim e scale for these speciation events is very brief com pared to the lifetimes of the individual species. Thus the evolutionary history of most species is characterized by long periods of stasis punctuated by relatively brief intervals of rapid evolutionary activity. Mayr's theory of allopatric speciation [2], which form s the basis of PE, suggests that the rapid evolutionary activity leading to speciation generally occurs in sm all isolated populations, w hereasm ore w idespread populations exhibit little evolutionary change over the sam e tim e scale. Understanding why the rates of evolutionary change di er so drastically in these two types of populations is one of the important problem s of evolutionary biology. A related problem is the study of patterns of extinctions in biological history, which seem to show scale-free behavior as discussed by Raup [3]. Stanley [4] has argued that these problem s, and most large-scale trends in evolutionary history, can be understood by using the species as a fundam ental unit of evolution.

Bak and Sneppen [5] have introduced a simple model aim ed at understanding these evolutionary patterns using the species as the fundam ental unit. The model is based on coevolution of species and exhibits \interm ittent dynam ics" - that is, species undergo long periods of little change, called stasis, which are punctuated by sudden bursts of activity called avalanches. It provides a natural explanation for the apparent scale-free behavior of the extinctions - the system evolves into a self-organized critical (SOC) state with avalanches (which are correlated with extinction events) occurring at all scales. The same model can also be used to understand the scalefree behavior of seem ingly unrelated phenom ena such as earthquakes, as noted by Ito [6]. A side from its applications to various problem s, the Bak-Sneppen m odel is of intrinsic interest, since it is one of the simplest models giving rise to SOC behavior.

The Bak-Sneppen model, as elaborated below, has

been extensively studied for regular networks [7]. However, as argued by W atts and Strogatz [8], most reallife networks are neither perfectly ordered nor completely disordered but fall under the category of $\mall+world$ " networks which interpolate smoothly between the two extrem es. Such networks are characterized by a high degree of local order, yet appear disordered on a large-scale because of the presence of shortcuts in the networks. Because of their wide applicability, there have recently been num erous papers characterizing the properties of such networks [9]. In the context of species interacting in an ecosystem, exam ples of food webs indicate that the interactions are better represented by small+world networks rather than by a sim ple ordered topology.

The aim of this Letter is to exam ine the dynamics of the Bak-Sneppen model on small-world networks. In our analysis, we nd it useful to de ne a site-dependent property which we call \connectance." We nd that the patterns of activity at each site are correlated with the connectance. In particular, the minim ally connected site in a given network shows activity characteristic of intermittent dynamics, whereas the maxim ally connected site shows uniform activity on the same time scale. Furthermore, for the maxim ally connected site we see a drastic reduction in the stasis times in going from the ordered topology to the small-world networks. We discuss the im plications of these results for speciation events.

We begin by recalling the de nition of the Bak-Sneppen model. Each species is represented by a site on a 1-d lattice with periodic boundary conditions. Each lattice site is connected to its 2k nearest neighbors by a bond (so that for n interacting sites, or species, we have nk bonds). W ith each site we associate a random number, called the barrier value, between 0 and 1; this number plays the role of a barrier against evolutionary change for that species. The dynam ics of the Bak-Sneppen m odel is carried out by the follow ing rule. At each time step, we select the site w ith the m inim al barrier value (denoted the m inim al site), and assign new barrier values to this site and its 2k neighboring sites. We refer to a reassignm ent of the barrier value at a site as activity at that site. This set of rules, called extrem aldynam ics, leads the system into a SOC state where the distribution of barrier values is uniform above a critical barrier value f_c [7].

The sm all-world networks are generated using the procedure outlined by W atts and Strogatz [8]. W e start with the 1-d lattice described above and \rewire" each bond with a probability p (0 p 1). The rewiring consists of taking a given bond and moving it so that, instead of connecting two neighbors, one end of the bond is moved so as to connect with another site chosen at random, with the constraint that double bonds are forbidden. The network so form ed can then be characterized by the functions L (p), which is de ned as the number of bonds in the shortest path between two sites, averaged over all pairs of sites, and C (p) which is a measure of clustering in the graph [8]. The network is said to fall in the smallworld regime if it satis are the conditions C (p)=C (0) 0. Table I shows the values of these and L (p)=L (0) two ratios for the networks we studied, all of which were characterized by n = 2000 and k = 2. A recent study on food webs [10] suggests that our choice of k is often realized in practice. Finally, for each site, we de ne the connectance q_i by

$$g_{i} = 1 \quad \frac{D_{i} \quad \min_{fjg} D_{j}}{\max_{fjg} D_{j} \quad \min_{fjg} D_{j}} ; p > 0 \quad (1)$$

$$D_{i} = \int_{j=1}^{X^{N}} d(i;j)$$
(2)

where d(i; j) is the m inim aldistance between sites i and j, that is, the m inim alnum ber of bonds which must be traversed in going from i to j.

We carried out our simulations for n = 2000 and for p = 0.0, 0.01, 0.04, and 1.0. For each disordered network ($p \in 0$) we studied the dynam ics for 10 di erent realizations. Remarkably, even though the topologies for a given p value were distinct, our calculated quantities all collapsed onto the same set of curves. The results we present were obtained by sam pling for 5 10^9 time steps after the stationary state characteristic of SOC had been reached. In the following, we discuss results for p = 0.0 and p = 0.01 since these capture the essential features in going from ordered to small-world networks.

In the Bak-Sneppen model, an avalanche is de ned as the sequence of time steps for which the minimal site has

TABLE I. The functions L (p)/L (0) and C (p)/C (0) as dened in the text, calculated for a system with n = 2000 sites and k = 2. For each p, we averaged over 25 di erent realizations.

р	L(p)/L(0)	C(p)/C(0)
0.01	0.14	0.97
0.04	0.06	0.89
1.00	0.02	0.04



FIG.1. Number of avalanches () and number of sites covered (full line without symbols; N_{cov}) by an avalanche vs. duration s of the avalanche, for p = 0.01. The in uence of short-cuts, or sm all-world behavior, is rejected in the slope changes between regions A and B. In region C, nite-size e ects dom inate. The inset shows the same plots for p = 0.0.

a barrier value sm aller than a threshold value f_0 . The avalanche thus ends when the minim al site has a barrier value greater than $f_0\,.$ For each p-value, we choose f_0 such that $f = f_c$ $f_0 = 0.01$. Fig. 1 shows the distribution of avalanche durations and the average num ber of sites covered (N_{cov}) by an avalanche of a given duration for p = 0.0 and p = 0.01. For p = 0.0 (inset) we see the expected power-law behavior with a cut-o for large avalanche sizes. For p = 0.01, on the other hand, we see novel features arising from the small-world properties of the network. In particular, we now see two power-law regimes, denoted by regions A and B.As the avalanche size grows, it increasingly \senses" the short-cuts in the system, and becom es m ore delocalized; this delocalization produces an increased slope in N_{cov} in going from regime A to regime B.A more detailed analysis of this behavior will be presented elsewhere [11].

In Fig. 2, we show the so-called nst-return plots, i.e., the distribution of waiting times between subsequent returns of activity, at the maximally connected sites for p = 0.0 and 0.01. All the plots show power-law behavior followed by a cut-o which corresponds to the maximal stasis time. We note that the maximal stasis time decreases by roughly two orders of magnitude in going from p = 0.0 to p = 0.01. Thus the transition from ordered to small-world networks is marked by a drastic reduction of the maximal stasis times. The inset shows the

rst-return plots for the maximally and the minimally connected sites in a p = 0.01 network. From these plots we see that the cut-o is correlated with the connectance: higher connectance leads to a smaller cut-o time for a given network.

A dram atic consequence of the reduction in stasis times



FIG.2. Distribution of rst-return (stasis) times for the maximally connected sites for p = 0.0 () and p = 0.01 (). Note that the cut-o, which corresponds to the maximal stasis time, is reduced by 2 orders of magnitude in going from p = 0.0 to p = 0.01. The inset shows the distribution of stasis times for the maximally () and the minimally (+) connected site in a p = 0.01 network.

can be seen in Fig. 3, where we show the activity plots at the maximally connected sites for p = 0.0 and p = 0.01. For p = 0:0, the cum ulative activity at a particular site shows a pattern of \punctuated equilibrium :" long periods of stasis interrupted by much shorter periods of activity. By contrast, the maxim ally connected site at p > 0 shows a uniform pattern of activity on the same time scale. This behavior results from the decrease in maximal stasis time (cf. Fig. 2) with increasing p. This relationship is clearly indicated in the inset, which shows that the same pattern now exhibits \punctuated equilibrium "behavior on a much smaller time scale. The correlation between maximal stasis time and connectance for a given network can be seen in Fig. 4, where the activity is plotted for the sites with maxim al and minim al connectance at p = 0.01. The m inim ally connected site shows a pattern of \punctuated equilibrium," whereas the site with maximal connectance exhibits uniform behavior. In short, the periods of stasis are signi cantly reduced when either p or g is increased.

The uniform activity pattern seen for p > 0 may appear to be at variance with the theory of punctuated equilibrium. But in fact, we argue that this behavior actually supports that theory in the form postulated by G ould and E ldredge [1]. We est clarify the correspondence between our idealized model and an ecosystem . As a lready noted, the sites correspond to species, while the network of connections between them can be viewed as representing the food web of the ecosystem . A ctivity at a site represents a signi cant change for the corresponding species population in the ecosystem; an accumulation of activity can lead to either speciation or extinction. Our



FIG.3. Calculated cum ulative activity at the maximally connected sites for (a) p = 0.0 and (b) p = 0.01. W hile (a) shows a \punctuated equilibrium " pattern, (b) appears uniform on the same time scale. The inset shows that actually (b) also exhibits a \punctuated equilibrium " pattern, but on a much reduced time scale.

m odel does not distinguish between the two, and we expect instances of both speciation and extinction to be proportional to the amount of activity. Finally we note that a bond between two species represents a dependency link between them, which is typically, but not exclusively, a predator-prey relationship.

We now discuss the correspondence between the activity patterns described above and the theory of punctuated equilibrium . We note that the staircase pattern seen in Fig. 3 (a) corresponds to an evolutionary process called anagenesis, in which the entire ancestral species evolves into a new species. But, as noted by Stanley [4], most speciation processes in nature are actually branching processes (called cladogenesis), in which a new species is created from a geographically isolated subpopulation of the ancestral species. During cladogenesis, the ancestral species shows little change (stasis), whereas the isolated sub-population undergoes rapid evolutionary change. This is the basic process involved in the theory ofpunctuated equilibrium . Thus, what is required is that the periods of stasis be greatly reduced for the isolated population, in comparison to the ancestral species. This behavior is precisely what is seen in Fig. 3: in these plots, increasing p results in m uch shorter periods of stasis and hence, an activity plot which has the appearance of uniform ity in time.

Thus, our scenario for the time sequence of allopatric speciation events, as motivated by our simulations, is the following. (1) A sub-population migrates to a peripherally isolated region, such as an island. We expect the island ecology to have a higher value of p for the interaction network; hence, the periods of stasis for the isolated sub-population should be much sm aller than those of the



FIG.4. Cumulative activity for sites with (a) maximal and (b) m inimal connectance for p = 0.01. The maximally connected site has a cumulative activity which is 40 times that of the m inimally connected site.

ancestral population. (2) The m igration event itself initiates a period of activity on the island [12], while stasis continues to prevail for the ancestral population. The com bination of these two e ects gives rise to rapid evolutionary change leading to speciation for the island population, and simultaneously stasis in the ancestral population.

Note that this model does not exclude other proposed mechanisms, such as the founder e ect proposed by Mayr [2], but instead is complementary to them . Our scenario focuses on the coevolutionary activity in di erent ecosystem s and thus on e ects external to the species populations, whereas the other mechanisms focus on the internal characteristics of the species populations. For example, the founder e ect requires that the peripherally isolated population be small, and therefore more am enable to rapid change. Speciation can, however, occur rapidly over a large area containing many millions of individuals. In particular, W illiam son [13] has shown that, in a group of A frican lakes in the Turkana basin, which shrank and became separated from the parental water bodies, the species underwent signi cant phylogenetic change over only 5 10^4 years, whereas the ancestral populations remained virtually unchanged in their parental waters. In this case, the observed speciation events cannot be explained by the founder e ect, as argued by W illiam son, but they do seem consistent with our proposed scenario.

O ur calculations show that, for a given network, the patterns of activity are strongly correlated with the connectance. How is this observation re ected in studies of real ecosystems? O ne possibility is the following: since the bonds in the network corresponds to dependency links, the species with high connectance have a high degree of dependency - i. e., they are specialists (stenotypic) - whereas species with low connectance are generalists (eurotypic). As already noted in Fig. 4, the cumulative activity is an order of magnitude larger for the maximally connected site than for the one with minimal connectance. Based on our model, we therefore expect stenotypic species to speciate or go extinct more offen than their eurotypic counterparts. This correlation between ecological specialization and speciation and extinction rates has been observed in num erous studies [14].

In sum m ary, we have extended the 1-d Bak-Sneppen m odel to small-world networks. This extension allows us to distinguish di erent sites in the network, and thereby the species in an ecosystem, based on a site-dependent property which we call connectance. We indicate the activity patterns are strongly correlated with the topology of the network and with the connectance. Finally, we discuss the possible in plications of these results for speciation events.

We thank Jens O. Andersen for useful discussions. This work has been supported by NASA through G rant NCC 8-152 and through NSF through G rant DMR 97-31511. C om putational support was provided by the O hio Supercom puter C enter and by the San D iego Supercom – puter C enter.

E-mail: rahul@mps.ohio-state.edu

- [L] S. Gould and N. Eldredge, Paleobiology 3, 114 (1977); Nature 366, 223 (1993).
- [2] E.M ayr Populations, Species and Evolution (C am bridge M ass., H arvard Univ. Press., 1970).
- [3] D.M.Raup, Science 231, 1528 (1986).
- [4] S.M. Stanley, M acroevolution, Pattern and Process (W. H.Freem an and Company, 1979).
- [5] P.Bak and K.Sneppen, Phys. Rev. Lett. 71, 4083 (1993).
- [6] K. Ito, Phys. Rev. E. 52, 3232 (1995).
- [7] M. Paczuski, S. Maslov, and P. Bak, Phys. Rev. E 53, 414 (1996) and references therein.
- [8] D.J.W atts, S.H. Strogatz, Nature 393, 440 (1998).
- [9] M. Barthelem y and L.A. N. Am aral, Phys. Rev. Lett. 82, 3180 (1999) See also cond-m at preprints 9903323, 9903357 and 9903411.
- [10] G. Sugihara, K. Schoenly, and A. Trom bla, Science 245, 48 (1989) see also M. E. J. Newman et. al. Proc. R. Soc. London B 260, 31 (1995).
- [11] E.Almaas, R.V.Kulkami, D.Stroud (in preparation).
- [12] See examples discussed by J.H.Brown in Complexity: metaphors, models, and reality edited by G A Cowan, D Pines, and D E M eltzer (Addison-W esley, 1994).
- [13] P.G.W illiam son, Nature 293, 437 (1981).
- [14] N. E klredge, M acroevolutionary D ynam ics (1989) In particular see E.S.Vrba, S.A fr. J. Sci. 76, 61 (1980).