

Topological Gauge Theory Of General Weytenbock Manifolds Of Dislocations In Crystals

Y. C. Huang^{1,3} B. L. Lin² S. Li³ M. X. Shao^{3,4}

¹Department of Applied Physics, Beijing Polytechnic University, Beijing, 100022, P. R. China

²Department of Transportation Management Engineering, Northeast Jiaotong University, Beijing, 100044, P. R. China

³Institute of Theoretical Physics, Chinese Academy of Science, Beijing, 100080, P. R. China

⁴Department of Physics, Beijing Normal University, Beijing, 100022, P. R. China

(¹Email: ychuang@bjpu.edu.cn)

Abstract

General Weytenbock material manifolds of dislocations in crystals are proposed, the reference, idealized and deformation states of the bodies in general case are uniformly described by the general manifolds, the topological gauge field theory of dislocations is given in general case, true distributions and evolution of dislocations in crystals are given by the formulas describing dislocations in terms of the general manifolds, furthermore, their properties are discussed.

1. Introduction

Practical crystals always contain a lot of dislocations and are filled with them, the defects strongly affect the properties of the crystals. Some pioneers even in the fifties had researched the relations between the defect crystals and differential geometry. K. Kondo used torsion of material manifolds to study dislocations¹, and Nye², Bilby³, Kroner⁴ etc studied the relations between the defects and torsion.

On the other hand, these defects are found having the features of elementary particles, and Kroner studied continuum theory of defects⁵. Due to the great success of Yang-Mills gauge field theory describing the interactions of elementary particles⁶, according to the research of gauge theory of gravitation, it can be discovered that geometry of Riemannian manifold essentially belongs to a kind of non-Abelian gauge theory⁷. Gauge theory and topological properties of dislocations were researched, and many important conclusions in the fields were obtained by Toulous and Kleiman⁸, Kleibert⁹,

Trebin¹⁰ and so on. Duan and his associates showed a unified approach to the study of defect mechanics and their topological and geometric properties by applying vielbein field theory and gauge field theory to dislocation and disclination continuum^{11;12}.

Sect.2 of the paper constructs the general Weitzenböck manifolds of dislocations in crystals, gives a unified description of these reference, idealized and deformation states; Sect.3 unifiedly studies topological gauge theory of dislocations, gives the formulas describing dislocations, their properties are discussed; Sect.4 gives the discussions of overcoming the defect; the last Sect. is summary.

2 General Weitzenböck Material Manifolds of Dislocations

The Weitzenböck manifold⁷ is the one whose $r_{ij}g_{jk} = 0$, $T_{jk}^i \neq 0$, $R_{jkl}^i = 0$, where g_{ij} , T_{jk}^i and R_{jkl}^i are, respectively, metric, torsion and curvature tensors of the manifold, when $T_{jk}^i = 0$, the manifold is reduced into Euclidean manifold. Since properties of dislocations correspond to properties of torsion tensor of a manifold¹⁻⁴, when considering dislocations, the corresponding manifold is thus taken as the Weitzenböck manifold. And the torsion tensor and metric tensor g_{ij} of a manifold are all defined locally on the manifold¹³, then we may naturally expect that the torsion tensor possesses different values, even zero, on different parts of the manifold, we call the manifold as the general Weitzenböck manifold, the corresponding different parts are just the corresponding submanifolds in the material manifold. In fact, the situations above just correspond to disappearance and appearance of dislocations for zero and non-zero of torsion tensor of the corresponding submanifold respectively. Therefore, we can use the above discussions to represent the complex situations of motion, disappearance and appearance of the dislocations.

Now we construct the general Weitzenböck material manifolds, relative to time parameter, of dislocations in crystals.

It is well known that Weitzenböck manifold is an ensemble of lots of small coordinate pieces homomorphic to some local Euclidean spaces, lying, however, to a certain degree anorphously, and the characters of the ensemble of lots of the small pieces can be described by means of a net connections of the small pieces. And the geometric properties of dislocations homomorphically correspond to the geometric properties of torsions of the corresponding material manifold.

Owing to the periodic regular distributions of the lattice particles of per-

fect crystal, the position coordinate x^a of lattice particles can be globally determined in Euclidean space in terms of their periodic rules, then we regard their corresponding material manifold as Euclidean manifold. Obviously, when dislocations appear, which means that we can not globally do the same in terms of their periodic rules, however, we can build up a lot of local coordinate systems in which the position coordinate x^a of any particles in the crystal can be locally defined.

In general, for a crystal with dislocations in 3-dimensional Euclidean space, we assume that the body is made of N lattice particles, for any lattice particle P_{i0} , we can always construct local coordinate system L_{i0} in local Euclidean space and presume that there are k lattice particles $P_{i1}; P_{i2}; \dots; P_{ik}$ which are close adjacent to P_{i0} , and the corresponding nonholonomic normalized frame fields are e^a ($a = 1; 2; 3$), then the distance vector of any two close adjacent particles is

$$dr = e^a dx^a \quad (1)$$

the metric tensor is

$$g_{ab} = e^a e^b = \delta_{ab} \quad (2)$$

and the square of the distance, i.e., the square of the line element is

$$ds^2 = dr \cdot dr = g_{ab} dx^a dx^b = dx^a dx^a \quad (3)$$

where dx^a is the corresponding nonholonomic coordinate difference.

Furthermore, for any particle P_{i1} ($i \neq 0$) discussed above, we can again construct a local coordinate system, and regard any such P_{i1} ($i \neq 0$) as a new P_{i0} ; do further the same as the above discussions about the $P_{i1}, \dots; P_{ik}$; up to all particles in the body.

For a open subset $V \subset \mathbb{R}^3$ of any local coordinate system discussed above, assume that there is an inverse homeomorphic map $\phi^{-1}: V \rightarrow M$ (where M is a family of open sets) and $\int_V = V_m$ (V_m is the volume of the crystal in Euclidean space) such that $\int M = M$ is topological space, which means that M is provided with family of pairs $\{M; \phi\}$. Further assume that given M_T and $M' (\in \mathcal{M})$ ($M_T \cap M' \neq \emptyset$) the map ϕ^{-1} from the subset $\phi^{-1}(M_T \cap M')$ of \mathbb{R}^3 to the subset $\phi^{-1}(M_T \cap M')$ of \mathbb{R}^3 is infinitely differentiable. Consequently, the aggregation of the all coordinate pieces form a general three dimensional differentiable material manifold¹³. Because the

general manifold inherits the topological and geometric properties of the dislocations in the body, and these properties correspond to properties of torsion of the manifold and can just be expressed by the torsion tensor of the manifold, therefore, there always exists the inverse map ϕ^{-1} such that M is the manifold that the properties of the dislocations of the crystal may be represented by means of the torsion tensors of the manifold. Because the torsion of the general manifold may have different values on different parts of the manifold, the different parts may correspond to Euclidean and Weyssenbock material submanifolds of the general material manifold. Evidently, the local Euclidean and Weyssenbock material submanifolds can be locally included in the general Weyssenbock material manifold, which just corresponds to the real distribution situations of dislocations of the body.

The relationship^{15;16} between the normalized nonholonomic and holonomic frames is

$$e^a = e_i^a e^i; a; i = 1; 2; 3 \quad (4)$$

where $e^i = dx^i$ is holonomic coframe of the manifold and e_i^a is the vielbein, then we have metric tensor of the manifold M

$$g_{ij} = e_i^a e_j^a \quad (5)$$

Using the vielbein theory^{7;14-16} we can have

$$dx^a = e_i^a dx^i \quad (6)$$

Substituting (6) into (3) and using (5), we have

$$ds^2 = dx^a dx^a = e_i^a dx^i e_j^a dx^j = g_{ij} dx^i dx^j \quad (7)$$

which means that the square of the line element of a local Euclidean system in the crystal is equal to that of the line element of the general material manifold. Using the above discussions, furthermore, we can study their stress and so on. Therefore, the above discussions is essential for our further study.

If we take time t_0 to label the reference state of time t_0 , when the body is experienced a deformation during time $4t = t - t_0$ the corresponding deformed state is labelled by time $t = t_0 + 4t$, in other words, at different time t , the lattice particles possess different distributions by which the creating and moving of the dislocations with time t are represented, the general material manifold with time parameter t is thus obtained.

In elastic and plastic mechanics, the reference, idealized and deformation states are chosen. We now give a unified description of the states by means of the general manifold. The reference state is usually chosen as perfect or idealized states of an ordered material. By the discussions above of translating the discreteness of a perfect crystal into the continuum, we obtain a Euclidean material manifold corresponding to the global vanishing torsion of the manifold. Usually the idealized state of the body is defined as an aggregation of lots of small pieces of the idealized material. In fact, the small pieces of the idealized material can always be viewed as having some ordered distributions of lattice particles, and our discussions about the piece distributions of the particles are general, i. e., our discussions are general, then the idealized state of the body is included in the states of the general Weytzenbock material manifold M . For elastic and plastic deformation states, they can be represented in microscope as the changes of the displacement and interaction of any adjacent lattice particles. Analogous to the above discussions, evidently the elastic and plastic deformation states are included in the general state of the general manifold M , especially plastic deformation has cooperative motions of the many defects and lattice particles. Now we generally consider reference state (or called standard state) and deformation state of M , the line elements of their manifolds are generally defined, respectively, as

$$ds_r^2 = ds_{t_0}^2 = g_{ij}(x(t_0); t_0) dx^i(t_0) dx^j(t_0) \quad (8)$$

and

$$ds_d^2 = ds_t^2 = g_{ij}(x(t); t) dx^i(t) dx^j(t) \quad (9)$$

Using (1) and (5), Eq.(9) can be generally rewritten as

$$ds_t^2 = e_i^a(x(t); t) e_j^b(x(t); t) dx^i(t) dx^j(t) \quad (10)$$

Similar to Ref.[11], the strain tensor is defined as

$$E_{ij} = E_{ij}(x(t); x(t_0); t; t_0) = \frac{1}{2} [g_{ij}(x(t); t) - g_{ij}(x(t_0); t_0)] \quad (11)$$

The general reference and deformation states and the strain tensor at any time t are useful for practical calculation, and it is convenient for acquiring different properties of a crystal, in fact, one usually compares reference states

with deformation states under different conditions and studies the evolution laws of the deformation states with time t .

It is important and useful to describe these different states in general case, we overcome the difficulty of self-consistence that generally describes these different states in terms of differential geometry, give true distributions and evolution of dislocations with variances of time in the crystals by means of the differential manifolds, and may give useful tool of describing the defects.

3. Topological Gauge Field Theory Of Dislocations

We now discuss the topological gauge field theory of the defects. In physics, any physical law doesn't depend on the choices of coordinates. For example, the square of the line element and Lagrangian of the system, there exist two kinds of indices i and a ($i, a = 1, 2, 3$) which is holonomic and nonholonomic indices respectively. For holonomic indices i , the coordinate transformations at a certain time t , i.e., at a certain state, are

$$x^{0i}(t) = x^{0i}(fx^j(t)g) \quad (12)$$

and

$$x^i(t) = x^i(fx^{0j}(t)g) \quad (13)$$

The transformation of the nonholonomic indices is the local $SO(3)$ gauge transformation¹¹

$$dx^{0a}(t) = S^{ab}(t)dx^b(t) \quad (14)$$

where

$$^{0ab} = \alpha S^{ac}S^{bd} \quad (15)$$

Using Eqs.(14), (15) and the discussions above, it is easy to prove that Eqs.(8-10) are all invariant under the two kinds of coordinate transformations.

In order to achieve the invariance of physical laws under the two kinds of coordinate transformations, the usual partial derivative should be substituted into two kinds of covariant derivatives as follows

$$r_i e_j^a = \partial_i e_j^a - \frac{k}{ij} e_k^a \quad (16)$$

and

$$D_i e_j^a = \partial_i e_j^a + \frac{!}{i}^{ab} e_j^b \quad (17)$$

where ω_{ij}^k and ω_i^{ab} are a affine connection and $SO(3)$ gauge potential (spin connection). Eqs.(16) and (17) satisfy the general and gauge covariant principles respectively, but the total Lagrangian of the system is invariant under the both transformations¹⁶:

Using the vielbein theory^{11;12;16} we have the torsion tensor of the general manifold as follows:

$$T_{ij}^k = \omega_{ij}^k - \omega_{ji}^k = e^{ka} (D_i e_j^a - D_j e_i^a) = e^{ka} T_{ij}^a \quad (18)$$

where T_{ij}^a is the torsion tensor with nonholonomic superscript a

$$T_{ij}^a = D_i e_j^a - D_j e_i^a \quad (19)$$

Analogous to Ref.[12,17], we may define the general tensor densities of dislocations of the body at any time t in the following

$${}^{ia}(\mathbf{x}(t);t) = \frac{\omega^{ilm}}{2 \sqrt{g(\mathbf{x}(t);t)}} T_{lm}^a(\mathbf{x}(t);t) \quad (20)$$

where $g = \det(g_{ij})$; $\omega^{123} = \omega^{132} = 1$, the coefficient $\frac{1}{2}$ is due to the tensor sums of subscripts l and m . Then Burgers vector of dislocation can be defined as follows

$$b^a = \int_{\Sigma} {}^{ia}(\mathbf{x}(t);t) \frac{q}{\sqrt{g(\mathbf{x}(t);t)}} d_i \quad (21)$$

where Σ is the surface including dislocations. The formulas (20) and (21) and $\mathbf{x}(t)$ are all the functions of the time t that labels the motions of the defects and deformations of a crystal, which includes the creations, motions and disappearances of the dislocations with evolution of time t in the general manifold, therefore, these variances can be represented by Burgers vector, or the tensor density of dislocations, these variances correspond to the variances that Eq.(20) and Eq.(21) are equal to different values at different time and position. Then it is given that true distributions and evolution of dislocations in the crystals by the formulas describing dislocations at any time t in the general manifold, and our theory is non-linear. Using gauge potential decompositions Duan and Zhang¹² gave the moving, geometric and topological descriptions of dislocations in terms of similar to Eqs. (20) and (21). Since the general Weytenbock manifold is very general, then the corresponding

further discussions of (20) and (21) are similar to those in Ref.[12], therefore, one can arrive in their all results of topology, geometry and kinematics of the defects, thus we shall not repeat here.

The discussions about elastic and plastic dynamics and the generalization to a four dimensional general pseudo-Weytenbock material manifold that concerns dissipative parts belonging to the fourth component of the vielbein fields will be written in a separated paper.

4. Summary

This paper proposes general Weytenbock material manifolds, relative to time parameter t , of dislocations in crystals, in the manifolds the local Euclidean and Weytenbock manifolds can be viewed as the submanifolds of the general Weytenbock material manifold, furthermore, by analyzing the microscope structures and change, with time t , of the general manifolds, we obtain the unified description of reference, idealized and deformation states in terms of the general manifolds, and the general manifold with evolution of time parameter t just represents the motions and the distributions of dislocations, and the topological gauge theory are acquired by leaving the physical laws invariant under the coordinate transformations. We overcome the difficulty of self-consistence that generally describes these different states in terms of differential geometry, give true distributions and evolution of dislocations in the crystals by the formulas describing dislocations to any time in general manifold, and their properties are discussed.

Acknowledgment

One, Huang, of the authors is grateful to Prof. Y. S. Duan and Prof. S. L. Zhang for help.

References

- ¹K .K ondo, M em oirs of the Unifying Study of the Basic P roblem s in Engi-
neering Sciences by M eans of G eom etry, Vol.1, D ivision D , P .6-17, P .70-
110, Tokoy (1955).
- ²J.F .N ye, A cta M etal, 1 (1953)153.
- ³B .A .B ilby and R .B ullough, and E .Sm ith, P roc.R .Soc., A 231 (1955)263.
- ⁴E .K roner and R .R ieder, Z .P hys., 145 (1956)424.
- ⁵E .K roner, Continuum theory of defects, in Less H ouches, Session X X X V ,
1980-P hysics of defects, North-H olland, A m sterdam , (1981).
- ⁶N .C .Y ang and R .L .M ills, P hys. Rev., 96 (1954)191.
- ⁷Y .S .D uan, G auge T heory of G ravitation, in P roc. Sym p on Yang-M ills
G auge T heories, Beijing, 1984.
- ⁸G .T oubus and M .K lem an, J. P hys. Lett. (Paris) , 37 (1976)L-249.
- ⁹H .K leinert, G auge F ields in Condensed M atter, W orld Scienti c P res, Singapore (1990).
- ¹⁰H .R .T reb in, A dv. P hys., 31 (1982)195.
- ¹¹Y .S .D uan and Z .P .D uan, Int. J. Engng. Sci. , 24 (1986)513.
- ¹²Y .S .D uan, and S .L .Z hang, Int. J. Engng. Sci., 28 (1990)689; 29 (1991)153.
- ¹³S .K obayashi, and K .N om izu, Foundations of di erential G eom etry, Inter-
science, New York, Vol.I (1963), Vol. II (1969).
- ¹⁴Y .C .H uang, Inner Structure of Torsion and Topological G auge F ield T he-
ory of D efects in Solid, Ph D Thesis, Lanzhou University, China, April
(1993).
- ¹⁵M .C arm ei, Classical F ields: General Relativity and G auge Theory, A
W iley-Interscience Publication, New York, 1982.
- ¹⁶Y .S .D uan et al, G en. Rel. G rav. 20 (1983)485.

- ¹⁷Y . C . Huang, Gauge eld theory of Riemann-Cartan manifolds of dislocations and disclinations, 7th International Conference on Mechanical Behaviour of Materials, Delft University Press, The Netherlands, P 848 (1995).