

Decomposing Finite Abelian Groups

Kevin K. H. Cheung

Michele Mosca^y

April 15, 2024

Abstract

This paper describes a quantum algorithm for efficiently decomposing finite Abelian groups. Such a decomposition is needed in order to apply the Abelian hidden subgroup algorithm. Such a decomposition (assuming the Generalized Riemann Hypothesis) also leads to an efficient algorithm for computing class numbers (known to be at least as difficult as factoring).

1 Introduction

The work by Shor [9] on factoring and finding discrete logarithms over $\mathbb{Z}_n$ can be generalized to solve the Abelian Hidden Subgroup Problem (see for example [10], [4], [7]). These algorithms find the hidden subgroup of a function $f : G \rightarrow S$, where $G = \mathbb{Z}_{N_1} \times \mathbb{Z}_{N_2} \times \dots \times \mathbb{Z}_{N_k}$ for some integers $N_1, N_2, \dots, N_k$. Any Abelian group G is isomorphic to such a product of cyclic groups, however it is not always known how to find such an isomorphism efficiently. Consider for example the group $\mathbb{Z}_N$, the multiplicative group of integers modulo N . This is an Abelian group we can compute in efficiently, yet no known classical algorithm can efficiently find its decomposition into a product of cyclic groups. Consider also the class group of a quadratic number field. This group is also Abelian, and finding its decomposition into a product of finite cyclic groups will give us the size of the group and therefore the class number of the quadratic number field. As Watrous [11] points out, assuming the generalized Riemann Hypothesis we can apply the algorithm in this paper and efficiently find class numbers (a problem known to be at least as hard as factoring).

In this paper, we show how we can make use of the solution to the Abelian Hidden Subgroup Problem to decompose a finite Abelian group. Such decompositions makes it possible to apply the Abelian Hidden Subgroup algorithm to a larger class of Abelian groups.

2 Integer Arithmetic Basics

A nonsingular integral matrix U is called unimodular if $\det U = \pm 1$. It is easy to check that U is unimodular if and only if U^{-1} is unimodular.

The following operations on a matrix are called elementary (unimodular) column operations:

1. exchanging two columns;
2. multiplying a column by -1 ;
3. adding an integral multiple of one column to another column.

^xPh.D. student, Department of Combinatorics and Optimization, Faculty of Mathematics, University of Waterloo, Waterloo, Ontario, N2L 3G1, Canada. E-mail: kkhcheun@math.uwaterloo.ca. Research of this author was supported by NSERC PGSB

^yDepartment of Combinatorics and Optimization, Faculty of Mathematics, University of Waterloo, Waterloo, Ontario, N2L 3G1, Canada. E-mail: mmosca@cacr.math.uwaterloo.ca. Research of this author was supported by NSERC

We define elementary row operations similarly.

Theorem 1 For any integral matrix A , one can find in polynomial time using elementary row and column operations unimodular matrices U and V such that $UAV = \begin{pmatrix} D & 0 \\ 0 & 0 \end{pmatrix}$ where $D = \text{Diag}(d_1, \dots, d_k)$ with positive integers such that $d_1 | d_2 | \dots | d_k$ and for each i , the product $d_1 \dots d_i$ is equal to the g.c.d. of the subdeterminants of A of order i . We call the matrix $\begin{pmatrix} D & 0 \\ 0 & 0 \end{pmatrix}$ the Smith normal form (abbreviated as SNF) of A .

Proof : See Kannan and Bachem [5].

3 Group Theory Basics

Recall that a group G is said to be Abelian if for all $a, b \in G$, $ab = ba$. In this paper, all groups are finite Abelian unless otherwise stated. G is said to be cyclic if there exists a $g \in G$, such that $G = \{a^n | n \in \mathbb{Z}\}$. Here, we call a a generator of G . $H \subseteq G$ is called a subgroup of G if H is a group under the operation induced by G . In this case, we write $H \leq G$. Let $a \in G$. The set $aH = \{ah | h \in H\}$ is called a coset of H in G determined by a .

Let $a \in G$. If $a^n = e$ for some $n \in \mathbb{N}$, then a is said to have finite order. The smallest such n is called the order of a , denoted by $\text{ord}(a)$. It is easy to see that the elements in $\{e, a, a^2, \dots, a^{n-1}\}$ form a subgroup. We call this subgroup the cyclic subgroup generated by a and we denote it by $\langle a \rangle$.

Let G_1, G_2 be groups such that $G_1 \cap G_2 = \{e\}$. The set $\{a_1a_2 | a_1 \in G_1, a_2 \in G_2\}$, denoted by $G_1 \times G_2$, is called the direct sum of G_1 and G_2 . Note that $G_1 \times G_2$ is a group under binary operation such that $(a_1b_1)(a_2b_2) = (a_1a_2)(b_1b_2)$.

Let G be a group and p be a prime number. Let $P \leq G$. Then P is called a Sylow p -subgroup of G if $|P| = p^k$ for some $k \in \mathbb{N}$ such that p^k divides $|G|$ but p^{k+1} does not.

We first quote a few classical results without proof. The interested reader can refer to a standard text on group theory.

Theorem 2 If N is a subgroup of an Abelian group G , then the set of cosets of N forms a group under the coset multiplication given by

$$aN \cdot bN = abN$$

for all $a, b \in G$. The group is denoted by G/N .

Theorem 3 Let N be a subgroup of a finite Abelian group G . If $a_1, \dots, a_k$ generates G , then $a_1N, \dots, a_kN$ generates G/N .

Theorem 4 A finite Abelian group can be expressed as a direct sum of its Sylow p -subgroups.

Theorem 5 Let K be a subgroup of $G = G_{p_1} \times \dots \times G_{p_l}$ where G_{p_i} is a Sylow p_i -subgroup for $i = 1, \dots, l$ and $p_1, \dots, p_l$ are distinct primes. Then there exists $K_{p_i} \leq G_{p_i}$, $i = 1, \dots, l$, such that $K = K_{p_1} \times \dots \times K_{p_l}$.

The next theorem is an important result on finite Abelian groups.

Theorem 6 (Fundamental Theorem of Finite Abelian Groups). Any finite Abelian group can be decomposed as a direct sum of cyclic subgroups of prime power order.

In this paper, we shall give an algorithm to find the decomposition.

The next theorem is an integral part of the algorithm.

Theorem 7 Given a generating set $\{a_1, \dots, a_k\}$ of a finite Abelian group G and a matrix M such that $a_1^{x_1} \dots a_k^{x_k} = e$ if and only if $x = (x_1, \dots, x_k)^T \in \text{intcol}(M)$ where $\text{intcol}(M)$ denotes the set of vectors obtainable by taking integer linear combinations of columns of M , we can find in polynomial time (in the size of M) $g_1, \dots, g_l$ with $1 \leq l \leq k$ such that $G = \langle g_1, \dots, g_l \rangle$.

Proof : (Adapted from Algorithm 4.1.3 in [3].) By Theorem 1, we can find in polynomial time unimodular matrices U and V such that $U^{-1} M V = \begin{pmatrix} D & 0 \\ 0 & 0 \end{pmatrix}$ where D is a diagonal matrix with diagonal entries $d_1, \dots, d_m$. Since V is unimodular, $\text{intcol}(M V) = \text{intcol}(M)$. Thus, $a_1^{x_1} \dots a_k^{x_k} = e$ if and only if $x = (x_1, \dots, x_k)^T \in \text{intcol}(M V)$. For each $i = 1, \dots, k$, set $a_i^0 = a_1^{U_{1i}} \dots a_k^{U_{ki}}$. Then

$$\begin{aligned} & a_1^{0x_1} \dots a_k^{0x_k} = e \\ & , \quad (x_1, \dots, x_k)^T \in \text{intcol}(U^{-1} M V) \\ & , \quad d_i \mid x_i \text{ for } i = 1, \dots, m; \text{ and } x_i = 0 \text{ for } i = m+1, \dots, k \end{aligned}$$

Since G is finite, we must have $m = k$. Otherwise, G will have an element of infinite order. Let j be the smallest index such that $d_j > 1$. Set $g_i = a_{i+j-1}^0$ for $i = 1, \dots, l$ where $l = m - j + 1$. It is clear that $g_1, \dots, g_l$ still generate G and g_i has order d_{i+j-1} for $i = 1, \dots, l$. Therefore, if $0 < x_i < \text{ord}(g_i)$, then $g_1^{x_1} \dots g_l^{x_l} \neq e$ implies that $x_i = 0$ for all i . Hence $G = \langle g_1, \dots, g_l \rangle$.

4 Hidden Subgroup Problem

Let $G = \mathbb{Z}_{N_1} \times \dots \times \mathbb{Z}_{N_l}$ where the N_j , $j = 1, \dots, l$ are prime powers. We are given $f : G \rightarrow S$ for some finite set S that is constant on cosets of some $K \leq G$ but distinct on each coset. (The case when distinct cosets are not mapped to distinct elements is addressed in Boneh and Lipton [2] and in the Appendix of [7]. Here, we need $m < \sum_{j=1}^l N_j$ where m is the maximum number of cosets that get mapped to the same output.) The hidden subgroup K is

$$\{k \in G \mid f(x) = f(x+k) \text{ for all } x \in G\}.$$

The Hidden Subgroup Problem is to find generators for K given only f and G .

There exist polynomial-time quantum algorithms to solve this problem.

Corollary 8 Let a be an element of a group G . The order r of a can be found in random quantum polynomial time.

Proof : Consider the function f from $\mathbb{Z}$ to the group G where $f(x) = a^x$. Then $f(x) = f(y)$ if and only if $x - y \in r\mathbb{Z}$. The hidden subgroup is $K = r\mathbb{Z}$ and a generator for K gives us the order r of a .

Using Corollary 8, one can deduce the result by Shor [9].

Theorem 9 Factoring can be solved in random quantum polynomial time.

In the next section, we shall show how to use the algorithm for finding hidden subgroup to decompose finite Abelian groups.

5 Decomposing Abelian Groups

By Theorem 6, we know that we can decompose a finite Abelian group into a direct sum of cyclic groups of prime power order. This problem was discussed briefly in [6]. We make four assumptions on the group G :

1. We have a unique binary representation for each element of G and we can efficiently recognize if a binary string represents an element of G or not.
2. Using the binary representation, for any $a \in G$, we can efficiently construct a quantum network for implementing $U_a : |y\rangle \mapsto |ay\rangle$.
3. We can efficiently find a generating set for G .
4. The orders of the generators are of prime power order.

To meet the third assumption, it suffices to have an upper bound of 2^k on the size of the groups we work with for some $k \leq (\log |G|)$ and that we can efficiently sample elements of G uniformly at random. (If we do not have such a bound, we can easily devise a procedure that tries an increasing sequence of values for k and still has expected running time in $O(\text{poly}(\log |G|))$.) Let K be a proper subgroup of G . Then there are at least two cosets of K . If we randomly sample an element x from G , then with probability at least $1/2$, the subgroup spanned by x and K will have size at least twice that of K because the elements xk for all $k \in K$ are in the span. Hence, it takes an expected number of at most $(1/(1/2))^k = 2^k$ samples to obtain a generating set G for G and therefore $2k + c \cdot k$ samples will find a generating set with probability in $1 - \epsilon$ for some $\epsilon \in (0,1)$ (by a Chernoff bound.)

Now we may assume that the order of the elements are of a prime power. Let a be an element in G with order p^q where $(p,q) = 1$, $p \notin 1$ and $q \notin 1$. Note that p and q can be determined efficiently as a result of Corollary 8 and Theorem 9. By the Euclidean algorithm, we can find r,s such that $rp + sq = 1$. Thus $(a^p)^r (a^q)^s = a$. Hence, replacing a with a^q and a^p still leaves us with a generating set. We repeat this procedure until each element in G has prime power order.

Since we know the order p^q of a , we can efficiently compute $a^{-1} = a^{p^q-1}$ and therefore efficiently perform the necessary uncomputation in order to satisfy the second assumption.

By Theorem 4, we have $G = G_{p_1} \times \dots \times G_{p_l}$ where p_i is a prime for all $i = 1, \dots, l$ and G_{p_i} is a Sylow p_i -subgroup of G . Let S_j be the set of all the elements in G having order a power of the prime p_j . For $a \in S_j$, let K_a denote the (cyclic) subgroup generated by a . By Theorem 5, we have $K_a = K_{p_1} \times \dots \times K_{p_l}$ where $K_{p_i} \leq G_{p_i}$ for all $i = 1, \dots, l$. Since $|K_a|$ is a power of p_j , we must have $K_a \leq G_{p_j}$. Thus $S_j \leq G_{p_j}$. Since G generates G , S_j generates G_{p_j} . Hence, we can first find the decomposition for each of the Sylow p -subgroups of G and then take their product to obtain a decomposition of G .

There are two primary reasons why we want to have the fourth assumption. One reason is that we want to minimize the amount of quantum computing resources required in any implementation. It is therefore advisable to decompose the problem whenever it is possible. The second reason is that working with p -groups can greatly simplify the amount of algebra one needs to perform to recover the generators. This latter point will be elaborated at the end of the section. In the meantime, we present the algorithm given in [6] which finds generators of a group G with prime power order.

Algorithm 10 Decompose Group $(a_1, \dots, a_k)$

Input:

A generating set $a_1, \dots, a_k$ of the group G .

The maximum order $q = p^f$ of the elements $a_1, \dots, a_k$.

Output:

A set of elements $g_1, \dots, g_l$, $1 \leq l \leq k$, from the group G .

Procedure:

1. Define $g : Z_q^k \rightarrow G$ by mapping $(x_1, \dots, x_k) \mapsto g(x) = a_1^{x_1} \dots a_k^{x_k}$. Find generators for the hidden subgroup K of Z_q^k as defined by the function g .
2. Compute a set $y_1, \dots, y_l \in Z_q^k = K$ of generators for $Z_q^k = K$.
3. Output $g(y_1), \dots, g(y_l)$.

To see the correctness of this algorithm, observe that the hidden subgroup K is the set $\{(x_1, \dots, x_k) \mid a_1^{x_1} \dots a_k^{x_k} = e\}$. We therefore have an isomorphism between $Z_q^k = K$ and G . If $y_1, \dots, y_l$ are generators for $Z_q^k = K$, then $g(y_1), \dots, g(y_l)$ are generators for G .

We now elaborate on how it is possible to find generators for $Z_q^k = K$. Observe that $e_1, \dots, e_k$ generate Z_q^k where e_i is a 0,1-vector with a 1 in the i th co-ordinate. Further, if we let $M = qI$ where I is the $k \times k$ identity matrix, then $x_1 e_1 + \dots + x_k e_k = 0$ (in Z_q^k) if and only if $x \in \text{intcol}(M)$. By Theorem 3, $e_1 + K, \dots, e_k + K$ generate $Z_q^k = K$. Note that

$$v_1(e_1 + K) + \dots + v_k(e_k + K) = K$$

if and only if $Iv \in K$ where I is the matrix $[e_1, \dots, e_k]$ and $v = (v_1, \dots, v_k)^T$. Let A be the matrix the columns of which generate K . Note that $Iv \in K$ if and only if there exists a vector x such that

$$\begin{aligned} Iv &= Ax \\ , \quad Iv &= IAx \\ , \quad I(v - Ax) &= 0 \\ , \quad v - Ax &\in \text{intcol}(M) \\ , \quad v &\in \text{intcol}(M - A): \end{aligned}$$

Applying Theorem 7 to $fe_i + K$ $j = 1, \dots, k$ and $M^{-1} = [M - A]$, we obtain $y_1, \dots, y_l \in Z_q^k = K$ such that $Z_q^k = K = \langle y_1, \dots, y_l \rangle$ as desired.

Technically, we need not work with each Sylow p -subgroup of G separately. Suppose $a_{i1}, \dots, a_{ik_1}$ generates the group G_{p_i} . Let $q_i = p_i^{r_i}$ be the maximum order of $a_{i1}, \dots, a_{ik_1}$. Define $g : Z_{q_1}^{k_1} \times \dots \times Z_{q_l}^{k_l} \rightarrow G$ by mapping $(x_{11}, \dots, x_{1k_1}, \dots, x_{l1}, \dots, x_{lk_l})$ to $g(x) = \prod_{i=1}^l \prod_{j=1}^{k_i} a_{ij}^{x_{ij}}$. Proceed as before. The only differences are that we need to build a huge quantum network to solve the hidden subgroup problem and that in computing generators for $Z_{q_1}^{k_1} \times \dots \times Z_{q_l}^{k_l} = K$ where K is the hidden subgroup K defined by g , we need to work with a huge matrix when applying Theorem 7 if the block structure of the matrix is not exploited. In practice, it is therefore desirable to avoid this approach. Furthermore, for each prime p , instead of using Z_q^k where $q = p^f$ is the maximum order of the elements a_j (i.e. $r = \max\{t_1, t_2, \dots, t_k\}$ where the order of a_j is p^{t_j}), we could use $Z_{p^{t_1}} \times Z_{p^{t_2}} \times \dots \times Z_{p^{t_k}}$.

R e f e r e n c e s

- [1] J.A.Beachy and W.D.Blair. Abstract algebra with a concrete introduction. Prentice-Hall Inc., New Jersey, 1990.
- [2] D.Boneh and R.J.Lipton. Quantum cryptanalysis of hidden linear functions (extended abstract). Lecture Notes on Computer Science, 963:424{437. Springer, Berlin, 1995.
- [3] H.Cohen. Advanced Topics in Computational Number Theory. Springer-Verlag, New York, 1991.
- [4] P.Hoyer. Conjugated operators in quantum algorithms. IMADA preprint, 1997.
- [5] R.Kannan and A.Bachem. Polynomial algorithms for computing the Smith and Hermite normal forms of an integer matrix. SIAM Journal on Computing, 8(4):499{507, 1979.
- [6] M.Mosca. Quantum Computer Algorithms. D.Phil Thesis, University of Oxford, 1999.
- [7] M.Mosca and A.Ekert. The hidden subgroup problem and eigenvalue estimation on a quantum computer. Lecture Notes in Computer Science, 1509:174{188, Springer, Berlin, 1999.
- [8] A.Schrijver. Theory of Linear and Integer Programming. Wiley and Sons, England, 1986.
- [9] P.W.Shor. Algorithms for quantum computations: Discrete logarithms and factoring. In S.Goldwasser, editor, Proceedings of the 35th Annual Symposium on Foundations of Computer Science, pages 124{134. IEEE Computer Society Press, November 1994.
- [10] U.Vazirani. UC Berkeley Course CS294-2 Quantum Computation Fall 1997.
- [11] J.Watrous. Quantum algorithms for solvable groups. preprint, 2000.