

Dichotomy Theorems for Alternation-Bounded Quantified Boolean Formulas^{*}

Edith Hemaspaandra

Department of Computer Science,
Rochester Institute of Technology,
Rochester, NY 14623, U.S.A.,
`eh@cs.rit.edu`

Abstract. In 1978, Schaefer proved his famous dichotomy theorem for generalized satisfiability problems. He defined an infinite number of propositional satisfiability problems, showed that all these problems are either in P or NP-complete, and gave a simple criterion to determine which of the two cases holds. This result is surprising in light of Ladner's theorem, which implies that there are an infinite number of complexity classes between P and NP-complete (under the assumption that P is not equal to NP).

Schaefer also stated a dichotomy theorem for quantified generalized Boolean formulas, but this theorem was only recently proven by Creignou, Khanna, and Sudan, and independently by Dalmau: Determining truth of quantified Boolean formulas is either PSPACE-complete or in P.

This paper looks at alternation-bounded quantified generalized Boolean formulas. In their unrestricted forms, these problems are the canonical problems complete for the levels of the polynomial hierarchy. In this paper, we prove dichotomy theorems for alternation-bounded quantified generalized Boolean formulas, by showing that these problems are either Σ_i^P -complete or in P, and we give a simple criterion to determine which of the two cases holds. This is the first result that obtains dichotomy for an infinite number of classes at once.

Keywords: quantified Boolean formulas, computational complexity, Boolean constraints, dichotomy, polynomial hierarchy

1 Introduction

In 1978, Schaefer proved his famous dichotomy theorem for generalized satisfiability problems. He defined an infinite number of propositional satisfiability problems (nowadays often called Boolean constraint satisfaction problems), showed that all these problems are either in P or NP-complete, and gave a simple criterion to determine which of the two cases holds. This result is surprising in light of Ladner's theorem, which implies that there are an infinite number of

^{*} Research supported in part by grant NSF-CCR-0311021

complexity classes between P and NP-complete (under the assumption that P is not equal to NP).

To make the discussion more concrete, we will quickly define what a constraint is and what a constraint problem is. Formal definitions can be found in Section 2. In this paper, we will be talking about *Boolean* constraints. See for example Feder and Vardi [FV98] for a discussion about general constraint satisfaction problems.

A *constraint* is a Boolean operator of fixed arity, specified as a Boolean function. For C a constraint of arity k , and x_1, \dots, x_k propositional variables (or constants), $C(x_1, \dots, x_k)$ is a *constraint application* of C . For example, $\lambda xy.(x \vee y)$ is a constraint, and $x_1 \vee x_2$ is a constraint application of this constraint. Each finite set of constraints \mathcal{C} gives rise to a satisfiability problem $\text{SAT}(\mathcal{C})$: $\text{SAT}(\mathcal{C})$ is the problem of, given a set of constraint applications of \mathcal{C} , determining whether this set has a satisfying assignment. We can view a set of constraint applications as a CNF formula. For example, 2CNF-SAT corresponds to $\text{SAT}(\{\lambda xy.(x \vee y), \lambda xy.(x \vee \bar{y}), \lambda xy.(\bar{x} \vee \bar{y})\})$.

Using constraint terminology, Schaefer's dichotomy theorem [Sch78] can now be formulated as follows: For any finite set of constraints \mathcal{C} , either $\text{SAT}(\mathcal{C})$ is in P, or $\text{SAT}(\mathcal{C})$ is NP-complete.

In recent years, dichotomy theorems (or dichotomy-like theorems) have been obtained for a number of other problems about logics. For example, such theorems have been obtained for the problem of determining whether a formula has exactly one satisfying assignment [Jub99], the problem of finding a satisfying assignment that satisfies a maximum number of constraint applications [Cre95], the problem of computing the number of satisfying assignments [CH96], the problem of finding the minimal satisfying assignment [KK03], the inverse satisfiability problem [KS98], the equivalence problem [BHRV02], the isomorphism problem [BHRV04], and the complexity of propositional circumscription [KK01]. Khanna, Sudan, Trevisan, and Williamson examined the approximability of some of these problems [KSTW01]. Consult the excellent monograph [CKS01] for an almost completely up-to-date overview of dichotomy theorems for Boolean constraint problems.

Schaefer also stated a dichotomy theorem for quantified generalized formulas (or, equivalently, quantified sets of constraint applications), but this theorem was only recently proven by Creignou, Khanna, and Sudan [CKS01], and independently by Dalmau [Dal97]: Depending on the underlying finite set of constraints, these problems are either PSPACE-complete or in P.

This paper looks at *alternation-bounded* quantified sets of constraint applications. In their unrestricted forms, alternation-bounded quantified Boolean formulas are the canonical problems complete for the levels of the polynomial hierarchy. In this paper, we prove dichotomy theorems for alternation-bounded quantified sets of constraint applications, by showing that these problems are either Σ_i^P -complete or in P, and we give a simple criterion to determine which of the two cases holds.

The importance of these results is two-fold. First of all, unlike all previous results, our result obtains dichotomy for an infinite number of classes at once (namely, we prove dichotomy for each level of the polynomial hierarchy). Secondly, Schaefer’s dichotomy theorem has proven very successful as a tool for proving NP-hardness. After all, his theorem supplies us with an infinite number of NP-complete variations of the already often-used satisfiability problem. We expect that our dichotomy theorems will likewise be useful in proving problems hard for higher levels of the polynomial hierarchy. Though there are not as many natural problems complete for higher levels of the polynomial hierarchy as for NP, there are in fact quite a few. See the survey by M. Schaefer and Umans [SU02].

The rest of this paper is organized as follows. In Section 2 we give the formal definitions of constraints, constraint applications, complexity classes, and the various constraint problems that we are interested in, and we will formally state Schaefer’s dichotomy theorem and the dichotomy theorem for quantified sets of constraint applications. In Section 3 we will prove the dichotomy theorems for alternation-bounded quantified constraint problems.

2 Preliminaries

2.1 Constraints

We will use the terminology and notation from [CKS01].¹

- Definition 1.**
1. A constraint C is a Boolean function from $\{0, 1\}^k$ to $\{0, 1\}$, where $k > 0$. k is the arity of C .
 2. If C is a constraint of arity k , and z_1, z_2, \dots, z_k are (not necessarily distinct) variables, then $C(z_1, z_2, \dots, z_k)$ is a constraint application of C .
 3. If C is a constraint of arity k , and for $1 \leq i \leq k$, z_i is a variable or a constant (0 or 1), then $C(z_1, z_2, \dots, z_k)$ is a constraint application of C with constants.

When we want to be explicit about the variables occurring in a set of constraint applications S , we will write $S(x_1, \dots, x_n)$, to denote that the variables of S are in $\{x_1, \dots, x_n\}$. If we also want to be explicit about constants, we will write $S(x_1, \dots, x_n, 0, 1)$.

Schaefer’s generalized satisfiability problems can now be defined formally, using constraint terminology.

Definition 2. Let \mathcal{C} be a finite set of constraints.

1. $\text{SAT}(\mathcal{C})$ is the problem of deciding whether a given set S of constraint applications of \mathcal{C} is satisfiable, i.e., whether there exists an assignment to the variables of S that satisfies every constraint application in S .

¹ It should be noted that not all papers use this notation. Many use the Schaefer notation instead. We have chosen to follow the [CKS01] notation, because we will use some of their constructions.

2. $\text{SAT}_c(\mathcal{C})$ is the problem of deciding whether given set S of constraint applications of \mathcal{C} with constants is satisfiable.

As mentioned in the introduction, Schaefer proved that all these problems are either in P or NP-complete. It is also easy to determine which of these two cases hold. This depends on simple properties of the constraints.

Definition 3. *Let C be a constraint.*

- C is 0-valid if $C(0, \dots, 0) = 1$.
- C is 1-valid if $C(1, \dots, 1) = 1$.
- C is Horn (or weakly negative) if C is equivalent to a CNF formula where each clause has at most one positive variable.
- C is anti-Horn (or weakly positive) if C is equivalent to a CNF formula where each clause has at most one negative variable.
- C is bijunctive if C is equivalent to a 2CNF formula.
- C is affine if C is equivalent to a XOR-CNF formula.
- C is complementive (or C-closed) if for every $s \in \{0, 1\}^k$, $C(s) = C(\bar{s})$, where k is the arity of C and $\bar{s} = (1-s_1)(1-s_2) \cdots (1-s_k)$ for $s = s_1 s_2 \cdots s_k$.

Let \mathcal{C} be a finite set of constraints. We say \mathcal{C} is 0-valid, 1-valid, Horn, anti-Horn, bijunctive, affine, or complementive if every constraint $C \in \mathcal{C}$ is 0-valid, 1-valid, Horn, anti-Horn, bijunctive, affine, or complementive, respectively.

Schaefer's theorem can now be stated as follows.

Theorem 4 (Schaefer [Sch78]). *Let \mathcal{C} be a finite set of constraints.*

1. If \mathcal{C} is 0-valid, 1-valid, Horn, anti-Horn, affine, or bijunctive, then $\text{SAT}(\mathcal{C})$ is in P; otherwise, $\text{SAT}(\mathcal{C})$ is NP-complete.
2. If \mathcal{C} is Horn, anti-Horn, affine, or bijunctive, then $\text{SAT}_c(\mathcal{C})$ is in P; otherwise, $\text{SAT}_c(\mathcal{C})$ is NP-complete.

2.2 Quantified constraint applications

QBF is the problem of deciding whether a given fully quantified Boolean formula is true. QBF is PSPACE-complete [SM73]. This problem remains PSPACE-complete if we restrict the Boolean formula to be in 3CNF [Sto77]. We use the following definition for quantified sets of constraint applications.

Definition 5 ([CKS01]). *Let \mathcal{C} be a finite set of constraints. A quantified \mathcal{C} expression [with constants] is an expression of the form $Q_1 x_1 Q_2 x_2 \dots Q_n x_n S(x_1, \dots, x_n)$, where S is a set of constraint applications of \mathcal{C} [with constants], and $Q_i \in \{\exists, \forall\}$ for all i .*

We now define the constraint analogs of QBF.

Definition 6 ([CKS01], Definition 3.9).

1. $\text{QSAT}(\mathcal{C})$ is the problem of deciding whether a given quantified \mathcal{C} expression is true.
2. $\text{QSAT}_c(\mathcal{C})$ is the problem of deciding whether a quantified \mathcal{C} expression with constants is true.

$\text{QSAT}(\mathcal{C})$ and $\text{QSAT}_c(\mathcal{C})$ exhibit dichotomy as well. Remarkably, if $\text{SAT}_c(\mathcal{C})$ is in P, then so are $\text{QSAT}(\mathcal{C})$ and $\text{QSAT}_c(\mathcal{C})$. In all other cases, $\text{SAT}_c(\mathcal{C})$ is NP-complete and $\text{QSAT}(\mathcal{C})$ and $\text{QSAT}_c(\mathcal{C})$ are PSPACE-complete.

Theorem 7 ([Sch78, CKS01, Dal97]). *Let \mathcal{C} be a finite set of constraints. If \mathcal{C} is Schaefer, then $\text{QSAT}(\mathcal{C})$ and $\text{QSAT}_c(\mathcal{C})$ are in P; otherwise, $\text{QSAT}(\mathcal{C})$ and $\text{QSAT}_c(\mathcal{C})$ are PSPACE-complete.*

The history behind this theorem is rather interesting. The dichotomy theorem for $\text{QSAT}_c(\mathcal{C})$ was stated without proof by Schaefer [Sch78]. Schaefer mentioned that the proof relies on the result that the set of true quantified 3CNF formulas is PSPACE-complete. Creignou et al. proved Theorem 7 in [CKS01]. The proofs of the PSPACE lower bounds for $\text{QSAT}_c(\mathcal{C})$ are similar to the NP-hardness proofs for $\text{SAT}_c(\mathcal{C})$. It is shown by Creignou et al. [CKS01], and independently by Dalmau [Dal97], that $\text{QSAT}_c(\mathcal{C})$ polynomial-time many-one reduces to $\text{QSAT}(\mathcal{C})$.

2.3 The Polynomial Hierarchy and Constraints

The polynomial-time hierarchy (polynomial hierarchy or PH for short) was defined by Meyer and Stockmeyer [MS72].

Definition 8 ([MS72]).

- $\Sigma_0^P = \Pi_0^P = \text{P}$.
- $\Sigma_{i+1}^P = \text{NP}^{\Sigma_i^P}$
- $\Pi_{i+1}^P = \text{coNP}^{\Sigma_i^P}$

QSAT_i is the set of all true fully quantified boolean formulas with $i - 1$ quantifier alternations, starting with an \exists quantifier. For all $i \geq 1$, QSAT_i is complete for Σ_i^P [SM73]. These problems remain Σ_i^P -complete if we restrict the Boolean formula to be in 3CNF for i odd and to 3DNF for i even [Wra77].

To generalize QSAT_i to arbitrary sets of constraints, it is important to realize that 3CNF formulas correspond to sets of constraint applications, but 3DNF formulas do not. Of course, a 3DNF formula is the negation of a 3CNF formula. For i even, we can view QSAT_i as the set of all *false* fully quantified boolean formulas of the form $\forall X_1 \exists X_2 \cdots \exists X_i \phi(X_1, \dots, X_k)$, where X_1, \dots, X_k are sets of variables. Restricting ϕ to 3CNF in this view of QSAT_i will still be Σ_i^P -complete. We can now generalize QSAT_i to arbitrary sets of constraints.

Definition 9. *Let \mathcal{C} be a finite set of constraints.*

- *For all $i \geq 1$, a $\Sigma_i(\mathcal{C})$ expression [with constants] is an expression of the form $\exists X_1 \forall X_2 \cdots Q_i X_i S(X_1, \dots, X_i)$, where S is a set of constraint applications of \mathcal{C} [with constants]. Here X_1, X_2, \dots are sets of variables.*

- For all $i \geq 1$, a $\Pi_i(\mathcal{C})$ expression [with constants] is an expression of the form $\forall X_1 \exists X_2 \cdots Q_i X_i S(X_1, \dots, X_i)$, where S is a set of constraint applications of \mathcal{C} [with constants].

Definition 10. Let \mathcal{C} be a set of constraints. Let $i \geq 1$.

1. For i odd, $\text{QSAT}_i(\mathcal{C})$ is the problem of deciding whether a given $\Sigma_i(\mathcal{C})$ expression is true, and $\text{QSAT}_{i,c}(\mathcal{C})$ is the problem of deciding whether a given $\Sigma_i(\mathcal{C})$ expression with constants is true.
2. For i even, $\text{QSAT}_i(\mathcal{C})$ is the problem of deciding whether a given $\Pi_i(\mathcal{C})$ expression is false, and $\text{QSAT}_{i,c}(\mathcal{C})$ is the problem of deciding whether a given $\Pi_i(\mathcal{C})$ expression with constants is false.

3 Dichotomy in the Polynomial Hierarchy

The main proof technique for lower bounds on constraint problems is to show that the problems can simulate an already-known-to-be hard problem. The freedom allowed in the simulations depends on the type of problem considered. For example, for satisfiability problems, we are allowed to introduce existentially quantified auxiliary variables. In [CKS01] terminology, this is known as a “perfect implementation.”

Definition 11 ([CKS01]).

1. A set of constraint applications $S(X, Y)$ perfectly implements constraint C iff $C(X) \equiv \exists Y S(X, Y)$.
2. A set of constraints \mathcal{D} perfectly implements constraint C iff there exists a set of constraint applications of \mathcal{D} that perfectly implements C .

Perfect implementations work well for satisfiability problems.

Lemma 12 ([CKS01], 5.12, 5.16).

1. If $\text{SAT}(\mathcal{C})$ is NP-hard and every constraint in \mathcal{C} can be perfectly implemented by \mathcal{D} , then $\text{SAT}(\mathcal{D})$ is also NP-hard.
2. If $\text{QSAT}(\mathcal{C})$ is PSPACE-hard and every constraint in \mathcal{C} can be perfectly implemented by \mathcal{D} , then $\text{QSAT}(\mathcal{D})$ is also PSPACE-hard.

It is easy to see that the same construction works for PH as well.

Lemma 13. For all $i \geq 2$, if $\text{QSAT}_i(\mathcal{C})$ is Σ_i^p -hard and every constraint in \mathcal{C} can be perfectly implemented by \mathcal{D} , then $\text{QSAT}_i(\mathcal{D})$ is also Σ_i^p -hard.

Proof. Much like the corresponding proof of Lemma 12 for $\text{QSAT}(\mathcal{C})$. Let $Q_1 X_1 Q_2 X_2 \cdots \exists X_i S(X_1, \dots, X_i)$ be a $\Sigma_i(\mathcal{C})$ expression if i is odd and a $\Pi_i(\mathcal{C})$ expression if i is even. For every constraint application $A(Y) \in S$, replace $A(Y)$ by a set of constraint applications $U(Y, Z)$ of \mathcal{D} such that $A(Y) \equiv \exists Z U(Y, Z)$. Make sure that the Z is a set of new variables, and that all introduced sets of new variables are disjoint. Let \hat{S} be the resulting set of constraint applications and let \hat{Z} be the set of all new variables. Then $Q_1 X_1 Q_2 X_2 \cdots \exists X_i S(X_1, \dots, X_i)$ is true iff $Q_1 X_1 Q_2 X_2 \cdots \exists X_i \exists \hat{Z} \hat{S}(X_1, \dots, X_i, \hat{Z})$ is true.

The dichotomy theorem for the case with constants now follows much in the same way as in the case for general quantified expressions.

Theorem 14. *Let \mathcal{C} be a finite set of constraints and let $i \geq 2$. If \mathcal{C} is Horn, anti-Horn, affine, or bijunctive, then $\text{QSAT}_{i,c}(\mathcal{C})$ is in P; otherwise, $\text{QSAT}_{i,c}(\mathcal{C})$ is Σ_i^P -complete.*

Proof. The polynomial-time cases follow immediately from the fact that if \mathcal{C} is Horn, anti-Horn, affine, or bijunctive, then even $\text{QSAT}_c(\mathcal{C})$ is in P (Theorem 7). It is also immediate that $\text{QSAT}_{i,c}(\mathcal{C})$ is in Σ_i^P .

It remains to show the Σ_i^P lower bounds. We closely follow the proof that $\text{QSAT}(\mathcal{C})$ is PSPACE-hard from [CKS01, Theorem 6.12].

Recall from Section 2.3 that the 3CNF version of QSAT_i is complete for Σ_i^P . In constraint terminology, $\text{QSAT}_i(\mathcal{D})$ is complete for Σ_i^P , where $\mathcal{D} = \{\lambda xyz.x \vee y \vee z, \lambda xyz.x \vee y \vee \bar{z}, \lambda xyz.x \vee \bar{y} \vee \bar{z}, \lambda xyz.\bar{x} \vee \bar{y} \vee \bar{z}\}$.

In addition, the constraint One-in-Three (which is defined as the ternary Boolean function that is true if and only if exactly one of its three arguments is true) can perfectly implement any ternary function [CKS01]. Using Lemma 13, it follows that $\text{QSAT}_i(\{\text{One-in-Three}\})$ is Σ_i^P -hard. If \mathcal{C} is not Horn, not anti-Horn, not affine, and not bijunctive, then $\mathcal{C} \cup \{\lambda x.\bar{x}, \lambda x.x\}$ perfectly implements One-in-Three [CKS01]. It follows from Lemma 13 that $\text{QSAT}_i(\mathcal{C} \cup \{\lambda x.\bar{x}, \lambda x.x\})$ is Σ_i^P -hard. This implies that $\text{QSAT}_{i,c}(\mathcal{C})$ is Σ_i^P -hard: Let $Q_1 X_1 Q_2 X_2 \cdots \exists X_i S(X_1, \dots, X_i)$ be a quantified $\mathcal{C} \cup \{\lambda x.\bar{x}, \lambda x.x\}$ expression. If there exists a variable x such that both x and \bar{x} are in S , then $S \equiv 0$. In that case, replace all of S by 0. Otherwise, for every variable x such that $x \in S$ and $\bar{x} \notin S$, replace this variable by 1, and remove x from S . For every variable x such that $\bar{x} \in S$ and $x \notin S$, replace x by 0 and remove \bar{x} from S . Call the resulting set of constraint applications \hat{S} . Then \hat{S} is a set of constraint applications of \mathcal{C} with constants, and $Q_1 X_1 Q_2 X_2 \cdots \exists X_i S(X_1, \dots, X_i)$ is true iff $Q_1 X_1 Q_2 X_2 \cdots \exists X_i \hat{S}(X_1, \dots, X_i)$ is true.

Far more effort is needed to prove the lower bounds for the case without constants. Indeed, the remainder of this paper is dedicated to establishing this result.

Theorem 15. *Let \mathcal{C} be a finite set of constraints and let $i \geq 2$. If \mathcal{C} is Horn, anti-Horn, affine, or bijunctive, then $\text{QSAT}_i(\mathcal{C})$ is in P; otherwise, $\text{QSAT}_i(\mathcal{C})$ is Σ_i^P -complete.*

Proof. The upper bounds follow from Theorem 14. For the remainder of this proof, suppose that \mathcal{C} is not Horn, not anti-Horn, not affine, and not bijunctive. We need to show that $\text{QSAT}_i(\mathcal{C})$ is Σ_i^P -hard. Without loss of generality, we assume that no constraint in \mathcal{C} is a constant function. (Since such constraints are bijunctive, we can simply remove them.)

We will prove that $\text{QSAT}_i(\mathcal{C})$ is Σ_i^P -hard by a case distinction that depends on whether or not \mathcal{C} is 0-valid, 1-valid, and/or complementive. In all cases, we will reduce $\text{QSAT}_{i,c}(\mathcal{C})$ to $\text{QSAT}_i(\mathcal{C})$.

\mathcal{C} is 0-valid and not complementive In this case, \mathcal{C} perfectly implements the constraint $\lambda xy.\bar{x} \vee y$ [CKS01, Lemma 5.41].

As a starting point, we will first review the reduction from $\text{QSAT}_c(\mathcal{C})$ to $\text{QSAT}(\mathcal{C})$ from [CKS01, Theorem 6.12] for the case that \mathcal{C} is 0-valid and not complementive. The main observation needed for this reduction is that $\forall y\{\bar{f} \vee y, \bar{y} \vee t\}$ is equivalent to $\bar{f} \wedge t$.

Let $Q_1x_1 \cdots Q_nx_n S(x_1, \dots, x_n, 0, 1)$ be a quantified \mathcal{C} expression with constants. Using the observation above, it is easy to see that this expression is equivalent to the quantified $\mathcal{C} \cup \{\lambda xy.\bar{x} \vee y\}$ expression $\exists f \exists t \forall y Q_1x_1 \cdots Q_nx_n [S(x_1, \dots, x_n, f, t) \cup \{\bar{f} \vee y, \bar{y} \vee t\}]$.

Thus, $\text{QSAT}(\mathcal{C} \cup \{\lambda xy.\bar{x} \vee y\})$ is PSPACE-hard. Since \mathcal{C} perfectly implements $\lambda xy.\bar{x} \vee y$, it follows by Lemma 12 that $\text{QSAT}(\mathcal{C})$ is PSPACE-hard.

Note that this construction does *not* prove that $\text{QSAT}_i(\mathcal{C})$ is Σ_i^p -complete, since the construction turns a $\Sigma_i(\mathcal{C})$ expression with constants into a $\Sigma_{i+2}(\mathcal{C})$ expression for i odd, and a $\Pi_i(\mathcal{C})$ expression with constants into a $\Sigma_{i+1}(\mathcal{C})$ expression for i even.

However, it is easy to see that we can place $\exists f \exists t \forall y$ anywhere in the quantifier string, as long as $\exists f$ and $\exists t$ precede $\forall y$. This implies that, as long as the original expression contains existential quantifiers followed by universal quantifiers, we obtain the required reduction. Formally, for $i > 2$, we reduce $\text{QSAT}_{i,c}(\mathcal{C})$ to $\text{QSAT}_i(\mathcal{C})$, by mapping $Q_1X_1 \cdots \exists X_{i-2} \forall X_{i-1} \exists X_i S(X_1, \dots, X_i, 0, 1)$ to $Q_1X_1 \cdots \exists X_{i-2} \exists f \exists t \forall y \forall X_{i-1} \exists X_i [S(X_1, \dots, X_i, f, t) \cup \{\bar{f} \vee y, \bar{y} \vee t\}]$.

Since \mathcal{C} is 0-valid, we know from Theorem 4 that, under the assumption that $P \neq NP$, $\text{QSAT}_{i,c}(\mathcal{C})$ is not reducible to $\text{QSAT}_i(\mathcal{C})$ for $i = 1$. It remains to handle the case that $i = 2$.

Let $\forall X_1 \exists X_2 S(X_1, X_2, 0, 1)$ be a $\Pi_2(\mathcal{C})$ expression with constants. We claim that this expression is equivalent to the following $\Pi_2(\mathcal{C} \cup \{\lambda xy.\bar{x} \vee y\})$ expression:

$$\forall X_1 \forall y \forall z \exists f \exists t \exists X_2 [S(X_1, X_2, f, t) \cup \{\bar{f} \vee y, \bar{z} \vee t\}].$$

For the proof, note that

$$\begin{aligned} & \forall X_1 \forall y \forall z \exists f \exists t \exists X_2 [S(X_1, X_2, f, t) \cup \{\bar{f} \vee y, \bar{z} \vee t\}] \\ & \quad \text{iff} \\ & \quad \forall X_1 \exists f \exists t \exists X_2 [S(X_1, X_2, f, t) \cup \{\bar{f} \vee 0, 1 \vee t\}], \\ & \quad \forall X_1 \exists f \exists t \exists X_2 [S(X_1, X_2, f, t) \cup \{\bar{f} \vee 0, 0 \vee t\}], \\ & \quad \forall X_1 \exists f \exists t \exists X_2 [S(X_1, X_2, f, t) \cup \{\bar{f} \vee 1, 1 \vee t\}], \text{ and} \\ & \quad \forall X_1 \exists f \exists t \exists X_2 [S(X_1, X_2, f, t) \cup \{\bar{f} \vee 1, 0 \vee t\}] \\ & \quad \text{iff} \\ & \quad \forall X_1 \exists f \exists t \exists X_2 [S(X_1, X_2, f, t) \cup \{\bar{f}, t\}] \\ & \quad \text{iff} \\ & \quad \forall X_1 \exists X_2 S(X_1, X_2, 0, 1) \end{aligned}$$

Note that this construction can be generalized to all $i \geq 2$, by mapping $Q_1 X_1 \cdots \exists X_{i-2} \forall X_{i-1} \exists X_i S(X_1, \dots, X_i, 0, 1)$ to $Q_1 X_1 \cdots \forall X_{i-1} \forall y \forall z \exists f \exists t \exists X_i [S(X_1, \dots, X_i, f, t) \cup \{\bar{f} \vee y, \bar{z} \vee t\}]$.

\mathcal{C} is 1-valid and not complementive In this case, we could simply state that the proof is similar to the proof of the case that \mathcal{C} is 0-valid and not complementive. But rather than making the reader work through the previous case to see that this is actually true, we will prove a theorem (Theorem 17) which relates satisfiability problems for sets of constraint applications of \mathcal{C} [with constants] to the satisfiability problems where the set of constraints is replaced by a type of “complement.” This theorem immediately implies the current case and will also be useful in the case that \mathcal{C} is complementive. We start with some definitions.

Definition 16. 1. Let C be a k -ary constraint. Define constraint C^c as follows. For all $s \in \{0, 1\}^k$, $C^c(s) = C(\bar{s})$, where, as in the definition of complementive, $\bar{s} = (1 - s_1)(1 - s_2) \cdots (1 - s_k)$ for $s = s_1 s_2 \cdots s_k$. Note that C is complementive iff $C = C^c$.

2. Let \mathcal{C} be a finite set of constraints. Define the set of constraints \mathcal{C}^c as $\mathcal{C}^c = \{C^c \mid C \in \mathcal{C}\}$.

3. For S a set of constraint applications of \mathcal{C} with constants, define S^c as $\{C^c(z_1, \dots, z_k) \mid C(z_1, \dots, z_k) \in S\}$, where each z_i is a variable or a constant.

Theorem 17. 1. For all $i \geq 1$, $\text{QSAT}_i(\mathcal{C}) \equiv_m^p \text{QSAT}_i(\mathcal{C}^c)$ and $\text{QSAT}_{i,c}(\mathcal{C}) \equiv_m^p \text{QSAT}_{i,c}(\mathcal{C}^c)$.

2. $\text{QSAT}(\mathcal{C}) \equiv_m^p \text{QSAT}(\mathcal{C}^c)$ and $\text{QSAT}_c(\mathcal{C}) \equiv_m^p \text{QSAT}_c(\mathcal{C}^c)$.

This theorem follows immediately from the following lemma.

Lemma 18. Let \mathcal{C} be a finite set of constraints and let $Q_1 x_1 \cdots Q_n x_n S(x_1, \dots, x_n, 0, 1)$ be a quantified \mathcal{C} expression with constants. Then $Q_1 x_1 \cdots Q_n x_n S(x_1, \dots, x_n, 0, 1)$ is true if and only if $Q_1 x_1 \cdots Q_n x_n S^c(x_1, \dots, x_n, 1, 0)$ is true.

Proof. The proof is by induction on n , the number of variables in S . For $n = 0$, by definition of S^c , $S(0, 1) = S^c(1, 0)$. Now let $n > 0$, and suppose the claim holds for $n - 1$.

If $Q_1 = \forall$, then $Q_1 x_1 \cdots Q_n x_n S(x_1, \dots, x_n, 0, 1)$ is true if and only if both $Q_2 x_2 \cdots Q_n x_n S(0, x_2, \dots, x_n, 0, 1)$ and $Q_2 x_2 \cdots Q_n x_n S(1, x_2, \dots, x_n, 0, 1)$ are true. By induction, this is the case if and only if both $Q_2 x_2 \cdots Q_n x_n S^c(1, x_2, \dots, x_n, 1, 0)$ and $Q_2 x_2 \cdots Q_n x_n S^c(0, x_2, \dots, x_n, 1, 0)$ are true, which holds if and only if $\forall x_1 \cdots Q_n x_n S^c(x_1, \dots, x_n, 1, 0)$ is true. The proof for $Q_1 = \exists$ is similar.

\mathcal{C} is 0-valid and complementive If \mathcal{C} is complementive, $C^c = C$ for all $C \in \mathcal{C}$. The following corollary follows immediately from Lemma 18.

Corollary 19. *Let \mathcal{C} be complementive. Let $S(x_1, \dots, x_n, 0, 1)$ be a set of constraint applications of \mathcal{C} with constants. Then $Q_1x_1Q_2x_2 \cdots Q_nx_nS(x_1, \dots, x_n, 0, 1)$ iff $Q_1x_1Q_2x_2 \cdots Q_nx_nS(x_1, \dots, x_n, 1, 0)$.*

From [CKS01, Lemma 5.41], we know that \mathcal{C} perfectly implements the constraint SymOR_1 , which is defined as the constraint $\lambda xyz.(\bar{x} \wedge (\bar{y} \vee z)) \vee (x \wedge (\bar{z} \vee y))$.

Using Corollary 19, we can now construct a reduction that is similar to the one described at the end of the case that \mathcal{C} is 0-valid and not complementive to reduce $\text{QSAT}_{i,c}(\mathcal{C})$ to $\text{QSAT}_i(\mathcal{C})$.

Let $Q_1X_1 \cdots \exists X_i S(X_1, \dots, X_i, 0, 1)$ be a $\Sigma_i(\mathcal{C})$ expression with constants if i is odd, and a $\Pi_i(\mathcal{C})$ expression with constants if i is even. We claim that this expression is true if and only if the following expression is true:

$$Q_1X_1 \cdots \forall X_{i-1} \forall x \forall y \forall z \exists f \exists t \exists X_i \\ [S(X_1, \dots, X_i, f, t) \cup \{\text{SymOR}_1(x, f, y), \text{SymOR}_1(x, z, t)\}].$$

For the proof, note that

$$Q_1X_1 \cdots \forall X_{i-1} \forall x \forall y \forall z \exists f \exists t \exists X_i \\ [S(X_1, \dots, X_i, f, t) \cup \{\text{SymOR}_1(x, f, y), \text{SymOR}_1(x, z, t)\}]$$

if and only if $Q_1X_1 \cdots \forall X_{i-1} \forall y \forall z \exists f \exists t \exists X_i [S(X_1, \dots, X_i, f, t) \cup \{\bar{f} \vee y, \bar{z} \vee t\}]$ and $Q_1X_1 \cdots \forall X_{i-1} \forall y \forall z \exists f \exists t \exists X_i [S(X_1, \dots, X_i, f, t) \cup \{\bar{y} \vee f, \bar{t} \vee z\}]$.

As in the 0-valid and not complementive case, this holds if and only if

$$Q_1X_1 \cdots \forall X_{i-1} \exists f \exists t \exists X_i [S(X_1, \dots, X_i, f, t) \cup \{\bar{f}, t\}] \text{ and} \\ Q_1X_1 \cdots \forall X_{i-1} \exists f \exists t \exists X_i [S(X_1, \dots, X_i, f, t) \cup \{\bar{t}, f\}] \\ \text{iff} \\ Q_1X_1 \cdots \forall X_{i-1} \exists X_i S(X_1, \dots, X_i, 0, 1) \text{ and} \\ Q_1X_1 \cdots \forall X_{i-1} \exists X_i S(X_1, \dots, X_i, 1, 0) \\ \text{iff (by complementivity)} \\ Q_1X_1 \cdots \forall X_{i-1} \exists X_i S(X_1, \dots, X_i, 0, 1)$$

\mathcal{C} is not 0-valid, not 1-valid, and complementive In this case, \mathcal{C} can perfectly implement $\lambda xy.x \oplus y$ [CKS01, proof of Lemma 5.24].

Using Corollary 19, it suffices to replace 0 by f , 1 by t and to add $\exists f \exists t \{f \oplus t\}$. However, this existential quantification has to be added at the start of the expression. For example, consider the (false) expression $\forall x \{x = 0\}$. Adding the existential quantification at the end of the quantifier string will give $\forall x \exists f \exists t \{x = f, f \oplus t\}$, which is true. (It doesn't matter whether t is set to 0 and f to 1 or vice-versa (by complementivity). However, we need to always look at the same assignment.) Adding the existential quantifiers at the start

of the expression gives a reduction from $\text{QSAT}_{i,c}(\mathcal{C})$ to $\text{QSAT}_i(\mathcal{C})$ for i odd. More formally, if i is odd, we map

$$\exists X_1 \forall X_2 \cdots \exists X_i S(X_1, \dots, X_i, 0, 1)$$

to

$$\exists f \exists t \exists X_1 \forall X_2 \cdots \exists X_i [S(X_1, \dots, X_i, f, t) \cup \{f \oplus t\}].$$

Note that this reduction also works for $i = 1$.

We will now show how to reduce $\text{QSAT}_{i,c}(\mathcal{C})$ to $\text{QSAT}_i(\mathcal{C})$ for i even. Let $S(X_1, \dots, X_i, 0, 1)$ be a set of constraint applications of \mathcal{C} with constants. We map

$$\forall X_1 \exists X_2 \cdots \exists X_i S(X_1, \dots, X_i, 0, 1)$$

to

$$\forall b \forall X_1 \exists X_2 \cdots \exists X_i \exists b' [S(X_1, \dots, X_i, b, b') \cup \{(b \oplus b')\}]$$

That this is indeed a reduction follows immediately from Corollary 19.

\mathcal{C} is not 0-valid, not 1-valid, and not complementive Let $A \in \mathcal{C}$ be not 0-valid, $B \in \mathcal{C}$ be not 1-valid, and $C \in \mathcal{C}$ be not complementive. Recall that we may assume that all constraints in \mathcal{C} are satisfiable. Let s_A be a satisfying assignment for A , let s_B be a satisfying assignment for B , and let s_C be a satisfying assignment for C such that $\overline{s_C}$ is not a satisfying assignment for C . Let $\widehat{A}(x, y)$ be a constraint application of A defined as follows: $\widehat{A}(x, y) = A(z_1, \dots, z_k)$, where $z_i = x$ if $(s_A)_i = 0$, and $z_i = y$ if $(s_A)_i = 1$. Define $\widehat{B}(x, y)$ from B and s_B and $\widehat{C}(x, y)$ from C and s_C in the same way. Then $\widehat{A}(0, 0) = 0$, $\widehat{A}(0, 1) = 1$, $\widehat{B}(0, 1) = 1$, $\widehat{B}(1, 1) = 0$, $\widehat{C}(0, 1) = 1$, and $\widehat{C}(1, 0) = 0$. Now consider the set of constraint applications $\{\widehat{A}(f, t), \widehat{B}(f, t), \widehat{C}(f, t)\}$. It is easy to see that this set perfectly implements $\overline{f} \wedge t$.

For our reductions, we need to replace 0 by f , 1 by t , and add $\exists f \exists t \{\overline{f} \wedge t\}$. Note that, unlike the previous case, we can add the existential quantifiers anywhere in the quantifier string, since $\exists f \exists t \{\overline{f} \wedge t\}$ completely fixes the truth assignment to f and t .

More formally, to reduce $\text{QSAT}_{i,c}(\mathcal{C})$ to $\text{QSAT}_i(\mathcal{C})$, map $Q_1 X_1 Q_2 X_2 \cdots \forall Q_{i-1} \exists X_i S(X_1, \dots, X_i, 0, 1)$ to $Q_1 X_1 Q_2 X_2 \cdots \forall Q_{i-1} \exists X_i \exists f \exists t [S(X_1, \dots, X_i, f, t) \cup \{\overline{f} \wedge t\}]$. This shows that $\text{QSAT}_i^p(\mathcal{C})$ is Σ_i^p -hard.

As in the previous case, this reduction will work for $i = 1$ as well. Thus, our proofs also imply Schaefer's dichotomy theorem.

References

- [BHRV02] E. Böhrer, E. Hemaspaandra, S. Reith, and H. Vollmer. Equivalence and isomorphism for Boolean constraint satisfaction. In *Proceedings of the 16th Annual Conference of the EACSL (CSL 2002)*, pages 412–426. Springer-Verlag *Lecture Notes in Computer Science* #2471, September 2002.

- [BHRV04] E. Böhrer, E. Hemaspaandra, S. Reith, and H. Vollmer. The complexity of Boolean constraint isomorphism. In *Proceedings of the 21st Symposium on Theoretical Aspects of Computer Science*. Springer-Verlag *Lecture Notes in Computer Science*, 2004. to appear.
- [CH96] N. Creignou and M. Hermann. Complexity of generalized satisfiability counting problems. *Information and Computation*, 125:1–12, 1996.
- [CKS01] N. Creignou, S. Khanna, and M. Sudan. *Complexity Classifications of Boolean Constraint Satisfaction Problems*. Monographs on Discrete Applied Mathematics. SIAM, 2001.
- [Cre95] N. Creignou. A dichotomy theorem for maximum generalized satisfiability problems. *Journal of Computer and System Sciences*, 51:511–522, 1995.
- [Dal97] V. Dalmau. Some dichotomy theorems on constant free Boolean formulas. Technical Report TR-LSI-97-43-R, Universitat Politècnica de Catalunya, 1997.
- [FV98] T. Feder and M. Vardi. Monadic SNP and constraint satisfaction: A study through datalog and group theory. *SIAM Journal on Computing*, 28(1):57–104, 1998.
- [Jub99] L. Juban. Dichotomy theorem for generalized unique satisfiability problem. In *Proceedings of the 12th Conference on Fundamentals of Computation Theory*, pages 327–337. Springer-Verlag *Lecture Notes in Computer Science #1684*, 1999.
- [KK01] L. Kirousis and P. Kolaitis. A dichotomy in the complexity of propositional circumscription. In *Proceedings of the 16th Symposium on Logic in Computer Science*, pages 71–80, 2001.
- [KK03] L. Kirousis and P. Kolaitis. The complexity of minimal satisfiability problems. *Information and Computation*, 187(1):20–39, 2003.
- [KS98] D. Kavvadias and M. Sideri. The inverse satisfiability problem. *SIAM Journal on Computing*, 28(1):152–163, 1998.
- [KSTW01] S. Khanna, M. Sudan, L. Trevisan, and D. Williamson. The approximability of constraint satisfaction problems. *SIAM Journal on Computing*, 30(6):1863–1920, 2001.
- [MS72] A. Meyer and L. Stockmeyer. The equivalence problem for regular expressions with squaring requires exponential space. In *Proceedings of the 13th IEEE Symposium on Switching and Automata Theory*, pages 125–129, 1972.
- [Sch78] T. Schaefer. The complexity of satisfiability problems. In *Proceedings of the 10th ACM Symposium on Theory of Computing*, pages 216–226, 1978.
- [SM73] L. Stockmeyer and A. Meyer. Word problems requiring exponential time: preliminary report. In *Proceedings of the 5th ACM Symposium on Theory of Computing*, pages 1–9, 1973.
- [Sto77] L. Stockmeyer. The polynomial-time hierarchy. *Theoretical Computer Science*, 3:1–22, 1977.
- [SU02] M. Schaefer and C. Umans. Completeness in the polynomial-time hierarchy. *SIGACT News*, 2002.
- [Wra77] C. Wrathall. Complete sets and the polynomial-time hierarchy. *Theoretical Computer Science*, 3:23–33, 1977.