arXiv:cs/0406010v1 [cs.DM] 6 Jun 2004

ANOTHER PROOF OF AN EXTENSION OF A CURIOUS IDENTITY

Yidong Sun
Department of Applied Mathematics, Dalian University of Technology, Dalian 116024, P. R. China
sydcom@eyou.com

Abstract

Based on Jensen formulae and the second kind of Chebyshev polynomials, another proof is presented for an extension of a curious binomial identity due to Z. W. Sun and K. J. Wu.

1. Introduction

Recently, Z. W. Sun and K. J. Wu [8] have presented an extension of a curious binomial identity [7], that is the following generalized version:

$$\begin{aligned}(x+(m+1)z)\sum_{k=0}^{m}(-1)^k\binom{x+y+kz}{m-k}\binom{y+k+kz}{k}\\ = \quad z\sum_{0\le i\le k\le m}(-1)^k\binom{k}{i}\binom{x+i}{m-k}(1+z)^{k+i}(1-z)^{k-i}+(x-m)\binom{x}{m}. \qquad (1)\end{aligned}$$

In 2002 Z. W. Sun [7] proved the case when $z=1$ using double recursion. Later four alternative proofs have been provided for the special case. A generating function proof was given by A. Panholzer and H. Prodinger [6]; D. Merlini and R. Sprugnoli [5] established it through Riordan arrays; S. B. Ekhad and M. Mohammed [4] proved it based on a WZ method. Later, W. Chu and L. V. D. Claudio [3] reobtained the identity by using Jensen formulae [2]. More recently, D. Callan [1] gave a combinatorial proof using weight-reversing involutions on suitable configurations involving dominos and colorings.

Motivated by the method [3], we give a simple proof of the identity (1).

2. Proof of (1)

Let

$$\begin{aligned}f(x,y,z) &= \sum_{k=0}^{m}(-1)^k\binom{x+y+kz}{m-k}\binom{y+k+kz}{k} \quad \text{and}\\ g(x,z) &= \sum_{0\le i\le k\le m}(-1)^k\binom{k}{i}\binom{x+i}{m-k}(1+z)^{k+i}(1-z)^{k-i}.\end{aligned}$$

Recall the Jensen formulae on binomial convolutions

$$\sum_{i=0}^{m}\binom{a+bi}{i}\binom{c-bi}{m-i}=\sum_{j=0}^{m}\binom{a+c-j}{m-j}b^j,$$

and the second kind of Chebyshev polynomials

$$U_n(t) \;=\; \frac{sin(n+1)\theta}{sin(\theta)}=\sum_{k=0}^{[n/2]}(-1)^k\binom{n-k}{k}(2t)^{n-2k}, \quad \text{by setting } t=cos(\theta).$$

Then we have

$$\begin{aligned}
f(x,y,z) &= \sum_{k=0}^{m}(-1)^k\binom{x+y+kz}{m-k}\binom{y+k+kz}{k}\\
&= (-1)^m\sum_{k=0}^{m}\binom{y+k(z+1)}{k}\binom{-1-x-y+m-k(z+1)}{m-k}\\
&= (-1)^m\sum_{j=0}^{m}\binom{-1-x+m-j}{m-j}(1+z)^j\\
&= \sum_{j=0}^{m}\binom{x}{m-j}(-1-z)^j.
\end{aligned}$$

Now we can express $g(x,z)$ as a simple form

$$\begin{aligned}
g(x,z) &= \sum_{k=0}^{m}\sum_{i=0}^{k}(-1)^k\binom{k}{i}\binom{x+i}{m-k}(1+z)^{k+i}(1-z)^{k-i}\\
&= \sum_{k=0}^{m}\sum_{i=0}^{k}(-1)^k(1+z)^{k+i}(1-z)^{k-i}\sum_{j=k}^{m}\binom{x}{m-j}\binom{i}{j-k}\binom{k}{i}\\
&= \sum_{j=0}^{m}\binom{x}{m-j}\sum_{k=0}^{j}\sum_{i=0}^{k}(-1)^k(1+z)^{k+i}(1-z)^{k-i}\binom{i}{j-k}\binom{k}{i}\\
&= \sum_{j=0}^{m}\binom{x}{m-j}\sum_{k=0}^{j}\sum_{i=0}^{k}(-1)^k(1+z)^{k+i}(1-z)^{k-i}\binom{k}{j-k}\binom{2k-j}{k+i-j}\\
&= \sum_{j=0}^{m}\binom{x}{m-j}(1+z)^j\sum_{k=0}^{j}(-1)^k\binom{k}{j-k}\sum_{i=0}^{k}\binom{2k-j}{k+i-j}(1+z)^{k+i-j}(1-z)^{k-i}\\
&= \sum_{j=0}^{m}\binom{x}{m-j}(1+z)^j\sum_{k=0}^{j}(-1)^k\binom{k}{j-k}2^{2k-j}\\
&= \sum_{j=0}^{m}\binom{x}{m-j}(-1-z)^jU_j(t)|_{t\to 1}\\
&= \sum_{j=0}^{m}(j+1)\binom{x}{m-j}(-1-z)^j.
\end{aligned}$$

Hence we obtain

$$\begin{aligned}
&(x+(m+1)z)f(x,y,z)-zg(x,z)\\
&= \sum_{j=0}^{m}(x+(m+1)z-(j+1)z)\binom{x}{m-j}(-1-z)^j\\
&= \sum_{j=0}^{m}(x-m+j)\binom{x}{m-j}(-1-z)^j-(m-j)\binom{x}{m-j}(-1-z)^{j+1}\\
&= \sum_{j=0}^{m}(1+m-j)\binom{x}{1+m-j}(-1-z)^j-(m-j)\binom{x}{m-j}(-1-z)^{j+1}\\
&= (1+m)\binom{x}{1+m}=(x-m)\binom{x}{m},
\end{aligned}$$

as desired (the second to last sum is telescoping).

References

[1] D. Callan, A combinatorial proof of Sun's "curious" identity, Integer: Electron. J. Combin. Number Theory 4 (2004), A05, 6pp.

[2] W. Chu, On an extension of a partition identity and its Abel analog, J. Math. Research & Exposition 6:4 (1986), 37-39, MR89b: 05021.

[3] W. Chu and L. V. D. Claudio, Jesen proof of a curious binomial identity, Integer: Electron. J. Combin. Number Theory 3 (2003), A20, 3pp.

[4] S. B. Ekhad and M. Mohammed, A WZ proof of a curious identity, Integer: Electron. J. Combin. Number Theory 3 (2003), A6, 2pp.

[5] D. Merlini and R. Sprugnoli, A Riordan array proof of a curious identity, Integer: Electron. J. Combin. Number Theory 2 (2002), A8, 3pp.

[6] A. Panholzer and H. Prodinger, A generating functions proof of a curious identity, Integer: Electron. J. Combin. Number Theory 2 (2002), A6, 3pp.

[7] Z. W. Sun, A curious identity involving binomial coefficients, Integer: Electron. J. Combin. Number Theory 2 (2002), A4, 8pp.

[8] Z. W. Sun and K. J. Wu, An extension of a curious binomial identity, arXiv:math.CO/0401057, submitted.