

INTEGRAL REPRESENTATION FOR A CLASS OF C^1 -CONVEX FUNCTIONALS

Gianni DAL MASO
Anneliese DEFRANCESCHI
Enrico VITALI

Abstract

In view of the applications to the asymptotic analysis of a family of obstacle problems, we consider a class of convex local functionals $F(u, A)$, defined for all functions u in a suitable vector valued Sobolev space and for all open sets A in $\mathbf{R}^n$. Sufficient conditions are given in order to obtain an integral representation of the form $F(u, A) = \int_A f(x, u(x)) d\mu + \nu(A)$, where μ and ν are Borel measures and f is convex in the second variable.

Introduction

This paper contains an integral representation theorem for a class of convex local functionals which arise in the study of the asymptotic behaviour of a sequence of minimum problems with obstacles for vector valued Sobolev functions.

Given an open subset Ω of $\mathbf{R}^n$ and $1 < p < +\infty$, let $W^{1,p}(\Omega, \mathbf{R}^m)$ be the usual space of Sobolev functions with values in $\mathbf{R}^m$, and let $\mathcal{A}(\Omega)$ be the family of all open subsets of Ω . The functionals $F: W^{1,p}(\Omega, \mathbf{R}^m) \times \mathcal{A}(\Omega) \rightarrow [0, +\infty]$ we are going to consider are assumed to satisfy the following properties:

- (i) (lower semicontinuity) for every $A \in \mathcal{A}(\Omega)$ the function $F(\cdot, A)$ is lower semicontinuous on $W^{1,p}(\Omega, \mathbf{R}^m)$;
- (ii) (measure property) for every $u \in W^{1,p}(\Omega, \mathbf{R}^m)$ the set function $F(u, \cdot)$ is (the trace of) a Borel measure on Ω ;
- (iii) (locality property) $F(u, A) = F(v, A)$ whenever $u, v \in W^{1,p}(\Omega, \mathbf{R}^m)$, $A \in \mathcal{A}(\Omega)$, and $u|_A = v|_A$;
- (iv) (C^1 -convexity) for every $A \in \mathcal{A}(\Omega)$ the function $F(\cdot, A)$ is convex on $W^{1,p}(\Omega, \mathbf{R}^m)$ and, in addition, $F(\varphi u + (1 - \varphi)v, A) \leq F(u, A) + F(v, A)$ for every $u, v \in W^{1,p}(\Omega, \mathbf{R}^m)$ and for every $\varphi \in C^1(\Omega) \cap W^{1,\infty}(\Omega)$ with $0 \leq \varphi \leq 1$ on Ω .

This set of conditions is motivated by the study of the limit behaviour, as $h \rightarrow \infty$, of a sequence of convex obstacle problems of the form

$$(0.1) \quad \min \left\{ \int_{\Omega} W(x, Du(x)) dx : u \in H_0^1(\Omega, \mathbf{R}^m), u(x) \in K_h(x) \text{ for q.e. } x \in A \right\},$$

where $W(x, \xi)$ is quadratic with respect to ξ , A is an open subset of Ω with $A \subset\subset \Omega$, and $K_h(x)$ is a closed convex subset of $\mathbf{R}^m$ for every $h \in \mathbf{N}$ and for every $x \in \Omega$. By using Γ -convergence techniques it is possible to prove (see [17]) that the limit problem can always be written in the form

$$\min \left\{ \int_{\Omega} W(x, Du(x)) dx + F(u, A) : u \in H_0^1(\Omega, \mathbf{R}^m) \right\},$$

with F satisfying the conditions considered above.

In this paper we are concerned only with the properties of F that can be deduced from (i)–(iv). The main result (Theorem 6.5) is that every functional F satisfying (i)–(iv) can be written in the form

$$(0.2) \quad F(u, A) = \int_A f(x, u(x)) d\mu + \nu(A),$$

where μ and ν are positive Borel measures and $f: \Omega \times \mathbf{R}^m \rightarrow [0, +\infty]$ is a Borel function, convex and lower semicontinuous in the second variable.

This result will be used in a forthcoming paper (see [17]) to provide a detailed description of the structure of the limits of sequences of obstacle problems of the form (0.1) under various assumptions on $W(x, \xi)$ and $K_h(x)$.

Conditions (i)–(iii) are not enough to obtain an integral representation of the form (0.2). Indeed, even convex functionals depending on the gradient, like $F(u, A) = \int_A |Du|^p dx$, satisfy (i)–(iii). Condition (iv) is the most important one, and is responsible for an integral representation of the form (0.2), i.e., without terms depending on the gradient. This notion of convexity, also used, e.g., in [27] and [16], is strictly related to the notion of C^1 -stability introduced by G. Bouchitté and M. Valadier in [7], whose results are frequently used in our paper.

For a general survey on integral representation theorems in L^p , $W^{1,p}$, and BV we refer to [9]. See also [1], [2], [6], [4], [21], [3] for more recent results.

In the scalar case (i.e., $m = 1$), integral representations on $W^{1,p}(\Omega)$ of the form (0.2), connected with limits of obstacle problems, can be found in [18], [15], [5], [12], [16] under suitable convexity conditions, and in [13] under monotonicity assumptions.

Although the final statement in the vector case is exactly the same as in the scalar case, the proof is completely different, since all arguments used in the papers mentioned above rely on the order structure of $\mathbf{R}$, involving truncations and monotonicity methods.

The main tools for the proof in the vector case are some technical results obtained in our previous paper [14], based on the theory of Lipschitz parametrization of convex sets developed in [24] and [26]. In particular we shall use the following result (Theorem 2.9 in [14]): given a finite number of functions $u_1, \dots, u_k$ in $W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$, their convex combinations with smooth coefficients form a dense subset in the set of all $W^{1,p}$ -selections of the polyhedral multivalued function $x \mapsto \text{co}\{u_1(x), \dots, u_k(x)\}$, where co denotes the convex hull.

The first step (Theorem 3.7) of our result deals with the integral representation of the functional F on the set of all $W^{1,p}$ -selections of such polyhedral multifunctions.

In Theorem 5.4 we extend the integral representation of F to all the functions of $W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ which satisfy, up to sets of capacity zero, a suitable “obstacle condition” of the form $u(x) \in K(x)$, which is necessary (but not sufficient) for the finiteness of the functional. We note, incidentally, that the main difficulty in the proof of our result lies in the fact that the functional F is not assumed to be finite everywhere,

in view of the applications to obstacle problems.

The restriction to $L^\infty(\Omega, \mathbf{R}^m)$, originated by the need of taking products of $W^{1,p}$ -functions, is dropped in Section 6. Moreover, the “obstacle condition”, given up to sets of capacity zero, is shown to be equivalent to the condition $u(x) \in K(x)$ almost everywhere with respect to a suitable measure (Proposition 6.3). This allows us to obtain the integral representation (0.2) for every $u \in W^{1,p}(\Omega, \mathbf{R}^m)$ and for every $A \in \mathcal{A}(\Omega)$.

In the last section (Theorem 7.3) we prove that, if F is quadratic or positively p -homogeneous, then so is f .

Acknowledgements

This work is part of the Project EURHomogenization – ERB4002PL910092 of the Program SCIENCE of the Commission of the European Communities, and of the Research Project “Irregular Variational Problems” of the Italian National Research Council. The second and third author were partially supported by a National Research Project of the Italian Ministry of Scientific Research.

1. Notation and preliminaries

Throughout this paper m, n are two fixed positive integers, p is a fixed real number, $1 < p < +\infty$, and Ω is an open subset of $\mathbf{R}^n$, possibly unbounded. We shall denote by $\mathcal{A}(\Omega)$ the family of the open subsets of Ω and by $\mathcal{B}(\Omega)$ the family of its Borel subsets. If $B \subseteq \mathbf{R}^n$ is a Borel set we denote its Lebesgue measure by $|B|$. The notation *a.e.* stands for almost everywhere with respect to the Lebesgue measure.

If d is a positive integer, for every $x \in \mathbf{R}^d$ and $r > 0$ we set $B_r(x) = \{y \in \mathbf{R}^d : |y - x| < r\}$, while $\overline{B}_r(x)$ denotes the closure of $B_r(x)$. The $(d-1)$ -dimensional simplex Σ_d is defined by

$$\Sigma_d = \{\lambda \in \mathbf{R}^d : \lambda^1 + \dots + \lambda^d = 1, \lambda^i \geq 0\},$$

where $\lambda = (\lambda^1, \dots, \lambda^d)$. If C is a convex subset of $\mathbf{R}^d$, we denote by $\text{ri } C$ its relative interior and by ∂C its relative boundary. In particular, $\text{ri } \Sigma_d = \{\lambda \in \mathbf{R}^d : \lambda^1 + \dots + \lambda^d = 1, \lambda^i > 0\}$ and $\partial \Sigma_d = \Sigma_d \setminus \text{ri } \Sigma_d$.

The space $L^p(\Omega, \mathbf{R}^m)$ is endowed with the usual norm

$$\|u\|_{L^p(\Omega, \mathbf{R}^m)} = \left(\int_{\Omega} |u|^p dx \right)^{1/p}.$$

Let $W^{1,p}(\Omega, \mathbf{R}^m)$ be the Banach space of all the functions $u \in L^p(\Omega, \mathbf{R}^m)$ with first order distributional derivative Du in $L^p(\Omega, \mathbf{R}^{mn})$, endowed with the norm

$$\|u\|_{W^{1,p}(\Omega, \mathbf{R}^m)} = (\|u\|_{L^p(\Omega, \mathbf{R}^m)}^p + \|Du\|_{L^p(\Omega, \mathbf{R}^{mn})}^p)^{1/p}.$$

The closure of $C_0^1(\Omega, \mathbf{R}^m)$ in $W^{1,p}(\Omega, \mathbf{R}^m)$ will be denoted by $W_0^{1,p}(\Omega, \mathbf{R}^m)$ ($\mathbf{R}^m$ will be omitted if $m = 1$).

Capacity. For every compact set $K \subseteq \Omega$ we define the capacity of K with respect to Ω by

$$\text{cap}(K, \Omega) = \inf \{ \|\varphi\|_{W^{1,p}(\Omega)}^p : \varphi \in C_0^\infty(\Omega), \varphi \geq 1 \text{ on } K \}.$$

The definition is extended to all subsets of Ω as external capacity in the usual way (see, for example, [11] and [31]).

Let E be a subset of $\mathbf{R}^n$. If a statement depending on $x \in \mathbf{R}^n$ holds for every $x \in E$ except for a set $N \subseteq E$ with capacity zero, then we say that it holds *quasi everywhere* (q.e.) on E .

A function $f: \Omega \rightarrow \mathbf{R}^m$ is said to be *quasi continuous* on Ω if for every $\varepsilon > 0$ there exists a set $E \subseteq \Omega$ with $\text{cap}(E, \Omega) < \varepsilon$ such that the restriction of f to $\Omega \setminus E$ is continuous. A subset A of Ω is said to be *quasi open* if for every $\varepsilon > 0$ there exists an open set A_ε with $\text{cap}(A_\varepsilon, \Omega) < \varepsilon$ such that $A \cup A_\varepsilon$ is an open set.

It is well known (see, for instance, [20]) that for every $u \in W^{1,p}(\Omega, \mathbf{R}^m)$ there exists a quasi continuous representative of u which is unique up to sets of capacity zero, and which is given by

$$\lim_{r \rightarrow 0^+} \frac{1}{|B_r(x)|} \int_{B_r(x)} u(y) dy$$

for q.e. $x \in \Omega$. Throughout this paper we shall use such a quasi continuous representative to individuate an element of $W^{1,p}(\Omega, \mathbf{R}^m)$. Moreover, we may also assume that the quasi continuous representative we are going to choose is Borel measurable.

It turns out that for every subset E of Ω

$$\text{cap}(E, \Omega) = \inf \{ \|u\|_{W^{1,p}(\Omega)}^p : u \in W_0^{1,p}(\Omega), u \geq 1 \text{ q.e. on } E \}.$$

Actually this infimum is attained by a unique function which is called the *capacitary potential* of E . It turns out that this function takes its values in $[0, 1]$.

A positive Borel measure μ on Ω is said to be *absolutely continuous* with respect to capacity if $\mu(B) = 0$ whenever $B \in \mathcal{B}(\Omega)$ and $\text{cap}(B, \Omega) = 0$.

If E is a subset of Ω and $F: E \rightarrow \mathbf{R}^m$ is a multivalued function, i.e., F maps E into the set of all subsets of $\mathbf{R}^m$, then we say that F is *lower semicontinuous at a point* x_0 of E if for every open subset G of $\mathbf{R}^m$ with $G \cap F(x_0) \neq \emptyset$ there exists a neighborhood U of x_0 such that for every $y \in U$ we have $G \cap F(y) \neq \emptyset$. We say that F is *upper semicontinuous in* x_0 if for every neighborhood G of $F(x_0)$ there exists a neighborhood U of x_0 such that $F(y) \subseteq G$ whenever $y \in U$. We say that F is *quasi lower semicontinuous* (resp. *quasi upper semicontinuous*) on Ω if for every $\varepsilon > 0$ there exists a set $E \subseteq \Omega$ with $\text{cap}(E, \Omega) < \varepsilon$ such that the restriction of F to $\Omega \setminus E$ is lower semicontinuous (resp. upper semicontinuous).

Measurability. Let $(X, \mathcal{M})$ be a measurable space. If μ is a positive measure on $(X, \mathcal{M})$ we denote by $\mathcal{M}_\mu$ the standard μ -completion of $\mathcal{M}$ and we still denote by μ the completed measure. If μ is σ -finite the $\mathcal{M}_\mu$ -measurability is equivalent to the μ -measurability in the Carathéodory sense. Moreover, $\widehat{\mathcal{M}}$ will denote the universal completion of $\mathcal{M}$, i.e., the intersection $\cap_\mu \mathcal{M}_\mu$ for all positive finite measures μ ; equivalently, the intersection can be extended to all positive σ -finite measures μ (see [10] Ch.III, parag. 4).

It is easy to verify that every quasi continuous function $u: \Omega \rightarrow \mathbf{R}$ is μ -measurable (i.e., $\mathcal{B}_\mu$ -measurable) for every positive Borel measure μ which is absolutely continuous with respect to capacity.

For convenience, we state here two results which will play an important role to prove the measurability of certain functions. They can be deduced from [10], Theorem III.23 and Theorem III.22, respectively.

Theorem 1.1. (Projection Theorem) *Let $(X, \mathcal{M})$ be a measurable space. If G is an element of $\mathcal{M} \otimes \mathcal{B}(\mathbf{R}^d)$, then the projection $\text{pr}_X(G)$ belongs to $\widehat{\mathcal{M}}$.*

Theorem 1.2. (Aumann-von Neumann Selection Theorem) *Let X be a topological space and let F be a multivalued function from X to $\mathbf{R}^d$. If the graph of F belongs to $\mathcal{B}(X) \otimes \mathcal{B}(\mathbf{R}^d)$, then there exists a $\widehat{\mathcal{B}}(X)$ -measurable function which is a selection of F on the set $\{x \in X : F(x) \neq \emptyset\}$.*

Lipschitz projections. Let us recall some results obtained in [14], Section 2.

Theorem 1.3. *Let $\mathcal{C}$ be the family of all non-empty, compact and convex subsets of $\mathbf{R}^m$. Then there exists a map $P: \mathbf{R}^m \times \mathcal{C} \rightarrow \mathbf{R}^m$ satisfying the following properties:*

- (i) *P is lipschitzian considering on $\mathcal{C}$ the Hausdorff metric;*
- (ii) *$P(\xi, C) \in C$ for every $\xi \in \mathbf{R}^m$, $C \in \mathcal{C}$, and $P(\xi, C) = \xi$ if $\xi \in C$;*
- (iii) *$d(\xi, C) \leq |\xi - P(\xi, C)| \leq \sqrt{3}d(\xi, C)$ for every $\xi \in \mathbf{R}^m$ and $C \in \mathcal{C}$.*

Corollary 1.4. *There exists a Lipschitz function $P_k: (\mathbf{R}^m)^{k+1} \rightarrow \mathbf{R}^m$ satisfying the following properties:*

- (i) *$P_k(\xi; \xi_1, \dots, \xi_k) \in \text{co}\{\xi_1, \dots, \xi_k\}$ for every $\xi, \xi_1, \dots, \xi_k \in \mathbf{R}^m$, and $P_k(\xi; \xi_1, \dots, \xi_k) = \xi$ if $\xi \in \text{co}\{\xi_1, \dots, \xi_k\}$;*
- (ii) *$d(\xi, \text{co}\{\xi_1, \dots, \xi_k\}) \leq |\xi - P_k(\xi; \xi_1, \dots, \xi_k)| \leq \sqrt{3}d(\xi, \text{co}\{\xi_1, \dots, \xi_k\})$ for every $\xi, \xi_1, \dots, \xi_k \in \mathbf{R}^m$.*

Remark 1.5. Let $u, u_1, \dots, u_k \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ and let $C_k(x) = \text{co}\{u_1(x), \dots, u_k(x)\}$. If $u(x) \in C_k(x)$ for a.e. $x \in \Omega$, then $u(x) \in C_k(x)$ for q.e. $x \in \Omega$. Indeed, since P_k is lipschitzian, the function $P_k(u; u_1, \dots, u_k)$ is quasi continuous and by assumption $P_k(u; u_1, \dots, u_k) = u$ a.e. on Ω . By well-known properties of Sobolev functions (see [20]) this implies $P_k(u; u_1, \dots, u_k) = u$ q.e. on Ω , hence $u(x) \in C_k(x)$ for q.e. $x \in \Omega$.

Remark 1.6. In connection with Theorem 1.3 and Corollary 1.4 we note that for every $u \in W^{1,p}(\Omega, \mathbf{R}^m)$, and for every Lipschitz function $f: \mathbf{R}^m \rightarrow \mathbf{R}$ such that $f \circ u \in L^p(\Omega)$, we have $f \circ u \in W^{1,p}(\Omega)$ and

$$|D_i(f \circ u)| \leq L |D_i u| \quad \text{a.e. on } \Omega \quad (i = 1, \dots, n),$$

where L is the Lipschitz constant of f . This result is classical if f is a C^1 function (see, for instance, [25], Theorem 3.1.9). In the general case it can be obtained by approximating f with a sequence of C^1 functions with Lipschitz constants bounded by L .

2. A class of C^1 -convex functionals: preliminary properties

Let us first introduce our class $\mathcal{F}$ of C^1 -convex local functionals.

Definition 2.1. Let $\mathcal{F}$ be the class of all functionals $F: W^{1,p}(\Omega, \mathbf{R}^m) \times \mathcal{A}(\Omega) \rightarrow [0, +\infty]$ satisfying the following properties:

- (i) (lower semicontinuity) for every $A \in \mathcal{A}(\Omega)$ the function $F(\cdot, A)$ is lower semicontinuous on $W^{1,p}(\Omega, \mathbf{R}^m)$;
- (ii) (measure property) for every $u \in W^{1,p}(\Omega, \mathbf{R}^m)$ the set function $F(u, \cdot)$ is the trace of a Borel measure on Ω ;
- (iii) (locality property) $F(u, A) = F(v, A)$ whenever $u, v \in W^{1,p}(\Omega, \mathbf{R}^m)$, $A \in \mathcal{A}(\Omega)$, and $u|_A = v|_A$;
- (iv) (C^1 -convexity) for every $A \in \mathcal{A}(\Omega)$ the function $F(\cdot, A)$ is convex on $W^{1,p}(\Omega, \mathbf{R}^m)$ and, in addition, $F(\varphi u + (1 - \varphi)v, A) \leq F(u, A) + F(v, A)$ for every $u, v \in W^{1,p}(\Omega, \mathbf{R}^m)$ and for every $\varphi \in C^1(\Omega) \cap W^{1,\infty}(\Omega)$, with $0 \leq \varphi \leq 1$ on Ω .

Example 2.2. Let $K: \Omega \rightarrow \mathbf{R}^m$ be any multifunction with closed convex values and let $F: W^{1,p}(\Omega, \mathbf{R}^m) \times \mathcal{A}(\Omega) \rightarrow [0, +\infty]$ be the “obstacle functional” defined by

$$F(u, A) = \begin{cases} 0, & \text{if } u(x) \in K(x) \text{ for q.e. } x \in A, \\ +\infty, & \text{otherwise.} \end{cases}$$

Then F satisfies all conditions of Definition 2.1, hence $F \in \mathcal{F}$. As mentioned in the introduction, it will be proved in [17] that all functionals which arise in the study of limits of obstacle problems of the form (0.1) still belong to the class $\mathcal{F}$.

Let μ and ν be two positive Borel measures on Ω and let $f: \Omega \times \mathbf{R}^m \rightarrow [0, +\infty]$ be a Borel function such that $f(x, \cdot)$ is convex and lower semicontinuous on $\mathbf{R}^m$ for μ -a.e. $x \in \Omega$. If μ is absolutely continuous with respect to capacity, then the functional

$$F(u, A) = \int_A f(x, u(x)) d\mu + \nu(A)$$

belongs to the class $\mathcal{F}$. In both examples the lower semicontinuity follows easily from well known properties of the quasi continuous representatives of Sobolev functions (see, for instance, [31], Lemma 2.6.4).

Remark 2.3. Given a functional F of the class $\mathcal{F}$, let us consider the following extension to $W^{1,p}(\Omega, \mathbf{R}^m) \times \mathcal{B}(\Omega)$:

$$(2.1) \quad F(u, B) = \inf\{F(u, A) : A \in \mathcal{A}(\Omega), B \subseteq A\}.$$

It turns out (see [19], Theorem 5.6) that condition (ii) of Definition 2.1 is equivalent to the assumption that, for every $u \in W^{1,p}(\Omega, \mathbf{R}^m)$, the extension (2.1) of $F(u, \cdot)$ is a Borel measure on Ω .

In the sequel, when dealing with Borel sets, we shall always consider the extension of F given by (2.1).

From property (iii) in Definition 2.1 and from (2.1) it follows that $F(u, B) = F(v, B)$ for every $B \in \mathcal{B}(\Omega)$ and for every u, v in $W^{1,p}(\Omega, \mathbf{R}^m)$ which coincide in a neighborhood of B .

Note that, if F is the obstacle functional of Example 2.2, then, in general, its extension given by (2.1) does not satisfy

$$F(u, B) = \begin{cases} 0, & \text{if } u(x) \in K(x) \text{ for q.e. } x \in B, \\ +\infty, & \text{otherwise,} \end{cases}$$

for every $B \in \mathcal{B}(\Omega)$. For instance, if $n = m = 1$ and $\Omega =]-2, 2[$, let us consider the obstacle functional $F: W^{1,p}(\Omega, \mathbf{R}^m) \times \mathcal{A}(\Omega) \rightarrow [0, +\infty]$ defined by

$$F(u, A) = \begin{cases} 0, & \text{if } u(x) \geq x^2 \text{ for q.e. } x \in A, \\ +\infty, & \text{otherwise.} \end{cases}$$

Then the extension (2.1) gives $F(1, [0, 1]) = +\infty$, although $1 \geq x^2$ for every $x \in [0, 1]$.

Remark 2.4. If F is a functional of $\mathcal{F}$ and $A \in \mathcal{A}(\Omega)$, then for every finite family $(u_i)_{i \in I}$ of elements of $W^{1,p}(\Omega, \mathbf{R}^m)$ and for every family $(\varphi^i)_{i \in I}$ of non-negative functions in $C^1(\Omega) \cap W^{1,\infty}(\Omega)$ such that $\sum_i \varphi^i = 1$ in Ω , we have $F(\sum_i \varphi^i u_i, A) \leq \sum_i F(u_i, A)$. Indeed, let $u_1, \dots, u_r \in W^{1,p}(\Omega, \mathbf{R}^m)$, $\varphi^1, \dots, \varphi^r \in C^1(\Omega) \cap W^{1,\infty}(\Omega)$, with $\varphi^i \geq 0$ and $\sum_{i=1}^r \varphi^i = 1$. It would be clear, by induction, that $F(\sum_{i=1}^r \varphi^i u_i, A) \leq \sum_{i=1}^r F(u_i, A)$ if we had $\varphi^i > \varepsilon$ for every $i = 1, \dots, r$ and for a suitable $\varepsilon > 0$. Since F is lower semicontinuous, we can reduce our problem to this case by considering the coefficients $\varphi_\varepsilon^i = (\varphi^i + \varepsilon)/(1 + r\varepsilon)$.

We also notice that, by using the definition (2.1) of F on Borel sets, property (iv) holds for $A \in \mathcal{B}(\Omega)$, too.

Given $F \in \mathcal{F}$, we now generalize to Borel sets the locality property (iii) for F (Proposition 2.6). As a consequence we can single out that part of the functional F which is absolutely continuous with respect to the capacity (Proposition 2.8).

For the proof of the locality property on Borel sets we need the following remark, which, for future convenience, we state in a slightly more general form than actually needed here.

Lemma 2.5. *Let $s > 0$ and $T_s: \mathbf{R}^m \rightarrow \mathbf{R}^m$ be the orthogonal projection onto the ball $\overline{B_s(0)}$, i.e.,*

$$T_s(\xi) = \frac{s}{|\xi| \vee s} \xi = \begin{cases} \xi, & \text{if } |\xi| \leq s, \\ s \frac{\xi}{|\xi|}, & \text{if } |\xi| \geq s. \end{cases}$$

If $u \in W^{1,p}(\Omega, \mathbf{R}^m)$, then $T_s \circ u \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ and the sequence $(T_s \circ u)$ converges, in the strong topology of $W^{1,p}(\Omega, \mathbf{R}^m)$, to 0 as s goes to 0^+ and to u as s goes to $+\infty$.

Proof. Since T_s is lipschitzian, for every fixed $u \in W^{1,p}(\Omega, \mathbf{R}^m)$ we have $T_s \circ u \in W^{1,p}(\Omega, \mathbf{R}^m)$ by Remark 1.6. We prove only the convergence as s tends to 0^+ , the other part being analogous. Since $(T_s \circ u)$ converges to 0 strongly in $L^p(\Omega, \mathbf{R}^m)$, there is only to verify the same kind of convergence for $(D(T_s \circ u))$. Let $\sigma \geq s$; since orthogonal projections have Lipschitz constant 1, the pointwise estimate in Remark 1.6 yields

$$\begin{aligned} \int_{\Omega} |D(T_s \circ u)|^p dx &= \int_{\{|u| \leq \sigma\}} |D(T_s \circ u)|^p dx + \int_{\{|u| > \sigma\}} |D(\frac{s}{\sigma}(T_\sigma \circ u))|^p dx \\ &\leq \int_{\{|u| \leq \sigma\}} |Du|^p dx + (\frac{s}{\sigma})^p \int_{\Omega} |Du|^p dx. \end{aligned}$$

The conclusion now follows taking first the limit as s tends to 0^+ and then the limit as σ tends to 0^+ . $\square$

Proposition 2.6. (Locality Property on Borel Sets) *Let $u, v \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ and $B \in \mathcal{B}(\Omega)$. If $u = v$ q.e. on B and $F(u, B), F(v, B) < +\infty$, then $F(u, B) = F(v, B)$.*

Proof. Step 1. Assume B is quasi open. For every $h \in \mathbf{N}$, let A_h be an open set with $\text{cap}(A_h, \Omega) < 1/h$ and such that $B_h = B \cup A_h$ is open. Let w_h be the capacitary potential of A_h and $u_h = u + w_h(v - u)$. It turns out that $u_h = v$ q.e. on B_h . Moreover, (u_h)

converges to u in $W^{1,p}(\Omega, \mathbf{R}^m)$ since (w_h) converges to 0 in $W^{1,p}(\Omega)$ and $0 \leq w_h \leq 1$ for every $h \in \mathbf{N}$.

Since by assumption $F(u, B) < +\infty$ and $F(v, B) < +\infty$, it follows that for every given $\varepsilon > 0$ there exist an open set A and a compact set K with $K \subseteq B \subseteq A$ such that $F(u, A \setminus K) < \varepsilon$ and $F(v, A \setminus K) < \varepsilon$.

By the lower semicontinuity of F on open sets we get

$$(2.2) \quad F(u, B) \leq F(u, A) \leq \liminf_{h \rightarrow \infty} F(u_h, A) \leq \liminf_{h \rightarrow \infty} [F(u_h, A \cap B_h) + F(u_h, A \setminus K)].$$

From the locality property of F on open sets it follows that

$$(2.3) \quad F(u_h, A \cap B_h) = F(v, A \cap B_h) \leq F(v, A) < F(v, B) + \varepsilon.$$

By approximating w_h in $W^{1,p}(\Omega)$ with a sequence of equibounded functions of $\mathcal{C}_0^1(\Omega)$, the semicontinuity and C^1 -convexity of F (properties (i) and (iv)) imply that

$$(2.4) \quad F(u_h, A \setminus K) \leq F(u, A \setminus K) + F(v, A \setminus K).$$

Hence, $F(u_h, A \setminus K) < 2\varepsilon$, and, by (2.2) and (2.3), $F(u, B) \leq F(v, B) + 3\varepsilon$. Since ε is arbitrary, we can conclude that $F(u, B) \leq F(v, B)$. Interchanging the roles of u and v , we obtain the opposite inequality. This proves the theorem when B is quasi open.

Step 2. Let now B be a Borel subset of Ω . For every $h \in \mathbf{N}$ let us define $B_h = \{x \in \Omega : |u(x) - v(x)| < 1/h\}$ and

$$u_h = u + \frac{1/h}{|v - u| \vee (1/h)} (v - u) = u + T_{1/h} \circ (v - u).$$

Clearly, $u_h = v$ q.e. on B_h . By Lemma 2.5 we have the convergence of u_h to u in $W^{1,p}(\Omega, \mathbf{R}^m)$. At this point we can introduce the sets A and K as in Step 1 and proceed in the same way replacing the locality property of F on the open sets with the locality property on the quasi open sets proved in Step 1. We have only to remark about the estimate (2.4). Let us notice that it is enough to consider $B \subset\subset \Omega$, hence we can choose $A \subset\subset \Omega$; for every $\Omega' \subset\subset \Omega$ the coefficient in the convex combination between u and v defining u_h can be approximated in $W^{1,p}(\Omega')$ by an equibounded sequence of functions of $C^1(\overline{\Omega'})$. As F is C^1 -convex and local on open sets, this suffices to get (2.4) as above. $\square$

Let us consider the function $\nu_0: \mathcal{B}(\Omega) \rightarrow [0, +\infty]$ defined by

$$(2.5) \quad \nu_0(B) = \inf\{F(v, B) : v \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)\}$$

for every $B \in \mathcal{B}(\Omega)$. Moreover, for every $A \in \mathcal{A}(\Omega)$ we define

$$\text{dom}F(\cdot, A) = \{u \in W^{1,p}(\Omega, \mathbf{R}^m) : F(u, A) < +\infty\}.$$

Then the following proposition holds.

Proposition 2.7. *For every $\Omega' \in \mathcal{A}(\Omega)$ with $\text{dom}F(\cdot, \Omega') \cap L^\infty(\Omega, \mathbf{R}^m) \neq \emptyset$, the restriction of ν_0 to $\mathcal{B}(\Omega')$ is a positive finite Borel measure.*

Proof. It is clear that ν_0 is an increasing function, $\nu_0(\emptyset) = 0$ and, in view of the definition of $F(u, \cdot)$ on $\mathcal{B}(\Omega)$, that $\nu_0(B) = \inf\{\nu_0(A) : A \in \mathcal{A}(\Omega'), A \supseteq B\}$. Hence, by Proposition 5.5 and Theorem 5.6 in [19], we have only to prove that ν_0 is superadditive, subadditive, and inner regular on $\mathcal{A}(\Omega')$. The superadditivity comes immediately from the definition of ν_0 and from the additivity of F in the second variable. Let us now prove that for every $A_1, A_2, A'_2 \in \mathcal{A}(\Omega')$ with $A'_2 \subset\subset A_2$ we have

$$(2.6) \quad \nu_0(A_1 \cup A'_2) \leq \nu_0(A_1) + \nu_0(A_2).$$

We can assume that $\nu_0(A_1) + \nu_0(A_2) < +\infty$. Then, for every $\varepsilon > 0$ there exist two functions u_1, u_2 in $W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ such that

$$\nu_0(A_1) + \frac{\varepsilon}{2} > F(u_1, A_1) \quad \nu_0(A_2) + \frac{\varepsilon}{2} > F(u_2, A_2).$$

Let $\varphi \in C_0^1(A_2)$, with $\varphi = 1$ on a neighborhood of $\overline{A'_2}$ and $0 \leq \varphi \leq 1$. We set $u = (1 - \varphi)u_1 + \varphi u_2$. By Remark 2.3 it follows that

$$\begin{aligned} \nu_0(A_1 \cup A'_2) &\leq F(u, A_1 \cup A'_2) \\ &\leq F(u_1, A_1 \setminus A_2) + F(u_2, \overline{A'_2}) + F((1 - \varphi)u_1 + \varphi u_2, (A_2 \setminus \overline{A'_2}) \cap A_1). \end{aligned}$$

Furthermore, the C^1 -convexity of F permits to estimate the last term in the above inequality by $F(u_1, (A_2 \setminus \overline{A'_2}) \cap A_1) + F(u_2, (A_2 \setminus \overline{A'_2}) \cap A_1)$. Hence,

$$\nu_0(A_1 \cup A'_2) \leq F(u_1, A_1) + F(u_2, A_2) < \nu_0(A_1) + \nu_0(A_2) + \varepsilon.$$

Thus we obtain (2.6). This inequality will give the subadditivity of ν_0 once inner regularity will be proved.

Since $\text{dom}F(\cdot, \Omega') \cap L^\infty(\Omega, \mathbf{R}^m) \neq \emptyset$, we can find $u \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ such that $F(u, \Omega') < +\infty$. Therefore, given $A \in \mathcal{A}(\Omega')$ and $\varepsilon > 0$ there exists $A'' \in \mathcal{A}(\Omega')$ with $A'' \subset\subset A$ and $F(u, A \setminus \overline{A''}) \leq \varepsilon$; it follows that $\nu_0(A \setminus \overline{A''}) \leq \varepsilon$. Let $A' \in \mathcal{A}(\Omega')$ such that $A'' \subset\subset A' \subset\subset A$. By (2.6) we have

$$\nu_0(A) \leq \nu_0(A') + J\nu_0(A \setminus \overline{A''}) \leq \nu_0(A') + \varepsilon.$$

We conclude that $\nu_0(A) = \sup\{\nu_0(A') : A' \in \mathcal{A}(\Omega'), A' \subset\subset A\}$, i.e., the inner regularity of ν_0 . $\square$

Proposition 2.8. *For every $A \in \mathcal{A}(\Omega)$ and $u \in \text{dom}F(\cdot, A) \cap L^\infty(\Omega, \mathbf{R}^m)$, the function $F(u, \cdot) - \nu_0(\cdot)$ is a positive Borel measure on A which is absolutely continuous with respect to capacity.*

Proof. Let $u \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ such that $F(u, A) < +\infty$. As ν_0 is a finite Borel measure on A (Proposition 2.7) and $\nu_0(\cdot) \leq F(u, \cdot)$ by (2.5), we conclude that $F(u, \cdot) - \nu_0(\cdot)$ is a positive Borel measure on A . Let us fix $B \in \mathcal{B}(A)$ with $\text{cap}(B, \Omega) = 0$. For every $v \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ with $F(v, B) < +\infty$, we have $v = u$ q.e. on B ; hence, by Proposition 2.6 we conclude that $F(v, B) = F(u, B)$. Since

$$\nu_0(B) = \inf\{F(v, B) : v \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m), F(v, B) < +\infty\}$$

it follows that $\nu_0(B) = F(u, B)$, i.e., $F(u, B) - \nu_0(B) = 0$. $\square$

We now conclude this section by giving a basic estimate for F on the convex hull of a finite number of functions in $W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$.

Proposition 2.9. *Let $u, u_1, \dots, u_k \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$. Assume that $u(x) \in \text{co}\{u_1(x), \dots, u_k(x)\}$ for a.e. $x \in \Omega$. Then*

$$F(u, B) \leq \sum_{i=1}^k F(u_i, B)$$

for every Borel set B in Ω .

Proof. In view of the definition of F on Borel sets, it is enough to prove the inequality for every open set B with $B \subset\subset \Omega$. Hence, let B be such a set. By means of the

Density Theorem 2.9 in [14], we can easily find a sequence of functions $\varphi_h: \mathbf{R}^n \rightarrow \Sigma_k$ such that $\varphi_h \in C^\infty(\mathbf{R}^n, \mathbf{R}^k)$ and

$$\sum_{i=1}^k \varphi_h^i u_i \xrightarrow[h]{} u$$

strongly in $W^{1,p}(A, \mathbf{R}^m)$, where A is a neighborhood of $\overline{B}$. Then, from the lower semicontinuity of F and the locality property on open sets we obtain

$$F(u, B) \leq \liminf_{h \rightarrow \infty} F\left(\sum_{i=1}^k \varphi_h^i u_i, B\right).$$

The conclusion follows now from the C^1 -convexity of F and from Remark 2.4. $\square$

3. Integral representation on moving polytopes

The aim of this section is the integral representation of the functionals in $\mathcal{F}$ when restricted to the pointwise convex hull of a finite number of functions in $W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ (see Theorem 3.7).

Let $F \in \mathcal{F}$, $k \in \mathbf{N}$, and $u_1, \dots, u_k \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ be fixed. Throughout this section we assume that $F(u_i, \Omega) < +\infty$ for every $i = 1, \dots, k$. We point out that our proof first produces a kind of integral representation of F on the functions of the form $u = \sum_{i=1}^k \psi^i u_i$, where $\psi: \Omega \rightarrow \Sigma_k$, with the integrand depending on the coefficient ψ rather than on the function u itself (Theorem 3.3). For a constant ψ , this result is contained in the following lemma.

Let ν_0 be the set function introduced in (2.5). Under the present assumptions, Proposition 2.7 tells us that ν_0 is a finite Borel measure on Ω . Let μ be a positive finite Borel measure on Ω with $\text{supp } \mu = \Omega$ such that μ is absolutely continuous with respect to capacity and $\mu(\cdot) \geq \sum_{i=1}^k (F(u_i, \cdot) - \nu_0(\cdot))$ (in view of Proposition 2.8, such a μ can be obtained, for example, by adding to $\sum_{i=1}^k (F(u_i, \cdot) - \nu_0(\cdot))$ the positive measure $f d\mathcal{L}^n$, where $f \in L^1(\Omega)$, $f > 0$ on Ω , and $\mathcal{L}^n$ is the n -dimensional Lebesgue measure).

Lemma 3.1. *For every $x \in \Omega$ and $\lambda \in \Sigma_k$ we define $u_\lambda(x) = \sum_{i=1}^k \lambda^i u_i(x)$ and*

$$(3.1) \quad g(x, \lambda) = \limsup_{r \rightarrow 0^+} \frac{F(u_\lambda, B_r(x)) - J\nu_0(B_r(x))}{\mu(B_r(x))}.$$

Then

- (i) *for every $x \in \Omega$ the function $g(x, \cdot)$ is convex and continuous in Σ_k ;*
- (ii) *for every $\lambda \in \Sigma_k$ the function $g(\cdot, \lambda)$ is Borel measurable on Ω ;*
- (iii) *$F(u_\lambda, B) = \int_B g(x, \lambda) d\mu + \nu_0(B)$ for every $\lambda \in \text{ri}\Sigma_k$ and $B \in \mathcal{B}(\Omega)$.*

Proof. From the definition of μ and ν_0 and from the convexity of F , it follows immediately that $0 \leq g \leq 1$ on $\Omega \times \text{ri}\Sigma_k$, and that $g(x, \cdot)$ is convex on $\text{ri}\Sigma_k$ for every $x \in \Omega$. Then, by Theorem 10.3 in [28], for every $x \in \Omega$ the function $g(x, \cdot)$ can be extended in one and only one way to a continuous convex function, still denoted by g , on the whole of Σ_k . Hence, $0 \leq g \leq 1$ on $\Omega \times \Sigma_k$ and (i) holds.

Let us proof (ii). If α is a positive Borel measure on Ω , the function $r \mapsto \alpha(B_r(x))$ is left continuous for every $x \in \Omega$. This implies that the upper limit which appears in (3.1) can equivalently be taken as $r \rightarrow 0^+$ with $r \in \mathbf{Q}$. Moreover, the function $x \mapsto \alpha(B_r(x))$ is lower semicontinuous for every $r > 0$ and hence Borel measurable, too. It follows that the function $g(\cdot, \lambda)$ is Borel measurable for every $\lambda \in \text{ri}\Sigma_k$. For $\lambda \in \partial\Sigma_k$, $g(\cdot, \lambda)$ is the pointwise limit of a sequence $g(\cdot, \lambda_n)$ with $\lambda_n \in \text{ri}\Sigma_k$; therefore, $g(\cdot, \lambda)$ is Borel measurable on Ω for every $\lambda \in \Sigma_k$.

By the Besicovitch differentiation theorem (see, e.g., [31], Section 1.3), we have

$$F(u_\lambda, B) = \int_B g(x, \lambda) d\mu + \nu_0(B)$$

for every $B \in \mathcal{B}(\Omega)$ and for every $\lambda \in \text{ri}\Sigma_k$. □

Before extending the previous result to non constant λ 's, we observe that the following selection lemma holds.

Lemma 3.2. *Let $u \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ such that $u(x) \in \text{co}\{u_1(x), \dots, u_k(x)\}$ for a.e. $x \in \Omega$. Then, there exists a $\widehat{\mathcal{B}}(\Omega)$ -measurable function $\psi: \Omega \rightarrow \Sigma_k$ such that*

$$(3.2) \quad u(x) = \sum_{i=1}^k \psi^i(x) u_i(x) \quad \text{for q.e. } x \in \Omega.$$

Proof. Let us fix some quasi continuous Borel measurable representatives of $u, u_1, \dots, u_k$ (see Section 1). Let $\Lambda(x) = \{\lambda \in \Sigma_k : u(x) = \sum_{i=1}^k \lambda^i u_i(x)\}$ for every $x \in \Omega$. Λ is a multivalued function from Ω to Σ_k with non-empty closed values for q.e. $x \in \Omega$ (see Remark 1.5). It is clear that $\text{graph } \Lambda \in \mathcal{B}(\Omega) \otimes \mathcal{B}(\mathbf{R}^k)$.

By the Aumann-von Neumann Selection Theorem 1.2 there exists a $\widehat{\mathcal{B}}(\Omega)$ -measurable selection ψ of the multivalued function Λ and, by the definition of $\Lambda(x)$, the function ψ satisfies (3.2). $\square$

Theorem 3.3. *Let $u \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ with $u(x) \in \text{co}\{u_1(x), \dots, u_k(x)\}$ for a.e. $x \in \Omega$. If g is the function given by Lemma 3.1 and $\psi: \Omega \rightarrow \Sigma_k$ is a $\widehat{\mathcal{B}}(\Omega)$ -measurable function such that (3.2) holds, then $g(\cdot, \psi(\cdot))$ is μ -measurable and*

$$(3.3) \quad F(u, A) = \int_A g(x, \psi(x)) d\mu + \nu_0(A)$$

for every $A \in \mathcal{A}(\Omega)$.

Let us explicitly notice that if u is as above, then $F(u, \Omega) < +\infty$ by Proposition 2.9. The proof of Theorem 3.3 heavily relies on the following approximation lemma, which essentially reduces the problem to the case of a constant ψ .

Lemma 3.4. *Let u and ψ be as in Theorem 3.3. Let $\lambda \in \text{ri}\Sigma_k$ with $d(\lambda, \partial\Sigma_k) = \eta > 0$, let $0 < \varepsilon < \eta$ and $B \in \mathcal{B}(\Omega)$ such that $|\psi(x) - \lambda| \leq \varepsilon$ for q.e. $x \in B$. Then,*

$$|F(u, B) - F(u_\lambda, B)| \leq \frac{\varepsilon}{\eta} \sum_{i=1}^k F(u_i, B).$$

Proof. Let us define $v = u + t(u_\lambda - u)$ on Ω , with $t = 1 + \eta/\varepsilon$. It turns out that $v(x) \in \text{co}\{u_1(x), \dots, u_k(x)\}$ for q.e. $x \in B$ and

$$u_\lambda = \frac{1}{t}v + (1 - \frac{1}{t})u \quad \text{on } \Omega.$$

In order to get from v a function which belongs to $\text{co}\{u_1, \dots, u_k\}$ a.e. on Ω , we consider the projection $w = P_k(v; u_1, \dots, u_k)$ as defined in Corollary 1.4. Set

$$z = \frac{1}{t}w + (1 - \frac{1}{t})u \quad \text{on } \Omega.$$

By Remark 1.6, the function w , and hence z , belongs to $W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$. Moreover, since $w = P_k(v; u_1, \dots, u_k) = v$ q.e. on B (see Corollary 1.4), we have $z = u_\lambda$ q.e. on B . By the convexity of F and Proposition 2.9

$$\begin{aligned} F(z, B) &\leq \frac{1}{t}F(w, B) + (1 - \frac{1}{t})F(u, B) \\ &\leq \frac{\varepsilon}{\eta} \sum_{i=1}^k F(u_i, B) + F(u, B). \end{aligned}$$

In view of the locality property of F on Borel sets (Proposition 2.6) we have $F(z, B) = F(u_\lambda, B)$; thus

$$F(u_\lambda, B) \leq F(u, B) + \frac{\varepsilon}{\eta} \sum_{i=1}^k F(u_i, B).$$

The inequality

$$F(u, B) \leq F(u_\lambda, B) + \frac{\varepsilon}{\eta} \sum_{i=1}^k F(u_i, B)$$

can be obtained analogously defining now $v = u_\lambda + t(u - u_\lambda)$ with $t = \eta/\varepsilon > 1$. $\square$

Proof of Theorem 3.3. Let us fix $A \in \mathcal{A}(\Omega)$.

Step 1. Assume that $\psi(x) \in \text{ri}\Sigma_k$ for every $x \in \Omega$.

Given $\eta > 0$, let us define $\Sigma_{k,\eta} = \{\lambda \in \Sigma_k : d(\lambda, \partial\Sigma_k) \geq \eta\}$ and $A_\eta = \psi^{-1}(\Sigma_{k,\eta}) \cap A$. For every $\varepsilon \in]0, \eta[$ we can fix a finite partition $(B_j)_{j \in J}$ of $\Sigma_{k,\eta}$ by means of Borel sets having diameter less than ε , and a family $(\lambda_j)_{j \in J}$ of elements of $\Sigma_{k,\eta}$ such that $\lambda_j \in B_j$ for every $j \in J$. Let us define $E_j = \psi^{-1}(B_j) \cap A$ for every $j \in J$. Since ψ is $\widehat{\mathcal{B}}(\Omega)$ -measurable, the sets A_η and E_j are in $\widehat{\mathcal{B}}(\Omega)$. According to the convention made in Section 1, for every $z \in W^{1,p}(\Omega, \mathbf{R}^m)$ the completion of the measures μ , ν_0 and $F(z, \cdot)$ will be still denoted by μ , ν_0 and $F(z, \cdot)$. By Lemma 3.4, for every $j \in J$

$$|F(u, E_j) - F(u_{\lambda_j}, E_j)| \leq \frac{\varepsilon}{\eta} \sum_{i=1}^k F(u_i, E_j).$$

This and Lemma 3.1 imply

$$\begin{aligned} F(u, A_\eta) - \nu_0(A_\eta) &= \sum_{j \in J} [F(u, E_j) - \nu_0(E_j)] \\ &\leq \sum_{j \in J} [F(u_{\lambda_j}, E_j) - \nu_0(E_j) + \frac{\varepsilon}{\eta} \sum_{i=1}^k F(u_i, E_j)] \\ &= \sum_{j \in J} \left[\int_{E_j} g(x, \lambda_j) d\mu + \frac{\varepsilon}{\eta} \sum_{i=1}^k F(u_i, E_j) \right]. \end{aligned}$$

Since $g(x, \cdot)$ is convex and bounded by 1 in Σ_k , it is Lipschitz continuous in $\Sigma_{k,\eta}$ with constant $1/\eta$; thus

$$\begin{aligned} F(u, A_\eta) - \nu_0(A_\eta) &\leq \sum_{j \in J} \left[\int_{E_j} g(x, \psi(x)) d\mu + \frac{1}{\eta} \int_{E_j} |\psi(x) - \lambda_j| d\mu + \frac{\varepsilon}{\eta} \sum_{i=1}^k F(u_i, E_j) \right] \\ &\leq \int_{A_\eta} g(x, \psi(x)) d\mu + \frac{\varepsilon}{\eta} \left[\mu(A_\eta) + \sum_{i=1}^k F(u_i, A_\eta) \right]. \end{aligned}$$

(Note that $g(\cdot, \psi(\cdot))$ is μ -measurable since g is Borel measurable and ψ is $\widehat{\mathcal{B}}(\Omega)$ -measurable). Since ε is arbitrary, we get

$$F(u, A_\eta) - \nu_0(A_\eta) \leq \int_{A_\eta} g(x, \psi(x)) d\mu.$$

Now, taking into account that $\psi(x) \in \text{ri}\Sigma_k$ for every $x \in \Omega$ and letting $\eta \rightarrow 0^+$, we obtain

$$F(u, A) - \nu_0(A) \leq \int_A g(x, \psi(x)) d\mu.$$

The reverse inequality can be obtained in a completely analogous way.

Step 2. We consider now the general case $\psi: \Omega \rightarrow \Sigma_k$.

Let $b_0 = \frac{1}{k}(e_1 + \dots + e_k)$ be the barycenter of Σ_k ($e_1, \dots, e_k$ are the elements of the standard basis of $\mathbf{R}^k$). For every $0 \leq \sigma \leq 1$ define $\psi_\sigma = b_0 + \sigma(\psi - b_0)$ and $u_\sigma = \sum_{i=1}^k \psi_\sigma^i u_i = u_0 + \sigma(u - u_0)$, where $u_0 = \frac{1}{k}(u_1 + \dots + u_k)$. If $0 \leq \sigma < 1$ then $\psi_\sigma(x) \in \text{ri}\Sigma_k$ for every $x \in \Omega$ (see [28], Theorem 6.1); therefore, by Step 1 we have

$$F(u_\sigma, A) = \int_A g(x, \psi_\sigma(x)) d\mu + \nu_0(A).$$

Observe now that the lower semicontinuity and the convexity of F imply that

$$\lim_{\sigma \rightarrow 1^-} F(u_\sigma, A) = F(u, A).$$

Moreover, the continuity of $g(x, \cdot)$ and the dominated convergence theorem yield

$$\lim_{\sigma \rightarrow 1^-} \int_A g(x, \psi_\sigma(x)) d\mu = \int_A g(x, \psi(x)) d\mu.$$

This concludes the proof. □

We point out that the values $u(x)$ of the function u enter the integral representation of Theorem 3.3 through the parameters $\psi^i(x)$ for which $u(x) = \sum_{i=1}^k \psi^i(x)u_i(x)$. When looking for an integrand depending directly on the values of u , the main difficulty we meet is that the expression of $u(x)$ as a convex combination of $u_1(x), \dots, u_k(x)$ is not necessarily unique. This problem is essentially overcome by the following lemma.

Lemma 3.5. *Let $u \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ and $u(x) \in \text{co}\{u_1(x), \dots, u_k(x)\}$ for a.e. $x \in \Omega$. Let g be the function defined in Lemma 3.1 and*

$$N = \{x \in \Omega : \exists \lambda, \lambda' \in \Sigma_k \text{ } u(x) = u_\lambda(x) = u_{\lambda'}(x) \text{ and } g(x, \lambda) \neq g(x, \lambda')\}$$

(N is defined up to sets of zero capacity). Then, $N \in \widehat{\mathcal{B}}(\Omega)$ and $\mu(N) = 0$.

Proof. As in the proof of Lemma 3.2 we can fix quasi continuous, Borel measurable representatives of $u, u_1, \dots, u_k$; the set N is now well defined all over Ω . Consider the multivalued map Γ from Ω into $\Sigma_k \times \Sigma_k$ defined by

$$\Gamma(x) = \{(\lambda, \lambda') \in \Sigma_k \times \Sigma_k : u(x) = u_\lambda(x) = u_{\lambda'}(x) \text{ and } g(x, \lambda) \neq g(x, \lambda')\}.$$

Arguing as in the proof of Lemma 3.2 and taking into account that g is Borel measurable on $\Omega \times \Sigma_k$, we obtain that $\text{graph} \Gamma \in \mathcal{B}(\Omega) \otimes \mathcal{B}(\mathbf{R}^k) \otimes \mathcal{B}(\mathbf{R}^k)$. By the Projection Theorem (Theorem 1.1) we get $N \in \widehat{\mathcal{B}}(\Omega)$. The Aumann-von Neumann Selection Theorem (Theorem 1.2) implies the existence of two $\widehat{\mathcal{B}}(\Omega)$ -measurable functions $\sigma_1, \sigma_2: \Omega \rightarrow \Sigma_k$ such that $(\sigma_1|_N, \sigma_2|_N)$ is a selection of Γ on N . Define for $j = 1, 2$

$$\psi_j = \begin{cases} \psi, & \text{on } \Omega \setminus N, \\ \sigma_j, & \text{on } N, \end{cases}$$

where $\psi: \Omega \rightarrow \Sigma_k$ is the $\widehat{\mathcal{B}}(\Omega)$ -measurable function given in Lemma 3.2. Then ψ_1 and ψ_2 are $\widehat{\mathcal{B}}(\Omega)$ -measurable functions such that

$$(3.4) \quad u(x) = \sum_{i=1}^k \psi_1^i(x)u_i(x) = \sum_{i=1}^k \psi_2^i(x)u_i(x) \quad \text{for q.e. } x \in \Omega,$$

$$(3.5) \quad g(x, \psi_1(x)) \neq g(x, \psi_2(x)) \quad \text{for every } x \in N.$$

By Theorem 3.3, (3.4) implies that

$$\int_A g(x, \psi_1(x)) d\mu = \int_A g(x, \psi_2(x)) d\mu$$

for every $A \in \mathcal{A}(\Omega)$. Hence,

$$g(x, \psi_1(x)) = g(x, \psi_2(x)) \quad \text{for } \mu\text{-a.e. } x \in \Omega.$$

Together with (3.5) this yields that $\mu(N) = 0$. □

For future convenience we single out a technical remark about measurability of functions.

Remark 3.6. (i) Let T be a Borel subset of $\mathbf{R}^m$. Given $g: \Omega \times \mathbf{R}^m \times T \rightarrow [0, +\infty]$, let us define $f: \Omega \times \mathbf{R}^m \rightarrow [0, +\infty]$ by setting $f(x, \xi) = \inf_{t \in T} g(x, \xi, t)$. If $g(x, \cdot, t)$ is continuous for every $(x, t) \in \Omega \times T$, uniformly with respect to $t \in T$, then $f(x, \cdot)$ is continuous for every $x \in \Omega$. Assume, in addition, that $g(\cdot, \xi, \cdot)$ is $\widehat{\mathcal{B}}(\Omega) \otimes \mathcal{B}(\mathbf{R}^m)$ -measurable for every $\xi \in \mathbf{R}^m$. Then $f(\cdot, \xi)$ is $\widehat{\mathcal{B}}(\Omega)$ -measurable for every $\xi \in \mathbf{R}^m$, hence f is $\widehat{\mathcal{B}}(\Omega) \otimes \mathcal{B}(\mathbf{R}^m)$ -measurable.

Indeed, given $\xi \in \mathbf{R}^m$ and $s \in \mathbf{R}$, the set

$$E_s = \{x \in \Omega : f(x, \xi) < s\} = \{x \in \Omega : \exists t \in T \ g(x, \xi, t) < s\}$$

is the projection on Ω of the set $\{(x, t) \in \Omega \times T : g(x, \xi, t) < s\}$. Since $(x, t) \mapsto g(x, \xi, t)$ is $\widehat{\mathcal{B}}(\Omega) \otimes \mathcal{B}(\mathbf{R}^m)$ -measurable, by the Projection Theorem (Theorem 1.1) we get $E_s \in \widehat{\mathcal{B}}(\Omega)$.

(ii) Let $f: \Omega \times \mathbf{R}^m \rightarrow [0, +\infty]$ be a $\widehat{\mathcal{B}}(\Omega) \otimes \mathcal{B}(\mathbf{R}^m)$ -measurable function. Then for every positive finite Borel measure μ on Ω there exists a set $N \in \mathcal{B}(\Omega)$ with $\mu(N) = 0$ such that $f|_{(\Omega \setminus N) \times \mathbf{R}^m}$ is a Borel function.

Indeed, for every $E \in \mathcal{B}_\mu(\Omega) \otimes \mathcal{B}(\mathbf{R}^m)$ there exists $N \in \mathcal{B}(\Omega)$ with $\mu(N) = 0$ such that $E \setminus (N \times \mathbf{R}^m) \in \mathcal{B}(\Omega) \otimes \mathcal{B}(\mathbf{R}^m)$.

Finally, we are able to prove the main result of this section, i.e., the integral representation on the pointwise convex combinations of a finite number of fixed functions.

Theorem 3.7. *Let $F \in \mathcal{F}$, $k \in \mathbf{N}$, $u_1, \dots, u_k \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$, and assume $F(u_i, \Omega) < +\infty$ for every $i = 1, \dots, k$. Let ν_0 be the positive finite Borel measure introduced in Proposition 2.7. Then, there exist a positive finite Borel measure μ on Ω , absolutely continuous with respect to capacity, and a function $f: \Omega \times \mathbf{R}^m \rightarrow [0, +\infty]$ with the following properties:*

- (i) *for every $x \in \Omega$ the function $f(x, \cdot)$ is convex and lower semicontinuous on $\mathbf{R}^m$;*
- (ii) *f is $\widehat{\mathcal{B}}(\Omega) \otimes \mathcal{B}(\mathbf{R}^m)$ -measurable;*
- (iii) *for every $u \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ such that $u(x) \in \text{co}\{u_1(x), \dots, u_k(x)\}$ for a.e. $x \in \Omega$, the function $f(\cdot, u(\cdot))$ is μ -measurable on Ω and $F(u, A) = \int_A f(x, u(x)) d\mu + \nu_0(A)$ for every $A \in \mathcal{A}(\Omega)$. Moreover, the restriction of $f(x, \cdot)$ to $\text{co}\{u_1(x), \dots, u_k(x)\}$ is continuous for μ -a.e. $x \in \Omega$.*

Proof. Let us fix quasi continuous Borel measurable representatives of $u_1, \dots, u_k$. For every $\lambda \in \Sigma_k$ and $x \in \Omega$ we set $u_\lambda(x) = \sum_{i=1}^k \lambda^i u_i(x)$ and $C_k(x) = \text{co}\{u_1(x), \dots, u_k(x)\}$. Let us define the function $f: \Omega \times \mathbf{R}^m \rightarrow [0, +\infty]$ as

$$f(x, \xi) = \begin{cases} \inf_{\substack{\lambda \in \Sigma_k \\ u_\lambda(x) = \xi}} g(x, \lambda), & \text{if } \xi \in C_k(x), \\ +\infty, & \text{otherwise,} \end{cases}$$

where g is the function introduced in Lemma 3.1.

Let us prove (i). Fix $x \in \Omega$; the multivalued function from $C_k(x)$ to Σ_k defined by $\xi \mapsto \{\lambda \in \Sigma_k : u_\lambda(x) = \xi\}$ has closed graph and compact range; hence it is upper semi-continuous. Moreover, by the continuity of $g(x, \cdot)$, the function $S \mapsto \inf\{g(x, \lambda) : \lambda \in S\}$, defined on the compact subsets of Σ_k , is continuous with respect to the Hausdorff metric and decreasing with respect to inclusion. Therefore, we can deduce the lower semicontinuity of $f(x, \cdot)$ on $C_k(x)$. This immediately implies the lower semicontinuity of $f(x, \cdot)$ on $\mathbf{R}^m$, while the convexity of $f(x, \cdot)$ can be easily verified directly. Hence, (i) holds true.

Let us prove (ii). For every $x \in \Omega$ let us consider the Moreau-Yosida transforms of $f(x, \cdot)$, defined by

$$f_s(x, \xi) = \inf_{\eta \in C_k(x)} [f(x, \eta) + s|\xi - \eta|] \quad (s \in \mathbf{N})$$

for every $\xi \in \mathbf{R}^m$. Since $f(x, \cdot)$ is lower semicontinuous on $\mathbf{R}^m$, for every $x \in \Omega$ and $\xi \in \mathbf{R}^m$ we have

$$(3.6) \quad f(x, \xi) = \sup_{s \in \mathbf{N}} f_s(x, \xi).$$

Let us prove that for every $\xi \in \mathbf{R}^m$, $f_s(\cdot, \xi)$ is $\widehat{\mathcal{B}}(\Omega)$ -measurable. Note that

$$\begin{aligned} f_s(x, \xi) &= \inf_{\eta \in C_k(x)} \inf_{\substack{\lambda \in \Sigma_k \\ u_\lambda(x) = \eta}} [g(x, \lambda) + s|\xi - \eta|] \\ &= \inf_{\lambda \in \Sigma_k} [g(x, \lambda) + s|\xi - u_\lambda(x)|]. \end{aligned}$$

Remark 3.6 shows that f_s is $\widehat{\mathcal{B}}(\Omega) \otimes \mathcal{B}(\mathbf{R}^m)$ -measurable; by (3.6) the same is true for f .

Let us now turn to the proof of (iii). Fix $u \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ such that $u(x) \in \text{co}\{u_1(x), \dots, u_k(x)\}$ for a.e. $x \in \Omega$ and choose a quasi continuous representative of u . Since such a representative is $\mathcal{B}_\mu(\Omega)$ -measurable (recall that μ is absolutely

continuous with respect to capacity), $f(\cdot, u(\cdot))$ is μ -measurable on Ω . By Lemma 3.2 and Theorem 3.3, for every $A \in \mathcal{A}(\Omega)$ we have

$$F(u, A) = \int_A g(x, \psi(x)) d\mu + \nu_0(A),$$

where $\psi: \Omega \rightarrow \Sigma_k$ is a $\widehat{\mathcal{B}}(\Omega)$ -measurable function such that (3.2) holds. By the definition of f we have

$$f(x, u(x)) = \inf_{\substack{\lambda \in \Sigma_k \\ u_\lambda(x) = u(x)}} g(x, \lambda) \quad \text{for q.e. } x \in \Omega.$$

Let N be the set given in Lemma 3.5. Then

$$f(x, u(x)) = g(x, \psi(x)) \quad \text{for q.e. } x \in \Omega \setminus N.$$

This proves (iii) since $\mu(N) = 0$. □

The following proposition shows that, given the measures μ and ν , the function f obtained in the integral representation theorem is essentially unique.

Proposition 3.8. *Let $F, u_1, \dots, u_k$ be as in Theorem 3.7. Let μ and ν be two positive finite Borel measures on Ω , with μ absolutely continuous with respect to capacity. Assume that two functions $f_1, f_2: \Omega \times \mathbf{R}^m \rightarrow [0, +\infty]$ satisfy conditions (i)–(iii) of Theorem 3.7 with ν_0 replaced by ν . Then $f_1(x, \xi) = f_2(x, \xi)$ for μ -a.e. $x \in \Omega$ and for every $\xi \in \text{co}\{u_1(x), \dots, u_k(x)\}$.*

Proof. From the finiteness of $F(u_i, \Omega)$ for $i = 1, \dots, k$ and from property (iii), we deduce that $f_1(\cdot, u_i(\cdot)) < +\infty$, $f_2(\cdot, u_i(\cdot)) < +\infty$ μ -a.e. on Ω . The convexity of $f_1(x, \cdot)$ and $f_2(x, \cdot)$ then guarantees that $f_1(x, \cdot)$ and $f_2(x, \cdot)$ are finite on $\text{co}\{u_1(x), \dots, u_k(x)\}$ for μ -a.e. $x \in \Omega$. By Theorem 10.1 in [28], it follows that $f_1(x, \cdot)$ and $f_2(x, \cdot)$ restricted to $\text{ri co}\{u_1(x), \dots, u_k(x)\}$ are continuous for μ -a.e. $x \in \Omega$. By (iii), for every $\lambda \in \Sigma_k \cap \mathbf{Q}^k$ we have

$$\int_A f_1(x, u_\lambda(x)) d\mu = \int_A f_2(x, u_\lambda(x)) d\mu$$

for every $A \in \mathcal{A}(\Omega)$; hence, there exists a set $N \in \mathcal{B}(\Omega)$, $\mu(N) = 0$ such that $f_1(x, u_\lambda(x)) = f_2(x, u_\lambda(x))$ for every $x \in \Omega \setminus N$ and $\lambda \in \Sigma_k \cap \mathbf{Q}^k$. Since the functions $f_1(x, \cdot)$ and $f_2(x, \cdot)$ restricted to $\text{ri co}\{u_1(x), \dots, u_k(x)\}$ are continuous, we have $f_1(x, \xi) = f_2(x, \xi)$ for every $x \in \Omega \setminus N$ and $\xi \in \text{ri co}\{u_1(x), \dots, u_k(x)\}$. By the continuity along line segments ([28], Corollary 7.5.1) $f_1(x, \xi) = f_2(x, \xi)$ for every $\xi \in \text{co}\{u_1(x), \dots, u_k(x)\}$. □

4. Auxiliary lemmas

We collect here some results we shall use in the next section.

Lemma 4.1. *Let X be a separable metric space and let $F: X \rightarrow \mathbf{R}$ be lower semicontinuous. Then there exists a countable subset D of X with the following property: for every $x \in X$ there exists a sequence (x_h) in D converging to x and such that $(F(x_h))$ converges to $F(x)$.*

Proof. It is enough to take a countable dense subset E of the epigraph of F and consider as D the projection of E onto X . $\square$

Lemma 4.2. *Let $d \in \mathbf{N}$ and X be a subset of $\mathbf{R}^d$. Let H be a Lipschitz multivalued function from X to $\mathbf{R}^m$ with non-empty, compact and convex values. Then, there exists a sequence (h_j) of Lipschitz functions from X to $\mathbf{R}^m$ such that*

$$H(x) = \text{cl}\{h_j(x) : j \in \mathbf{N}\}$$

for every $x \in X$, where cl denotes the closure in $\mathbf{R}^m$.

Proof. Let (ξ_j) be a dense sequence in $\mathbf{R}^m$; for every $x \in X$, define $h_j(x) = P(\xi_j, H(x)) \in H(x)$, where P is the projection map given in Theorem 1.3. Since P and H are both lipschitzian, so is h_j .

Given $\xi \in H(x)$ and $\varepsilon > 0$ there exists $\xi_j \in \mathbf{R}^m$ such that $|\xi - \xi_j| < \varepsilon$. If L denotes a Lipschitz constant for P , then

$$|\xi - h_j(x)| = |P(\xi, H(x)) - P(\xi_j, H(x))| \leq L|\xi - \xi_j| < L\varepsilon;$$

we conclude that $\xi \in \text{cl}\{h_j(x) : j \in \mathbf{N}\}$. $\square$

We shall now state a result due to G. Bouchitté and M. Valadier concerning the commutativity property for the operations of integration and infimum. To this aim we need the following notion of C^1 -convexity which is essentially the notion of C^1 -stability introduced in [7].

Definition 4.3. Given a positive Radon measure λ on Ω and a set $\mathcal{H}$ of λ -measurable functions from Ω into $\mathbf{R}^m$, we say that $\mathcal{H}$ is C^1 -convex if for every finite family $(u_i)_{i \in I}$ of elements of $\mathcal{H}$ and for every family $(\alpha_i)_{i \in I}$ of non-negative functions of $C^1(\Omega) \cap W^{1,\infty}(\Omega)$ such that $\sum_i \alpha_i = 1$ in Ω , we have that $\sum_i \alpha_i u_i$ belongs to $\mathcal{H}$.

Let λ be a positive Radon measure on Ω and $\mathcal{H}$ be a family of λ -measurable functions from Ω to $\mathbf{R}^m$. Then, there exists a closed valued λ -measurable multifunction $\Gamma: \Omega \rightarrow \mathbf{R}^m$ (i.e., such that $\Gamma^{-1}(C) = \{x \in \Omega : \Gamma(x) \cap C \neq \emptyset\}$ is λ -measurable for every closed subset C of $\mathbf{R}^m$) with the following properties (see [29], Proposition 14):

- (i) for every $w \in \mathcal{H}$ we have $w(x) \in \Gamma(x)$ for λ -a.e. $x \in \Omega$;
- (ii) if $\Phi: \Omega \rightarrow \mathbf{R}^m$ is a closed valued λ -measurable multifunction such that for every $w \in \mathcal{H}$, $w(x) \in \Phi(x)$ for λ -a.e. $x \in \Omega$, then $\Gamma(x) \subseteq \Phi(x)$ for λ -a.e. $x \in \Omega$.

This multifunction Γ is unique up to λ -equivalence and will be denoted by $\lambda\text{-ess sup}_{w \in \mathcal{H}} \{w(\cdot)\}$.

The next theorem is taken from [7], Theorem 1.

Theorem 4.4. Let λ be a positive Radon measure on Ω and let $\mathcal{H}$ be a C^1 -convex family of λ -measurable functions from Ω into $\mathbf{R}^m$. Let $f: \Omega \times \mathbf{R}^m \rightarrow]-\infty, +\infty]$ be a $\mathcal{B}_\lambda(\Omega) \otimes \mathcal{B}(\mathbf{R}^m)$ -measurable function such that $f(x, \cdot)$ is convex on $\mathbf{R}^m$ for λ -a.e. $x \in \Omega$. Suppose that $f(\cdot, u(\cdot)) \in L^1(\Omega, \lambda)$ for every $u \in \mathcal{H}$ and let $\Gamma(x) = \lambda\text{-ess sup}_{u \in \mathcal{H}} \{u(x)\}$. Then

$$\inf_{u \in \mathcal{H}} \int_{\Omega} f(x, u(x)) d\lambda = \int_{\Omega} \inf_{z \in \Gamma(x)} f(x, z) d\lambda.$$

The following technical result, proven in [14], Lemma 4.2, will be crucial in the next section.

Lemma 4.5. Let (w_k) be a sequence of functions in $W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ converging in $L^\infty(\Omega, \mathbf{R}^m)$ to a function $w \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$. Then there exists a sequence (v_k) in $W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ such that $v_k(x) \in \text{co}\{w_1(x), \dots, w_k(x)\}$ for a.e. $x \in \Omega$ and (v_k) converges to w strongly in $W^{1,p}(\Omega, \mathbf{R}^m)$.

Lemma 4.6. *Let λ be a positive Borel measure on Ω . Let (γ_h) and γ be non-negative functions in $L^1(\Omega, \lambda)$ satisfying*

$$(4.1) \quad \gamma(x) \leq \liminf_{h \rightarrow \infty} \gamma_h(x) \quad \text{for } \lambda\text{-a.e. } x \in \Omega,$$

$$(4.2) \quad \int_{\Omega} \gamma \, d\lambda \geq \limsup_{h \rightarrow \infty} \int_{\Omega} \gamma_h \, d\lambda.$$

Then, (γ_h) converges to γ strongly in $L^1(\Omega, \lambda)$.

Proof. Let us note that, by the Fatou Lemma, (4.2) ensures that

$$(4.3) \quad \int_{\Omega} \gamma \, d\lambda = \lim_{h \rightarrow \infty} \int_{\Omega} \gamma_h \, d\lambda.$$

In view of (4.1) we have

$$\gamma \leq \liminf_{h \rightarrow \infty} (\gamma_h \wedge \gamma) \leq \limsup_{h \rightarrow \infty} (\gamma_h \wedge \gamma) \leq \gamma \quad \text{on } \Omega.$$

Thus, the dominated convergence theorem guarantees that $(\gamma_h \wedge \gamma)$ converges to γ in $L^1(\Omega, \lambda)$, and, in particular

$$(4.4) \quad \int_{\Omega} \gamma_h \wedge \gamma \, d\lambda \rightarrow \int_{\Omega} \gamma \, d\lambda.$$

By noticing that $\gamma_h + \gamma = (\gamma_h \wedge \gamma) + (\gamma_h \vee \gamma)$, (4.3) and (4.4) permit to conclude that

$$\int_{\Omega} \gamma_h \vee \gamma \, d\lambda \rightarrow \int_{\Omega} \gamma \, d\lambda;$$

hence $(\gamma_h \vee \gamma)$ converges to γ in $L^1(\Omega, \lambda)$, being $\gamma_h \vee \gamma \geq \gamma$. Now, the conclusion can be obtained by using again the relation $\gamma_h = (\gamma_h \wedge \gamma) + (\gamma_h \vee \gamma) - \gamma$ on Ω . $\square$

We conclude this section with a Dini-type lemma for which we refer, e.g., to [14], Lemma 4.3.

Lemma 4.7. *Let E be a compact subset of $\mathbf{R}^n$ and (H_k) be an increasing (with respect to inclusion) sequence of lower semicontinuous multifunctions from E to $\mathbf{R}^m$ with closed values. Let $u \in C^0(E, \mathbf{R}^m)$ such that*

$$u(x) \in \text{cl}\left(\bigcup_{k=1}^{\infty} H_k(x)\right) \quad \text{for every } x \in E.$$

Then, for every $r > 0$ there exists $h \in \mathbf{N}$ such that $B_r(u(x)) \cap H_k(x) \neq \emptyset$ for every $k \geq h$ and for every $x \in E$.

5. Integral representation on $W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$

The main result of this section is the integral representation of the functionals of the class $\mathcal{F}$ on the bounded functions of $W^{1,p}(\Omega, \mathbf{R}^m)$ (Theorem 5.4).

Given $F \in \mathcal{F}$, let us introduce the least closed valued multifunction having the elements of $\text{dom}F(\cdot, \Omega)$ among its selections.

Proposition 5.1. *Let $F \in \mathcal{F}$ and let A be an open subset of Ω with $\text{dom}F(\cdot, A) \neq \emptyset$. Then there exists a closed valued multifunction K_A from A to $\mathbf{R}^m$, unique up to sets of capacity zero, such that*

- (i) *for every $u \in \text{dom}F(\cdot, A)$ we have $u(x) \in K_A(x)$ for q.e. $x \in A$;*
- (ii) *if H is a closed valued multifunction from A to $\mathbf{R}^m$ such that for every $u \in \text{dom}F(\cdot, A)$ we have $u(x) \in H(x)$ for q.e. $x \in A$, then $K_A(x) \subseteq H(x)$ for q.e. $x \in A$.*

Moreover, K_A satisfies the following properties:

- (iii) *K_A is quasi lower semicontinuous and $K_A(x)$ is convex for q.e. $x \in A$;*
- (iv) *if (u_k) is a countable dense subset of $\text{dom}F(\cdot, A)$, then*

$$K_A(x) = \text{cl}\{u_k(x) : k \in \mathbf{N}\} = \text{cl}\left(\bigcup_{k=1}^{\infty} C_k(x)\right) \quad \text{for q.e. } x \in A,$$

where $C_k(x) = \text{co}\{u_1(x), \dots, u_k(x)\}$.

Proof. The same argument applied in Proposition 3.3 in [14] works now for the subset $\{u|_A : u \in \text{dom}F(\cdot, A)\}$ of $W^{1,p}(A, \mathbf{R}^m)$. $\square$

Remark 5.2. Let A and A' be open subsets of Ω , with $\text{dom}F(\cdot, A) \neq \emptyset$ and $\text{dom}F(\cdot, A') \neq \emptyset$. If K_A and $K_{A'}$ are the multifunctions given by the previous proposition, then $K_A = K_{A'}$ q.e. on $A \cap A'$.

It is enough to give the proof in the case $A' \subseteq A$. Since $\text{dom}F(\cdot, A) \subseteq \text{dom}F(\cdot, A')$, the inclusion $K_A(x) \subseteq K_{A'}(x)$ for q.e. $x \in A'$ follows immediately from property (i) satisfied by $K_{A'}$ and property (ii) applied to K_A and

$$H(x) = \begin{cases} K_{A'}(x), & \text{if } x \in A', \\ \mathbf{R}^m, & \text{if } x \in A \setminus A'. \end{cases}$$

To get the opposite inclusion let us choose $u_0 \in \text{dom}F(\cdot, A)$. Fix now $u \in \text{dom}F(\cdot, A')$ and $A'' \in \mathcal{A}(\Omega)$ with $A'' \subset\subset A'$. If φ is a function in $C_0^1(A')$, with $\varphi = 1$ on A'' and $0 \leq \varphi \leq 1$, by the C^1 -convexity and the locality property of F on open sets, we have

$$\begin{aligned} F(\varphi u + (1 - \varphi)u_0, A) &\leq F(\varphi u + (1 - \varphi)u_0, A') + F(\varphi u + (1 - \varphi)u_0, A \setminus \text{supp}\varphi) \\ &\leq F(u, A') + F(u_0, A') + F(u_0, A \setminus \text{supp}\varphi) < +\infty. \end{aligned}$$

Therefore, $\varphi u + (1 - \varphi)u_0 \in \text{dom}F(\cdot, A)$ so that $u(x) \in K_A(x)$ for q.e. $x \in A''$. By the arbitrariness of A'' we deduce that $u(x) \in K_A(x)$ for q.e. $x \in A'$. By applying property (ii) we conclude that $K_{A'}(x) \subseteq K_A(x)$ for q.e. $x \in A'$.

Lemma 5.3. *Let $s > 0$ and let T_s be the orthogonal projection onto the ball $\overline{B}_s(0)$ defined in Lemma 2.5. Then for every $u, v \in W^{1,p}(\Omega, \mathbf{R}^m)$ and $A \in \mathcal{A}(\Omega)$*

$$F(u + T_s \circ (v - u), A) \leq F(u, A) + F(v, A).$$

Proof. It is enough to consider the case $A \subset\subset \Omega$. Let $\varphi \in C_0^1(\Omega)$, $\varphi = 1$ on A , $0 \leq \varphi \leq 1$. By Lemma 3.6 in [14] there exists a sequence (ψ_h) of functions in $C^\infty(\mathbf{R}^n)$ such that $0 \leq \psi_h \leq 1$ and $(\psi_h \varphi(v - u))$ converges to $T_s \circ [\varphi(v - u)]$ weakly in $W^{1,p}(\Omega, \mathbf{R}^m)$ as h goes to ∞ . Since $F(\cdot, A)$ is weakly lower semicontinuous on $W^{1,p}(\Omega, \mathbf{R}^m)$ (recall that $F(\cdot, A)$ is convex) we get

$$\begin{aligned} F(u + T_s \circ [\varphi(v - u)], A) &\leq \liminf_{h \rightarrow \infty} F(u + \psi_h \varphi(v - u), A) \\ &\leq F(u, A) + F(v, A), \end{aligned}$$

where in the last estimate we have used the C^1 -convexity of F . Now the conclusion can be obtained by applying the locality property of F on open sets. $\square$

Theorem 5.4. *Let $F \in \mathcal{F}$ with $\text{dom}F(\cdot, \Omega) \cap L^\infty(\Omega, \mathbf{R}^m) \neq \emptyset$. Let ν_0 be the positive finite Borel measure introduced in Proposition 2.7 and let $K = K_\Omega$ be the closed valued multifunction from Ω to $\mathbf{R}^m$ given by Proposition 5.1 for $A = \Omega$. Then, there exist a positive finite Borel measure μ on Ω , absolutely continuous with respect to capacity, and a Borel function $f: \Omega \times \mathbf{R}^m \rightarrow [0, +\infty]$ with the following properties:*

(i) *for every $x \in \Omega$, the function $f(x, \cdot)$ is convex and lower semicontinuous on $\mathbf{R}^m$;*

(ii) for every $u \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ and for every $A \in \mathcal{A}(\Omega)$

$$F(u, A) = \begin{cases} \int_A f(x, u(x)) d\mu + \nu_0(A), & \text{if } u(x) \in K(x) \text{ for } q.e. x \in A, \\ +\infty, & \text{otherwise.} \end{cases}$$

Proof. Step 1. Let (u_i) be a sequence of functions in $\text{dom}F(\cdot, \Omega) \cap L^\infty(\Omega, \mathbf{R}^m)$ which will be specified in Step 2. We construct now the measure μ and the integrand f (see (5.3)) satisfying (i) and we prove that for every $A \in \mathcal{A}(\Omega)$ and for every $k \in \mathbf{N}$ we have

$$(5.1) \quad F(u, A) = \int_A f(x, u(x)) d\mu + \nu_0(A)$$

whenever $u \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ and $u(x) \in \text{co}\{u_1(x), \dots, u_k(x)\}$ for a.e. $x \in A$.

Fix quasi continuous Borel measurable representatives of (u_i) . For every $x \in \Omega$ define $C_k(x) = \text{co}\{u_1(x), \dots, u_k(x)\}$. By Theorem 3.7, for every $k \in \mathbf{N}$ there exist a positive finite Borel measure μ_k on Ω , absolutely continuous with respect to capacity, and a function $f_k: \Omega \times \mathbf{R}^m \rightarrow [0, +\infty]$ such that

- (a) for every $x \in \Omega$ the function $f_k(x, \cdot)$ is convex and lower semicontinuous on $\mathbf{R}^m$; moreover, the restriction of $f_k(x, \cdot)$ to $C_k(x)$ is continuous for μ_k -a.e. $x \in \Omega$;
- (b) f_k is $\widehat{\mathcal{B}}(\Omega) \otimes \mathcal{B}(\mathbf{R}^m)$ -measurable;
- (c) for every $u \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ such that $u(x) \in C_k(x)$ for a.e. $x \in \Omega$ the function $f_k(\cdot, u(\cdot))$ is μ -measurable on Ω and for every $A \in \mathcal{A}(\Omega)$

$$(5.2) \quad F(u, A) = \int_A f_k(x, u(x)) d\mu_k + \nu_0(A).$$

By a standard cut-off argument we obtain that (5.2) still holds if $u(x) \in C_k(x)$ for a.e. $x \in A$.

Let μ be a positive finite Borel measure on Ω absolutely continuous with respect to capacity and such that $\mu_k \ll \mu$ for every $k \in \mathbf{N}$ (for instance, take $\mu(B) = \sum_{k=1}^{\infty} 2^{-k} \frac{\mu_k(B)}{\mu_k(\Omega)}$ for every $B \in \mathcal{B}(\Omega)$). Define

$$g_k(x, \xi) = f_k(x, \xi) \frac{d\mu_k}{d\mu}(x),$$

where $d\mu_k/d\mu$ is a fixed (μ -measurable) representative of the Radon-Nikodym derivative of μ_k with respect to μ . By Proposition 3.8, there exists a set $N \in \mathcal{B}(\Omega)$ with $\mu(N) = 0$

such that $g_k(x, \xi) = g_{k+1}(x, \xi)$ for every $k \in \mathbf{N}$, $x \in \Omega \setminus N$ and $\xi \in C_k(x)$. Hence, we can define $g: \Omega \times \mathbf{R}^m \rightarrow [0, +\infty]$ as

$$g(x, \xi) = \begin{cases} g_k(x, \xi), & \text{if } x \in \Omega \setminus N \text{ and } \xi \in C_k(x) \text{ for some } k \in \mathbf{N}, \\ +\infty, & \text{otherwise.} \end{cases}$$

Since $C_k(x) = \{\sum_{i=1}^k \lambda^i u_i(x) : \lambda \in \Sigma_k\}$, by Theorem III.9 and Proposition III.13 in [10], the graph of C_k belongs to $\mathcal{B}(\Omega) \otimes \mathcal{B}(\mathbf{R}^m)$. Recalling the definition of g_k it follows that g is $\widehat{\mathcal{B}}(\Omega) \otimes \mathcal{B}(\mathbf{R}^m)$ -measurable. An easy check gives the convexity of $g(x, \cdot)$ on $\mathbf{R}^m$ for every $x \in \Omega$.

Now, for every $x \in \Omega$ let us set $h(x, \cdot) = \text{sc}^- g(x, \cdot)$, where $\text{sc}^- g(x, \cdot)$ denotes the lower semicontinuous envelope of $g(x, \cdot)$. It turns out that

$$h(x, \xi) = \sup_{s \in \mathbf{N}} g_s(x, \xi),$$

where $g_s(x, \xi) = \inf_{\eta \in \mathbf{R}^m} [g(x, \eta) + s|\xi - \eta|]$. By Remark 3.6, for every $s \in \mathbf{N}$ there exists a set $Z_s \in \mathcal{B}(\Omega)$, with $\mu(Z_s) = 0$, such that $g_s|_{(\Omega \setminus Z_s) \times \mathbf{R}^m}$ is a Borel function. Set $Z = \bigcup_{s \in \mathbf{N}} Z_s$; then $\mu(Z) = 0$ and $h|_{(\Omega \setminus Z) \times \mathbf{R}^m}$ is Borel measurable. Now we are in a position to define the function f as

$$(5.3) \quad f(x, \cdot) = \begin{cases} h(x, \cdot) = \text{sc}^- g(x, \cdot) & \text{if } x \in \Omega \setminus Z \\ 0 & \text{if } x \in Z. \end{cases}$$

Then, f is a Borel function on $\Omega \times \mathbf{R}^m$ and satisfies (i) (see [28], Theorem 7.4).

Let us prove that for μ -a.e. $x \in \Omega$

$$(5.4) \quad f(x, \cdot) = g_k(x, \cdot) \quad \text{on } C_k(x).$$

Let us fix $x \in \Omega \setminus (N \cup Z)$ and $k \in \mathbf{N}$. Let $H(x)$ be the affine hull of $\bigcup_{k=1}^{\infty} C_k(x)$. As the sequence $(C_k(x))$ is increasing, there exists $h \geq k$ such that the interior of $C_h(x)$ relative to $H(x)$ is non-empty. Since $g(x, \cdot) = g_h(x, \cdot)$ on $C_h(x)$, and the restriction of $g_h(x, \cdot)$ to $C_h(x)$ is continuous, we have $f(x, \cdot) = g_h(x, \cdot)$ on the interior of $C_h(x)$ relative to $H(x)$. As $C_h(x)$ is a polytope, the restriction of $f(x, \cdot)$ to $C_h(x)$ is continuous (see [28], Theorem 10.2), hence $f(x, \cdot) = g_h(x, \cdot)$ on $C_h(x)$. Since $C_k(x) \subseteq C_h(x)$ and $g_k(x, \cdot) = g_h(x, \cdot)$ on $C_k(x)$, we conclude that (5.4) is satisfied.

Let us now prove (5.1). Fix $k \in \mathbf{N}$, $A \in \mathcal{A}(\Omega)$ and $u \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ with $u(x) \in C_k(x)$ for a.e. $x \in A$. For every $0 < \sigma < 1$ let us define $u_\sigma = \sigma u + (1 - \sigma)u_0$,

with $u_0 = \frac{1}{k} \sum_{i=1}^k u_i$. Then $u_\sigma(x) \in \text{ri}C_k(x)$ for q.e. $x \in A$. Therefore, by (5.4) and the integral representation formula (5.2) satisfied by f_k , we get

$$F(u_\sigma, A) = \int_A f(x, u_\sigma(x)) d\mu + \nu_0(A).$$

Since every lower semicontinuous proper convex function is continuous along line segments (see [28], Corollary 7.5.1) it turns out that

$$\begin{aligned} \lim_{\sigma \rightarrow 1^-} F(u_\sigma, A) &= F(u, A) \\ \lim_{\sigma \rightarrow 1^-} \int_A f(x, u_\sigma(x)) d\mu &= \int_A f(x, u(x)) d\mu \quad \text{for every } x \in \Omega. \end{aligned}$$

We thus obtain (5.1).

Step 2. We choose now a suitable sequence (u_i) , dense in $\text{dom}F(\cdot, \Omega) \cap L^\infty(\Omega, \mathbf{R}^m)$, to which Step 1 will be applied.

Let $\mathcal{D}$ be a countable base for the open subsets of Ω , closed under finite unions. For every $A \in \mathcal{D}$ we can apply Lemma 4.1 to $F(\cdot, A)$ on $\text{dom}F(\cdot, \Omega) \cap L^\infty(\Omega, \mathbf{R}^m)$ (with the $W^{1,p}(\Omega, \mathbf{R}^m)$ topology); this yields the existence of a set $\mathcal{G}_A \subseteq \text{dom}F(\cdot, \Omega) \cap L^\infty(\Omega, \mathbf{R}^m)$ such that for every $u \in \text{dom}F(\cdot, \Omega) \cap L^\infty(\Omega, \mathbf{R}^m)$ there exists a sequence (u_h) in $\mathcal{G}_A$ satisfying

$$\begin{aligned} u_h &\rightarrow u \quad \text{strongly in } W^{1,p}(\Omega, \mathbf{R}^m), \\ F(u_h, A) &\rightarrow F(u, A) \quad \text{in } \mathbf{R}. \end{aligned}$$

Let (u_i) be an enumeration of $\bigcup_{A \in \mathcal{D}} \mathcal{G}_A$; starting from (u_i) we then construct by means of Step 1 a Borel function $f: \Omega \times \mathbf{R}^m \rightarrow [0, +\infty]$ satisfying (i) and (5.1).

Step 3. Let us prove that for every $u \in \text{dom}F(\cdot, \Omega) \cap L^\infty(\Omega, \mathbf{R}^m)$ and for every $A \in \mathcal{A}(\Omega)$

$$(5.5) \quad F(u, A) \geq \int_A f(x, u(x)) d\mu + \nu_0(A).$$

Fix $u \in \text{dom}F(\cdot, \Omega) \cap L^\infty(\Omega, \mathbf{R}^m)$ and $A \in \mathcal{D}$. By Step 2 it is possible to extract a sequence (u_{i_h}) from $\{u_i : i \in \mathbf{N}\}$ such that

$$\begin{aligned} u_{i_h} &\rightarrow u \quad \text{q.e. in } \Omega \quad (\text{hence } \mu\text{-a.e.}), \\ F(u_{i_h}, A) &\rightarrow F(u, A) \quad \text{in } \mathbf{R}. \end{aligned}$$

Therefore, by (5.1)

$$F(u, A) = \lim_{h \rightarrow \infty} \int_A f(x, u_{i_h}(x)) d\mu + \nu_0(A);$$

by the Fatou Lemma and the lower semicontinuity of $f(x, \cdot)$ we get (5.5) for every $A \in \mathcal{D}$. The result for an arbitrary $A \in \mathcal{A}(\Omega)$ can be obtained by approximation, since each $A \in \mathcal{A}(\Omega)$ is the union of an increasing sequence of elements of $\mathcal{D}$ (recall that $\mathcal{D}$ is closed under finite unions).

Step 4. It is now easy to prove that for every $u \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ and $A \in \mathcal{A}(\Omega)$ the inequality (5.5) holds.

Given $A \in \mathcal{A}(\Omega)$ and $u \in \text{dom}F(\cdot, A) \cap L^\infty(\Omega, \mathbf{R}^m)$, let $G \subset\subset A$ and $\varphi \in C_0^1(A)$ with $\varphi = 1$ on G and $0 \leq \varphi \leq 1$. Set $u_\varphi = \varphi u + (1 - \varphi)w$, where w belongs to $\text{dom}F(\cdot, \Omega) \cap L^\infty(\Omega, \mathbf{R}^m)$, which is non-empty by assumption. By the convexity and the locality property of F we have $F(u_\varphi, \Omega) < +\infty$. Therefore, Step 3 applied to u_φ yields

$$F(u_\varphi, G) \geq \int_G f(x, u_\varphi(x)) d\mu + \nu_0(G);$$

since $\varphi = 1$ on G we get

$$F(u, G) \geq \int_G f(x, u(x)) d\mu + \nu_0(G).$$

As $G \subset\subset A$ is arbitrary, the conclusion is easily achieved.

Step 5. Let $K = K_\Omega$ be the closed valued multifunction from Ω to $\mathbf{R}^m$ given by Proposition 5.1 for $A = \Omega$. The aim is now to prove that for every $A \in \mathcal{A}(\Omega)$ and $u \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ such that $u(x) \in K(x)$ for q.e. $x \in A$, we have

$$(5.6) \quad F(u, A) \leq \int_A f(x, u(x)) d\mu + \nu_0(A).$$

Recall that for every $k \in \mathbf{N}$ and $x \in \Omega$ we have $C_k(x) = \text{co}\{u_1(x), \dots, u_k(x)\}$, where (u_i) is the sequence given in Step 2. By Lemmas 2.5 and 5.3, $\text{dom}F(\cdot, \Omega) \cap L^\infty(\Omega, \mathbf{R}^m)$, which is non-empty by assumption, is dense in $\text{dom}F(\cdot, \Omega)$. Hence, (u_i) is dense in $\text{dom}F(\cdot, \Omega)$ and, by Proposition 5.1, $K(x) = \text{cl}(\bigcup_{k=1}^\infty C_k(x))$ for q.e. $x \in \Omega$.

Fix $u \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ and $A \in \mathcal{A}(\Omega)$ such that $u(x) \in K(x)$ for q.e. $x \in A$. Clearly, we can assume that the right-hand side in (5.6) is finite. Moreover, we

can consider open sets $A \subset\subset \Omega$ with smooth boundary, so that there exists an extension operator $W^{1,p}(A) \rightarrow W^{1,p}(\Omega)$.

In a first moment we work with the additional assumption that $u(x)$ is in the closure of $\bigcup_{k=1}^{\infty} C_k(x)$ “uniformly” for $x \in A$; more precisely, given a sequence (r_h) of positive numbers decreasing to 0, we require that for every $h \in \mathbf{N}$ there exists $n_h \in \mathbf{N}$ such that

$$(5.7) \quad B_{r_h/2}(u(x)) \cap C_k(x) \neq \emptyset \quad \text{for q.e. } x \in A$$

for every $k \geq n_h$.

To achieve (5.6) we look for a sequence (v_h) of functions in $W^{1,p}(A, \mathbf{R}^m) \cap L^\infty(A, \mathbf{R}^m)$ and a strictly increasing sequence (k_h) of positive integers such that $v_h \in C_{k_h}(x)$ for q.e. $x \in A$, (v_h) converges to u in $W^{1,p}(A, \mathbf{R}^m)$ and

$$(5.8) \quad \limsup_{h \rightarrow \infty} \int_A f(x, v_h(x)) d\mu \leq \int_A f(x, u(x)) d\mu.$$

Indeed, as ∂A is smooth, we can assume that (v_h) is a sequence in $W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ converging in $W^{1,p}(\Omega, \mathbf{R}^m)$ to a function v such that $v = u$ a.e. on A . By the lower semicontinuity of $F(\cdot, A)$ and the integral representation (5.1) obtained in Step 1, we can then conclude

$$\begin{aligned} F(u, A) &= F(v, A) \leq \liminf_{h \rightarrow \infty} F(v_h, A) \\ &\leq \limsup_{h \rightarrow \infty} \left(\int_A f(x, v_h(x)) d\mu + \nu_0(A) \right) \leq \int_A f(x, u(x)) d\mu + \nu_0(A). \end{aligned}$$

Let us first construct a sequence (w_h) of functions in $W^{1,p}(A, \mathbf{R}^m) \cap L^\infty(A, \mathbf{R}^m)$ and a strictly increasing sequence (k_h) of positive integers, with the following properties:

$$(5.9) \quad \begin{aligned} w_h(x) &\in C_{k_h}(x) && \text{for q.e. } x \in A \\ w_h &\rightarrow u && \text{uniformly on } A \\ \limsup_{h \rightarrow \infty} \int_A f(x, w_h(x)) d\mu &\leq \int_A f(x, u(x)) d\mu. \end{aligned}$$

To this aim let us prove that for every $h \in \mathbf{N}$ there exists $k_h \in \mathbf{N}$ such that

$$(5.10) \quad \inf_{w \in \mathcal{H}_k^h} \int_A f(x, w(x)) d\mu < \int_A f(x, u(x)) d\mu + r_h,$$

for every $k \geq k_h$, where

$$\mathcal{H}_k^h = \{w \in W^{1,p}(A, \mathbf{R}^m) \cap L^\infty(A, \mathbf{R}^m) : w(x) \in \overline{B}_{r_h}(u(x)) \cap C_k(x) \text{ for q.e. } x \in A\}.$$

Let us fix $h \in \mathbf{N}$ and let n_h be as in (5.7). For every fixed $k \geq n_h$ we want to apply Theorem 4.4 to the set $\mathcal{H}_k^h$. For this purpose let us verify that

$$(5.11) \quad \mu\text{-ess sup}_{w \in \mathcal{H}_k^h} \{w(x)\} = \overline{B}_{r_h}(u(x)) \cap C_k(x) \quad \text{for q.e. } x \in A.$$

Let $\Xi = \{(\xi, \xi_1, \dots, \xi_k) \in (\mathbf{R}^m)^{k+1} : d(\xi, \text{co}\{\xi_1, \dots, \xi_k\}) \leq r_h/2\}$ and let H be the multivalued function from Ξ to $\mathbf{R}^m$ defined by

$$H(\xi, \xi_1, \dots, \xi_k) = \overline{B}_{r_h}(\xi) \cap \text{co}\{\xi_1, \dots, \xi_k\}.$$

By Theorem 1 in [24], H is lipschitzian. Hence, we can apply Lemma 4.2 to H obtaining a sequence (h_j) of Lipschitz functions from Ξ to $\mathbf{R}^m$ such that

$$H(\xi, \xi_1, \dots, \xi_k) = \text{cl}\left(\bigcup_{j=1}^{\infty} \{h_j(\xi, \xi_1, \dots, \xi_k)\}\right).$$

Since $B_{r_h/2}(u(x)) \cap C_k(x) \neq \emptyset$ for q.e. $x \in A$, we can define $z_j = h_j(u, u_1, \dots, u_k)$ q.e. on A for every $j \in \mathbf{N}$. By Remark 1.6., $z_j \in W^{1,p}(A, \mathbf{R}^m) \cap L^\infty(A, \mathbf{R}^m)$. Thus $z_j \in \mathcal{H}_k^h$, and

$$\overline{B}_{r_h}(u(x)) \cap C_k(x) = H(u(x), u_1(x), \dots, u_k(x)) = \text{cl}\left(\bigcup_{j=1}^{\infty} \{z_j(x)\}\right)$$

for q.e. $x \in A$. Hence, (5.11) holds.

Moreover, since every $w \in \mathcal{H}_k^h$ is a convex combination of $u_1, \dots, u_k$, we have

$$\int_A f(x, w(x)) d\mu \leq \int_A \sum_{i=1}^k f(x, u_i(x)) d\mu < +\infty.$$

We can now apply Theorem 4.4; by (5.11), for every $k \geq n_h$ we have

$$(5.12) \quad \inf_{w \in \mathcal{H}_k^h} \int_A f(x, w(x)) d\mu = \int_A \inf_{\xi \in C_k^h(x)} f(x, \xi) d\mu,$$

where $C_k^h(x) = \overline{B}_{r_h}(u(x)) \cap C_k(x)$. Since $u(x) \in K(x)$ for q.e. $x \in A$, in view of the continuity property along line segments for a proper, lower semicontinuous convex function, for q.e. $x \in A$ we can approximate $u(x)$ by a sequence $(\xi_k(x))$ in $\text{ri}K(x)$ such that

$$f(x, u(x)) = \lim_{k \rightarrow \infty} f(x, \xi_k(x)).$$

As $\text{ri}K(x) \subseteq \bigcup_{k=1}^{\infty} C_k(x)$ (see [28], Theorem 6.3), we can suppose that $\xi_k(x) \in C_k(x)$ for every $k \in \mathbf{N}$. Thus, for every $h \in \mathbf{N}$ we have

$$(5.13) \quad \inf_{k \in \mathbf{N}} \inf_{\xi \in C_k^h(x)} f(x, \xi) \leq f(x, u(x)) \quad \text{for q.e. } x \in A.$$

Moreover, for every $k \geq n_h$ the set $C_k^h(x)$ is non-empty for q.e. $x \in A$. Therefore, the convexity of f ensures that

$$\int_A \inf_{\xi \in C_k^h(x)} f(x, \xi) d\mu \leq \int_A \sum_{i=1}^k f(x, u_i(x)) d\mu < +\infty;$$

by (5.13) and by the monotone convergence theorem it follows that

$$\inf_{k \in \mathbf{N}} \int_A \inf_{\xi \in C_k^h(x)} f(x, \xi) d\mu \leq \int_A f(x, u(x)) d\mu.$$

This inequality, together with (5.12), proves (5.10).

Let (k_h) be the sequence given in (5.10) which we can assume to be strictly increasing. For every $h \in \mathbf{N}$, by (5.10) there exists a function $w_h \in W^{1,p}(A, \mathbf{R}^m) \cap L^\infty(A, \mathbf{R}^m)$ such that

$$\begin{aligned} w_h(x) &\in \overline{B}_{r_h}(u(x)) \cap C_{k_h}(x) \quad \text{for q.e. } x \in A, \\ \int_A f(x, w_h(x)) d\mu &\leq \int_A f(x, u(x)) d\mu + r_h. \end{aligned}$$

It is easy to verify that (w_h) satisfies the properties in (5.9).

Let us set now $\gamma_h = f(\cdot, w_h(\cdot))$ and $\gamma = f(\cdot, u(\cdot))$. We claim that

$$(5.14) \quad \gamma_h \rightarrow \gamma \quad \text{strongly in } L^1(A, \mu).$$

Indeed, as (w_h) converges to u q.e. on A , by the lower semicontinuity of $f(x, \cdot)$ we get $\gamma(x) \leq \liminf_{h \rightarrow \infty} \gamma_h(x)$. By (5.9) and Lemma 4.6, it follows that (γ_h) converges to γ in the strong topology of $L^1(A, \mu)$.

In view of (5.14) it is not restrictive to assume that for every $h \in \mathbf{N}$

$$(5.15) \quad \int_A |\gamma_h - \gamma| d\mu \leq \frac{1}{2^h}.$$

At this point let us apply Lemma 4.5 to the sequence $(w_j)_{j \geq h}$ for every $h \in \mathbf{N}$. We obtain a sequence $(v_{h,j})_{j \geq h}$ of functions in $W^{1,p}(A, \mathbf{R}^m) \cap L^\infty(A, \mathbf{R}^m)$ such that

$$\begin{aligned} v_{h,j}(x) &\in \text{co}\{w_h(x), w_{h+1}(x), \dots, w_j(x)\} \quad \text{for q.e. } x \in A, \\ v_{h,j} &\rightarrow u \quad \text{strongly in } W^{1,p}(A, \mathbf{R}^m) \text{ as } j \rightarrow \infty. \end{aligned}$$

By a standard argument we can find a strictly increasing sequence (j_h) of positive integers such that (v_{h,j_h}) converges to u in $W^{1,p}(A, \mathbf{R}^m)$. Define $v_h = v_{h,j_h}$ for every $h \in \mathbf{N}$. Then $v_h \in W^{1,p}(A, \mathbf{R}^m) \cap L^\infty(A, \mathbf{R}^m)$ and

$$\begin{aligned} v_h(x) &\in \text{co}\{w_h(x), \dots, w_{j_h}(x)\} \quad \text{for q.e. } x \in A, \\ v_h &\rightarrow u \quad \text{strongly in } W^{1,p}(A, \mathbf{R}^m). \end{aligned}$$

In particular, a suitable sequence (k_h) exists such that $v_h(x) \in C_{k_h}(x)$ for q.e. $x \in A$. Now we only need to verify that (5.8) holds for the sequence (v_h) just obtained. By Lemma 3.2 we can write $v_h(x) = \sum_{i=h}^{k_h} \psi_h^i(x) w_i(x)$ for q.e. $x \in A$, where $\psi_h: A \rightarrow \Sigma_{k_h-h+1}$ are μ -measurable. Let us now make use of the convexity of f , together with (5.15):

$$\begin{aligned} \int_A f(x, v_h(x)) d\mu &\leq \sum_{i=h}^{k_h} \int_A \psi_h^i(x) |\gamma_i(x) - \gamma(x)| d\mu + \int_A \gamma(x) d\mu \\ &\leq \sum_{i=h}^{k_h} \frac{1}{2^i} + \int_A \gamma(x) d\mu \leq \int_A \gamma(x) d\mu + \frac{1}{2^h}. \end{aligned}$$

This implies

$$\limsup_{h \rightarrow \infty} \int_A f(x, v_h(x)) d\mu \leq \int_A f(x, u(x)) d\mu.$$

Finally, let us remove the additional assumption (5.7). Fix $G \subset\subset A$ and a sequence (r_h) of positive real numbers decreasing to 0. For every $\varepsilon > 0$ there exists an open set $A_\varepsilon \subseteq \Omega$, with $\text{cap}(A_\varepsilon, \Omega) < \varepsilon$, such that $u_i|_{\Omega \setminus A_\varepsilon}$ and $u|_{\Omega \setminus A_\varepsilon}$ are continuous for every $i \in \mathbf{N}$. In particular, the multifunction C_k is continuous on $\Omega \setminus A_\varepsilon$ with respect

to the Hausdorff metric. By Lemma 4.7 for every $h \in \mathbf{N}$ there exists $n_h^\varepsilon \in \mathbf{N}$ such that $B_{r_h/2}(u(x)) \cap C_k(x) \neq \emptyset$ for every $k \geq n_h^\varepsilon$ and for every $x \in G \setminus A_\varepsilon$. Let z_ε be the capacitary potential of A_ε and $u_\varepsilon = (1 - z_\varepsilon)u + z_\varepsilon u_1$, where u_1 is the first term of the sequence (u_i) . Then one can easily check that (u_ε) converges to u in $W^{1,p}(\Omega, \mathbf{R}^m)$, that $u_\varepsilon \in K(x)$ for q.e. $x \in G$, and that for every $h \in \mathbf{N}$ there exists $n_h^\varepsilon \in \mathbf{N}$ such that $B_{r_h/2}(u_\varepsilon(x)) \cap C_k(x) \neq \emptyset$ for q.e. $x \in G$ and for every $k \geq n_h^\varepsilon$. Therefore we can apply the previous result for u_ε and G in place of u and A ; this gives

$$F(u_\varepsilon, G) \leq \int_G f(x, u_\varepsilon(x)) d\mu + \nu_0(G).$$

Since

$$\int_G f(x, u_\varepsilon(x)) d\mu \leq \int_A [(1 - z_\varepsilon(x))f(x, u(x)) + z_\varepsilon(x)f(x, u_1(x))] d\mu$$

and $\int_A f(x, u(x)) d\mu < +\infty$ by assumption, the lower semicontinuity of F and the dominated convergence theorem imply

$$F(u, G) \leq \liminf_{\varepsilon \rightarrow 0^+} F(u_\varepsilon, G) \leq \int_A f(x, u(x)) d\mu + \nu_0(A).$$

Taking the supremum for $G \subset\subset A$ we get (5.6).

Step 6. In view of Step 4 and Step 5 we get

$$F(u, A) = \int_A f(x, u(x)) d\mu + \nu_0(A)$$

for every $u \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ with $u(x) \in K(x)$ for q.e. $x \in A$. Property (ii) now follows by taking into account that if $u \in \text{dom} F(\cdot, A)$ then $u(x) \in K(x)$ for q.e. $x \in A$ by Remark 5.2. $\square$

6. Integral representation on $W^{1,p}(\Omega, \mathbf{R}^m)$

We now eliminate (Theorem 6.1) the restrictive condition $u \in L^\infty(\Omega, \mathbf{R}^m)$ considered in the previous section. Furthermore, Proposition 6.3 will allow us to treat in a unified way both cases of the representation formula established in Theorem 5.4(ii). Thus, we achieve (Theorem 6.5) the conclusive integral representation theorem, which is the main result of the paper.

Given $F \in \mathcal{F}$ let us define

$$(6.1) \quad \bar{\nu}(B) = \inf\{F(u, B) : u \in W^{1,p}(\Omega, \mathbf{R}^m)\},$$

for every $B \in \mathcal{B}(\Omega)$. It is easily seen that the proof of Proposition 2.7 still works for the set function $\bar{\nu}$ on every $\Omega' \in \mathcal{A}(\Omega)$ with $\text{dom}F(\cdot, \Omega') \neq \emptyset$. Therefore, on such sets, $\bar{\nu}$ is a positive finite Borel measure.

Theorem 6.1. *Let $F \in \mathcal{F}$ and assume $\text{dom}F(\cdot, \Omega) \neq \emptyset$. Then the conclusions of Theorem 5.4 still hold with u (in item (ii)) ranging all over $W^{1,p}(\Omega, \mathbf{R}^m)$ and ν_0 replaced by the measure $\bar{\nu}$ defined in (6.1).*

Proof. For every $v \in W^{1,p}(\Omega, \mathbf{R}^m)$ and for every $B \in \mathcal{B}(\Omega)$ let us define

$$\begin{aligned} X_v &= \{u \in W^{1,p}(\Omega, \mathbf{R}^m) : u - v \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)\}, \\ \nu_v(B) &= \inf\{F(u, B) : u \in X_v\}. \end{aligned}$$

By a suitable application of Theorem 5.4, it turns out that for every $v \in \text{dom}F(\cdot, \Omega)$ there exist a positive finite Borel measure μ_v on Ω , absolutely continuous with respect to capacity, and a Borel function $f_v : \Omega \times \mathbf{R}^m \rightarrow [0, +\infty]$ such that

- (i) for every $x \in \Omega$ the function $f_v(x, \cdot)$ is convex and lower semicontinuous on $\mathbf{R}^m$;
- (ii) for every $u \in X_v$ and for every $A \in \mathcal{A}(\Omega)$

$$(6.2) \quad F(u, A) = \begin{cases} \int_A f_v(x, u(x)) d\mu_v + \nu_v(A), & \text{if } u(x) \in K(x) \text{ for q.e. } x \in A, \\ +\infty, & \text{otherwise,} \end{cases}$$

where $K = K_\Omega$ is the closed valued multifunction from Ω to $\mathbf{R}^m$ given by Proposition 5.1 for $A = \Omega$.

Step 1. Let us show first that for every $v \in \text{dom}F(\cdot, \Omega)$, $u \in W^{1,p}(\Omega, \mathbf{R}^m)$, and $A \in \mathcal{A}(\Omega)$ we have

$$(6.3) \quad F(u, A) < +\infty \quad \text{if and only if} \quad \begin{cases} u(x) \in K(x) & \text{for q.e. } x \in A, \\ \int_A f_v(x, u(x)) d\mu_v + \nu_v(A) < +\infty. \end{cases}$$

By the definition of K and Remark 5.2, if $F(u, A) < +\infty$ then $u(x) \in K(x)$ for q.e. $x \in A$. Hence, let us assume that $u(x) \in K(x)$ for q.e. $x \in A$ and prove that $F(u, A) < +\infty$ if and only if $\int_A f_v(x, u(x)) d\mu_v + \nu_v(A) < +\infty$.

For every $k \in \mathbf{N}$, let $T_k: \mathbf{R}^m \rightarrow \mathbf{R}^m$ be the orthogonal projection onto the ball $\overline{B}_k(0)$; by Lemma 2.5, for every $w \in W^{1,p}(\Omega, \mathbf{R}^m)$ the function $T_k \circ w$ belongs to $W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$, and the sequence $(T_k \circ w)$ converges to w in the strong topology of $W^{1,p}(\Omega, \mathbf{R}^m)$ as k tends to ∞ .

For every $k \in \mathbf{N}$ let us set $u_k = v + T_k \circ (u - v)$. By (6.2) we have

$$(6.4) \quad F(u_k, A) = \int_A f_v(x, u_k(x)) d\mu_v + \nu_v(A).$$

Assume now $F(u, A) < +\infty$. By (6.4) and Lemma 5.3 we have

$$\int_A f_v(x, u_k(x)) d\mu_v + \nu_v(A) \leq F(u, A) + F(v, A).$$

Since, up to a subsequence, (u_k) converges to u q.e. on Ω , the Fatou Lemma and the lower semicontinuity of $f_v(x, \cdot)$ ensure that

$$\int_A f_v(x, u(x)) d\mu_v + \nu_v(A) \leq F(v, A) + F(u, A) < +\infty.$$

Conversely, assume $\int_A f_v(x, u(x)) d\mu_v + \nu_v(A) < +\infty$. For every $k \in \mathbf{N}$, by (6.2), (6.4), and by the convexity of $f_v(x, \cdot)$ we get

$$(6.5) \quad \begin{aligned} F(u_k, A) &\leq \int_A f_v(x, u(x)) d\mu_v + \int_A f_v(x, v(x)) d\mu_v + \nu_v(A) = \\ &= \int_A f_v(x, u(x)) d\mu_v + F(v, A). \end{aligned}$$

Hence, by the lower semicontinuity of $F(\cdot, A)$ we conclude that $F(u, A) < +\infty$.

Step 2. Let us fix $A \in \mathcal{A}(\Omega)$ and $u, v \in \text{dom} F(\cdot, \Omega)$. We claim that

$$(6.6) \quad F(u, A) = \int_A f_v(x, u(x)) d\mu_v + \nu_v(A).$$

Let us show first that for every $w \in X_u$ with $w(x) \in \text{co}\{u(x), v(x)\}$ for q.e. $x \in \Omega$, we have

$$(6.7) \quad F(w, A) \leq \int_A f_v(x, w(x)) d\mu_v + \nu_v(A).$$

Let us fix $w \in X_u$ with $w(x) \in \text{co}\{u(x), v(x)\}$ for q.e. $x \in \Omega$, and let $u_k = v + T_k \circ (w - v)$ for $k \in \mathbf{N}$. By the lower semicontinuity of $F(\cdot, A)$ and by (6.2)

$$(6.8) \quad F(w, A) \leq \liminf_{k \rightarrow \infty} F(u_k, A) = \liminf_{k \rightarrow \infty} \int_A f_v(x, u_k(x)) d\mu_v + \nu_v(A).$$

Note that (u_k) converges to w q.e. on Ω . Since $u_k(x)$ is on the segment with endpoints $u(x)$ and $v(x)$, by the convexity of f_v it turns out that $f_v(x, u_k(x)) \leq f_v(x, v(x)) + f_v(x, u(x))$ for q.e. $x \in \Omega$. From (6.3) we have $\int_A f_v(x, v(x)) d\mu_v < +\infty$ and $\int_A f_v(x, u(x)) d\mu_v < +\infty$. Hence, by the continuity property of $f_v(x, \cdot)$ along line segments ([28], Corollary 7.5.1) and the dominated convergence theorem

$$(6.9) \quad \lim_{k \rightarrow \infty} \int_A f_v(x, u_k(x)) d\mu_v = \int_A f_v(x, w(x)) d\mu_v.$$

In view of (6.8), this implies (6.7).

From (6.7) with $w = u$ we obtain

$$F(u, A) \leq \int_A f_v(x, u(x)) d\mu_v + \nu_v(A).$$

Let us now prove the opposite inequality.

If in (6.7) we apply (6.2) to represent $F(w, A)$ we obtain

$$\int_A f_u(x, w(x)) d\mu_u + \nu_u(A) \leq \int_A f_v(x, w(x)) d\mu_v + \nu_v(A).$$

By exchanging now the roles of u and v we obtain that for every $w \in X_v$ with $w(x) \in \text{co}\{u(x), v(x)\}$ for q.e. $x \in \Omega$

$$\int_A f_v(x, w(x)) d\mu_v + \nu_v(A) \leq \int_A f_u(x, w(x)) d\mu_u + \nu_u(A).$$

Now, if we take $w = v + T_k \circ (u - v)$ and argue as for (6.9), by the Fatou Lemma we get

$$\begin{aligned} \int_A f_v(x, u(x)) d\mu_v + \nu_v(A) &\leq \liminf_{k \rightarrow \infty} \int_A f_u(x, v(x) + T_k(u(x) - v(x))) d\mu_u + \nu_u(A) \\ &= \int_A f_u(x, u(x)) d\mu_u + \nu_u(A) = F(u, A). \end{aligned}$$

Step 3. For every $v \in \text{dom}F(\cdot, \Omega)$, $A \in \mathcal{A}(\Omega)$, and $u \in \text{dom}F(\cdot, A)$ it turns out that

$$(6.10) \quad F(u, A) = \int_A f_v(x, u(x)) d\mu_v + \nu_v(A).$$

This follows by applying the same argument used in Step 4 of Theorem 5.4.

Step 4. Since $\text{dom}F(\cdot, \Omega) \neq \emptyset$, there exists a function v for which (6.10) holds for every $A \in \mathcal{A}(\Omega)$ and $u \in \text{dom}F(\cdot, A)$. Finally, we obtain that for every $u \in W^{1,p}(\Omega, \mathbf{R}^m)$ and $A \in \mathcal{A}(\Omega)$

$$(6.11) \quad F(u, A) = \begin{cases} \int_A f_v(x, u(x)) d\mu_v + \nu_v(A), & \text{if } u(x) \in K(x) \text{ for q.e. } x \in A, \\ +\infty, & \text{otherwise.} \end{cases}$$

Indeed, if $u(x) \in K(x)$ for q.e. $x \in A$ and $u \notin \text{dom}F(\cdot, A)$, then by (6.3) we have $\int_A f_v(x, u(x)) d\mu_v + \nu_v(A) = +\infty$.

So far we have proved the integral representation by means of any of the measures ν_v with $v \in \text{dom}F(\cdot, \Omega)$. We claim that for every $v \in \text{dom}F(\cdot, \Omega)$ and $B \in \mathcal{B}(\Omega)$

$$(6.12) \quad \nu_v(B) = \inf\{F(u, B) : u \in W^{1,p}(\Omega, \mathbf{R}^m)\}.$$

Let $B \in \mathcal{B}(\Omega)$ and $u \in W^{1,p}(\Omega, \mathbf{R}^m)$ with $F(u, B) < +\infty$. In view of the definition of F on Borel sets, by (6.11) we have

$$F(u, B) = \int_B f_v(x, u(x)) d\mu_v + \nu_v(B) \geq \nu_v(B);$$

hence, $\inf\{F(u, B) : u \in W^{1,p}(\Omega, \mathbf{R}^m), F(u, B) < +\infty\} \geq \nu_v(B)$. By the definition of $\nu_v(B)$, this implies (6.12). $\square$

The following proposition shows that, given the measures μ and ν , the function f obtained in the integral representation theorem is essentially unique.

Proposition 6.2. *Let $F \in \mathcal{F}$ with $\text{dom}F(\cdot, \Omega) \neq \emptyset$ and let $K = K_\Omega$ be the closed valued multifunction from Ω to $\mathbf{R}^m$ given by Proposition 5.1 for $A = \Omega$. Let μ and ν be two positive finite Borel measures on Ω , with μ absolutely continuous with respect to capacity, and let $f_1, f_2: \Omega \times \mathbf{R}^m \rightarrow [0, +\infty]$ be two Borel functions such that $f_1(x, \cdot)$ and $f_2(x, \cdot)$ are convex and lower semicontinuous on $\mathbf{R}^m$ for μ -a.e. $x \in \Omega$. Assume that for every $A \in \mathcal{A}(\Omega)$ and for every $u \in \text{dom}F(\cdot, A)$ we have $F(u, A) = \int_A f_i(x, u(x)) d\mu + \nu(A)$ for $i = 1, 2$. Then $f_1(x, \xi) = f_2(x, \xi)$ for μ -a.e. $x \in \Omega$ and for every $\xi \in K(x)$.*

Proof. By a translation we can easily reduce the problem to the case $F(0, \Omega) < +\infty$. Let (u_i) and $C_k(x)$ be as in the proof of Theorem 5.4. By Proposition 2.9 we have

$$F(u, A) = \int_A f_1(x, u(x)) d\mu + \nu(A) = \int_A f_2(x, u(x)) d\mu + \nu(A)$$

for every $A \in \mathcal{A}(\Omega)$ and for every $u \in W^{1,p}(\Omega, \mathbf{R}^m)$ with $u(x) \in C_k(x)$ for q.e. $x \in \Omega$. By Proposition 3.8 it turns out that $f_1(x, \xi) = f_2(x, \xi)$ for μ -a.e. $x \in \Omega$ and for every $\xi \in C_k(x)$. Hence the equality holds for every $\xi \in \text{ri}K(x) \subseteq \bigcup_{k=1}^{\infty} C_k(x)$, and, therefore, for every $\xi \in K(x)$ by the continuity along line segments (see [28], Corollary 7.5.1). $\square$

Proposition 6.3. *Let $K(x)$ be a closed and convex valued multifunction from Ω to $\mathbf{R}^m$ for which there exists a sequence (u_k) of functions in $W^{1,p}(\Omega, \mathbf{R}^m)$ such that $K(x) = \text{cl}\{u_k(x) : k \in \mathbf{N}\}$ for q.e. $x \in \Omega$. Then, there exists a positive finite Borel measure ρ on Ω , absolutely continuous with respect to capacity, such that for every $A \in \mathcal{A}(\Omega)$ and for every $u \in W^{1,p}(\Omega, \mathbf{R}^m)$ the following conditions are equivalent:*

- (i) $u(x) \in K(x)$ for q.e. $x \in A$,
- (ii) $u(x) \in K(x)$ for ρ -a.e. $x \in A$.

Proof. It is not restrictive to assume that $0 \in K(x)$ for q.e. $x \in \Omega$. Moreover, we can suppose that $u_k \in W^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$ for every $k \in \mathbf{N}$. Indeed, if $T_h \circ u_k$, with $h \in \mathbf{N}$, denotes the truncation introduced in Lemma 2.5, it turns out that $K(x) = \text{cl}\{(T_h \circ u_k)(x) : h, k \in \mathbf{N}\}$, since $K(x)$ is closed and convex and $(T_h \circ u_k)_h$ converges to u_k strongly in $W^{1,p}(\Omega, \mathbf{R}^m)$, as h tends to ∞ .

Let us note that, by a standard cut-off argument, it is enough to consider the case $A = \Omega$. Moreover, (i) clearly implies (ii) as ρ is absolutely continuous with respect to capacity.

Step 1. Here we prove that (ii) implies (i) for $u \in W_0^{1,p}(\Omega, \mathbf{R}^m)$ under the additional assumption that $\partial\Omega$ is smooth.

Let us define the convex sets

$$\begin{aligned} \mathcal{K} &= \{u \in W_0^{1,p}(\Omega, \mathbf{R}^m) : u(x) \in K(x) \text{ for q.e. } x \in \Omega\}, \\ \mathcal{K}_k &= \{u \in W_0^{1,p}(\Omega, \mathbf{R}^m) : u(x) \in K(x) + \frac{1}{k}B_1(0) \text{ for q.e. } x \in \Omega\}, \end{aligned}$$

for every $k \in \mathbf{N}$. Since $W_0^{1,p}(\Omega, \mathbf{R}^m)$ is separable, the set $\mathcal{K}_k$ is the intersection of a countable family of closed half-spaces of $W_0^{1,p}(\Omega, \mathbf{R}^m)$. Hence, there exists a sequence $(\mu_{k,h})_h$ in $W^{-1,p'}(\Omega, \mathbf{R}^m)$, with $p' = \frac{p}{p-1}$, and a sequence $(a_{k,h})_h$ in $\mathbf{R}$ such that

$$\mathcal{K}_k = \bigcap_{h \in \mathbf{N}} \{u \in W_0^{1,p}(\Omega, \mathbf{R}^m) : \langle \mu_{k,h}, u \rangle \geq a_{k,h}\},$$

where $\langle \cdot, \cdot \rangle$ denotes the duality pairing between $W^{-1,p'}(\Omega, \mathbf{R}^m)$ and $W_0^{1,p}(\Omega, \mathbf{R}^m)$.

Denote by $\mathcal{M}(\Omega, \mathbf{R}^m)$ the space of all $\mathbf{R}^m$ -valued Radon measures on Ω with bounded total variation. We say that an element $T \in W^{-1,p'}(\Omega, \mathbf{R}^m)$ belongs to $\mathcal{M}(\Omega, \mathbf{R}^m)$ if there exists $\mu \in \mathcal{M}(\Omega, \mathbf{R}^m)$ such that

$$\langle T, \varphi \rangle = \int_{\Omega} \varphi d\mu$$

for every $\varphi \in C_0^\infty(\Omega, \mathbf{R}^m)$. In this case T and μ will be identified.

Let us prove that $\mu_{k,h} \in \mathcal{M}(\Omega, \mathbf{R}^m)$ for every $h, k \in \mathbf{N}$. Fix $\varphi \in C_0^\infty(\Omega, \mathbf{R}^m)$ with $\|\varphi\|_\infty \leq 1$. Since $u_1 + \frac{1}{k}\varphi$ and $u_1 - \frac{1}{k}\varphi$ belong to $\mathcal{K}_k$, we have

$$-k(\langle \mu_{k,h}, u_1 \rangle - a_{k,h}) \leq \langle \mu_{k,h}, \varphi \rangle \leq k(\langle \mu_{k,h}, u_1 \rangle - a_{k,h}).$$

Therefore, there exists $C_{k,h} > 0$ such that $|\langle \mu_{k,h}, \varphi \rangle| \leq C_{k,h} \|\varphi\|_\infty$ for every $\varphi \in C_0^\infty(\Omega, \mathbf{R}^m)$. Hence $\mu_{k,h}|_{C_0^\infty(\Omega, \mathbf{R}^m)}$ can be uniquely extended to a continuous linear functional on the space of continuous functions on Ω vanishing on $\partial\Omega$. We conclude by the Riesz representation theorem.

Since $\mathcal{K} = \bigcap_{k \in \mathbf{N}} \mathcal{K}_k$, we can assert that there exists a sequence (μ_h) in $W^{-1,p'}(\Omega, \mathbf{R}^m) \cap \mathcal{M}(\Omega, \mathbf{R}^m)$ and a sequence (a_h) in $\mathbf{R}$ such that

$$(6.13) \quad \mathcal{K} = \bigcap_{h \in \mathbf{N}} \{u \in W_0^{1,p}(\Omega, \mathbf{R}^m) : \langle \mu_h, u \rangle \geq a_h\}.$$

Moreover, as $\mu_h \in W^{-1,p'}(\Omega, \mathbf{R}^m) \cap \mathcal{M}(\Omega, \mathbf{R}^m)$, by [22] and [8], Lemma 2 we have that $|\mu_h|$ is absolutely continuous with respect to capacity. For every $B \in \mathcal{B}(\Omega)$ define

$$\rho(B) = \sum_{h=1}^{\infty} 2^{-h} \frac{|\mu_h|(B)}{|\mu_h|(\Omega)}$$

(clearly, we can assume that $|\mu_h|(\Omega) > 0$ for every $h \in \mathbf{N}$). Then ρ is a positive finite Borel measure on Ω absolutely continuous with respect to capacity. Let g_h be the Radon-Nikodym derivative of μ_h with respect to ρ . Then $g_h \in L^1(\Omega, \rho)$. By Corollary 6 in [8], for every $u \in W_0^{1,p}(\Omega, \mathbf{R}^m) \cap L^\infty(\Omega, \mathbf{R}^m)$, we have $u \cdot g_h \in L^1(\Omega, \rho)$ and

$$(6.14) \quad \langle \mu_h, u \rangle = \int_{\Omega} u \cdot g_h d\rho.$$

Let us now prove that (ii) implies (i). Fix $u \in W_0^{1,p}(\Omega, \mathbf{R}^m)$ with $u(x) \in K(x)$ for ρ -a.e. $x \in \Omega$. Assume first that u belongs to $L^\infty(\Omega, \mathbf{R}^m)$.

For every $h \in \mathbf{N}$ and $v \in \mathcal{K} \cap L^\infty(\Omega, \mathbf{R}^m)$, by (6.14) we have

$$a_h \leq \langle \mu_h, v \rangle = \int_{\Omega} v \cdot g_h \, d\rho,$$

hence

$$(6.15) \quad a_h \leq \inf_{v \in \mathcal{K} \cap L^\infty} \int_{\Omega} v \cdot g_h \, d\rho.$$

In view of the fact that the functions u_k are in $L^\infty(\Omega, \mathbf{R}^m)$, it turns out that $K(x) = \rho\text{-ess sup}_{v \in \mathcal{K} \cap L^\infty} \{v(x)\}$ for ρ -a.e. $x \in \Omega$; then Theorem 4.4 yields

$$(6.16) \quad \inf_{v \in \mathcal{K} \cap L^\infty} \int_{\Omega} v \cdot g_h \, d\rho = \int_{\Omega} \inf_{\xi \in K(x)} \xi \cdot g_h(x) \, d\rho.$$

By the assumption $u(x) \in K(x)$ for ρ -a.e. $x \in \Omega$, by (6.15) and (6.16), it follows that

$$a_h \leq \int_{\Omega} u(x) \cdot g_h(x) \, d\rho = \langle \mu_h, u \rangle$$

for every $h \in \mathbf{N}$. By (6.13) this proves that $u \in \mathcal{K}$, i.e., $u(x) \in K(x)$ for q.e. $x \in \Omega$.

Consider now a general $u \in W_0^{1,p}(\Omega, \mathbf{R}^m)$; let us note that, since $K(x)$ is convex and $0 \in K(x)$ for q.e. $x \in \Omega$, the condition $u(x) \in K(x)$ for ρ -a.e. $x \in \Omega$ implies that $(T_h \circ u)(x) \in K(x)$ for ρ -a.e. $x \in \Omega$ and for every $h \in \mathbf{N}$. The previous step and the q.e. convergence of $(T_h \circ u)$ to u allow us to conclude as $K(x)$ is closed.

Step 2. Let us now prove that (ii) implies (i) for every $u \in W^{1,p}(\Omega, \mathbf{R}^m)$ without assuming the smoothness of the boundary of Ω .

Let (Ω_h) be a sequence of open subsets of Ω with $\Omega_h \subset\subset \Omega_{h+1}$, $\bigcup_h \Omega_h = \Omega$, and $\partial\Omega_h$ smooth. Let φ_h be a $C_0^1(\Omega_h)$ function with $\varphi_h = 1$ on Ω_{h-1} and $0 \leq \varphi_h \leq 1$. Define

$$K_h(x) = \text{cl}\{\varphi_h(x)u_k(x) : k \in \mathbf{N}\}$$

for q.e. $x \in \Omega_h$. By Step 1 there exists a positive finite Borel measure ρ_h on Ω_h , absolutely continuous with respect to capacity, and such that for every $u \in W_0^{1,p}(\Omega, \mathbf{R}^m)$ the condition $u(x) \in K_h(x)$ for q.e. $x \in \Omega_h$ is equivalent to the condition $u(x) \in K_h(x)$ for ρ_h -a.e. $x \in \Omega_h$. We can consider ρ_h as a measure on Ω by setting $\rho_h(B) = \rho_h(B \cap \Omega_h)$ for $B \in \mathcal{B}(\Omega)$. Let us define

$$\rho(B) = \sum_{h=1}^{\infty} 2^{-h} \frac{\rho_h(B)}{\rho_h(\Omega)}$$

for every $B \in \mathcal{B}(\Omega)$. Then ρ is a positive finite Borel measure on Ω which is absolutely continuous with respect to capacity.

Let us fix $u \in W^{1,p}(\Omega, \mathbf{R}^m)$ with $u(x) \in K(x)$ for ρ -a.e. $x \in \Omega$. Then, $\varphi_h(x)u(x) \in K_h(x)$ for ρ_h -a.e. $x \in \Omega_h$, so that $\varphi_h(x)u(x) \in K_h(x)$ for q.e. $x \in \Omega_h$. Since $\varphi_h = 1$ on Ω_{h-1} , we obtain that $u(x) \in K(x)$ for q.e. $x \in \Omega_{h-1}$. As h is arbitrary, we conclude that $u(x) \in K(x)$ for q.e. $x \in \Omega$. $\square$

Lemma 6.4. *Let $F \in \mathcal{F}$ and define Ω_0 to be the union of all $A \in \mathcal{A}(\Omega)$ such that $\text{dom}F(\cdot, A) \neq \emptyset$. Then $\text{dom}F(\cdot, A) \neq \emptyset$ for every $A \in \mathcal{A}(\Omega)$ with $A \subset\subset \Omega_0$.*

Proof. By induction we can reduce ourselves to prove that, whenever A_1, A_2 are open subsets of Ω with $\text{dom}F(\cdot, A_1) \neq \emptyset$ and $\text{dom}F(\cdot, A_2) \neq \emptyset$, then $\text{dom}F(\cdot, A) \neq \emptyset$ for every open set $A \subset\subset A_1 \cup A_2$.

Let A_1, A_2 and A be as above, and let $A' \subset\subset A_1$ with $A \subset\subset A' \cup A_2$. We shall show that $\text{dom}F(\cdot, A' \cup A_2) \neq \emptyset$, which clearly implies $\text{dom}F(\cdot, A) \neq \emptyset$. Consider a function $\varphi \in C_0^1(A_1)$ with $\varphi = 1$ on A' and $0 \leq \varphi \leq 1$. By assumption we can find $u \in \text{dom}F(\cdot, A_1)$ and $v \in \text{dom}F(\cdot, A_2)$; define $w = \varphi u + (1 - \varphi)v$. Then, by the usual properties of the class $\mathcal{F}$ it is easy to see that $F(w, A' \cup A_2) < +\infty$. $\square$

Theorem 6.5. *Let Ω be an open subset of $\mathbf{R}^n$ (not necessarily bounded) and let $F \in \mathcal{F}$. Then, there exist a positive finite Borel measure μ on Ω , absolutely continuous with respect to capacity, a positive Borel measure ν on Ω , and a Borel function $f: \Omega \times \mathbf{R}^m \rightarrow [0, +\infty]$ with the following properties:*

- (i) *for every $x \in \Omega$ the function $f(x, \cdot)$ is convex and lower semicontinuous on $\mathbf{R}^m$;*
- (ii) *for every $u \in W^{1,p}(\Omega, \mathbf{R}^m)$ and for every $A \in \mathcal{A}(\Omega)$*

$$(6.17) \quad F(u, A) = \int_A f(x, u(x)) d\mu + \nu(A).$$

Proof. Let $\overline{\nu}$ be the function on $\mathcal{B}(\Omega)$ defined in (6.1). Let us set

$$\nu(A) = \sup\{\overline{\nu}(A') : A' \in \mathcal{A}(\Omega), A' \subset\subset A\}$$

for every $A \in \mathcal{A}(\Omega)$. Clearly ν is increasing with respect to the inclusion, and $\nu(\emptyset) = 0$. Moreover, for every $A \in \mathcal{A}(\Omega)$

$$(6.18) \quad \nu(A) < +\infty \Rightarrow A \subseteq \Omega_0.$$

Clearly ν is inner regular on $\mathcal{A}(\Omega)$; therefore, by Proposition 5.5 and Theorem 5.6 in [19], to prove that ν can be extended to a Borel measure on Ω it suffices to show that ν is subadditive and superadditive on $\mathcal{A}(\Omega)$.

Let $A_1, A_2 \in \mathcal{A}(\Omega)$, and note that

$$(6.19) \quad \nu(A_1 \cup A_2) = \sup\{\bar{\nu}(A'_1 \cup A'_2) : A'_i \in \mathcal{A}(\Omega), A'_i \subset\subset A_i \ (i = 1, 2)\}.$$

If $\nu(A_1)$ and $\nu(A_2)$ are finite, then $A_1, A_2 \subseteq \Omega_0$ by (6.18). Since, by Lemma 6.4, $\bar{\nu}$ is a measure on every $\Omega' \subset\subset \Omega_0$, from (6.19) it follows that $\nu(A_1 \cup A_2) \leq \nu(A_1) + \nu(A_2)$. In a similar way we get superadditivity.

This allows us to conclude that the set function $\nu: \mathcal{B}(\Omega) \rightarrow [0, +\infty]$ defined by

$$\nu(B) = \inf\{\nu(A) : A \in \mathcal{A}(\Omega), B \subseteq A\}$$

is a Borel measure on Ω and that $\nu(B) = \bar{\nu}(B)$ for every $B \in \mathcal{B}(\Omega)$ with $B \subset\subset \Omega_0$.

Let us construct now μ and f . Let (Ω_h) be a sequence of open subsets of Ω_0 with a smooth boundary such that $\Omega_h \subset\subset \Omega_{h+1}$ and $\Omega_0 = \cup_h \Omega_h$. In particular, $\text{dom}F(\cdot, \Omega_h) \neq \emptyset$ by Lemma 6.4. Since there exists an extension operator from $W^{1,p}(\Omega_h, \mathbf{R}^m)$ to $W^{1,p}(\Omega, \mathbf{R}^m)$, it is possible to apply Theorem 6.1 to each Ω_h using $W^{1,p}(\Omega, \mathbf{R}^m)$ instead of $W^{1,p}(\Omega_h, \mathbf{R}^m)$. Let K_{Ω_h} be the multifunction defined in Proposition 5.1 for $A = \Omega_h$; then there exist a positive finite Borel measure μ_h on Ω_h , absolutely continuous with respect to capacity, and a Borel function $f_h: \Omega_h \times \mathbf{R}^m \rightarrow [0, +\infty]$ such that

- (a) for every $x \in \Omega_h$ the function $f_h(x, \cdot)$ is convex and lower semicontinuous on $\mathbf{R}^m$;
- (b) for every $u \in W^{1,p}(\Omega, \mathbf{R}^m)$ and $A \in \mathcal{A}(\Omega_h)$

$$F(u, A) = \begin{cases} \int_A f_h(x, u(x)) d\mu_h + \bar{\nu}(A), & \text{if } u(x) \in K_{\Omega_h}(x) \text{ for q.e. } x \in A, \\ +\infty, & \text{otherwise.} \end{cases}$$

Moreover, we have a uniqueness property for the integrand as stated in Proposition 6.2.

By Proposition 6.3, for every $h \in \mathbf{N}$ there exists a positive finite Borel measure ρ_h on Ω_h , absolutely continuous with respect to capacity, such that for every $u \in W^{1,p}(\Omega, \mathbf{R}^m)$ the condition $u(x) \in K_{\Omega_h}(x)$ for q.e. $x \in \Omega_h$ is equivalent to the condition

$u(x) \in K_{\Omega_h}(x)$ for ρ_h -a.e. $x \in \Omega_h$. Let us consider μ_h and ρ_h as measures on Ω by setting $\mu_h(B) = \mu_h(B \cap \Omega_h)$ and $\rho_h(B) = \rho_h(B \cap \Omega_h)$ for every $B \in \mathcal{B}(\Omega)$. Define

$$\mu(B) = \sum_{h=1}^{\infty} 2^{-h} \frac{(\mu_h + \rho_h)(B)}{(\mu_h + \rho_h)(\Omega)},$$

$$g_h(x, \xi) = \begin{cases} f_h(x, \xi) \frac{d\mu_h}{d\mu}(x), & \text{if } x \in \Omega_h \text{ and } \xi \in K_{\Omega_h}(x), \\ +\infty, & \text{otherwise,} \end{cases}$$

where $d\mu_h/d\mu$ is a fixed Borel representative of the Radon-Nikodym derivative. Then μ is a positive finite Borel measure on Ω , absolutely continuous with respect to capacity. Since for every $h \in \mathbf{N}$ there is a sequence (v_i) in $W^{1,p}(\Omega, \mathbf{R}^m)$ such that $K_{\Omega_h}(x) = \text{cl}\{v_i(x) : i \in \mathbf{N}\}$, by Theorem III.9 and Proposition III.13 in [10], the graph of K_{Ω_h} belongs to $\mathcal{B}(\Omega_h) \times \mathcal{B}(\mathbf{R}^m)$. Therefore $g_h : \Omega_h \times \mathbf{R}^m \rightarrow [0, +\infty]$ is a Borel function, and for every $x \in \Omega_h$, the function $g_h(x, \cdot)$ is convex and lower semicontinuous on $\mathbf{R}^m$.

By recalling that $K_{\Omega_h} = K_{\Omega_{h+1}}$ q.e. on Ω_h (see Remark 5.2), and by using the uniqueness property of the integrand mentioned above, we easily obtain that $g_h(x, \cdot) = g_{h+1}(x, \cdot)$ for μ -a.e. $x \in \Omega_h$.

Therefore, there exists a Borel function $f : \Omega \times \mathbf{R}^m \rightarrow [0, +\infty]$ satifying (i) and such that for every $h \in \mathbf{N}$

$$f(x, \cdot) = g_h(x, \cdot) \quad \text{on } \mathbf{R}^m \text{ for } \mu\text{-a.e. } x \in \Omega_h.$$

Let us now prove (ii). Fix $u \in W^{1,p}(\Omega, \mathbf{R}^m)$ and $A \in \mathcal{A}(\Omega)$. If $A \setminus \Omega_0 \neq \emptyset$, then $\nu(A) = +\infty$ by (6.18). On the other hand, by the definition of Ω_0 we have $F(u, A) = +\infty$ for every $u \in W^{1,p}(\Omega, \mathbf{R}^m)$. Therefore

$$F(u, A) = \int_A f(x, u(x)) d\mu + \nu(A).$$

Let now $A \subseteq \Omega_0$ and $A' \subset\subset A$. Then $A' \subseteq \Omega_h$ for a suitable $h \in \mathbf{N}$. In view of the properties of the measure ρ_h , from the definition of g_h and f we easily obtain

$$F(u, A') = \int_{A'} g_h(x, u(x)) d\mu + \overline{\nu}(A') = \int_{A'} f(x, u(x)) d\mu + \overline{\nu}(A').$$

Therefore (ii) follows from the definition of ν taking the supremum for $A' \subset\subset A$. $\square$

Remark 6.6. Let $F \in \mathcal{F}$ and Ω_0 be as in Lemma 6.4. By Proposition 5.1 and Remark 5.2, there exists a closed valued multifunction K from Ω_0 to $\mathbf{R}^m$, unique up to sets of capacity zero, such that

$$(6.20) \quad K(x) = K_A(x) \quad \text{for q.e. } x \in A$$

whenever $A \in \mathcal{A}(\Omega)$ and $\text{dom}F(\cdot, A) \neq \emptyset$. Moreover, $K(x)$ is non-empty and convex for q.e. $x \in \Omega_0$.

It is clear that the function f constructed in the proof of Theorem 6.5 satisfies the additional condition $f(x, \xi) = +\infty$ for μ -a.e. $x \in \Omega_0$ and for every $\xi \notin K(x)$. This is not necessarily true for every function f which satisfies conditions (i) and (ii) of Theorem 6.5. Let us consider, for instance, the functional

$$F(u, A) = \begin{cases} 0, & \text{if } u = 0 \text{ a.e. on } A, \\ +\infty, & \text{otherwise,} \end{cases}$$

in the case $n = m = 1$ and $\Omega = \mathbf{R}$. Then, clearly, $K(x) = \{0\}$ for q.e. $x \in \mathbf{R}$ and

$$(6.21) \quad F(u, A) = \int_A f(x, u(x)) dx,$$

with

$$f(x, \xi) = \begin{cases} 0, & \text{if } \xi = 0, \\ +\infty, & \text{if } \xi \neq 0. \end{cases}$$

But (6.21) holds also with $f(x, \xi) = a(x)|\xi|^2$, where $a: \mathbf{R} \rightarrow [0, +\infty[$ is any finite valued Borel function such that $\int_A a(x) dx = +\infty$ for every open subset A of $\mathbf{R}$ (see [23], Section 43, Exercise 7).

Remark 6.7. If $\text{dom}F(\cdot, \Omega) = \emptyset$ and ν is not necessarily finite, the uniqueness result of Proposition 6.2 still holds, with an obvious localization of the proof, in the weaker form:

$$f_1(x, \xi) = f_2(x, \xi) \quad \text{for } \mu\text{-a.e. } x \in \Omega_0 \text{ and for every } \xi \in K(x),$$

where Ω_0 is defined in Lemma 6.4 and $K(x)$ is now defined by (6.20).

7. Quadratic functionals

In this section we show how certain algebraic properties of the functional F are inherited by the integrand which appears in the representation of F according to Theorem 6.5. We recall that a cone in a vector space X (with vertex at 0) is a set K such that $tx \in K$ for every $t > 0$ and for every $x \in K$.

Definition 7.1. Let X be a real vector space and let $p \in \mathbf{R}$. We say that a function $f: X \rightarrow [0, +\infty]$ is:

- (i) *positively homogeneous of degree p* on a cone K if $F(tx) = t^p F(x)$ for every $t > 0$ and for every $x \in K$;
- (ii) a (non-negative) *quadratic form* (with extended real values) on X if there exist a linear subspace Y of X and a symmetric bilinear form $B: Y \times Y \rightarrow \mathbf{R}$ such that

$$F(x) = \begin{cases} B(x, x), & \text{if } x \in Y, \\ +\infty, & \text{if } x \in X \setminus Y. \end{cases}$$

We shall refer to Y as the *domain* of F .

Remark 7.2. In the previous definition it is not restrictive to assume that B is defined over all of $X \times X$. Indeed, let Z be an algebraic complement of Y in X and denote by $P: X \rightarrow Y$ the canonical projection on Y associated to the pair (Y, Z) . Then, it is enough to consider the extension $(x, y) \mapsto B(Px, Py)$ defined for every $(x, y) \in X \times X$. As a consequence, if X is finite dimensional and $\dim X = m$, then there exists an $m \times m$ symmetric matrix (a_{ij}) such that $F(x) = \sum_{i,j=1}^m a_{ij} x^i x^j$ for every $x \in Y$, where $x^1, \dots, x^m$ denote the components of x with respect to a fixed basis of X .

Theorem 7.3. Let $F \in \mathcal{F}$ and $p \in \mathbf{R}$. Let f, μ, ν be as in Theorem 6.5 and let Ω_0 and K be as in Remark 6.6. Assume that $f(x, \xi) = +\infty$ for μ -a.e. $x \in \Omega_0$ and for every $\xi \notin K(x)$. Then the following properties hold:

- (i) if $F(\cdot, A)$ is positively homogeneous of degree p on $W^{1,p}(\Omega, \mathbf{R}^m)$ for every $A \in \mathcal{A}(\Omega)$, then $K(x)$ is a closed convex cone for q.e. $x \in \Omega_0$, and $f(x, \cdot)$ is positively homogeneous of degree p on $K(x)$ for μ -a.e. $x \in \Omega_0$; if, in addition, $p \neq 0$, then $\nu(B) = 0$ for every $B \in \mathcal{B}(\Omega_0)$;
- (ii) if $F(\cdot, A)$ is a quadratic form on $W^{1,p}(\Omega, \mathbf{R}^m)$ for every $A \in \mathcal{A}(\Omega)$, then $\nu = 0$, $K(x)$ is a linear subspace of $\mathbf{R}^m$ for q.e. $x \in \Omega$, and for μ -a.e. $x \in \Omega$ the function $f(x, \cdot)$ is a quadratic form on $\mathbf{R}^m$ with domain $K(x)$.

Proof. Proof of (i). For every $A \in \mathcal{A}(\Omega)$ the positive homogeneity of degree p implies that $tu \in \text{dom} F(\cdot, A)$ whenever $t > 0$ and $u \in \text{dom} F(\cdot, A)$. Recalling the definition and properties of K_A given in Proposition 5.1, it is easy to see that, if $\text{dom} F(\cdot, A) \neq \emptyset$, then there exists a set $N \subseteq \Omega$ with $\text{cap}(N) = 0$ and such that for every $x \in A \setminus N$, $q \in \mathbf{Q}^+ \setminus \{0\}$

and $\xi \in K_A(x)$ we have $q\xi \in K_A(x)$. Since $K_A(x)$ is closed, it follows that $K_A(x)$ is a cone for every $x \in A \setminus N$. The convexity of $K_A(x)$ is proved in Proposition 5.1(iii). By the definition (6.20) of $K(x)$ and by the definition of Ω_0 we conclude that $K(x)$ is a closed convex cone for q.e. $x \in \Omega_0$.

Let us now prove that $f(x, \cdot)$ is positively homogeneous of degree p on $K(x)$ for μ -a.e. $x \in \Omega_0$. Let us first consider the case $p = 0$. If $A \in \mathcal{A}(\Omega)$ and $u \in \text{dom}F(\cdot, A)$, then the function $t \mapsto F(tu, A)$ from $[0, 1]$ into $[0, +\infty]$ is convex and lower semicontinuous; moreover, $F(tu, A) = F(u, A) < +\infty$ for every $t > 0$. Therefore, $F(0, A) = \lim_{t \rightarrow 0^+} F(tu, A) = F(u, A)$ for every $A \in \mathcal{A}(\Omega)$ and $u \in \text{dom}F(\cdot, A)$. This shows that $0 \in K(x)$ for q.e. $x \in \Omega_0$. By the uniqueness of the integrand stated in Remark 6.7, we conclude that $f(x, \xi) = f(x, 0)$ for μ -a.e. $x \in \Omega_0$ and for every $\xi \in K(x)$.

Assume now $p \neq 0$. Let $A \in \mathcal{A}(\Omega)$ and $w \in \text{dom}F(\cdot, A)$. Since $\nu(A) \leq F(tw, A) = t^p F(w, A)$ for every $t > 0$, taking the limit as $t \rightarrow 0^+$ or $t \rightarrow +\infty$ according to whether $p > 0$ or $p < 0$, we get $\nu(A) = 0$. In view of the positive homogeneity of $F(\cdot, A)$ and (6.17) we have

$$F(u, A) = \int_A \frac{1}{t^p} f(x, tu(x)) d\mu$$

for every $A \in \mathcal{A}(\Omega)$, $u \in W^{1,p}(\Omega, \mathbf{R}^m)$ and $t > 0$. Now, by the uniqueness of the integrand (Remark 6.7), we have $f(x, \xi) = (1/t^p)f(x, t\xi)$ for μ -a.e. $x \in \Omega_0$ and for every $\xi \in K(x)$.

Proof of (ii). Assume that $F(\cdot, A)$ is a quadratic form for every $A \in \mathcal{A}(\Omega)$. Then $F(0, A) = 0$ for every $A \in \mathcal{A}(\Omega)$, hence $\Omega_0 = \Omega$, $\nu(B) = 0$ for every $B \in \mathcal{B}(\Omega)$, and $f(x, 0) = 0$ for μ -a.e. $x \in \Omega$. Directly from Definition 7.1 it follows that $\text{dom}F(\cdot, A)$ is a linear space; in particular, $u + v$ and $-u$ belong to $\text{dom}F(\cdot, A)$ whenever $u, v \in \text{dom}F(\cdot, A)$. As in the first part of (i), it can be shown that for q.e. $x \in \Omega$, $\xi + \eta$ and $-\xi$ belong to $K(x)$ if $\xi, \eta \in K(x)$. Since $K(x)$ is a cone (part (i)), this guarantees that $K(x)$ is a linear subspace of $\mathbf{R}^m$ for q.e. $x \in \Omega$.

If X is a (real) vector space, it is well known (Fréchet-Von Neumann-Jordan Theorem, see, for instance, [30]) that a function $F: X \rightarrow [0, +\infty]$ is a quadratic form if and only if $F(0) = 0$, F is positively homogeneous of degree 2, and satisfies the following “parallelogram identity”:

$$F(\xi + \eta) + F(\xi - \eta) = 2F(\xi) + 2F(\eta)$$

for every $\xi, \eta \in X$. Since $f(x, 0) = 0$ for μ -a.e. $x \in \Omega$ and $f(x, t\xi) = t^2 f(x, \xi)$ for μ -a.e. $x \in \Omega$ and for every $t > 0$, $\xi \in K(x)$ (see part (i)), to complete the proof of (ii) it remains only to show that $f(x, \cdot)$ satisfies the parallelogram identity on $K(x)$ for μ -a.e. $x \in \Omega$. Define the functional $G: [W^{1,p}(\Omega, \mathbf{R}^m)]^2 \times \mathcal{A}(\Omega) \rightarrow [0, +\infty]$ as

$$G(u, v, A) = F(u + v, A) + F(u - v, A) = 2F(u, A) + 2F(v, A).$$

Since $\nu = 0$, from (6.17) we obtain

$$(7.1) \quad G(u, v, A) = \int_A 2[f(x, u(x)) + f(x, v(x))] d\mu = \int_A [f(x, u(x) + v(x)) + f(x, u(x) - v(x))] d\mu$$

for every $A \in \mathcal{A}(\Omega)$. Since $[W^{1,p}(\Omega, \mathbf{R}^m)]^2$ can be identified with $W^{1,p}(\Omega, \mathbf{R}^{2m})$, we can apply Remark 6.7 to the functional G , with the set $K(x) \times K(x)$ playing the role of $K(x)$ for q.e. $x \in \Omega$. Therefore, (7.1) gives that

$$2(f(x, \xi) + f(x, \eta)) = f(x, \xi + \eta) + f(x, \xi - \eta)$$

for μ -a.e. $x \in \Omega_0$ and for every $(\xi, \eta) \in K(x) \times K(x)$. □

Corollary 7.4. *Let $F \in \mathcal{F}$. Assume that $F(\cdot, A)$ is a quadratic form on $W^{1,p}(\Omega, \mathbf{R}^m)$ for every $A \in \mathcal{A}(\Omega)$. Then there exist:*

- (i) *a positive finite Borel measure μ on Ω , absolutely continuous with respect to capacity,*
- (ii) *a symmetric $m \times m$ matrix (a_{ij}) of Borel functions from Ω into $\mathbf{R}$ such that $\sum_{i,j=1}^m a_{ij}(x) \xi^i \xi^j \geq 0$ for q.e. $x \in \Omega$ and for every $\xi \in \mathbf{R}^m$,*
- (iii) *for every $x \in \Omega$ a linear subspace $V(x)$ of $\mathbf{R}^m$,*

with the following properties: for every $u \in W^{1,p}(\Omega, \mathbf{R}^m)$ and $A \in \mathcal{A}(\Omega)$

- (a) *if $F(u, A) < +\infty$, then $u(x) \in V(x)$ for q.e. $x \in A$;*
- (b) *if $u(x) \in V(x)$ for q.e. $x \in A$, then $F(u, A) = \int_A \sum_{i,j=1}^m a_{ij}(x) u^i(x) u^j(x) d\mu$.*

Proof. The conclusion follows from Theorems 6.5 and 7.3(ii), and from Remark 7.2. □

References

- [1] ALBERTI G.: Integral representation of local functionals. Preprint Scuola Normale, Pisa, 1990.
- [2] ALBERTI G., BUTTAZZO G.: Integral representation of functionals defined on Sobolev spaces. *Composite Media and Homogenization Theory (Trieste, 1990)*, 1-12, Birkhäuser, Boston, 1991.
- [3] AMBROSIO L., DAL MASO G.: On the relaxation in $BV(\Omega; \mathbf{R}^m)$ of quasi-convex integrals. *J. Funct. Anal.*, to appear.
- [4] AMBROSIO L., PALLARA D.: Integral representation of relaxed functionals on $BV(\mathbf{R}^n; \mathbf{R}^k)$ and polyhedral approximation. Preprint Scuola Normale, Pisa, submitted to *Indiana Univ. Math. J.*
- [5] ATTOUCH H., PICARD C.: Variational inequalities with varying obstacles: the general form of the limit problem. *J. Funct. Anal.* **50** (1983), 329-386.
- [6] BOUCHITTÉ G., DAL MASO G.: Integral representation and relaxation of convex local functionals on $BV(\Omega)$. Preprint SISSA, Trieste, 1991.
- [7] BOUCHITTÉ G., VALADIER M.: Integral representation of convex functionals on a space of measures. *J. Funct. Anal.* **80** (1988), 398-420.
- [8] BREZIS H., BROWDER F.E.: Some properties of higher order Sobolev spaces. *J. Math. Pures Appl. (9)* **61** (1982), 245-259.
- [9] BUTTAZZO G.: Semicontinuity, Relaxation and Integral Representation Problems in the Calculus of Variations. Pitman Res. Notes in Math. 207, Longman, Harlow, 1989.
- [10] CASTAING C., VALADIER M.: Convex Analysis and Measurable Multifunctions. Lecture Notes in Math. 580, Springer-Verlag, Berlin, 1977.
- [11] CHOQUET G.: Lectures on Analysis. Vol. I. Mathematics Lecture Note Series, Benjamin, New York, 1969.
- [12] DAL MASO G.: Asymptotic behaviour of minimum problems with bilateral obstacles. *Ann. Mat. Pura Appl. (4)* **129** (1981), 327-366.
- [13] DAL MASO G.: On the integral representation of certain local functionals. *Ricerche Mat.* **32** (1983), 85-113.
- [14] DAL MASO G., DEFRANCESCHI A., VITALI E.: A characterization of C^1 -convex sets in Sobolev spaces. *Manuscripta Math.*, to appear.
- [15] DAL MASO G., LONGO P.: Γ -limits of obstacles. *Ann. Mat. Pura Appl. (4)* **128** (1980), 1-50.
- [16] DAL MASO G., PADERNI G.: Integral representation of some convex local functionals. *Ricerche Mat.* **36** (1987), 197-214.
- [17] DEFRANCESCHI A., VITALI E.: Paper in preparation.
- [18] DE GIORGI E., DAL MASO G., LONGO P.: Γ -limiti di ostacoli. *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (8)* **68** (1980), 481-487.
- [19] DE GIORGI E., LETTA G.: Une notion générale de convergence faible pour des fonctions croissantes d'ensemble. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **4** (1977), 61-99.
- [20] FEDERER H., ZIEMER W.P.: The Lebesgue set of a function whose distribution derivatives are p th power summable. *Indiana Univ. Math. J.* **22** (1982), 139-158.
- [21] FONSECA I., RYBKA P.: Relaxation of multiple integrals in the space $BV(\Omega; \mathbf{R}^p)$. Research Report, Carnegie Mellon University, Pittsburgh, 1991.
- [22] GRUN-REHOMME M.: Caractérisation du sous-différentiel d'intégrandes convexes dans les espaces de Sobolev. *J. Math. Pures Appl.* **56** (1977), 149-156.
- [23] HALMOS P.R.: Measure Theory. Van Nostrand, Princeton, 1950.
- [24] LOJASIEWICZ S., Jr.: Parametrizations of convex sets. *J. Approx. Theory*, to appear.
- [25] MORREY C.B., Jr.: Multiple Integrals in the Calculus of Variations. Springer-Verlag, Berlin, 1966.

- [26] ORNELAS A.: Parametrization of Carathéodory multifunctions. *Rend. Sem. Mat. Univ. Padova* **83** (1990), 33-44.
- [27] PICARD C. Problème biharmonique avec obstacles variables. Thèse, Univ. Paris-Sud, 1984.
- [28] ROCKAFELLAR R.T.: Convex Analysis. Princeton University Press, Princeton, 1970.
- [29] VALADIER M.: Multi-applications mesurables à valeurs convexes compactes. *J. Math. Pures Appl.* **50** (1971), 265-297.
- [30] YOSIDA K.: Functional Analysis. Springer-Verlag, Berlin, 1965.
- [31] ZIEMER W.P.: Weakly Differentiable Functions. Springer-Verlag, Berlin, 1989.

Gianni DAL MASO
SISSA
Via Beirut 2/4
I-34014 TRIESTE

Anneliese DEFRANCESCHI
Università di Trento
Dip. di Matematica
I-38050 POVO
(TRENTO)

Enrico VITALI
Università di Parma
Dip. di Matematica
Via D'Azeglio 85/A
I-43100 PARMA